



**HAL**  
open science

# Geometric realizations of the accordion complex of a dissection

Thibault Manneville, Vincent Pilaud

► **To cite this version:**

Thibault Manneville, Vincent Pilaud. Geometric realizations of the accordion complex of a dissection. *Discrete and Computational Geometry*, 2019, 61 (3), pp.507-540. 10.1007/s00454-018-0004-2. hal-02343579

**HAL Id: hal-02343579**

**<https://hal.science/hal-02343579v1>**

Submitted on 2 Nov 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# GEOMETRIC REALIZATIONS OF THE ACCORDION COMPLEX OF A DISSECTION

THIBAUT MANNEVILLE AND VINCENT PILAUD

ABSTRACT. Consider  $2n$  points on the unit circle and a reference dissection  $D_\circ$  of the convex hull of the odd points. The accordion complex of  $D_\circ$  is the simplicial complex of non-crossing subsets of the diagonals with even endpoints that cross a connected subset of diagonals of  $D_\circ$ . In particular, this complex is an associahedron when  $D_\circ$  is a triangulation and a Stokes complex when  $D_\circ$  is a quadrangulation. In this paper, we provide geometric realizations (by polytopes and fans) of the accordion complex of any reference dissection  $D_\circ$ , generalizing known constructions arising from cluster algebras.

KEYWORDS. Permutahedra · Zonotopes · Associahedra ·  $\mathbf{g}$ -,  $\mathbf{c}$ - and  $\mathbf{d}$ -vectors.

The  $(n-3)$ -dimensional *associahedron* is a simple polytope whose face poset is isomorphic to the reverse inclusion poset of non-crossing subsets of diagonals of a convex  $n$ -gon. Introduced in early works of D. Tamari [Tam51] and J. Stasheff [Sta63], it was first realized as a convex polytope by M. Haiman [Hai84] and C. Lee [Lee89], and later constructed by more systematic methods developed by several authors, in particular [GKZ08, Lod04, HL07, CSZ15]. Various relevant generalizations of the associahedron were introduced and studied, in particular secondary polytopes and fiber polytopes [GKZ08, BFS90], generalized associahedra [FZ03b, CFZ02, HLT11, Ste13, Hoh] in connection to cluster algebras [FZ02, FZ03a], graph associahedra [CD06, Pos09, FS05, Zel06, Pil13, MP17], or brick polytopes [PS12, PS15].

In a different context, Y. Baryshnikov [Bar01] introduced the simplicial complex of crossing-free subsets of the set of diagonals of a polygon that are in some sense compatible with a reference quadrangulation  $Q_\circ$ . Although the precise definition of compatibility is a bit technical in [Bar01], it turns out that a diagonal is compatible with  $Q_\circ$  if and only if it crosses a connected subset of diagonals of  $Q_\circ$  that we call *accordion* of  $Q_\circ$ . We thus call Y. Baryshnikov's simplicial complex the *accordion complex*  $\mathcal{AC}(Q_\circ)$ . A polytopal realization of  $\mathcal{AC}(Q_\circ)$  was announced in [Bar01], but the explicit construction and its proof were never published as far as we know. Revisiting some combinatorial and algebraic properties of  $\mathcal{AC}(Q_\circ)$ , F. Chapoton [Cha16, Intro.p.4] raised three explicit challenges: first prove that the oriented dual graph of  $\mathcal{AC}(Q_\circ)$  has a lattice structure extending the Tamari and Cambrian lattices [MHPS12, Rea06]; second construct geometric realizations of  $\mathcal{AC}(Q_\circ)$  as fans and polytopes generalizing the known constructions of the associahedron; third show that the facets of  $\mathcal{AC}(Q_\circ)$  are in bijection with other combinatorial objects called serpent nests [Cha16, Sect. 4].

In [GM16], A. Garver and T. McConville defined and studied the accordion complex  $\mathcal{AC}(D_\circ)$  of any reference dissection  $D_\circ$  (their presentation slightly differs as they use a compatibility condition on the dual tree of the dissection  $D_\circ$ , but the simplicial complex is the same). In this context, they settled F. Chapoton's lattice question, using lattice quotients of a lattice of biclosed sets. In this paper, we present geometric realizations of  $\mathcal{AC}(D_\circ)$  for any reference dissection  $D_\circ$ , providing in particular an answer to F. Chapoton's geometric question. In fact, we present three methods to realize  $\mathcal{AC}(D_\circ)$  based on constructions of the classical associahedron.

Our first method is based on the  $\mathbf{g}$ -vector fan. It belongs to a series of constructions of the (generalized) associahedra initiated by S. Shnider and S. Sternberg [SS93], popularised by J.-L. Loday [Lod04], developed by C. Hohlweg, C. Lange and H. Thomas [HL07, HLT11] using works of N. Reading and D. Speyer [Rea06, Rea07, RS09], and revisited by S. Stella [Ste13] and by V. Pilaud, F. Santos, and C. Stump [PS12, PS15]. It was recently extended by C. Hohlweg, V. Pilaud, and S. Stella [HPS18] to construct an associahedron parametrized by any initial triangulation. Here, we first extend to the  $D_\circ$ -accordion complex  $\mathcal{AC}(D_\circ)$  the  $\mathbf{g}$ -vectors and  $\mathbf{c}$ -vectors

defined in the context of cluster algebras by S. Fomin and A. Zelevinski [FZ07]. Note that  $\mathbf{c}$ -vectors were already implicitly considered in [GM16], while  $\mathbf{g}$ -vectors are new in this context. When  $D_\circ$  is a triangulation, our definitions coincide with those given in terms of triangulations and laminations for cluster algebras from surfaces by S. Fomin and D. Thurston [FT12]. We then show that the  $\mathbf{g}$ -vectors with respect to the dissection  $D_\circ$  support a complete simplicial fan  $\mathcal{F}^{\mathbf{g}}(D_\circ)$  realizing the  $D_\circ$ -accordion complex  $\mathcal{AC}(D_\circ)$ . Finally, we construct a  $D_\circ$ -accordiohedron  $\text{Acco}(D_\circ)$  realizing the  $\mathbf{g}$ -vector fan  $\mathcal{F}^{\mathbf{g}}(D_\circ)$  by deleting inequalities from the facet description of the  $D_\circ$ -zonotope  $\text{Zono}(D_\circ)$  obtained as the Minkowski sum of all  $\mathbf{c}$ -vectors. See Figure 7 for an illustration of  $D_\circ$ -accordiohedra.

Our second method is based on the  $\mathbf{d}$ -vector fan. This construction is inspired from the original cluster fan of S. Fomin and A. Zelevinsky [FZ03a] later realized as a polytope by F. Chapoton, S. Fomin and A. Zelevinsky [CFZ02], and from the generalization of C. Ceballos, F. Santos and G. Ziegler [CSZ15] to construct a compatibility fan and an associahedron from any initial triangulation. For any reference dissection  $D_\circ$ , we associate to each diagonal a  $\mathbf{d}$ -vector which records the crossings of this diagonal with those of  $D_\circ$ . We show that the  $\mathbf{d}$ -vectors support a complete simplicial fan realizing the  $D_\circ$ -accordion complex  $\mathcal{AC}(D_\circ)$  if and only if  $D_\circ$  contains no even interior cell. The polytopality of the resulting fan remains open in general, but was shown for arbitrary triangulations in [CSZ15].

Finally, our third method is based on projections of associahedra. Namely, for any dissection  $D_\circ$  and triangulation  $T_\circ$  such that  $D_\circ \subseteq T_\circ$ , the accordion complex  $\mathcal{AC}(D_\circ)$  is a subcomplex of the simplicial associahedron  $\mathcal{AC}(T_\circ)$ . It turns out that the  $\mathbf{g}$ -vector fan  $\mathcal{F}^{\mathbf{g}}(D_\circ)$  is then a section of the  $\mathbf{g}$ -vector fan  $\mathcal{F}^{\mathbf{g}}(T_\circ)$  by a coordinate subspace. Therefore, the accordion complex  $\mathcal{AC}(D_\circ)$  is realized by a projection of the associahedron  $\text{Asso}(T_\circ)$  of [HPS18]. This point of view provides a complementary perspective on accordion complexes that leads on the one hand to more concise but less instructive proofs of combinatorial and geometric properties of the accordion complex (pseudomanifold,  $\mathbf{g}$ -vector fan, accordiohedron), and on the other hand to natural extensions to coordinate sections of the  $\mathbf{g}$ -vector fan in arbitrary cluster algebras.

As recently observed in [GM16, PPP17, PPS17, BDM<sup>+</sup>17], accordion complexes are prototypes of support  $\tau$ -tilting complexes introduced in [AIR14], for certain associative algebras called gentle algebras. In this context,  $\mathbf{g}$ -vectors have a deep algebraic meaning and still define a  $\mathbf{g}$ -vector fan. Although this fan is still polytopal for finite support  $\tau$ -tilting complexes, it is not in general obtained by deleting inequalities in the facet description of a zonotope. We refer to [PPP17, Part 4] for details.

The paper is organized as follows. Section 1 introduces the accordion complex and accordion lattice of a dissection  $D_\circ$ . We essentially follow the definitions and arguments of A. Garver and T. McConville [GM16], except that we prefer to work on the dissection  $D_\circ$  rather than on its dual graph. Section 2 is devoted to the generalization of the  $\mathbf{g}$ -vector fan and the associahedra of [HL07, HPS18]. Section 3 discusses the generalization of the construction of the  $\mathbf{d}$ -vector fan and associahedra of [FZ03a, CSZ15]. Finally, Section 4 shows that the accordion complex is realized by a projection of a well-chosen associahedron and presents related questions on cluster algebras, subcomplexes of the cluster complex, and sections of the  $\mathbf{g}$ -vector fan.

## 1. THE ACCORDION COMPLEX AND THE ACCORDION LATTICE

In this section, we define the accordion complex  $\mathcal{AC}(D_\circ)$  of a dissection  $D_\circ$ , show that it is a pseudomanifold, and define an orientation of its dual graph. Our definitions and proofs are essentially translations of the arguments of A. Garver and T. McConville [GM16] given in terms of the dual tree of the dissection  $D_\circ$ . However our presentation in terms of dissections is more convenient for our purposes.

**1.1. The accordion complex.** Let  $P$  be a convex polygon. We call *diagonals* of  $P$  the segments connecting two vertices of  $P$ . This includes both the internal diagonals and the external diagonals (or boundary edges) of  $P$ . A *dissection* of  $P$  is a set  $D$  of non-crossing internal diagonals of  $P$ . The *cells* of  $D$  are the closures of the connected components of  $P$  minus the diagonals of  $D$ . A *triangulation* (resp. *quadrangulation*) is a dissection whose cells are all triangles (resp. quadrangles).




FIGURE 1. A dissection  $D$  (left) and three accordions whose zigzags are bolded (middle and right).

We denote by  $\bar{D}$  the dissection  $D$  together with all boundary edges of  $P$ . A *cut* of  $D$  is the subset of  $\bar{D}$  intersected by a line crossing two boundary edges of  $P$ . An *accordion* is a connected cut. By definition, an accordion is a tree and contains two boundary edges of  $P$ . The *zigzag* of an accordion  $A$  is the chain obtained by deleting all degree 1 vertices of  $A$ . A *subaccordion* of  $D$  is a connected subset of  $D$  intersected by a segment in the interior of  $P$ . Note that any subaccordion of an accordion  $A$  consists of the diagonals of  $A$  between two internal diagonals of  $A$ . Note that we include boundary edges of  $P$  in the accordions of  $D$ , but not in the subaccordions nor in the zigzags of  $D$ . See Figure 1.

We consider  $2n$  points on the unit circle labeled clockwise by  $1_\circ, 2_\bullet, 3_\circ, 4_\bullet, \dots, (2n-1)_\circ, (2n)_\bullet$ . We say that  $1_\circ, \dots, (2n-1)_\circ$  are the *hollow vertices* while  $2_\bullet, \dots, (2n)_\bullet$  are the *solid vertices*. The *hollow polygon* is the convex hull  $P_\circ$  of  $1_\circ, \dots, (2n-1)_\circ$  while the *solid polygon* is the convex hull  $P_\bullet$  of  $2_\bullet, \dots, (2n)_\bullet$ . We simultaneously consider *hollow diagonals*  $\delta_\circ$  (with two hollow vertices) and *solid diagonals*  $\delta_\bullet$  (with two solid vertices), but we never consider diagonals with one hollow vertex and one solid vertex. Similarly, we consider *hollow dissections*  $D_\circ$  (of the hollow polygon, with only hollow diagonals) and *solid dissections*  $D_\bullet$  (of the solid polygon, with only solid diagonals), but never mix hollow and solid diagonals in a dissection. To help distinguish them, hollow (resp. solid) vertices and diagonals appear red (resp. blue) in all pictures.

We fix an arbitrary reference hollow dissection  $D_\circ$ . A solid diagonal  $\delta_\bullet$  is a  *$D_\circ$ -accordion diagonal* if the hollow diagonals of  $\bar{D}_\circ$  crossed by  $\delta_\bullet$  form an accordion of  $D_\circ$ . In other words,  $\delta_\bullet$  cannot enter and exit a cell of  $D_\circ$  using two non-incident diagonals. For example, note that for any hollow diagonal  $i_\circ j_\circ \in \bar{D}_\circ$ , the solid diagonals  $(i-1)_\bullet(j-1)_\bullet$  and  $(i+1)_\bullet(j+1)_\bullet$  are  $D_\circ$ -accordion diagonals (here and throughout, labels are considered modulo  $2n$ ). In particular, all boundary edges of the solid polygon are  $D_\circ$ -accordion diagonals. A  *$D_\circ$ -accordion dissection* is a set of non-crossing internal  $D_\circ$ -accordion diagonals. We define the  *$D_\circ$ -accordion complex* to be the simplicial complex  $\mathcal{AC}(D_\circ)$  of  $D_\circ$ -accordion dissections.

**Example 1.** As a running example, we consider the reference dissection  $D_\circ^{\text{ex}}$  of Figure 2 (left). Examples of maximal  $D_\circ^{\text{ex}}$ -accordion dissections are given in Figure 2 (right). The  $D_\circ^{\text{ex}}$ -accordion complex is illustrated in Figure 3 (left).




FIGURE 2. A hollow dissection  $D_\circ^{\text{ex}}$ , a solid  $D_\circ^{\text{ex}}$ -accordion diagonal whose corresponding hollow accordion is bolded, and two maximal solid  $D_\circ^{\text{ex}}$ -accordion dissections.




FIGURE 3. The  $D_0^{\text{ex}}$ -accordion complex (left) and the  $D_0^{\text{ex}}$ -accordion lattice (right), oriented from bottom to top, for the reference hollow dissection  $D_0^{\text{ex}}$  of Figure 2 (left).

**Example 2.** Special reference hollow dissections  $D_\circ$  give rise to special accordion complexes  $\mathcal{AC}(D_\circ)$ :

- ◊ If  $D_\circ$  is the empty dissection with the whole hollow polygon as unique cell, then the  $D_\circ$ -accordion complex  $\mathcal{AC}(D_\circ)$  is reduced to the empty  $D_\circ$ -accordion dissection.
- ◊ If  $D_\circ$  has a unique internal diagonal, then the  $D_\circ$ -accordion complex  $\mathcal{AC}(D_\circ)$  consists of only two points.
- ◊ For a hollow triangulation  $T_\circ$ , all solid diagonals are  $T_\circ$ -accordions, so that the  $T_\circ$ -accordion complex  $\mathcal{AC}(T_\circ)$  is the simplicial associahedron.
- ◊ For a hollow quadrangulation  $Q_\circ$ , a solid diagonal is a  $Q_\circ$ -accordion if and only if it does not cross two opposite edges of a quadrangle of  $Q_\circ$ . The  $Q_\circ$ -accordion complex  $\mathcal{AC}(Q_\circ)$  is thus the Stokes complex defined by Y. Baryshnikov [Bar01] and studied by F. Chapoton [Cha16].

**Remark 3.** Following the original definition of the non-crossing complex of A. Garver and T. McConville [GM16], the accordion complex could equivalently be defined in terms of the dual tree  $D_\circ^*$  of  $D_\circ$  (with one node in each cell of  $D$  and one edge connecting two adjacent cells). More precisely, the duality provides the following dictionary between the two definitions:

present paper		A. Garver and T. McConville [GM16]
reference dissection $D_\circ$	$\longleftrightarrow$	embedded tree $D_\circ^*$
diagonal $u_\bullet v_\bullet$ of $P_\bullet$	$\longleftrightarrow$	path connecting the leaves $u_\bullet^*$ and $v_\bullet^*$ of $D_\circ^*$
$D_\circ$ -accordion diagonal	$\longleftrightarrow$	arc (path where any two consecutive edges belong to the boundary of a face of the complement of $D_\circ^*$ in the unit disk)
$D_\circ$ -subaccordion	$\longleftrightarrow$	segment
$D_\circ$ -accordion complex	$\longleftrightarrow$	non-crossing complex of $D_\circ^*$

The **g**-, **c**- and **d**-vectors defined in Section 2.1 could as well be defined in terms of  $D_\circ^*$ . In fact, **c**-vectors were already implicitly considered in [GM16], while **g**- and **d**-vectors are new in this context. For this paper, we find more convenient to work directly with dissections, in particular in Sections 3 and 4.

**1.2. Two structural observations.** Before studying the accordion complex in details in Section 1.3, we present two simple structural observations. For this, let us recall two classical notions on simplicial complexes. The *join* of two simplicial complexes  $\Delta, \Delta'$  with disjoint ground sets  $X, X'$  is the simplicial complex  $\Delta * \Delta'$  with ground set  $X \sqcup X'$  whose faces are disjoint unions of faces

of  $\Delta$  with faces of  $\Delta'$ . For a face  $D$  in a simplicial complex  $\Delta$  on  $X$ , the *link* of  $D$  is the simplicial complex on  $X \setminus D$  whose faces are the subsets  $D'$  of  $X \setminus D$  such that  $D \cup D'$  is a face of  $\Delta$ .

**Proposition 4.** *If the reference hollow dissection  $D_\circ$  has a cell containing  $p$  boundary edges of the hollow polygon  $P_\circ$ , then the  $D_\circ$ -accordion complex  $\mathcal{AC}(D_\circ)$  is the join of  $p$  accordion complexes.*

*Proof.* Assume that  $D_\circ$  has a cell  $C_\circ$  containing  $p$  boundary edges of the hollow polygon  $P_\circ$ . Let  $C_\circ^1, \dots, C_\circ^p$  denote the  $p$  (possibly empty) connected components of the hollow polygon minus  $C_\circ$ . For  $i \in [p] := \{1, \dots, p\}$ , let  $D_\circ^i$  denote the dissection formed by the cell  $C_\circ$  together with the cells of  $D_\circ$  contained in the closure of  $C_\circ^i$ . Observe that for  $i \neq j$ , the internal diagonals of  $D_\circ^i$  are not incident to the internal diagonals of  $D_\circ^j$ . Thus, no  $D_\circ$ -accordion can contain internal diagonals from distinct dissections  $D_\circ^i$  and  $D_\circ^j$ . Therefore, the set of  $D_\circ$ -accordion diagonals is the union of the sets of  $D_\circ^i$ -accordion diagonals for  $i \in [p]$ . Moreover, for  $i \neq j$ , the  $D_\circ^i$ -accordion diagonals do not cross the  $D_\circ^j$ -accordion diagonals. It follows that the  $D_\circ$ -accordion complex is the join of the  $D_\circ^i$ -accordion complexes:  $\mathcal{AC}(D_\circ) = \mathcal{AC}(D_\circ^1) * \dots * \mathcal{AC}(D_\circ^p)$ .  $\square$

**Remark 5.** In view of Proposition 4, we can do the following reductions:

- (i) If a non-triangular cell of  $D_\circ$  has two consecutive boundary edges  $\gamma_\circ, \delta_\circ$  of the hollow polygon, then contracting  $\gamma_\circ$  and  $\delta_\circ$  to a single boundary edge preserves the  $D_\circ$ -accordion complex.
- (ii) If a cell of  $D_\circ$  has two non-consecutive boundary edges of the hollow polygon, then the  $D_\circ$ -accordion complex is a join of smaller accordion complexes.

In all the examples of the paper, we therefore only consider dissections where any non-triangular cell of  $D_\circ$  has at most one boundary edge. All of our constructions work in general, but are just obtained as products or joins of the non-degenerate situation.

**Proposition 6.** *The links in an accordion complex are joins of accordion complexes.*

*Proof.* Consider a  $D_\circ$ -accordion dissection  $D_\bullet$  with cells  $C_\bullet^1, \dots, C_\bullet^p$ . Let  $D_\circ^i$  denote the hollow dissection obtained from  $D_\circ$  by contracting all hollow boundary edges which do not cross  $C_\bullet^i$ . Then a diagonal  $\delta_\bullet$  of a cell  $C_\bullet^i$  is a  $D_\circ$ -accordion diagonal if and only if it is a  $D_\circ^i$ -accordion diagonal. Moreover, for  $i \neq j$ , the diagonals of  $C_\bullet^i$  do not cross the diagonals of  $C_\bullet^j$ . It follows that the link of  $D_\bullet$  in  $\mathcal{AC}(D_\circ)$  is isomorphic to the join  $\mathcal{AC}(D_\circ^1) * \dots * \mathcal{AC}(D_\circ^p)$ .  $\square$

**1.3. Pseudo-manifold.** We now prove that the accordion complex  $\mathcal{AC}(D_\circ)$  is a *pseudomanifold*, *i.e.* that it is:

- (i) *pure*: all maximal  $D_\circ$ -accordion dissections have the same number of diagonals as  $D_\circ$ , and
- (ii) *thin*: any codimension 1 simplex of  $\mathcal{AC}(D_\circ)$  is contained in exactly two maximal  $D_\circ$ -accordion dissections.

We follow the arguments of A. Garver and T. McConville [GM16] (except that they work on the dual tree of the dissection  $D_\circ$ ). A much more concise but less instructive proof of the pseudomanifold property will be derived from geometric considerations in Remark 60.

Recall that we denote by  $\overline{D}_\circ$  the set formed by  $D_\circ$  together with all boundary edges of the hollow polygon. An *angle*  $u_\circ v_\circ w_\circ$  of  $\overline{D}_\circ$  is a pair  $\{u_\circ v_\circ, v_\circ w_\circ\}$  of two consecutive diagonals of  $\overline{D}_\circ$  around a common vertex  $v_\circ$ , called the *apex*. Note that  $\overline{D}_\circ$  has  $2|D_\circ| + n = 2|\overline{D}_\circ| - n$  angles. Observe also that an accordion  $A_\circ$  of  $D_\circ$  can be seen as a sequence of  $|A_\circ| - 1$  angles where two consecutive angles are separated by a diagonal of  $A_\circ$ . We say that a solid vertex  $p_\bullet$  belongs to an angle  $u_\circ v_\circ w_\circ$  if it lies in the cone generated by the edges  $v_\circ u_\circ$  and  $v_\circ w_\circ$  of the angle. The main observation is given in the following statement.

**Lemma 7.** *Let  $D_\bullet$  be a maximal  $D_\circ$ -accordion dissection, and let  $p_\bullet, q_\bullet, r_\bullet, s_\bullet$  denote four consecutive vertices of a cell  $C_\bullet$  of  $D_\bullet$  (with possibly  $p_\bullet = s_\bullet$  if  $C_\bullet$  is a triangle). Then  $p_\bullet$  and  $s_\bullet$  belong to the same angle of the accordion of  $\overline{D}_\circ$  which is crossed by  $q_\bullet r_\bullet$ .*

*Proof.* Let  $A_\circ$  be the accordion of  $\overline{D}_\circ$  which is crossed by  $q_\bullet r_\bullet$ . Assume that  $p_\bullet$  and  $s_\bullet$  belong to distinct angles of  $A_\circ$ . Then they are separated by a diagonal  $\varepsilon_\circ$  of  $A_\circ$ . Therefore, there are two boundary edges  $q_\bullet r_\bullet$  and  $u_\bullet v_\bullet$  of  $C_\bullet$  with distinct vertices such that the hollow diagonal  $\varepsilon_\circ$  separates the vertices  $q_\bullet, u_\bullet$  from the vertices  $r_\bullet, v_\bullet$ . Let  $\gamma_\circ^1, \dots, \gamma_\circ^i = \varepsilon_\circ, \dots, \gamma_\circ^a$  (resp.  $\delta_\circ^1, \dots, \delta_\circ^j = \varepsilon_\circ, \dots, \delta_\circ^b$ )






FIGURE 4. Two maximal  $D_0$ -accordion dissection  $D_•$  (left) and  $D'_•$  (right) related by the flip of  $\delta_•$  to  $\delta'_•$ . The angles of  $D_0$  closed by  $\delta_•$  and  $\delta'_•$  are shaded. The flip is oriented from  $D_•$  to  $D'_•$ .

denote the diagonals of  $D_0$  crossed by  $q_•r_•$  from  $q_•$  to  $r_•$  (resp. crossed by  $u_•v_•$  from  $u_•$  to  $v_•$ ). Then the hollow diagonals  $\gamma_0^1, \dots, \gamma_0^i = \varepsilon_0 = \delta_0^j, \dots, \delta_0^b$  which are crossed by  $q_•v_•$  also form an accordion. It follows that  $D_•$  is not maximal as we can still include  $q_•v_•$ .  $\square$

Consider now an angle  $u_0v_0w_0$  of  $\overline{D_0}$ . In any maximal  $D_0$ -accordion dissection  $D_•$ , the set  $X_•$  of diagonals of  $\overline{D_•}$  that cross both  $u_0v_0$  and  $v_0w_0$  is non-empty (since it contains the boundary edge  $(v-1)_•(v+1)_•$ ) and totally ordered (since the diagonals of  $D_•$  do not cross). Let  $\delta_•$  be the largest diagonal of  $X_•$  (meaning the farthest from  $v_0$ ). We say that the diagonal  $\delta_•$  *closes* the angle  $u_0v_0w_0$ . Note that each angle of  $\overline{D_0}$  is closed by precisely one diagonal of  $\overline{D_•}$ . The following lemma is stated and proved in [GM16] in terms of the dual tree  $D_0^*$  of the dissection  $D_0$ .

**Lemma 8** ([GM16]). *For any maximal  $D_0$ -accordion dissection  $D_•$ , each internal diagonal  $\delta_•$  of  $D_•$  closes two angles of  $\overline{D_0}$  (one apex on each side of  $\delta_•$ ) while each boundary edge of the solid polygon closes one angle of  $\overline{D_0}$ . Therefore the accordion complex  $\mathcal{AC}(D_0)$  is pure of dimension  $|D_0|$ .*

*Proof.* The first sentence is a consequence of Lemma 7: for any four consecutive vertices  $p_•, q_•, r_•, s_•$  of a cell of  $\overline{D_•}$ , the diagonal  $q_•r_•$  closes the unique angle of the accordion of  $\overline{D_0}$  crossed by  $q_•r_•$  that contains the vertices  $p_•$  and  $s_•$ . Therefore,  $q_•r_•$  closes precisely two angles (resp. one angle) of  $D_0$  if it is an internal diagonal (resp. a boundary edge of the solid polygon). We finally obtain by double-counting that  $2|D_0| + n = |\{\text{angles of } \overline{D_0}\}| = 2|D_•| + n$  and thus  $|D_•| = |D_0|$  for any maximal  $D_0$ -accordion dissection  $D_•$ .  $\square$

We are now ready to prove that the  $D_0$ -accordion complex is thin, *i.e.* that each internal diagonal of a maximal  $D_0$ -accordion dissection can be flipped into a unique other internal diagonal to form a new maximal  $D_0$ -accordion dissection. Here and throughout the paper,  $X \triangle Y$  denotes the symmetric difference of two sets  $X, Y$  defined by  $X \triangle Y := (X \setminus Y) \cup (Y \setminus X)$ .

The following notations are illustrated in Figure 4. Let  $D_•$  be a maximal  $D_0$ -accordion dissection and  $\delta_•$  be a diagonal of  $D_•$ . Let  $u_0$  and  $v_0$  be the apices of the angles of  $D_0$  closed by  $\delta_•$ , let  $\mu_•$  and  $\nu_•$  denote the edges of the cells of  $D_•$  containing  $\delta_•$ , which separate  $\delta_•$  from  $u_0$  and  $v_0$  respectively, and let  $Q_•$  denote the quadrilateral defined by the four vertices of  $\mu_•$  and  $\nu_•$ . Note that  $\delta_•$  is a diagonal of  $Q_•$ , and let  $\delta'_•$  denote the other diagonal.

**Lemma 9** ([GM16]). *With the previous notations, the collection of diagonals  $D'_• := D_• \triangle \{\delta_•, \delta'_•\}$  is a maximal  $D_0$ -accordion dissection, and  $D_•$  and  $D'_•$  are the only maximal  $D_0$ -accordion dissections containing  $D_• \setminus \{\delta_•\}$ . In other words, the accordion complex  $\mathcal{AC}(D_0)$  is thin.*

*Proof.* We first observe that  $\delta'_•$  is a  $D_0$ -accordion diagonal, since the edges of  $\overline{D_0}$  crossed by  $\delta'_•$  are obtained by merging three subaccordions of  $D_0$ : the subaccordion formed by the diagonals of  $\overline{D_0}$  crossed by  $\mu_•$  but not  $\delta_•$  nor  $\nu_•$ , the subaccordion formed by the diagonals of  $\overline{D_0}$  crossed by  $\delta_•, \mu_•$  and  $\nu_•$ , and the subaccordion formed by the diagonals of  $\overline{D_0}$  crossed by  $\nu_•$  but not  $\delta_•$  nor  $\mu_•$ .

Moreover,  $\delta_\bullet$  and  $\delta'_\bullet$  are the only  $D_\circ$ -accordion diagonals compatible with  $D_\bullet \setminus \{\delta_\bullet\}$ . Indeed, any other such diagonal would cross  $\delta_\bullet$  and  $\delta'_\bullet$  (by maximality of  $D_\bullet$  and  $D'_\bullet$ ), and thus also the subaccordion  $A_\circ$  of  $D_\circ$  crossed by  $\delta_\bullet$  and  $\delta'_\bullet$  (because it cannot cross  $\mu$  and  $\nu$ ). But it would then improperly intersect the two cells of  $D_\circ$  containing precisely one diagonal of  $A_\circ$ .  $\square$

The  $D_\circ$ -*accordion flip graph* is the dual graph  $\mathcal{AFG}(D_\circ)$  of the  $D_\circ$ -accordion complex: its vertices are the maximal  $D_\circ$ -accordion dissections, and its edges are the *flips* between them, *i.e.* the pairs  $\{D_\bullet, D'_\bullet\}$  of maximal  $D_\circ$ -accordion dissections with  $D_\bullet \setminus \{\delta_\bullet\} = D'_\bullet \setminus \{\delta'_\bullet\}$ . See Figure 3 (right).

**1.4. The accordion lattice.** We now define a natural orientation on the  $D_\circ$ -accordion flip graph. We use the same notations as in Lemma 9 (see also Figure 4), where  $D_\bullet \setminus \{\delta_\bullet\} = D'_\bullet \setminus \{\delta'_\bullet\}$  and  $\delta_\bullet, \delta'_\bullet$  are the two diagonals of the quadrilateral defined by  $\mu_\bullet, \nu_\bullet$ . Observe that one of the path  $\mu_\bullet \delta_\bullet \nu_\bullet$  and  $\mu_\bullet \delta'_\bullet \nu_\bullet$  forms a  $\Sigma$  while the other forms a  $Z$ , see Figure 4. We then orient the flip from the dissection containing the  $\Sigma$  to that containing the  $Z$ . See Figure 3 (right) for an illustration of  $D_\circ$ -accordion oriented flip graph (where the graph is oriented from bottom to top).

A. Garver and T. McConville introduced a natural closure on sets of  $D_\circ$ -subaccordions, and showed that the inclusion poset of biclosed sets of  $D_\circ$ -subaccordions is a well-behaved lattice (namely, semidistributive, congruence-uniform and polygonal). Then, they introduced a lattice quotient map from biclosed sets of  $D_\circ$ -subaccordions to maximal  $D_\circ$ -accordion dissections, which imply the following statement.

**Theorem 10** ([GM16]). *The  $D_\circ$ -accordion oriented flip graph is the Hasse diagram of a lattice, that we call the  $D_\circ$ -accordion lattice and denote by  $\mathcal{AL}(D_\circ)$ .*

In particular, the  $D_\circ$ -accordion oriented flip graph is connected and acyclic, and has a unique source  $D_\bullet^- := \{(i-1)_\bullet(j-1)_\bullet \mid i_\circ j_\circ \in D_\circ\}$  (obtained by slightly rotating  $D_\circ$  counterclockwise) and a unique sink  $D_\bullet^+ := \{(i+1)_\bullet(j+1)_\bullet \mid i_\circ j_\circ \in D_\circ\}$  (obtained by slightly rotating  $D_\circ$  clockwise).

**Example 11.** Following Example 2, note that special reference hollow dissections  $D_\circ$  give rise to special accordion lattices  $\mathcal{AL}(D_\circ)$ , as it was already observed in [GM16]:

- ◊ For a fan triangulation  $F_\circ$  (*i.e.* where all internal diagonals are incident to a common vertex), the  $F_\circ$ -accordion lattice  $\mathcal{AL}(F_\circ)$  is the famous Tamari lattice [Tam51, MHPS12] defined equivalently by slope increasing flips on triangulations of a convex polygon, by right rotations on binary trees, or by flips on Dyck paths.
- ◊ In general, accordion lattices of accordion triangulations (*i.e.* with no interior triangle) precisely correspond to type *A* Cambrian lattices defined by N. Reading [Rea06].
- ◊ For an arbitrary triangulation  $T_\circ$  (with or without interior triangle), the  $T_\circ$ -accordion oriented flip graph  $\mathcal{AFG}(A_\circ)$  is a particular instance of the oriented exchange graphs of 2-acyclic quivers defined by T. Brüstle, G. Dupont and M. Pérotin [BDP14]. These oriented exchange graphs are far more general and their transitive closures are in general not lattices.
- ◊ For a quadrangulation  $Q_\circ$ , the  $Q_\circ$ -accordion lattice  $\mathcal{AL}(Q_\circ)$  is the Stokes poset on  $Q_\circ$ -compatible quadrangulations studied by F. Chapoton [Cha16].

The following statement is a direct consequence of Proposition 4.

**Proposition 12.** *If the reference hollow dissection  $D_\circ$  has a cell containing  $p$  boundary edges of the hollow polygon  $P_\circ$ , then the  $D_\circ$ -accordion lattice  $\mathcal{AL}(D_\circ)$  is a Cartesian product of  $p$  accordion lattices.*

*Proof.* Consider the dissections  $D_\circ^1, \dots, D_\circ^p$  as in the proof of Proposition 4. Since any increasing flip in  $\mathcal{AL}(D_\circ)$  is an increasing flip in one of the  $\mathcal{AL}(D_\circ^i)$ , we obtain that the  $D_\circ$ -accordion lattice is the Cartesian product of the  $D_\circ^i$ -accordion lattices:  $\mathcal{AL}(D_\circ) = \mathcal{AL}(D_\circ^1) \times \dots \times \mathcal{AL}(D_\circ^p)$ .  $\square$

In particular, if two consecutive boundary edges  $\gamma_\circ, \delta_\circ$  of the hollow polygon belong to the same non-triangular cell of  $D_\circ$ , then contracting  $\gamma_\circ$  and  $\delta_\circ$  to a single boundary edge preserves the  $D_\circ$ -accordion lattice. This shows the following statement conjectured for quadrangulations in [Cha16] and proved in [BMP16].



**Corollary 13.** *Consider an accordion dissection  $A_\circ$ , i.e. a dissection where each cell has at most 2 edges which are internal diagonals of  $P_\circ$ . Then the  $A_\circ$ -accordion lattice is a Cambrian lattice.*

**Remark 14.** Call *cell-sequence* of a dissection the sequence whose  $i$ th entry is its number of  $(i+2)$ -cells. For example, the dissection of Figure 2 (left) has cell-sequence  $3, 1, 0^\infty$  and all  $(p+2)$ -angulations of a  $(pm+2)$ -gon have cell-sequence  $0^{p-1}, m, 0^\infty$ . Observe that the flip preserves the cell-sequence. Thus, all maximal  $D_\circ$ -accordion dissections have the same cell-sequence as  $D_\circ$ .

We conclude this section with a reciprocity result on accordion dissections.

**Proposition 15.** *Let  $D_\circ$  be a hollow dissection and  $D_\bullet$  be a solid dissection. Then  $D_\bullet$  is a maximal  $D_\circ$ -accordion dissection if and only if  $D_\circ$  is a maximal  $D_\bullet$ -accordion dissection.*

*Proof.* Since  $D_\bullet^- := \{(i-1)\bullet(j-1)\bullet \mid i_\circ j_\circ \in D_\circ\}$  and  $D_\bullet^+ := \{(i+1)\bullet(j+1)\bullet \mid i_\circ j_\circ \in D_\circ\}$  are both  $D_\circ$ -accordion dissections, we already know that  $D_\circ$  is a  $D_\bullet^-$ -accordion dissection. Observe now in Figure 4 that if  $D_\bullet$  and  $D'_\bullet$  are maximal  $D_\circ$ -accordion dissections connected by a flip, then  $D_\circ$  is a  $D_\bullet$ -accordion dissection if and only if it is a  $D'_\bullet$ -accordion dissection. Indeed, if  $\delta_\bullet$  belongs to the zigzag of the  $D_\bullet$ -accordion  $A_\bullet$  of a hollow diagonal  $\delta_\circ$ , then  $\delta_\circ$  crosses both  $\mu_\bullet$  and  $\nu_\bullet$ , but then  $\delta_\circ$  also crosses  $\delta'_\bullet$ , and thus  $\delta_\circ$  crosses the  $D'_\bullet$ -accordion  $A_\bullet \triangle \{\delta_\bullet, \delta'_\bullet\}$ . Since the  $D_\circ$ -accordion flip graph is connected, we obtain that  $D_\circ$  is a  $D_\bullet$ -accordion dissection for any maximal  $D_\circ$ -accordion dissection  $D_\bullet$ . Finally, maximality follows since all maximal  $D_\circ$ -accordion dissections have  $|D_\circ|$  diagonals. The equivalence follows by symmetry.  $\square$

## 2. THE $\mathbf{g}$ -VECTOR FAN

In this Section, we construct accordiohedra using  $\mathbf{g}$ - and  $\mathbf{c}$ -vectors. Our construction is in the same spirit as the Cambrian fans of N. Reading and D. Speyer [Rea06, Rea07, RS09] and their polytopal realizations by C. Hohlweg, C. Lange and H. Thomas [HL07, HLT11], recently extended in [HPS18] to any initial triangulation, acyclic or not. A different approach to the  $\mathbf{g}$ -vector fan together with an alternative polytopal realization will be presented in Section 4.

**2.1.  $\mathbf{g}$ - and  $\mathbf{c}$ -vectors.** Consider a hollow dissection  $D_\circ$  and a solid dissection  $D_\bullet$  that are maximal accordion dissections of each other (see Proposition 15), and let  $\delta_\circ \in D_\circ$  and  $\delta_\bullet \in D_\bullet$ . When  $\delta_\circ$  crosses  $\delta_\bullet$ , we let  $\mu_\circ$  and  $\nu_\circ$  be the other diagonals of  $\bar{D}_\circ$  crossed by  $\delta_\bullet$  in the two cells of  $D_\circ$  containing  $\delta_\circ$ . We say that  $\delta_\bullet$  *slaloms* on  $\delta_\circ$  if  $\mu_\circ \delta_\circ \nu_\circ$  forms a path, and we define  $\varepsilon_\circ(\delta_\circ \in D_\circ \mid \delta_\bullet)$  to be 1,  $-1$ , or 0 depending on whether  $\mu_\circ \delta_\circ \nu_\circ$  forms a  $\mathbf{Z}$ , a  $\mathbf{\Sigma}$ , or a  $\mathbf{V}$ . Similarly we let  $\mu_\bullet$  and  $\nu_\bullet$  be the other diagonals of  $\bar{D}_\bullet$  crossed by  $\delta_\circ$  in the two cells of  $D_\bullet$  containing  $\delta_\bullet$ , we say that  $\delta_\circ$  slaloms on  $\delta_\bullet$  if  $\mu_\bullet \delta_\bullet \nu_\bullet$  forms a path, and we define  $\varepsilon_\bullet(\delta_\circ \mid \delta_\bullet \in D_\bullet)$  to be 1,  $-1$ , or 0 depending on whether  $\mu_\bullet \delta_\bullet \nu_\bullet$  forms a  $\mathbf{\Sigma}$ , a  $\mathbf{Z}$ , or a  $\mathbf{V}$ . Note that the sign convention for  $\varepsilon_\circ(\delta_\circ \in D_\circ \mid \delta_\bullet)$  and  $\varepsilon_\bullet(\delta_\circ \mid \delta_\bullet \in D_\bullet)$  is opposite: the reciprocity already observed in Proposition 15 naturally reverses the orientation. More informally, we exchange the role of hollow and solid dissections by looking at the picture from the opposite side of the blackboard, which of course reverses the orientation. Finally, if  $\delta_\circ$  and  $\delta_\bullet$  do not cross, then we let  $\varepsilon_\circ(\delta_\circ \in D_\circ \mid \delta_\bullet) = \varepsilon_\bullet(\delta_\circ \mid \delta_\bullet \in D_\bullet) = 0$ . Let  $(\mathbf{e}_{\delta_\circ})_{\delta_\circ \in D_\circ}$  denote the canonical basis of  $\mathbb{R}^{D_\circ}$ . As in [HPS18], we define the following vectors:

- (i) the  **$\mathbf{g}$ -vector** of  $\delta_\bullet$  with respect to  $D_\circ$  is  $\mathbf{g}(D_\circ \mid \delta_\bullet) := \sum_{\delta_\circ \in D_\circ} \varepsilon_\circ(\delta_\circ \in D_\circ \mid \delta_\bullet) \mathbf{e}_{\delta_\circ}$ . We also define  $\mathbf{g}(D_\circ \mid D_\bullet) := \{\mathbf{g}(D_\circ \mid \delta_\bullet) \mid \delta_\bullet \in D_\bullet\}$ .
- (ii) the  **$\mathbf{c}$ -vector** of  $\delta_\bullet \in D_\bullet$  with respect to  $D_\circ$  is  $\mathbf{c}(D_\circ \mid \delta_\bullet \in D_\bullet) := \sum_{\delta_\circ \in D_\circ} \varepsilon_\bullet(\delta_\circ \mid \delta_\bullet \in D_\bullet) \mathbf{e}_{\delta_\circ}$ . We denote by  $\mathbf{c}(D_\circ \mid D_\bullet) := \{\mathbf{c}(D_\circ \mid \delta_\bullet \in D_\bullet) \mid \delta_\bullet \in D_\bullet\}$  the set of  $\mathbf{c}$ -vectors of the diagonals of  $D_\bullet$  and by  $\mathbf{C}(D_\circ) := \bigcup_{D_\bullet} \mathbf{c}(D_\circ \mid D_\bullet)$  the set of all  $\mathbf{c}$ -vectors with respect to  $D_\circ$ .

**Example 16.** Consider the hollow dissection  $D_\circ^{\text{ex}} = \{3_\circ 7_\circ, 3_\circ 13_\circ, 9_\circ 13_\circ\}$  and the rightmost solid dissection  $D_\bullet^{\text{ex}} = \{2_\bullet 6_\bullet, 2_\bullet 10_\bullet, 10_\bullet 14_\bullet\}$  of Figure 2. Then we have for example

- $\diamond \varepsilon_\circ(3_\circ 13_\circ \in D_\circ^{\text{ex}} \mid 2_\bullet 10_\bullet) = 1$  since the path  $1_\circ - 3_\circ - 13_\circ - 9_\circ$  forms a  $\mathbf{Z}$ ,
- $\diamond \varepsilon_\circ(9_\circ 13_\circ \in D_\circ^{\text{ex}} \mid 2_\bullet 10_\bullet) = -1$  since the path  $3_\circ - 13_\circ - 9_\circ - 11_\circ$  forms a  $\mathbf{\Sigma}$ , and
- $\diamond \varepsilon_\circ(3_\circ 13_\circ \in D_\circ^{\text{ex}} \mid 2_\bullet 6_\bullet) = 0$  since  $3_\circ$  connects  $1_\circ, 13_\circ, 7_\circ$  as a  $\mathbf{V}$ .

Moreover, we have

$$\begin{aligned} \mathbf{g}(D_\circ^{\text{ex}} | 2_\bullet 6_\bullet) &= \mathbf{e}_{3_\circ 7_\circ}, & \mathbf{c}(D_\circ^{\text{ex}} | 2_\bullet 6_\bullet \in D_\bullet^{\text{ex}}) &= \mathbf{e}_{3_\circ 7_\circ}, \\ \mathbf{g}(D_\circ^{\text{ex}} | 2_\bullet 10_\bullet) &= \mathbf{e}_{3_\circ 13_\circ} - \mathbf{e}_{9_\circ 13_\circ}, & \mathbf{c}(D_\circ^{\text{ex}} | 2_\bullet 10_\bullet \in D_\bullet^{\text{ex}}) &= \mathbf{e}_{3_\circ 13_\circ}, \\ \mathbf{g}(D_\circ^{\text{ex}} | 10_\bullet 14_\bullet) &= -\mathbf{e}_{9_\circ 13_\circ}, & \mathbf{c}(D_\circ^{\text{ex}} | 10_\bullet 14_\bullet \in D_\bullet^{\text{ex}}) &= -\mathbf{e}_{3_\circ 13_\circ} - \mathbf{e}_{9_\circ 13_\circ}. \end{aligned}$$

**Example 17.** For any hollow diagonal  $i_\circ j_\circ \in D_\circ$ , we have

$$\begin{aligned} \mathbf{g}(D_\circ | (i-1)_\bullet (j-1)_\bullet) &= -\mathbf{e}_{i_\circ j_\circ}, & \mathbf{c}(D_\circ | (i-1)_\bullet (j-1)_\bullet \in D_\bullet^-) &= -\mathbf{e}_{i_\circ j_\circ}, \\ \mathbf{g}(D_\circ | (i+1)_\bullet (j+1)_\bullet) &= \mathbf{e}_{i_\circ j_\circ}, & \mathbf{c}(D_\circ | (i+1)_\bullet (j+1)_\bullet \in D_\bullet^+) &= \mathbf{e}_{i_\circ j_\circ}. \end{aligned}$$

**Remark 18.** For a hollow triangulation  $T_\circ$ , our definitions of  $\mathbf{g}$ - and  $\mathbf{c}$ -vectors coincide with the shear coordinates of S. Fomin and D. Thurston [FT12], defined in the much more general context of cluster algebras on surfaces [FST08].

**Remark 19.** Consider the quiver  $Q(D_\circ)$  of the reference dissection  $D_\circ$ , with one node on each internal diagonal of  $D_\circ$  and one arrow between two diagonals counter-clockwise consecutive around a cell of  $D_\circ$ . Let  $W(D_\circ)$  be the reflection group whose Dynkin diagram is the underlying graph of  $Q(D_\circ)$ . Then all  $\mathbf{g}$ -vectors of the  $D_\circ$ -accordion diagonals are weights of  $W(D_\circ)$  and all  $\mathbf{c}$ -vectors of  $\mathbf{C}(D_\circ)$  are roots of  $W(D_\circ)$ .

**Remark 20.** Informally, the  $\mathbf{g}$ - and  $\mathbf{c}$ -vectors can be interpreted as follows:

- (i) The  $\mathbf{g}$ -vector  $\mathbf{g}(D_\circ | \delta_\bullet)$  has coordinate 1 and  $-1$  alternating along the zigzag of the accordion crossed by  $\delta_\bullet$  in  $D_\circ$ , and coordinate 0 on all other diagonals of  $D_\circ$ .
- (ii) The  $\mathbf{c}$ -vector  $\mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet)$  is, up to a sign, the characteristic vector of the diagonals of the subaccordion of  $D_\circ$  crossed by both diagonals  $\mu_\bullet$  and  $\nu_\bullet$  of Lemma 9 (see also Figure 4). Thus, any  $\mathbf{c}$ -vector is either *positive* (only non-negative coordinates) or *negative* (only non-positive coordinates).

In fact, the  $\mathbf{g}$ -vectors are clearly in bijection with the accordions and with the zigzags in  $D_\circ$ . In contrast, many pairs  $(\delta_\bullet, D_\bullet)$  produce the same  $\mathbf{c}$ -vector  $\mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet)$ . For example, if two dissections  $D_\bullet, D'_\bullet$  contain  $\delta_\bullet$  and have the same cells incident to  $\delta_\bullet$ , then  $\mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet) = \mathbf{c}(D_\circ | \delta_\bullet \in D'_\bullet)$ . The set of  $\mathbf{c}$ -vectors  $\mathbf{C}(D_\circ)$  without repetitions can be understood as follows.

**Lemma 21.** *There are bijections between:*

- ◊ the negative (resp. positive)  $\mathbf{c}$ -vectors of  $\mathbf{C}(D_\circ)$ ,
- ◊ the subaccordions of  $D_\circ$ ,
- ◊ the  $D_\circ$ -accordion diagonals not in the source dissection  $D_\bullet^- := \{(i-1)_\bullet (j-1)_\bullet | i_\circ j_\circ \in D_\circ\}$  (resp. not in the sink dissection  $D_\bullet^+ := \{(i+1)_\bullet (j+1)_\bullet | i_\circ j_\circ \in D_\circ\}$ ).

*Proof.* By Remark 20 (ii), the support of any  $\mathbf{c}$ -vector is a subaccordion of  $D_\circ$ . Reciprocally, let  $A_\circ$  be a subaccordion of  $D_\circ$ , let  $C_\circ$  and  $C'_\circ$  denote the two cells of  $D_\circ$  containing exactly one diagonal of  $A_\circ$ , and let  $p_\circ, q_\circ, r_\circ, s_\circ$  (resp.  $p'_\circ, q'_\circ, r'_\circ, s'_\circ$ ) denote the four consecutive vertices in clockwise order around  $C_\circ$  (resp. around  $C'_\circ$ ) such that  $q_\circ r_\circ$  (resp.  $q'_\circ r'_\circ$ ) is the diagonal of  $A_\circ$  in  $C_\circ$  (resp. in  $C'_\circ$ ). Let  $\delta_\bullet := (s-1)_\bullet (s'-1)_\bullet$ ,  $\mu_\bullet := (p+1)_\bullet (s'-1)_\bullet$  and  $\nu_\bullet := (p'+1)_\bullet (s-1)_\bullet$  and consider any  $D_\circ$ -accordion dissection  $D_\bullet$  containing  $\{\mu_\bullet, \delta_\bullet, \nu_\bullet\}$ . Then  $A_\circ$  is precisely the support of the negative  $\mathbf{c}$ -vector  $\mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet)$ . Finally, we have associated to the subaccordion  $A_\circ$  of  $D_\circ$  a  $D_\circ$ -diagonal  $\delta_\bullet = (s-1)_\bullet (s'-1)_\bullet$  which cannot be in  $D_\bullet^-$  as otherwise  $s_\circ s'_\circ$  would cross  $q_\circ r_\circ$ . Reciprocally,  $A_\circ$  is precisely the set of diagonals of  $D_\circ$  crossed by  $\delta_\bullet$  and not incident to  $s_\circ$  or  $s'_\circ$ .  $\square$

The  $\mathbf{g}$ -vectors and  $\mathbf{c}$ -vectors are connected in the following two statements, inspired and motivated by classical analogues in cluster algebra theory.

**Proposition 22.** *For any maximal  $D_\circ$ -accordion dissection  $D_\bullet$ , the set of  $\mathbf{g}$ -vectors  $\mathbf{g}(D_\circ | D_\bullet)$  and the set of  $\mathbf{c}$ -vectors  $\mathbf{c}(D_\circ | D_\bullet)$  form dual bases.*

*Proof.* Let  $\langle \cdot | \cdot \rangle$  denote the standard Euclidean inner product of  $\mathbb{R}^{D_\circ}$ . Given two solid diagonals  $\gamma_\bullet, \delta_\bullet$  of  $D_\bullet$ , we want to compute  $\langle \mathbf{g}(D_\circ | \gamma_\bullet) | \mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet) \rangle$ . By Remark 20 (i), the  $\mathbf{g}$ -vector  $\mathbf{g}(D_\circ | \gamma_\bullet)$  has coordinate  $\pm 1$  alternating along the zigzag  $Z_\circ$  of the accordion crossed by  $\gamma_\bullet$  in  $D_\circ$ , and coordinate 0 on all other diagonals of  $D_\circ$ . Moreover, by Remark 20 (ii), the




FIGURE 5. Illustration of the proof of Proposition 22. The red hollow diagonals form the zigzag of  $\gamma_\bullet$ , and the bolded ones are slaloming on  $\delta_\bullet$ . There are an even number of bolded diagonals when  $\gamma_\bullet \neq \delta_\bullet$  (left) and an odd number when  $\gamma_\bullet = \delta_\bullet$  (right).

$\mathbf{c}$ -vector  $\mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet)$  has coordinate  $\pm 1$  on the diagonals of  $D_\circ$  which slalom on  $\delta_\bullet$  in  $D_\bullet$ , and coordinate 0 on all other diagonals of  $D_\circ$ . We thus need to understand how the diagonals of  $Z_\circ$  slalom on  $\delta_\bullet$  in  $D_\bullet$ . See Figure 5 for a schematic illustration. Observe that there is an even (resp. odd) number of hollow diagonals of  $Z_\circ$  that slalom on  $\delta_\bullet$  when  $\delta_\bullet \neq \gamma_\bullet$  (resp. when  $\delta_\bullet = \gamma_\bullet$ ). Moreover, since they are non-crossing, all hollow diagonals of  $Z_\circ$  slaloming on  $\delta_\bullet$  do it the same way (either all as a  $\Sigma$  or all as a  $Z$ ). Finally, when  $\gamma_\bullet = \delta_\bullet$ , consider the first hollow diagonal  $\delta_\circ$  of the zigzag  $Z_\circ$  which slaloms on  $\delta_\bullet$ . Then  $\delta_\circ$  slaloms on  $\delta_\bullet$  in the opposite way as  $\delta_\bullet$  slaloms on  $\delta_\circ$ . This shows that

$$\langle \mathbf{g}(D_\circ | \gamma_\bullet) | \mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet) \rangle = \sum_{\delta_\circ \in D_\circ} \varepsilon_\circ(\delta_\circ \in D_\circ | \gamma_\bullet) \cdot \varepsilon_\bullet(\delta_\circ | \delta_\bullet \in D_\bullet) = \mathbb{1}_{\gamma_\bullet = \delta_\bullet},$$

since we sum an even number of alternating  $\pm 1$  when  $\gamma_\bullet \neq \delta_\bullet$ , and an odd number of alternating  $\pm 1$  starting by a 1 when  $\gamma_\bullet = \delta_\bullet$ . In other words,  $\mathbf{g}(D_\circ | D_\bullet)$  and  $\mathbf{c}(D_\circ | D_\bullet)$  form dual bases.  $\square$

**Proposition 23.** *Let  $D_\circ$  be a hollow dissection and  $D_\bullet$  be a solid dissection such that  $D_\circ$  and  $D_\bullet$  are maximal accordion dissections of each other (see Proposition 15). Then*

$$\mathbf{g}(D_\circ | D_\bullet) = -\mathbf{c}(D_\bullet | D_\circ)^t \quad \text{and} \quad \mathbf{c}(D_\circ | D_\bullet) = -\mathbf{g}(D_\bullet | D_\circ)^t,$$

where we consider the sets of  $\mathbf{g}$ -vectors  $\mathbf{g}(D_\circ | D_\bullet)$  and  $\mathbf{c}$ -vectors  $\mathbf{c}(D_\circ | D_\bullet)$  as matrices in  $\mathbb{R}^{D_\circ \times D_\bullet}$ , and  $M^t$  denotes the transpose of a matrix  $M$ .

*Proof.* We immediately derive from the definitions that for any  $\delta_\circ \in D_\circ$  and  $\delta_\bullet \in D_\bullet$ ,

$$\mathbf{g}(D_\circ | D_\bullet)_{(\delta_\circ, \delta_\bullet)} = \varepsilon_\circ(\delta_\circ \in D_\circ | \delta_\bullet) = -\varepsilon_\bullet(\delta_\bullet | \delta_\circ \in D_\circ) = -\mathbf{c}(D_\bullet | D_\circ)_{(\delta_\bullet, \delta_\circ)},$$

which shows  $\mathbf{g}(D_\circ | D_\bullet) = -\mathbf{c}(D_\bullet | D_\circ)^t$ . The other equality follows by exchanging  $D_\circ$  and  $D_\bullet$ .  $\square$

**Corollary 24.** *For any maximal  $D_\circ$ -accordion dissection  $D_\bullet$ , we have the following [sign coherence](#):*

- (i) *for any  $\delta_\bullet \in D_\bullet$ , all coordinates of the  $\mathbf{c}$ -vector  $\mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet)$  have the same sign,*
- (ii) *for any  $\delta_\circ \in D_\circ$ , the  $\delta_\circ$ -coordinates of all  $\mathbf{g}$ -vectors  $\mathbf{g}(D_\circ | \delta_\bullet)$  for  $\delta_\bullet \in D_\bullet$  have the same sign.*

*Proof.* Point (i) was already seen in Remark 20 (ii), and Point (ii) follows by Proposition 23.  $\square$

**2.2.  $\mathbf{c}$ -vector fan and  $D_\circ$ -zonotope.** Define the  $\mathbf{c}$ -vector fan of  $D_\circ$  to be the complete polyhedral fan  $\mathcal{F}^c(D_\circ)$  given by the arrangement of the linear hyperplanes orthogonal to the  $\mathbf{c}$ -vectors of  $\mathbf{C}(D_\circ)$ . Be careful: in contrast to the  $\mathbf{g}$ - and  $\mathbf{d}$ -vector fans defined later, the  $\mathbf{c}$ -vectors are not the rays of  $\mathcal{F}^c(D_\circ)$  but the normal vectors of the hyperplanes supporting the facets of  $\mathcal{F}^c(D_\circ)$ .

We call  $D_\circ$ -zonotope the Minkowski sum  $\text{Zono}(D_\circ)$  of all  $\mathbf{c}$ -vectors:

$$\text{Zono}(D_\circ) := \sum_{\mathbf{c} \in \mathbf{C}(D_\circ)} \mathbf{c}.$$

The normal fan of the  $D_\circ$ -zonotope  $\text{Zono}(D_\circ)$  is the  $\mathbf{c}$ -vector fan  $\mathcal{F}^{\mathbf{c}}(D_\circ)$ . Note that the  $\mathbf{c}$ -vector fan is not always simplicial, and thus the  $D_\circ$ -zonotope  $\text{Zono}(D_\circ)$  is not always simple. See Figure 7.

**Example 25.** Consider an accordion dissection  $A_\circ$  (where each cell has at most 2 edges which are internal diagonals of  $P_\circ$ ). Label its internal diagonals by  $\delta_\circ^1, \dots, \delta_\circ^{|A_\circ|}$  such that  $\delta_\circ^k$  and  $\delta_\circ^{k+1}$  belong to the same cell of  $A_\circ$  for all  $k$ . Identifying  $\mathbf{e}_{\delta_\circ^k}$  to the simple root  $\mathbf{f}_k - \mathbf{f}_{k+1}$  of type  $A_{|A_\circ|}$ , the  $\mathbf{c}$ -vectors of  $\mathbf{C}(A_\circ)$  are all roots  $\pm(\mathbf{f}_i - \mathbf{f}_j) = \pm \sum_{i \leq k \leq j} \mathbf{e}_{\delta_\circ^k}$  of type  $A_{|A_\circ|}$ . Therefore, the  $\mathbf{c}$ -vector fan is the type  $A_{|A_\circ|}$  Coxeter fan and the  $A_\circ$ -zonotope is a permutahedron. More precisely,

$$\text{Zono}(A_\circ) = \sum_{k \in [|A_\circ|+1]} k(|A_\circ| + 1 - k) [-\mathbf{e}_{\delta_\circ^k}, \mathbf{e}_{\delta_\circ^k}] = 2 \text{Perm}(|A_\circ|) - (|A_\circ| + 2) \sum_{i \in [|A_\circ|+1]} \mathbf{f}_i,$$

where  $\text{Perm}(|A_\circ|) := \text{conv} \{ \sum_{i \in [|A_\circ|+1]} \sigma(i) \mathbf{f}_i \mid \sigma \in \mathfrak{S}_{|A_\circ|+1} \}$  is the classical permutahedron.

The vertices of  $\text{Zono}(D_\circ)$  correspond to *separable* subsets of  $\mathbf{C}(D_\circ)$ , *i.e.* those which can be strictly separated from their complement by a hyperplane. Although we could work out all facets of  $\text{Zono}(D_\circ)$ , we will only need the following specific inequalities.

**Proposition 26.** *For any  $D_\circ$ -accordion diagonal  $\gamma_\bullet$ , the  $D_\circ$ -zonotope  $\text{Zono}(D_\circ)$  has a facet defined by the inequality*

$$\langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{x} \rangle \leq \omega(D_\circ | \gamma_\bullet),$$

where  $\omega(D_\circ | \gamma_\bullet)$  is the  *$D_\circ$ -height* of  $\gamma_\bullet$ , *i.e.* the number of  $D_\circ$ -accordion diagonals that cross  $\gamma_\bullet$ .

*Proof.* Let  $\omega(D_\circ | \gamma_\bullet)$  denote the maximum of  $\langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{x} \rangle$  over  $\text{Zono}(D_\circ)$ . As  $\text{Zono}(D_\circ)$  is the Minkowski sum of all  $\mathbf{c}$ -vectors, we have

$$\omega(D_\circ | \gamma_\bullet) = \sum_{\substack{\mathbf{c} \in \mathbf{C}(D_\circ) \\ \langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c} \rangle > 0}} \langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c} \rangle.$$

By Remark 20, we have  $\langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c} \rangle \in \{-1, 0, 1\}$  for any  $\mathbf{c} \in \mathbf{C}(D_\circ)$ . We thus just need to count the distinct  $\mathbf{c}$ -vectors  $\mathbf{c}$  such that  $\langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c} \rangle > 0$ . It turns out that it is more convenient and equivalent (since  $\mathbf{C}(D_\circ) = -\mathbf{C}(D_\circ)$ ) to count the distinct  $\mathbf{c}$ -vectors  $\mathbf{c}$  such that  $\langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c} \rangle < 0$ . For that, let  $Z_\circ$  denote the zigzag of the accordion crossed by  $\gamma_\bullet$  in  $D_\circ$ , and decompose  $Z_\circ = Z_\circ^- \sqcup Z_\circ^+$  such that  $\mathbf{g}(D_\circ | \gamma_\bullet) = \mathbb{1}_{Z_\circ^+} - \mathbb{1}_{Z_\circ^-}$  (where  $\mathbb{1}_{X_\circ} := \sum_{\delta_\circ \in X_\circ} \mathbf{e}_{\delta_\circ}$  for  $X_\circ \subseteq D_\circ$ ).

Let  $\delta_\bullet$  be a  $D_\circ$ -accordion diagonal. Let  $A_\circ^-$  (resp.  $A_\circ^+$ ) denote the accordion crossed by  $\delta_\bullet = u_\bullet v_\bullet$  in  $D_\circ$  and not including  $(u+1)_\circ$  or  $(v+1)_\circ$  (resp.  $(u-1)_\circ$  or  $(v-1)_\circ$ ). Let  $\mathbf{c}^-(\delta_\bullet) := -\mathbb{1}_{A_\circ^-}$  and  $\mathbf{c}^+(\delta_\bullet) := \mathbb{1}_{A_\circ^+}$ . Recall from Lemma 21 that the negative (resp. positive)  $\mathbf{c}$ -vectors of  $\mathbf{C}(D_\circ)$  are given by  $\mathbf{c}^-(\delta_\bullet)$  (resp.  $\mathbf{c}^+(\delta_\bullet)$ ) for all  $D_\circ$ -accordion diagonal  $\delta_\bullet$  not in  $D_\bullet^-$  (resp.  $D_\bullet^+$ ). We let the reader check that:

- ◇ If  $\gamma_\bullet$  and  $\delta_\bullet$  do not cross and have no common endpoint, both  $|Z_\circ \cap A_\circ^-|$  and  $|Z_\circ \cap A_\circ^+|$  are even. Thus  $\langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c}^-(\delta_\bullet) \rangle = \langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c}^+(\delta_\bullet) \rangle = 0$ .
- ◇ If  $\gamma_\bullet$  and  $\delta_\bullet$  have a common endpoint, and  $\gamma_\bullet \delta_\bullet$  form a counterclockwise angle, then  $|Z_\circ \cap A_\circ^-|$  is even while  $Z_\circ \cap A_\circ^+$  is empty or starts and ends in  $Z_\circ^+$ . Thus  $\langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c}^-(\delta_\bullet) \rangle = 0$  while  $\langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c}^+(\delta_\bullet) \rangle \geq 0$ . The situation is similar if  $\gamma_\bullet \delta_\bullet$  form a clockwise angle.
- ◇ If  $\gamma_\bullet$  and  $\delta_\bullet$  cross,  $Z_\circ \cap A_\circ^-$  and  $Z_\circ \cap A_\circ^+$  are empty or start and end both in  $Z_\circ^-$  or both in  $Z_\circ^+$ . Thus, either  $\langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c}^-(\delta_\bullet) \rangle < 0$  and  $\langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c}^+(\delta_\bullet) \rangle \geq 0$  or conversely.

We conclude from this case analysis that

$$\omega(D_\circ | \gamma_\bullet) = |\{ \mathbf{c} \in \mathbf{C}(D_\circ) \mid \langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c} \rangle < 0 \}| = |\{ D_\circ\text{-accordion diagonals crossing } \gamma_\bullet \}|.$$

Finally, the inequality  $\langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{x} \rangle \leq \omega(D_\circ | \gamma_\bullet)$  defines a priori a face  $\mathbf{F}(\gamma_\bullet)$  of the zonotope  $\text{Zono}(D_\circ)$ . This face  $\mathbf{F}(\gamma_\bullet)$  is the Minkowski sum of the  $\mathbf{c}$ -vectors of  $\mathbf{C}(D_\circ)$  orthogonal to  $\mathbf{g}(D_\circ | \gamma_\bullet)$ . Proposition 22 ensures that any  $D_\circ$ -accordion dissection  $D_\bullet$  containing  $\gamma_\bullet$  already provides  $|D_\bullet| - 1$  linearly independent such  $\mathbf{c}$ -vectors  $\mathbf{c}(D_\circ | \delta_\bullet) \in D_\bullet$  for  $\delta_\bullet \in D_\bullet \setminus \{\gamma_\bullet\}$ . We obtain that  $\mathbf{F}(\gamma_\bullet)$  has dimension  $|D_\bullet| - 1 = |D_\circ| - 1$  and is therefore a facet of the zonotope  $\text{Zono}(D_\circ)$ .  $\square$

Define the half-space and the hyperplane corresponding to a solid  $D_\circ$ -accordion diagonal  $\gamma_\bullet$  by

$$\mathbf{H}^\leq(D_\circ | \gamma_\bullet) := \{\mathbf{x} \in \mathbb{R}^{D_\circ} \mid \langle \mathbf{g}(D_\circ | \gamma_\bullet) | \mathbf{x} \rangle \leq \omega(D_\circ | \gamma_\bullet)\},$$

and

$$\mathbf{H}^=(D_\circ | \gamma_\bullet) := \{\mathbf{x} \in \mathbb{R}^{D_\circ} \mid \langle \mathbf{g}(D_\circ | \gamma_\bullet) | \mathbf{x} \rangle = \omega(D_\circ | \gamma_\bullet)\}.$$

**2.3.  $\mathbf{g}$ -vector fan and  $D_\circ$ -accordiohedron.** In this section, we give a geometric realization of the  $D_\circ$ -accordion complex. We start by realizing this simplicial complex as a complete simplicial fan in  $\mathbb{R}^{D_\circ}$ . We denote by  $\mathbb{R}_{\geq 0}\mathbf{R}$  the nonnegative span of a set  $\mathbf{R}$  of vectors in  $\mathbb{R}^{D_\circ}$ .

**Theorem 27.** *The collection of cones*

$$\mathcal{F}^{\mathbf{g}}(D_\circ) := \{\mathbb{R}_{\geq 0}\mathbf{g}(D_\circ | D_\bullet) \mid D_\bullet \text{ any } D_\circ\text{-accordion dissection}\}$$

*forms a complete simplicial fan, that we call the  $\mathbf{g}$ -vector fan of  $D_\circ$ .*

The proof uses the following characterization of complete simplicial fans [DRS10, Coro. 4.5.20]. We will provide as well an alternative proof in Remark 60 based on sections of Cambrian fans.

**Proposition 28.** *Consider a pseudomanifold  $\Delta$  on a finite vertex set  $X$  and a set of vectors  $\mathbf{R} := (\mathbf{r}_x)_{x \in X}$  of  $\mathbb{R}^d$ . For  $D \in \Delta$ , define the cone  $\mathbf{R}_D := \{\mathbf{r}_x \mid x \in D\}$ . The collection of cones  $\{\mathbb{R}_{\geq 0}\mathbf{R}_D \mid D \in \Delta\}$  forms a complete simplicial fan if and only if*

- (1) *there exists a facet  $D$  of  $\Delta$  such that  $\mathbf{R}_D$  is a basis of  $\mathbb{R}^d$  and such that the open cones  $\mathbb{R}_{> 0}\mathbf{R}_D$  and  $\mathbb{R}_{> 0}\mathbf{R}_{D'}$  are disjoint for any facet  $D'$  of  $\Delta$  distinct from  $D$ ;*
- (2) *for two adjacent facets  $D, D'$  of  $\Delta$  with  $D \setminus \{x\} = D' \setminus \{x'\}$ , there is a linear dependence*

$$\alpha \mathbf{r}_x + \alpha' \mathbf{r}_{x'} + \sum_{y \in D \cap D'} \beta_y \mathbf{r}_y = 0$$

*on  $\mathbf{R}_{D \cup D'}$  where the coefficients  $\alpha$  and  $\alpha'$  have the same sign. (When these conditions hold, these coefficients do not vanish and the linear dependence is unique up to rescaling.)*

*Proof of Theorem 27.* By Corollary 24, the cone  $\mathbb{R}_{\geq 0}\mathbf{g}(D_\circ | D_\bullet^-)$  is the only cone of  $\mathcal{F}^{\mathbf{g}}(D_\circ)$  intersecting the interior of the positive orthant  $(\mathbb{R}_{\geq 0})^{D_\circ}$ . Consider now two adjacent maximal  $D_\circ$ -accordion dissections  $D_\bullet, D'_\bullet$ . Let  $\delta_\bullet \in D_\bullet$  and  $\delta'_\bullet \in D'_\bullet$  be such that  $D_\bullet \setminus \{\delta_\bullet\} = D'_\bullet \setminus \{\delta'_\bullet\}$ , and let  $\mu_\bullet$  and  $\nu_\bullet$  be the other diagonals as in Lemma 9 (see also Figure 4). Note that a diagonal of  $D_\circ$  crosses none of (resp. one of, resp. both) the diagonals  $\delta_\bullet, \delta'_\bullet$  if and only if it crosses none of (resp. one of, resp. both) the diagonals  $\mu_\bullet, \nu_\bullet$ . The same holds for a  $Z$  or a  $\Sigma$  of  $D_\circ$ . Therefore, we have the linear dependence  $\mathbf{g}(D_\circ | \delta_\bullet) + \mathbf{g}(D_\circ | \delta'_\bullet) = \mathbf{g}(D_\circ | \mu_\bullet) + \mathbf{g}(D_\circ | \nu_\bullet)$ . This shows that  $\mathcal{F}^{\mathbf{g}}(D_\circ)$  satisfies the two conditions of Proposition 28, and thus concludes the proof.  $\square$

**Remark 29.** The linear dependence  $\mathbf{g}(D_\circ | \delta_\bullet) + \mathbf{g}(D_\circ | \delta'_\bullet) = \mathbf{g}(D_\circ | \mu_\bullet) + \mathbf{g}(D_\circ | \nu_\bullet)$  relating the  $\mathbf{g}$ -vectors of two adjacent maximal  $D_\circ$ -accordion dissections  $D_\bullet, D'_\bullet$  with  $D_\bullet \setminus \{\delta_\bullet\} = D'_\bullet \setminus \{\delta'_\bullet\}$  shows that  $\det(\mathbf{g}(D_\circ | D_\bullet)) = -\det(\mathbf{g}(D_\circ | D'_\bullet))$ . Since the initial cone  $\mathbb{R}_{\geq 0}\mathbf{g}(D_\circ | D_\bullet^-)$  is generated by the coordinate vectors (see Example 17), we obtain that  $\det(\mathbf{g}(D_\circ | D_\bullet)) = \pm 1$  for all  $D_\circ$ -accordion dissection  $D_\bullet$ , so that the  $\mathbf{g}$ -vector fan  $\mathcal{F}^{\mathbf{g}}(D_\circ)$  is always *smooth*.

By Proposition 22, any non-maximal cone of  $\mathcal{F}^{\mathbf{g}}(D_\circ)$  is supported by a hyperplane orthogonal to a  $\mathbf{c}$ -vector of  $\mathbf{C}(D_\circ)$ . We thus obtain the following consequence.

**Corollary 30.** *The  $\mathbf{g}$ -vector fan  $\mathcal{F}^{\mathbf{g}}(D_\circ)$  coarsens the  $\mathbf{c}$ -vector fan  $\mathcal{F}^{\mathbf{c}}(D_\circ)$ .*

**Example 31.** Following Example 2, we observe that special reference dissections give rise to the following relevant fans:

- ◊ For an accordion triangulation  $A_\circ$  (*i.e.* with no interior triangle), the  $\mathbf{g}$ -vector fan  $\mathcal{F}^{\mathbf{g}}(A_\circ)$  coincides with a type  $A$  Cambrian fan of  $N$ . Reading and D. Speyer [RS09].
- ◊ For an arbitrary triangulation  $T_\circ$  (with or without interior triangle), the  $\mathbf{g}$ -vector fan  $\mathcal{F}^{\mathbf{g}}(T_\circ)$  was recently constructed in [HPS18].

**Example 32.** Figure 6 illustrates the  $\mathbf{g}$ -vector fans  $\mathcal{F}^{\mathbf{g}}(D_\circ)$  for various reference dissections  $D_\circ$ : the fan, the snake, and the cyclic triangulation of the hexagon, and a dissection of the heptagon. More precisely, we have represented the stereographic projection of the fans from the point  $[1, 1, 1]$ .




FIGURE 6. Stereographic projections of the  $\mathbf{g}$ -vector fans  $\mathcal{F}^{\mathbf{g}}(D_{\circ})$  for various reference hollow dissections  $D_{\circ}$ . See Figure 9 for alternative simplicial fan realizations of these accordion complexes.

Therefore, the external face of the projection corresponds to the  $D_{\circ}$ -accordion dissection  $D_{\bullet}^{-}$ . We have labeled all vertices of the projection (*i.e.* the rays of the fan) by the corresponding  $D_{\circ}$ -accordion diagonals.

We now provide a first polytopal realization of the  $\mathbf{g}$ -vector fan  $\mathcal{F}^{\mathbf{g}}(D_{\circ})$  (see also Section 4). This fan has a maximal cone for each maximal  $D_{\circ}$ -accordion dissection and a ray for each  $D_{\circ}$ -accordion diagonal. For a maximal  $D_{\circ}$ -accordion dissection  $D_{\bullet}$ , we define a point  $\mathbf{p}(D_{\circ} | D_{\bullet}) \in \mathbb{R}^{D_{\circ}}$  by

$$\mathbf{p}(D_{\circ} | D_{\bullet}) := \sum_{\delta_{\bullet} \in D_{\bullet}} \omega(D_{\circ} | \delta_{\bullet}) \cdot \mathbf{c}(D_{\circ} | \delta_{\bullet} \in D_{\bullet}),$$

where  $\omega(D_{\circ} | \delta_{\bullet})$  still denotes the  $D_{\circ}$ -height of  $\delta_{\bullet}$  defined as the number of  $D_{\circ}$ -accordion diagonals that cross  $\delta_{\bullet}$ . We will need the following two technical lemmas in the proof of Theorem 35.

**Lemma 33.** *For any maximal  $D_{\circ}$ -accordion dissection  $D_{\bullet}$ , the point  $\mathbf{p}(D_{\circ} | D_{\bullet})$  is the intersection of all hyperplanes  $\mathbf{H}^{\circ}(D_{\circ} | \delta_{\bullet})$  with  $\delta_{\bullet} \in D_{\bullet}$ .*



*Proof.* Observe first that the hyperplanes  $\mathbf{H}^=(D_\circ | \delta_\bullet)$  with  $\delta_\bullet \in D_\bullet$  have a unique intersection point, since  $\mathbf{g}(D_\circ | D_\bullet)$  is a basis. Moreover, since  $\mathbf{g}(D_\circ | D_\bullet)$  and  $\mathbf{c}(D_\circ | D_\bullet)$  form dual bases by Proposition 22, we have for any  $\gamma_\bullet \in D_\bullet$ :

$$\begin{aligned} \langle \mathbf{g}(D_\circ | \gamma_\bullet) | \mathbf{p}(D_\circ | D_\bullet) \rangle &= \sum_{\delta_\bullet \in D_\bullet} \omega(D_\circ | \delta_\bullet) \cdot \langle \mathbf{g}(D_\circ | \gamma_\bullet) | \mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet) \rangle \\ &= \sum_{\delta_\bullet \in D_\bullet} \omega(D_\circ | \delta_\bullet) \cdot \mathbb{1}_{\gamma_\bullet = \delta_\bullet} = \omega(D_\circ | \gamma_\bullet). \quad \square \end{aligned}$$

**Lemma 34.** *If  $D_\bullet, D'_\bullet$  are two adjacent maximal  $D_\circ$ -accordion dissections, and  $\delta_\bullet \in D_\bullet$  and  $\delta'_\bullet \in D'_\bullet$  are such that  $D_\bullet \setminus \{\delta_\bullet\} = D'_\bullet \setminus \{\delta'_\bullet\}$ , then*

$$\mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet) = -\mathbf{c}(D_\circ | \delta'_\bullet \in D'_\bullet) \quad \text{and} \quad \mathbf{p}(D_\circ | D'_\bullet) - \mathbf{p}(D_\circ | D_\bullet) \in \mathbb{Z}_{<0} \cdot \mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet).$$

*Proof.* Let  $D_\bullet, D'_\bullet$  be two adjacent maximal  $D_\circ$ -accordion dissections, let  $\delta_\bullet \in D_\bullet$  and  $\delta'_\bullet \in D'_\bullet$  be such that  $D_\bullet \setminus \{\delta_\bullet\} = D'_\bullet \setminus \{\delta'_\bullet\}$ , and let  $\mu_\bullet$  and  $\nu_\bullet$  be the other diagonals as in Lemma 9 (see also Figure 4). A quick case analysis then shows that

$$\mathbf{c}(D_\circ | \gamma_\bullet \in D'_\bullet) = \begin{cases} \mathbf{c}(D_\circ | \gamma_\bullet \in D_\bullet) & \text{for all diagonal } \gamma_\bullet \in D_\bullet \setminus \{\delta_\bullet, \mu_\bullet, \nu_\bullet\}, \\ -\mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet) & \text{if } \gamma_\bullet = \delta'_\bullet, \\ \mathbf{c}(D_\circ | \gamma_\bullet \in D_\bullet) + \mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet) & \text{if } \gamma_\bullet \in \{\mu_\bullet, \nu_\bullet\}. \end{cases}$$

Summing the contribution of all  $\mathbf{c}$ -vectors with their coefficients  $\omega(D_\circ | \gamma_\bullet)$ , we obtain

$$\mathbf{p}(D_\circ | D'_\bullet) - \mathbf{p}(D_\circ | D_\bullet) = (\omega(D_\circ | \mu_\bullet) + \omega(D_\circ | \nu_\bullet) - \omega(D_\circ | \delta_\bullet) - \omega(D_\circ | \delta'_\bullet)) \cdot \mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet).$$

Finally, note that any diagonal of  $P_\bullet$  that crosses one of (resp. both) the diagonals  $\mu_\bullet, \nu_\bullet$  also crosses one of (resp. both) the diagonals  $\delta_\bullet, \delta'_\bullet$ . Moreover,  $\delta_\bullet$  and  $\delta'_\bullet$  cross each other but do not cross  $\mu_\bullet$  and  $\nu_\bullet$ . It follows that  $\omega(D_\circ | \mu_\bullet) + \omega(D_\circ | \nu_\bullet) - \omega(D_\circ | \delta_\bullet) - \omega(D_\circ | \delta'_\bullet) \leq -2 < 0$ .  $\square$

**Theorem 35.** *The  $\mathbf{g}$ -vector fan is the normal fan of the  $D_\circ$ -accordiohedron  $\text{Acco}(D_\circ)$  defined equivalently as*

- $\diamond$  the convex hull of the points  $\mathbf{p}(D_\circ | D_\bullet)$  for all maximal  $D_\circ$ -accordion dissection  $D_\bullet$ , or
- $\diamond$  the intersection of the half-spaces  $\mathbf{H}^\leq(D_\circ | \gamma_\bullet)$  for all  $D_\circ$ -accordion diagonals  $\gamma_\bullet$ .

*Thus, the polar dual of  $\text{Acco}(D_\circ)$  is a polytopal realization of the  $D_\circ$ -accordion complex  $\mathcal{AC}(D_\circ)$ .*

The proof of Theorem 35 is based on the following characterization of polytopal realizations of a complete simplicial fan, whose proof can be found *e.g.* in [HLT11, Thm. 4.1].

**Theorem 36.** *Given a complete simplicial fan  $\mathcal{F}$  in  $\mathbb{R}^d$ , consider for each ray  $\mathbf{r}$  of  $\mathcal{F}$  a half-space  $\mathbf{H}_\mathbf{r}^\leq$  of  $\mathbb{R}^d$  containing the origin and defined by a hyperplane  $\mathbf{H}_\mathbf{r}^\perp$  orthogonal to  $\mathbf{r}$ . For each maximal cone  $C$  of  $\mathcal{F}$ , let  $\mathbf{a}(C) \in \mathbb{R}^d$  be the intersection of all hyperplanes  $\mathbf{H}_\mathbf{r}^\perp$  with  $\mathbf{r} \in C$ . Then the following assertions are equivalent:*

- (i) *The vector  $\mathbf{a}(C') - \mathbf{a}(C)$  points from  $C$  to  $C'$  for any two adjacent maximal cones  $C, C'$  of  $\mathcal{F}$ .*
- (ii) *The polytopes*

$$\text{conv} \{ \mathbf{a}(C) \mid C \text{ maximal cone of } \mathcal{F} \} \quad \text{and} \quad \bigcap_{\mathbf{r} \text{ ray of } \mathcal{F}} \mathbf{H}_\mathbf{r}^\leq$$

*coincide and their normal fan is  $\mathcal{F}$ .*

*Proof of Theorem 35.* The  $\mathbf{g}$ -vector fan  $\mathcal{F}^\mathbf{g}(D_\circ)$  has a ray  $\mathbf{g}(D_\circ | \delta_\bullet)$  for each  $D_\circ$ -accordion diagonal  $\delta_\bullet$  and a maximal cone  $C(D_\bullet) = \mathbb{R}_{\geq 0} \mathbf{g}(D_\circ | D_\bullet)$  for each maximal  $D_\circ$ -accordion dissection  $D_\bullet$ . Consider the half-spaces  $\mathbf{H}^\leq(D_\circ | \gamma_\bullet)$  for all  $D_\circ$ -accordion diagonals  $\gamma_\bullet$ . Lemma 33 ensures that the point  $\mathbf{a}(C(D_\bullet))$  coincides with  $\mathbf{p}(D_\circ | D_\bullet)$  for each maximal  $D_\circ$ -accordion dissection  $D_\bullet$ . Finally, Lemma 34 shows that the conditions of application of Theorem 36 are fulfilled.  $\square$

**Example 37.** Following Example 2, observe that special reference hollow dissections give rise to the following relevant polytopes, illustrated in Figure 7:

- $\diamond$  For a fan triangulation  $T_\circ$ , the  $T_\circ$ -accordiohedron  $\text{Acco}(T_\circ)$  is the classical associahedron constructed by S. Shnider and S. Sternberg [SS93] and J.-L. Loday [Lod04].




FIGURE 7. The zonotope  $Zono(D_o)$ ,  $D_o$ -accordiohedron  $Acco(D_o)$  and parallelepiped  $Para(D_o)$  for different reference dissections  $D_o$ . The first column is J.-L. Loday’s associahedron [Lod04], the second column is one of C. Hohlweg and C. Lange’s associahedra [HL07], the third column appeared in a discussion in C. Ceballos, F. Santos and G. Ziegler’s survey on associahedra [CSZ15, Fig. 3] and was explained in C. Hohlweg, V. Pilaud and S. Stella’s recent paper [HPS18], and the last column is a Stokes complex discussed by F. Chapoton in [Cha16] and illustrated in Figure 3.

- ◇ The  $A_o$ -accordiohedra  $Acco(A_o)$  for all accordion triangulations  $A_o$  are precisely the associahedra constructed by C. Hohlweg and C. Lange in [HL07].
- ◇ For a triangulation  $T_o$  with an interior triangle, the  $T_o$ -accordiohedron  $Acco(T_o)$  was recently constructed in [HPS18]. For example, for the triangulation of the hexagon with an interior triangle, this associahedron appeared as a mysterious realization in [CSZ15].
- ◇ For a quadrangulation  $Q_o$ , the  $Q_o$ -accordiohedron  $Acco(Q_o)$  is a realization of the Stokes polytope announced by Y. Baryshnikov [Bar01] and discussed by F. Chapoton in [Cha16].

We conclude this section by an immediate consequence of Theorem 35. To our knowledge, this property of accordion complexes was not observed before. However, using the connection between accordion complexes and support  $\tau$ -tilting complexes [GM16, PPP17, PPS17, BDM<sup>+</sup>17], it can also be obtained from [DIJ17, Thm. 1.7].

**Corollary 38.** *For any reference dissection  $D_o$ , the  $D_o$ -accordion complex  $\mathcal{AC}(D_o)$  is shellable.*

**2.4. Some properties of  $\text{Acco}(D_\circ)$ .** We conclude this section by pointing out some relevant combinatorial and geometric properties and observations on the  $D_\circ$ -accordiohedron.

**Proposition 39.** *The graph of the  $D_\circ$ -accordiohedron  $\text{Acco}(D_\circ)$  linearly oriented in the direction  $-\mathbb{1} := -\sum_{\delta_\circ \in D_\circ} \mathbf{e}_{\delta_\circ}$  is the Hasse diagram of the accordion lattice  $\mathcal{AL}(D_\circ)$ .*

*Proof.* Consider two adjacent maximal  $D_\circ$ -accordion dissections  $D_\bullet, D'_\bullet$  such that the flip from  $D_\bullet$  to  $D'_\bullet$  is increasing. Let  $\delta_\bullet \in D_\bullet$  and  $\delta'_\bullet \in D'_\bullet$  be such that  $D_\bullet \setminus \{\delta_\bullet\} = D'_\bullet \setminus \{\delta'_\bullet\}$ . As observed in Remark 20 (ii), the  $\mathbf{c}$ -vector  $\mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet)$  is the characteristic vector  $\mathbb{1}_{A_\circ}$  of the set  $A_\circ$  of diagonals of  $D_\circ$  crossed by both  $\delta_\bullet$  and  $\delta'_\bullet$ . Applying Lemma 34, we therefore obtain that

$$\langle -\mathbb{1} \mid \mathbf{p}(D_\circ | D'_\bullet) - \mathbf{p}(D_\circ | D_\bullet) \rangle = \langle -\mathbb{1} \mid \lambda \cdot \mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet) \rangle = \lambda \cdot \langle -\mathbb{1} \mid \mathbb{1}_{A_\circ} \rangle = -\lambda \cdot |A_\circ|,$$

for some  $\lambda \in \mathbb{Z}_{<0}$ . Thus, the linear functional  $-\mathbb{1}$  indeed orients the edge  $[\mathbf{p}(D_\circ | D_\bullet), \mathbf{p}(D_\circ | D'_\bullet)]$  from  $\mathbf{p}(D_\circ | D_\bullet)$  to  $\mathbf{p}(D_\circ | D'_\bullet)$ .  $\square$

**Remark 40.** Since the  $\mathbf{c}$ -vector fan  $\mathcal{F}^{\mathbf{c}}(D_\circ)$  refines the  $\mathbf{g}$ -vector fan  $\mathcal{F}^{\mathbf{g}}(D_\circ)$ , there is a natural projection  $\pi$  from the vertices of the  $D_\circ$ -zonotope  $\text{Zono}(D_\circ)$  to that of the  $D_\circ$ -accordiohedron  $\text{Acco}(D_\circ)$ . In analogy to the acyclic case, one could hope to obtain the accordion lattice as a lattice quotient through this projection. However, the transitive closure of the graph of the  $D_\circ$ -zonotope  $\text{Zono}(D_\circ)$  oriented in the direction  $-\mathbb{1}$  is not a lattice in general (the first counter-example is the dissection with a central square surrounded by 4 triangles). As shown in [GM16], the right objects are not the separable subsets of  $\mathbf{c}$ -vectors (*i.e.* the vertices of  $\text{Zono}(D_\circ)$ ) but the biclosed subsets of  $\mathbf{c}$ -vectors.

**Proposition 41.** *The accordiohedron  $\text{Acco}(D_\circ)$  has precisely  $|D_\circ|$  pairs of parallel facets.*

*Proof.* Two facets of  $\text{Acco}(D_\circ)$  are parallel if and only if the corresponding  $\mathbf{g}$ -vectors are opposite. We therefore want to prove that the pairs of opposite coordinate vectors are the only pairs of opposite  $\mathbf{g}$ -vectors. Assume by contradiction that there exist two hollow diagonals  $\delta_\circ, \delta'_\circ \in D_\circ$  and two solid  $D_\circ$ -diagonals  $\delta_\bullet, \delta'_\bullet$  such that  $\mathbf{g}(D_\circ | \delta_\bullet)$  and  $\mathbf{g}(D_\circ | \delta'_\bullet)$  have non-zero opposite coordinate both on  $\delta_\circ$  and  $\delta'_\circ$ . Then both  $\delta_\bullet$  and  $\delta'_\bullet$  cross both  $\delta_\circ$  and  $\delta'_\circ$ . But this implies that they both slalom on  $\delta_\circ$  (and on  $\delta'_\circ$ ) in the same way. Contradiction.  $\square$

Recall from Example 17 that the  $\mathbf{g}$ -vectors of the diagonals of  $D_\bullet^-$  (resp.  $D_\bullet^+$ ) are the coordinate vectors (resp. negative of the coordinate vectors). Consider the  *$D_\circ$ -parallelepiped*

$$\text{Para}(D_\circ) := \{ \mathbf{x} \in \mathbb{R}^{D_\circ} \mid \langle \mathbf{g}(D_\circ | \delta_\bullet) \mid \mathbf{x} \rangle \leq \omega(D_\circ | \delta_\bullet) \text{ for all } \delta_\bullet \in D_\bullet^- \cup D_\bullet^+ \}$$

defined by the inequalities of the  $D_\circ$ -zonotope  $\text{Zono}(D_\circ)$  corresponding to the positive and negative basis vectors. Our next statement follows from Proposition 41 and is illustrated in Figure 7.

**Corollary 42.** *For any  $D_\circ$ , we have matriochka polytopes:  $\text{Zono}(D_\circ) \subseteq \text{Acco}(D_\circ) \subseteq \text{Para}(D_\circ)$ .*

In fact, each polytope in this chain is obtained by deleting facets from the previous one.

Consider now an isometry  $\sigma$  of the plane that preserves the hollow polygon  $P_\circ$  and the solid polygon  $P_\bullet$ . For any diagonals and dissections  $\delta_\bullet \in D_\bullet$  and  $\delta_\circ \in D_\circ$ , we have

- ◇  $\delta_\bullet$  is a  $D_\circ$ -accordion diagonal  $\iff \sigma(\delta_\bullet)$  is a  $\sigma(D_\circ)$ -accordion diagonal,
- ◇  $D_\bullet$  is a  $D_\circ$ -accordion dissection  $\iff \sigma(D_\bullet)$  is a  $\sigma(D_\circ)$ -accordion dissection,
- ◇ if  $\Sigma : \mathbb{R}^{D_\circ} \rightarrow \mathbb{R}^{\sigma(D_\circ)}$  denotes the isometry defined by  $(\Sigma(\mathbf{x}))_{\sigma(\delta_\circ)} := \varepsilon(\sigma) \cdot \mathbf{x}_{\delta_\circ}$ , (where  $\varepsilon(\sigma) = 1$  if  $\sigma$  is direct and  $-1$  if  $\sigma$  is indirect), then we have

$$\begin{aligned} \mathbf{g}(\sigma(D_\circ) | \sigma(\delta_\bullet)) &= \Sigma(\mathbf{g}(D_\circ | \delta_\bullet)), & \mathbf{c}(\sigma(D_\circ) | \sigma(\delta_\bullet) \in \sigma(D_\bullet)) &= \Sigma(\mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet)), \\ \omega(\sigma(D_\circ) | \sigma(\delta_\bullet)) &= \omega(D_\circ | \delta_\bullet), & \text{and} & \quad \mathbf{p}(\sigma(D_\circ) | \sigma(D_\bullet)) &= \Sigma(\mathbf{p}(D_\circ | D_\bullet)). \end{aligned}$$

This immediately implies the following statement.

**Proposition 43.** *Any  $P_\circ$ -preserving isometry  $\sigma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  induces an isometry  $\Sigma : \mathbb{R}^{D_\circ} \rightarrow \mathbb{R}^{\sigma(D_\circ)}$  with  $\Sigma(\text{Zono}(D_\circ)) = \text{Zono}(\sigma(D_\circ))$ ,  $\Sigma(\text{Acco}(D_\circ)) = \text{Acco}(\sigma(D_\circ))$  and  $\Sigma(\text{Para}(D_\circ)) = \text{Para}(\sigma(D_\circ))$ .*

We say that a dissection  $D$  is  $\sigma$ -invariant when  $\sigma(D) = D$ . Assume now that  $\sigma$  is a rotation and  $D_\circ$  is  $\sigma$ -invariant. We call  $\sigma$ -invariant  $D_\circ$ -accordion complex the simplicial complex  $\mathcal{AC}^\sigma(D_\circ)$  whose vertices are the crossing-free  $\sigma$ -orbits of  $D_\circ$ -accordion diagonals, and whose faces are sets of such orbits whose union is crossing-free. In other words, the faces of  $\mathcal{AC}^\sigma(D_\circ)$  are  $\sigma$ -invariant  $D_\circ$ -accordion dissections, seen as sets of  $\sigma$ -orbits of diagonals.

**Lemma 44.** *The  $\sigma$ -invariant  $D_\circ$ -accordion complex  $\mathcal{AC}^\sigma(D_\circ)$  is a pseudomanifold.*

*Proof.* Assume first that  $\sigma$  is the central symmetry. In this case, there are two possible types of orbits: the long  $D_\circ$ -accordion diagonals and the centrally symmetric pairs of  $D_\circ$ -accordion diagonals. One can check that any facet of  $\mathcal{AC}^\sigma(D_\circ)$  has a long diagonal if and only if  $D_\circ$  has, and has as many centrally symmetric pairs of diagonals as  $D_\circ$ . Finally, any orbit in any facet of  $\mathcal{AC}^\sigma(D_\circ)$  can be flipped: long diagonals can already be flipped in  $\mathcal{AC}(D_\circ)$ , and a centrally symmetric pair of diagonals can be flipped by flipping one after the other its two diagonals in  $\mathcal{AC}(D_\circ)$ .

Finally, the general statement follows from this special case. Indeed, if  $\sigma$  is not a central symmetry, let  $C_\circ$  denote the cell of  $D_\circ$  containing the center of  $P_\circ$ , let  $u_\circ$  be a vertex of  $C_\circ$ , let  $\underline{D}_\circ$  be the set of diagonals of  $D_\circ$  whose endpoints are between  $u_\circ$  and  $\sigma(u_\circ)$ , and let  $\rho$  be the central symmetry around the middle of  $u_\circ\sigma(u_\circ)$ . Then  $\mathcal{AC}^\sigma(D_\circ)$  is isomorphic to  $\mathcal{AC}^\rho(\underline{D}_\circ \cup \rho(\underline{D}_\circ))$ .  $\square$

Let  $\Sigma : \mathbb{R}^{D_\circ} \rightarrow \mathbb{R}^{D_\circ}$  denote the isometry defined by  $(\Sigma(\mathbf{x}))_{\sigma(\delta_\circ)} := \mathbf{x}_{\delta_\circ}$  and  $\text{Fix}(\Sigma)$  denote the linear subspace of fixed points of  $\Sigma$ . According to the previous discussion, a maximal  $D_\circ$ -accordion dissection  $D_\bullet$  is  $\sigma$ -invariant if and only if  $\mathbf{p}(D_\circ | D_\bullet) \in \text{Fix}(\Sigma)$ . We obtain the following statement.

**Proposition 45.** *For a  $\sigma$ -invariant dissection  $D_\circ$ , the polytope  $\text{Acco}^\sigma(D_\circ)$  defined equivalently as*

- $\diamond$  *the convex hull of  $\mathbf{p}(D_\circ | D_\bullet)$  for all  $\sigma$ -invariant maximal  $D_\circ$ -accordion dissections  $D_\bullet$ ,*
- $\diamond$  *the intersection of the  $D_\circ$ -accordiohedron  $\text{Acco}(D_\circ)$  with the fixed space  $\text{Fix}(\Sigma)$ ,*

*is a polytopal realization of the  $\sigma$ -invariant accordion complex  $\mathcal{AC}^\sigma(D_\circ)$ .*

*Proof.* Denote by  $P = \text{conv}\{\mathbf{p}(D_\circ | D_\bullet) \mid \sigma\text{-invariant maximal } D_\circ\text{-accordion dissections } D_\bullet\}$  and by  $Q = \text{Acco}(D_\circ) \cap \text{Fix}(\Sigma)$ . The inclusion  $P \subseteq Q$  is clear since  $D_\bullet$  is  $\sigma$ -invariant if and only if  $\mathbf{p}(D_\circ | D_\bullet) \in \text{Fix}(\Sigma)$ . We now prove the reverse inclusion. For that, consider an arbitrary  $\sigma$ -invariant maximal  $D_\circ$ -accordion dissection  $D_\bullet$ . Its corresponding point  $\mathbf{p}(D_\circ | D_\bullet)$  is a common vertex of  $P$  and  $Q$ . Moreover, any edge  $e$  of  $Q$  incident to  $\mathbf{p}(D_\circ | D_\bullet)$  is the intersection of  $\text{Fix}(\Sigma)$  with a face  $F$  of  $\text{Acco}(D_\circ)$  that corresponds to a  $\sigma$ -invariant  $D_\circ$ -dissection. Since  $\mathcal{AC}^\sigma(D_\circ)$  is a pseudomanifold, this dissection can be refined into another maximal  $\sigma$ -invariant  $D_\circ$ -accordion dissection  $D'_\bullet$ . The point  $\mathbf{p}(D_\circ | D'_\bullet)$  belongs to  $F$  and to  $\text{Fix}(\Sigma)$  and thus to  $e$ . We conclude that if  $v$  is a common vertex of  $P$  and  $Q$ , then so are all neighbors of  $v$  in the graph of  $Q$ . Propagating this property, we obtain that all vertices of  $Q$  are also vertices of  $P$ , so that  $P = Q$ . Finally, there is a clear injection from the  $\sigma$ -invariant accordion complex  $\mathcal{AC}^\sigma(D_\circ)$  to the boundary complex of  $P = Q$ , thus a bijection (since these complexes are two spheres with the same vertex set).  $\square$

### 3. THE $\mathbf{d}$ -VECTOR FAN

In this section, we discuss the generalization to the  $D_\circ$ -accordion complex of another classical geometric realization of the associahedron coming from the theory of cluster algebras [FZ02, FZ03a, CFZ02, CSZ15]. Namely, we define compatibility vectors in analogy with the denominator vectors of cluster variables, and we characterize the reference dissections  $D_\circ$  for which these vectors support a complete simplicial fan realizing the  $D_\circ$ -accordion complex.

**3.1.  $\mathbf{d}$ -vectors.** Fix a dissection  $D_\circ$  of the hollow  $n$ -gon. For a hollow diagonal  $\delta_\circ = i_\circ j_\circ$  and a solid diagonal  $\delta_\bullet$ , we denote by

$$(\delta_\circ | \delta_\bullet) := \begin{cases} -1 & \text{if } \delta_\bullet = (i-1)_\bullet(j-1)_\bullet, \\ 0 & \text{if } \delta_\bullet \text{ and } (i-1)_\bullet(j-1)_\bullet \text{ do not cross,} \\ 1 & \text{if } \delta_\bullet \text{ and } (i-1)_\bullet(j-1)_\bullet \text{ cross.} \end{cases}$$

For any  $D_\circ$ -accordion diagonal  $\delta_\bullet$ , the **d-vector** of  $\delta_\bullet$  with respect to  $D_\circ$  is the vector

$$\mathbf{d}(D_\circ | \delta_\bullet) = \sum_{\delta_\circ \in D_\circ} (\delta_\circ | \delta_\bullet) \mathbf{e}_{\delta_\circ}.$$

In other words, our **d-vector**  $\mathbf{d}(D_\circ | \delta_\bullet)$  records the compatibility of the diagonal  $\delta_\bullet$  with the dissection  $D_\circ^-$ . For a  $D_\circ$ -accordion dissection  $D_\bullet$ , we define  $\mathbf{d}(D_\circ | D_\bullet) := \{\mathbf{d}(D_\circ | \delta_\bullet) \mid \delta_\bullet \in D_\bullet\}$ .

**Example 46.** Consider the hollow dissection  $D_\circ^{\text{ex}} = \{3_\circ 7_\circ, 3_\circ 13_\circ, 9_\circ 13_\circ\}$  and the rightmost solid dissection  $D_\bullet^{\text{ex}} = \{2_\bullet 6_\bullet, 2_\bullet 10_\bullet, 10_\bullet 14_\bullet\}$  of Figure 2. Its **d-vectors** are given by

$$\mathbf{d}(D_\circ^{\text{ex}} | 2_\bullet 6_\bullet) = -\mathbf{e}_{3_\circ 7_\circ}, \quad \mathbf{d}(D_\circ^{\text{ex}} | 2_\bullet 10_\bullet) = \mathbf{e}_{9_\circ 13_\circ}, \quad \text{and} \quad \mathbf{d}(D_\circ^{\text{ex}} | 10_\bullet 14_\bullet) = \mathbf{e}_{3_\circ 13_\circ} + \mathbf{e}_{9_\circ 13_\circ}.$$

**3.2. d-vector fan.** We now consider the set of cones

$$\{\mathbb{R}_{\geq 0} \mathbf{d}(D_\circ | D_\bullet) \mid D_\bullet \text{ any } D_\circ\text{-accordion dissection}\}$$

generated by the **d-vectors** of the  $D_\circ$ -accordion dissections. We want to characterize the reference hollow dissections  $D_\circ$  for which these cones form a complete simplicial fan realizing the  $D_\circ$ -accordion complex. We start with a negative result. An *even interior cell* of a dissection  $D$  is a cell with an even number of edges which are all internal diagonals of  $D$ .

**Proposition 47.** *If the reference hollow dissection  $D_\circ$  contains an even interior cell, then the **d-vectors** cannot realize the  $D_\circ$ -accordion complex.*

*Proof.* Assume that  $D_\circ$  contains an even interior cell  $C_\circ$ . Denote its vertices by  $i_\circ^1, \dots, i_\circ^{2p}$  (in clockwise order) and its edges  $\delta_\circ^k := i_\circ^k i_\circ^{k+1}$  for  $k \in [2p]$  (where  $i_\circ^{2p+1} = i_\circ^1$  by convention). Denote by  $D_\circ^k$  the set of diagonals of  $D_\circ$  separated from  $C_\circ$  by  $\delta_\circ^k$  (including  $\delta_\circ^k$  itself), and let  $D_\bullet^k := \{(i-1)_\bullet (j-1)_\bullet \mid i_\circ j_\circ \in D_\circ^k\}$ . Consider the solid diagonals  $\delta_\bullet^k := (i^k+1)_\bullet (i^{k+1}+1)_\bullet$  for  $k \in [2p]$ . Observe that  $\delta_\bullet^k$  only crosses diagonals of  $D_\bullet^{k-1}$  and  $D_\bullet^k$ , and that  $\delta_\bullet^k$  and  $\delta_\bullet^{k+1}$  cross precisely the same diagonals of  $D_\bullet^k$ . Since the cell is even, it ensures that the **d-vectors** of the diagonals  $\delta_\bullet^k$  for  $k \in [2p]$  satisfy the linear dependence

$$\sum_{\substack{k \in [2p] \\ k \text{ even}}} \mathbf{d}(D_\circ | \delta_\bullet^k) = \sum_{\substack{k \in [2p] \\ k \text{ odd}}} \mathbf{d}(D_\circ | \delta_\bullet^k).$$

However, as already mentioned in Section 1.4, the diagonals  $\delta_\bullet^k$  for  $k \in [2p]$  all belong to the  $D_\circ$ -accordion dissection  $D_\bullet^+ := \{(i+1)_\bullet (j+1)_\bullet \mid i_\circ j_\circ \in D_\circ\}$ . Therefore, the cone  $\mathbb{R}_{\geq 0} \mathbf{d}(D_\circ | D_\bullet^+)$  is degenerate, so that the **d-vectors** cannot realize the  $D_\circ$ -accordion complex.  $\square$

**Example 48.** Consider a hollow octagon and the reference dissection  $D_\circ := \{1_\circ 5_\circ, 5_\circ 9_\circ, 9_\circ 13_\circ, 13_\circ 1_\circ\}$  with an interior square cell  $1_\circ 5_\circ 9_\circ 13_\circ$ . Then we have

$$\begin{aligned} \mathbf{d}(D_\circ | 2_\bullet 6_\bullet) &= \mathbf{e}_{1_\circ 5_\circ} + \mathbf{e}_{5_\circ 9_\circ} & \mathbf{d}(D_\circ | 6_\bullet 10_\bullet) &= \mathbf{e}_{5_\circ 9_\circ} + \mathbf{e}_{9_\circ 13_\circ} \\ \mathbf{d}(D_\circ | 10_\bullet 14_\bullet) &= \mathbf{e}_{9_\circ 13_\circ} + \mathbf{e}_{13_\circ 1_\circ} & \mathbf{d}(D_\circ | 14_\bullet 2_\bullet) &= \mathbf{e}_{13_\circ 1_\circ} + \mathbf{e}_{1_\circ 5_\circ} \end{aligned}$$

so that there is already a linear dependence

$$\mathbf{d}(D_\circ | 2_\bullet 6_\bullet) + \mathbf{d}(D_\circ | 10_\bullet 14_\bullet) = \mathbf{d}(D_\circ | 6_\bullet 10_\bullet) + \mathbf{d}(D_\circ | 14_\bullet 2_\bullet)$$

among the **d-vectors** of the  $D_\circ$ -accordion dissection  $D_\bullet^+ = \{2_\bullet 6_\bullet, 6_\bullet 10_\bullet, 10_\bullet 14_\bullet, 14_\bullet 2_\bullet\}$ .

On the negative side, we have seen that the presence of even interior cells prohibits the **d-vectors** from forming a complete simplicial fan. The positive side is that the even interior cells are the only obstructions.

**Theorem 49.** *The collection of cones*

$$\mathcal{F}^{\mathbf{d}}(D_\circ) := \{\mathbb{R}_{\geq 0} \mathbf{d}(D_\circ | D_\bullet) \mid D_\bullet \text{ any } D_\circ\text{-accordion dissection}\}$$

*forms a complete simplicial fan, that we call the **d-vector fan** of  $D_\circ$ , if and only if  $D_\circ$  contains no even interior cell.*




FIGURE 8. Illustration of the notations and of the different cases in the proof of Theorem 49.

*Proof.* We use the characterization of complete simplicial fans presented in Proposition 28.

Observe first that  $\mathbf{d}(D_\circ | D_\circ^-) = (\mathbb{R}_{\leq 0})^{D_\circ}$  is the only cone of  $\mathcal{F}^{\mathbf{d}}(D_\circ)$  intersecting the interior of the negative orthant  $(\mathbb{R}_{\leq 0})^{D_\circ}$ . Therefore,  $\mathcal{F}^{\mathbf{d}}(D_\circ)$  fulfills Condition (1) in Proposition 28.

To check Condition (2), consider two adjacent maximal  $D_\circ$ -accordion dissections  $D_\bullet$  and  $D'_\bullet$  and let  $\delta_\bullet \in D_\bullet$  and  $\delta'_\bullet \in D'_\bullet$  be such that  $D_\bullet \setminus \{\delta_\bullet\} = D'_\bullet \setminus \{\delta'_\bullet\}$ . Let  $\mu_\bullet$  and  $\nu_\bullet$  be the diagonals of  $\overline{D_\bullet} \cap \overline{D'_\bullet}$  as in Lemma 9 (see also Figure 4). In other words,  $\mu_\bullet$  and  $\nu_\bullet$  are incident to both  $\delta_\bullet$  and  $\delta'_\bullet$ , and they are crossed by the hollow diagonal which intersect  $\delta_\bullet$  and  $\delta'_\bullet$ . Let  $\gamma_\circ = i_\circ j_\circ$  be such a hollow diagonal crossing  $\delta_\bullet, \delta'_\bullet, \mu_\bullet$  and  $\nu_\bullet$ , and let  $\gamma_\bullet = (i-1)_\bullet(j-1)_\bullet$ . We now distinguish three cases:

- ◊ Assume that  $\gamma_\bullet$  still crosses  $\mu_\bullet$  and  $\nu_\bullet$ . In this case, any diagonal of  $D_\bullet^-$  crossing both (resp. either)  $\delta_\bullet$  and (resp. or)  $\delta'_\bullet$  also crosses both (resp. either)  $\mu_\bullet$  and (resp. or)  $\nu_\bullet$ . See Figure 8 (left). Therefore, the  $\mathbf{d}$ -vectors of  $D_\bullet \cup D'_\bullet$  satisfy the linear dependence

$$\mathbf{d}(D_\circ | \delta_\bullet) + \mathbf{d}(D_\circ | \delta'_\bullet) = \mathbf{d}(D_\circ | \mu_\bullet) + \mathbf{d}(D_\circ | \nu_\bullet).$$

- ◊ Assume that  $\gamma_\bullet$  crosses neither  $\mu_\bullet$  nor  $\nu_\bullet$ . Then  $\gamma_\bullet$  is incident to both  $\mu_\bullet$  and  $\nu_\bullet$ , and therefore is either  $\delta_\bullet$  or  $\delta'_\bullet$ , say  $\gamma_\bullet = \delta_\bullet$ . Then  $\mathbf{d}(\gamma_\circ | \delta_\bullet) = -1$  while  $\mathbf{d}(\gamma_\circ | \delta'_\bullet) = 1$  (since  $\delta'_\bullet$  crosses  $\delta_\bullet = \gamma_\bullet$ ), so that  $\mathbf{d}(\gamma_\circ | \delta_\bullet) + \mathbf{d}(\gamma_\circ | \delta'_\bullet) = 0$ . Moreover, we have  $\mathbf{d}(\gamma_\circ | \delta'_\bullet) = 0$  for any diagonal  $\varepsilon_\bullet \in D_\bullet \cap D'_\bullet$  since  $\delta_\bullet = \gamma_\bullet$  cannot cross  $\varepsilon_\bullet$  as they both belongs to  $D_\bullet$ . Therefore, the set  $\{\mathbf{d}(D_\circ | \delta_\bullet) + \mathbf{d}(D_\circ | \delta'_\bullet)\} \cup \mathbf{d}(D_\circ | D_\bullet \cap D'_\bullet)$  contains  $|D_\circ|$  vectors of  $\mathbb{R}^{D_\circ}$  whose  $\gamma_\circ$ -coordinate all vanish, so that it admits a linear dependence.

- ◊ Otherwise, we can assume that  $\gamma_\bullet$  crosses  $\mu_\bullet$  but not  $\nu_\bullet$ . Then  $\gamma_\bullet$  has a common endpoint with  $\nu_\bullet$  and  $\delta_\bullet$  (or  $\delta'_\bullet$ , but we then permute notations). Changing our initial choice of  $\gamma_\circ$ , we can assume that no diagonal of  $D_\bullet^-$  separates  $\gamma_\bullet$  from  $\delta_\bullet$ . We now denote clockwise
  - by  $\nu_\bullet =: \lambda_\bullet^0, \lambda_\bullet^1, \dots, \lambda_\bullet^\ell := \delta_\bullet$  the edges of the cell  $C_\bullet$  of  $D_\bullet$  containing  $\nu_\bullet$  and  $\delta_\bullet$ ,
  - by  $\gamma_\bullet =: \gamma_\bullet^0, \gamma_\bullet^1, \dots, \gamma_\bullet^k$  the edges of the cell  $C_\bullet^-$  of  $D_\bullet^-$  containing  $\gamma_\bullet$  and crossed by  $\delta_\bullet$ .

These notations are illustrated on Figure 8. We still distinguish two subcases as in Figure 8:

- If  $\gamma_\bullet^i$  crosses  $\lambda_\bullet^i$  for all  $i$  as in Figure 8 (middle), then  $\ell = k$  and we have the linear dependence

$$2\mathbf{d}(D_\circ | \delta_\bullet) + \mathbf{d}(D_\circ | \delta'_\bullet) = \mathbf{d}(D_\circ | \mu_\bullet) + \sum_{i \in [\ell-1]} (-1)^{(i-1)} \mathbf{d}(D_\circ | \lambda_\bullet^i).$$

It is essential here that  $\ell = k$  is even. This is guarantied by the assumption that  $D_\circ$  (and thus  $D_\bullet^-$ ) has no even interior cell, since  $C_\bullet^-$  is an interior cell of  $D_\bullet^-$  of size  $k$ .

- Otherwise, we are in a situation similar to Figure 8 (right). Considering the maximal index  $m$  such that  $\gamma_\bullet^i$  crosses  $\lambda_\bullet^i$  for all  $i \leq m$ , and we have the linear dependence

$$\mathbf{d}(D_\circ | \delta_\bullet) + \mathbf{d}(D_\circ | \delta'_\bullet) = \mathbf{d}(D_\circ | \mu_\bullet) + \sum_{i \in [m]} (-1)^{(i-1)} \mathbf{d}(D_\circ | \lambda_\bullet^i). \quad \square$$






FIGURE 9. Stereographic projections of the  $\mathbf{d}$ -vector fans  $\mathcal{F}^{\mathbf{d}}(D_{\circ})$  for various reference hollow dissections  $D_{\circ}$ . See Figure 6 for alternative simplicial fan realizations of these accordion complexes.

**Example 50.** Following Example 2, we observe that special reference dissections give rise to the following relevant fans:

- ◇ For a snake triangulation  $\Sigma_{\circ}$ , the  $\mathbf{d}$ -vector fan  $\mathcal{F}^{\mathbf{d}}(\Sigma_{\circ})$  coincides with the type  $A$  cluster fan of S. Fomin and A. Zelevinsky [FZ03a].
- ◇ For any triangulation  $T_{\circ}$ , the  $\mathbf{d}$ -vector fan  $\mathcal{F}^{\mathbf{d}}(T_{\circ})$  was already constructed in [CSZ15].
- ◇ For a quadrangulation  $Q_{\circ}$  with no interior quadrangle (equivalently, with no cross), we obtain an alternative realization of the Stokes complexes studied in [Bar01, Cha16]. This was observed by A.-H. Bateni, T. Manneville and V. Pilaud in [BMP16].

Figure 9 illustrates the  $\mathbf{d}$ -vector fans  $\mathcal{F}^{\mathbf{d}}(D_{\circ})$  for the same reference dissections  $D_{\circ}$  as in Figure 6. More precisely, we have represented the stereographic projection of the fans from the point  $[-1, -1, -1]$ . Therefore, the external face of the projection corresponds to the  $D_{\circ}$ -accordion dissection  $D_{\circ}^{-}$ . We have labeled all vertices of the projection (*i.e.* the rays of the fan) by the corresponding  $D_{\circ}$ -accordion diagonals. Compare with Figure 6.

**Remark 51.** To prove that the  $\mathbf{d}$ -vector fan  $\mathcal{F}^{\mathbf{d}}(D_{\circ})$  is polytopal, we would need to find suitable hyperplanes orthogonal to their rays in order to apply Theorem 36. For the  $\mathbf{g}$ -vector fan, these

hyperplanes were defined using the height function  $\omega(D_\circ | \delta_\bullet)$ . It would be natural to use the same height function for the  $\mathbf{d}$ -vector fan as well. Unfortunately, for this choice of height function, we can only prove Condition (i) of Theorem 36 when  $D_\circ$  is a triangulation (see also [CSZ15]). We were not able to find suitable right hand sides for any dissection  $D_\circ$ .

**Remark 52.** Our  $\mathbf{d}$ -vectors record the compatibility with the dissection  $D_\bullet^-$ . A priori, we could compute compatibility vectors with respect to any other maximal  $D_\circ$ -accordion dissection  $D_\bullet^{\text{ini}}$ . Experiments suggest that the  $\mathbf{d}$ -vector construction provides a complete simplicial fan as long as neither  $D_\circ$  nor  $D_\bullet^{\text{ini}}$  contain no even interior cell. We checked it for reference quadrangulations with at most 5 diagonals. The linear dependences involved seem however much more complicated than those of the proof of Theorem 49 (in particular, they may involve  $\mathbf{d}$ -vectors of diagonals not included in the cells containing  $\delta_\bullet$  and  $\delta'_\bullet$ ).

#### 4. SECTIONS AND PROJECTIONS

Recall that for a fan  $\mathcal{F}$  of  $\mathbb{R}^d$  and a linear subspace  $V$  of  $\mathbb{R}^d$ , the *section* of  $\mathcal{F}$  by  $V$  is the fan  $\mathcal{F}|_V := \{C \cap V \mid C \in \mathcal{F}\}$ . For a polytope  $P \subseteq \mathbb{R}^d$  and a projection  $\pi : \mathbb{R}^d \rightarrow V$ , the normal fan of the projected polytope  $\pi(P)$  is the section of the normal fan of  $P$  by  $V$  [Zie95, Lem. 7.11]. We now consider sections of the  $\mathbf{g}$ - and  $\mathbf{d}$ -vector fans by coordinate subspaces. For two dissections  $D_\circ \subset D'_\circ$ , we naturally identify  $\mathbb{R}^{D_\circ}$  with the subspace spanned by  $\{\mathbf{e}_{\delta_\circ} \mid \delta_\circ \in D_\circ\}$  in  $\mathbb{R}^{D'_\circ}$ .

**4.1. Coordinate sections of the  $\mathbf{d}$ -vector fan.** We start by presenting sections of the  $\mathbf{d}$ -vector fan which are not very surprising. The following lemma is immediate from the definition of  $\mathbf{d}$ -vectors.

**Lemma 53.** *Consider two dissections  $D_\circ \subset D'_\circ$ , and a  $D'_\circ$ -accordion diagonal  $\delta_\bullet$ . Then we have  $\mathbf{d}(D_\circ | \delta_\bullet) \in \mathbb{R}^{D_\circ}$  if and only if  $\delta_\bullet$  does not cross any diagonal of  $\{(i-1)_\bullet(j-1)_\bullet \mid i_\circ j_\circ \in D'_\circ \setminus D_\circ\}$ .*

**Corollary 54.** *For two dissections  $D_\circ \subset D'_\circ$ , the face complex of the section of the  $\mathbf{d}$ -vector fan  $\mathcal{F}^{\mathbf{d}}(D'_\circ)$  by  $\mathbb{R}^{D_\circ}$  is isomorphic to the link of the dissection  $\{(i-1)_\bullet(j-1)_\bullet \mid i_\circ j_\circ \in D'_\circ \setminus D_\circ\}$  in the  $D'_\circ$ -accordion complex  $\mathcal{AC}(D'_\circ)$ .*

**4.2. Coordinate sections of the  $\mathbf{g}$ -vector fan.** More relevant are the sections of the  $\mathbf{g}$ -vector fan. They provide an alternative approach to polytopal realizations of the accordion complex based on projected associahedra. This approach relies on the following crucial observation.

**Lemma 55.** *Consider two dissections  $D_\circ \subset D'_\circ$ , and a  $D'_\circ$ -accordion diagonal  $\delta_\bullet$ . Then we have  $\mathbf{g}(D'_\circ | \delta_\bullet) \in \mathbb{R}^{D_\circ}$  if and only if  $\delta_\bullet$  is a  $D_\circ$ -accordion diagonal. Moreover, in this case, the  $\mathbf{g}$ -vectors  $\mathbf{g}(D_\circ | \delta_\bullet)$  and  $\mathbf{g}(D'_\circ | \delta_\bullet)$  coincide.*

*Proof.* Let  $\delta_\circ \in D'_\circ \setminus D_\circ$ . By definition, a  $D'_\circ$ -accordion diagonal  $\delta_\bullet$  does not slalom on  $\delta_\circ$  if and only if the  $\delta_\circ$ -coordinate of  $\mathbf{g}(D_\circ | \delta_\bullet)$  vanishes. Thus,  $\delta_\bullet$  is a  $D_\circ$ -accordion diagonal if and only if the  $\delta_\circ$ -coordinate of  $\mathbf{g}(D'_\circ | \delta_\bullet)$  vanishes for all  $\delta_\circ \in D'_\circ \setminus D_\circ$ .  $\square$

Based on this lemma, we obtain in the following statements an alternative realization on the  $\mathbf{g}$ -vector fan, which is illustrated on Figure 10.

**Theorem 56.** *For two dissections  $D_\circ \subset D'_\circ$ , the  $\mathbf{g}$ -vector fan  $\mathcal{F}^{\mathbf{g}}(D_\circ)$  is precisely the set of cones  $\{C \in \mathcal{F}^{\mathbf{g}}(D'_\circ) \mid C \subset \mathbb{R}^{D_\circ}\}$  and coincides with the section of the  $\mathbf{g}$ -vector fan  $\mathcal{F}^{\mathbf{g}}(D'_\circ)$  by  $\mathbb{R}^{D_\circ}$ .*

*Proof.* Lemma 55 immediately implies that  $\mathcal{F}^{\mathbf{g}}(D_\circ) = \{C \in \mathcal{F}^{\mathbf{g}}(D'_\circ) \mid C \subset \mathbb{R}^{D_\circ}\}$ . A priori, it is a subfan of the section  $\mathcal{F}^{\mathbf{g}}(D'_\circ)|_{\mathbb{R}^{D_\circ}} = \{C \cap \mathbb{R}^{D_\circ} \mid C \in \mathcal{F}^{\mathbf{g}}(D'_\circ)\}$ . However, since  $\mathcal{F}^{\mathbf{g}}(D_\circ)$  is already a complete simplicial fan of  $\mathbb{R}^{D_\circ}$ , it coincides with  $\mathcal{F}^{\mathbf{g}}(D'_\circ)|_{\mathbb{R}^{D_\circ}}$ .  $\square$

**Theorem 57.** *For two dissections  $D_\circ \subset D'_\circ$ , the  $\mathbf{g}$ -vector fan  $\mathcal{F}^{\mathbf{g}}(D_\circ)$  is realized by the orthogonal projection of the  $D'_\circ$ -accordiohedron  $\text{Acco}(D'_\circ)$  on  $\mathbb{R}^{D_\circ}$ , which is equivalently described by:*

- ◊ the convex hull of the points  $\sum_{\delta_\bullet \in D_\bullet} \omega(D'_\circ | \delta_\bullet) \cdot \mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet)$  for all  $D_\circ$ -accordion dissections  $D_\bullet$ ,
- ◊ the intersection of the half-spaces  $\{\mathbf{x} \in \mathbb{R}^{D_\circ} \mid \langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{x} \rangle \leq \omega(D'_\circ | \delta_\circ)\}$  for all  $D_\circ$ -accordion diagonals  $\gamma_\bullet$ .

*Proof.* Since  $\mathcal{F}^{\mathbf{g}}(D'_\circ)$  is the normal fan of  $\text{Acco}(D'_\circ)$ , Theorem 56 implies that  $\mathcal{F}^{\mathbf{g}}(D_\circ) = \mathcal{F}^{\mathbf{g}}(D'_\circ)|_{\mathbb{R}^{D_\circ}}$  is the normal fan of the orthogonal projection of  $\text{Acco}(D'_\circ)$  on  $\mathbb{R}^{D_\circ}$  [Zie95, Lem. 7.11]. We therefore just need to prove the given vertex and facet descriptions of this projection. First, since  $\mathcal{F}^{\mathbf{g}}(D_\circ) = \mathcal{F}^{\mathbf{g}}(D'_\circ)|_{\mathbb{R}^{D_\circ}}$ , the inequalities of the projection of  $\text{Acco}(D'_\circ)$  on  $\mathbb{R}^{D_\circ}$  are just the inequalities of  $\text{Acco}(D'_\circ)$  whose normal vectors are in  $\mathbb{R}^{D_\circ}$ . Finally, the vertex description follows from the inequality description using the same argument as in Lemma 33.  $\square$

**Remark 58.** The projection of the accordiohedron  $\text{Acco}(D'_\circ)$  on  $\mathbb{R}^{D_\circ}$  differs from the accordiohedron  $\text{Acco}(D_\circ)$ : they have both  $\mathcal{F}^{\mathbf{g}}(D_\circ)$  as normal fan, but their precise geometry is different.

**Corollary 59.** *For any hollow dissection  $D_\circ$ , the  $\mathbf{g}$ -vector fan  $\mathcal{F}^{\mathbf{g}}(D_\circ)$  is realized by a projection of an associahedron of [HPS18].*

*Proof.* Apply Theorem 57 to any triangulation  $T_\circ$  that refines  $D_\circ$ .  $\square$

**Remark 60.** Approaching accordion complexes as coordinate sections of  $\mathbf{g}$ -vector fans actually provides more concise (but also less instructive) proofs for Sections 1.3 and 2.3. Namely, consider any dissection  $D_\circ$  and let  $T_\circ$  be a triangulation that refines  $D_\circ$ . The sign coherence property for triangulations (see Corollary 24) shows that the section  $\mathcal{F}^{\mathbf{g}}(T_\circ)|_{\mathbb{R}^{D_\circ}} = \{C \cap \mathbb{R}^{D_\circ} \mid C \in \mathcal{F}^{\mathbf{g}}(T_\circ)\}$  actually coincides with  $\{C \in \mathcal{F}^{\mathbf{g}}(T_\circ) \mid C \subset \mathbb{R}^{D_\circ}\}$ . Therefore, this gives an alternative concise proof that the collection of cones  $\{C \in \mathcal{F}^{\mathbf{g}}(T_\circ) \mid C \subset \mathbb{R}^{D_\circ}\}$  forms a complete simplicial fan. Moreover, this fan has the same combinatorics as the  $D_\circ$ -accordion complex  $\mathcal{AC}(D_\circ)$  by Lemma 55. We conclude directly that  $\mathcal{AC}(D_\circ)$  is a pseudomanifold realized by the fan  $\{C \in \mathcal{F}^{\mathbf{g}}(T_\circ) \mid C \subset \mathbb{R}^{D_\circ}\}$  and by the orthogonal projection of the associahedron  $\text{Asso}(T_\circ)$  on  $\mathbb{R}^{D_\circ}$ .

**4.3. Cluster algebra analogues.** The perspective on accordion complexes developed in this section also opens the door to generalizations on arbitrary cluster algebras (finite type or not). Namely, consider an arbitrary cluster  $X_\circ = (x_\circ^1, \dots, x_\circ^m)$  in an arbitrary cluster algebra  $\mathcal{A}$ . For any cluster variable  $y \in \mathcal{A}$ , we denote by  $\mathbf{g}(X_\circ \mid y) \in \mathbb{R}^m$  and  $\mathbf{d}(X_\circ \mid y) \in \mathbb{R}^m$  the  $\mathbf{g}$ - and  $\mathbf{d}$ -vectors of  $y$  computed with respect to  $X_\circ$ , see [FZ02, FZ07]. Fix a non-empty proper subset  $I$  of  $[m]$ . We consider two natural subcomplexes of the cluster complex of  $\mathcal{A}$ :

- ◊ the subcomplex  $\Delta^{\mathbf{d}}(X_\circ, I)$  induced by the variables  $y$  such that  $\mathbf{d}(X_\circ \mid y)_i = 0$  for all  $i \in I$ ,
- ◊ the subcomplex  $\Delta^{\mathbf{g}}(X_\circ, I)$  induced by the variables  $y$  such that  $\mathbf{g}(X_\circ \mid y)_i = 0$  for all  $i \in I$ .

It is well-known that the subcomplex  $\Delta^{\mathbf{d}}(X_\circ, I)$  is the cluster complex obtained by freezing all variables  $x_i$  for  $i \in I$ . For example in type  $A$ , it is a join of simplicial associahedra and it can therefore be realized by a product of smaller associahedra. In contrast, we do not know whether the subcomplex  $\Delta^{\mathbf{g}}(X_\circ, I)$  has been investigated. The present paper dealt with the type  $A$  situation.

**Example 61.** Let  $T_\circ$  be a triangulation, with internal diagonals labeled by  $1, \dots, m$ . Consider the corresponding type  $A_m$  cluster  $X_\circ$ . Then for any non-empty proper subset  $I$  of  $[m]$ , the




FIGURE 10. Projecting accordiohedra on coordinate planes yields smaller accordiohedra.

subcomplex  $\Delta^{\mathfrak{g}}(X_{\circ}, I)$  is isomorphic to the  $D_{\circ}$ -accordion complex, where  $D_{\circ}$  is the dissection obtained by deleting in  $T_{\circ}$  the diagonals labeled by  $I$ .

**Example 62.** Example 61 extends to cluster algebras on surfaces [FST08, FT12], using accordions of dissections of surfaces.

The following statement extends Theorem 56 to arbitrary cluster algebras.

**Theorem 63.** *The subset  $\{C \in \mathcal{F}^{\mathfrak{g}}(X_{\circ}) \mid C \subseteq \mathbb{R}^{[m] \setminus I}\}$  of the  $\mathfrak{g}$ -vector fan  $\mathcal{F}^{\mathfrak{g}}(X_{\circ})$  of  $X_{\circ}$  coincides with the section  $\mathcal{F}^{\mathfrak{g}}(X_{\circ})|_{\mathbb{R}^{[m] \setminus I}} = \{C \cap \mathbb{R}^{[m] \setminus I} \mid C \in \mathcal{F}^{\mathfrak{g}}(X_{\circ})\}$ .*

*Proof.* The inclusion  $\{C \in \mathcal{F}^{\mathfrak{g}}(X_{\circ}) \mid C \subseteq \mathbb{R}^{[m] \setminus I}\} \subseteq \mathcal{F}^{\mathfrak{g}}(X_{\circ})|_{\mathbb{R}^{[m] \setminus I}}$  is clear. For the reverse inclusion, we use the sign coherence property of  $\mathfrak{g}$ -vectors in cluster algebras, which was conjectured in [FZ07, Conj. 6.13] and proved in [GHKK, Thm. 5.1] in general. This property implies that the coordinate plane  $\mathbb{R}^{[m] \setminus I}$  intersects any cone  $C$  of  $\mathcal{F}^{\mathfrak{g}}(X_{\circ})$  in a face  $C'$ . This shows that  $C \cap \mathbb{R}^{[m] \setminus I} = C'$  belongs to  $\{C \in \mathcal{F}^{\mathfrak{g}}(X_{\circ}) \mid C \subseteq \mathbb{R}^{[m] \setminus I}\}$ .  $\square$

**Corollary 64.** *The subcomplex  $\Delta^{\mathfrak{g}}(X_{\circ}, I)$  induced by the variables  $y$  such that  $\mathfrak{g}(X_{\circ} \mid y)_i = 0$  for all  $i \in I$  is a pseudomanifold.*

Moreover, extending the result of C. Hohlweg, C. Lange and H. Thomas [HLT11] in the acyclic case, C. Hohlweg, V. Pilaud and S. Stella recently constructed a polytope  $\text{Asso}(X_{\circ})$  realizing the  $\mathfrak{g}$ -vector fan  $\mathcal{F}^{\mathfrak{g}}(X_{\circ})$  in [HPS18]. We can use this associahedron to realize the subcomplex  $\Delta^{\mathfrak{g}}(X_{\circ}, I)$  as a convex polytope, extending Theorem 57.

**Corollary 65.** *The orthogonal projection of  $\text{Asso}(X_{\circ})$  on  $\mathbb{R}^{[m] \setminus I}$  is a realization of  $\Delta^{\mathfrak{g}}(X_{\circ}, I)$ .*

Finally, when oriented in the suitable direction  $v$  (the sum of the positive roots, or equivalently the sum of the fundamental weights), the graph of the generalized associahedron  $\text{Asso}(X_{\circ})$  is the Hasse diagram of a Cambrian lattice [Rea06]. One can similarly orient the graph of the projection of  $\text{Asso}(X_{\circ})$  on  $\mathbb{R}^{[m] \setminus I}$  in the direction of the projection of  $v$  on  $\mathbb{R}^{[m] \setminus I}$ . Is the resulting graph the Hasse diagram of a lattice? Combining the results of [GM16] with that of the present paper shows that this property holds in type  $A$ . We also computationally verified the statement in types  $B_4$ ,  $B_5$ ,  $D_4$  and  $D_5$ . Following [GM16] it seems promising to construct first a lattice structure on biclosed sets of  $\mathfrak{c}$ -vectors, and to obtain then the graph of the projection of  $\text{Asso}(X_{\circ})$  on  $\mathbb{R}^{[m] \setminus I}$  as the Hasse diagram of a lattice quotient.

To conclude, let us mention that the ideas developed in this section have also inspired further investigation of sections of  $\mathfrak{g}$ -vector fans of support  $\tau$ -tilting complexes of associative algebras, see [PPS17] and [PPP17, Sect. 4.2.6].

ACKNOWLEDGEMENTS

We thank C. Hohlweg and S. Stella for many helpful discussions on realizations of the associahedron [HPS18] which were the starting point of this paper. We are grateful to F. Chapoton for various conversations on quadrangulations and Stokes posets, and to A. Garver and T. McConville for introducing us with the accordion complexes during FPSAC'16. Their works [Cha16, GM16] were highly inspiring and motivating. We also thank N. Thiery for a question which led to the approach of Section 4.2, and to P.-G. Plamondon for discussions on the generalization to cluster algebras presented in Section 4.3. Finally, we are grateful to two anonymous referees for their attentive reading and their suggestions on the content and presentation which largely improved our original draft.

REFERENCES

[AIR14] Takahide Adachi, Osamu Iyama, and Idun Reiten.  $\tau$ -tilting theory. *Compos. Math.*, 150(3):415–452, 2014.

[Bar01] Yuliy Baryshnikov. On Stokes sets. In *New developments in singularity theory (Cambridge, 2000)*, volume 21 of *NATO Sci. Ser. II Math. Phys. Chem.*, pages 65–86. Kluwer Acad. Publ., Dordrecht, 2001.

- [BDM<sup>+</sup>17] Thomas Brüstle, Guillaume Douville, Kaveh Mousavand, Hugh Thomas, and Emine Yıldırım. On the combinatorics of gentle algebras. Preprint, [arXiv:1707.07665](https://arxiv.org/abs/1707.07665), 2017.
- [BDP14] Thomas Brüstle, Grégoire Dupont, and Matthieu Pérotin. On maximal green sequences. *Int. Math. Res. Not. IMRN*, (16):4547–4586, 2014.
- [BFS90] Louis J. Billera, Paul Filliman, and Bernd Sturmfels. Constructions and complexity of secondary polytopes. *Adv. Math.*, 83(2):155–179, 1990.
- [BMP16] Amir-Hossein Bateni, Thibault Manneville, and Vincent Pilaud. A note on quadrangulations and Stokes complexes. In preparation, 2016.
- [CD06] Michael P. Carr and Satyan L. Devadoss. Coxeter complexes and graph-associahedra. *Topology Appl.*, 153(12):2155–2168, 2006.
- [CFZ02] Frédéric Chapoton, Sergey Fomin, and Andrei Zelevinsky. Polytopal realizations of generalized associahedra. *Canad. Math. Bull.*, 45(4):537–566, 2002.
- [Cha16] Frédéric Chapoton. Stokes posets and serpent nests. *Discrete Math. Theor. Comput. Sci.*, 18(3), 2016.
- [CSZ15] Cesar Ceballos, Francisco Santos, and Günter M. Ziegler. Many non-equivalent realizations of the associahedron. *Combinatorica*, 35(5):513–551, 2015.
- [DIJ17] Laurent Demonet, Osamu Iyama, and Gustavo Jasso.  $\tau$ -tilting finite algebras,  $\mathbf{g}$ -vectors and brick- $\tau$ -rigid correspondence. *Int. Math. Res. Not. IMRN*, pages Art. ID rnx135, 17, 2017.
- [DRS10] Jesus A. De Loera, Jörg Rambau, and Francisco Santos. *Triangulations: Structures for Algorithms and Applications*, volume 25 of *Algorithms and Computation in Mathematics*. Springer Verlag, 2010.
- [FS05] Eva Maria Feichtner and Bernd Sturmfels. Matroid polytopes, nested sets and Bergman fans. *Port. Math. (N.S.)*, 62(4):437–468, 2005.
- [FST08] Sergey Fomin, Michael Shapiro, and Dylan Thurston. Cluster algebras and triangulated surfaces I. Cluster complexes. *Acta Math.*, 201(1):83–146, 2008.
- [FT12] Sergey Fomin and Dylan Thurston. Cluster algebras and triangulated surfaces. part II: Lambda lengths. Preprint, [arXiv:1210.5569](https://arxiv.org/abs/1210.5569), 2012.
- [FZ02] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. I. Foundations. *J. Amer. Math. Soc.*, 15(2):497–529, 2002.
- [FZ03a] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. II. Finite type classification. *Invent. Math.*, 154(1):63–121, 2003.
- [FZ03b] Sergey Fomin and Andrei Zelevinsky.  $Y$ -systems and generalized associahedra. *Ann. of Math. (2)*, 158(3):977–1018, 2003.
- [FZ07] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. IV. Coefficients. *Compos. Math.*, 143(1):112–164, 2007.
- [GHKK] Mark Gross, Paul Hacking, Sean Keel, and Maxim Kontsevich. Canonical bases for cluster algebras. *J. Amer. Math. Soc.*
- [GKZ08] Israel Gelfand, Mikhail Kapranov, and Andrei Zelevinsky. *Discriminants, resultants and multidimensional determinants*. Modern Birkhäuser Classics. Birkhäuser Boston Inc., Boston, MA, 2008. Reprint of the 1994 edition.
- [GM16] Alexander Garver and Thomas McConville. Oriented flip graphs and noncrossing tree partitions. Preprint, [arXiv:1604.06009](https://arxiv.org/abs/1604.06009), 2016.
- [Hai84] Mark Haiman. Constructing the associahedron. Unpublished manuscript, 11 pages, available at <http://www.math.berkeley.edu/~mhaiman/ftp/assoc/manuscript.pdf>, 1984.
- [HL07] Christophe Hohlweg and Carsten Lange. Realizations of the associahedron and cyclohedron. *Discrete Comput. Geom.*, 37(4):517–543, 2007.
- [HLT11] Christophe Hohlweg, Carsten Lange, and Hugh Thomas. Permutahedra and generalized associahedra. *Adv. Math.*, 226(1):608–640, 2011.
- [Hoh] Christophe Hohlweg. Permutahedra and associahedra. Pages 129–159 in [MHPS12].
- [HPS18] Christophe Hohlweg, Vincent Pilaud, and Salvatore Stella. Polytopal realizations of finite type  $\mathbf{g}$ -vector fans. *Adv. Math.*, 328:713–749, 2018.
- [Lee89] Carl W. Lee. The associahedron and triangulations of the  $n$ -gon. *European J. Combin.*, 10(6):551–560, 1989.
- [Lod04] Jean-Louis Loday. Realization of the Stasheff polytope. *Arch. Math. (Basel)*, 83(3):267–278, 2004.
- [MHPS12] Folkert Müller-Hoissen, Jean Marcel Pallo, and Jim Stasheff, editors. *Associahedra, Tamari Lattices and Related Structures. Tamari Memorial Festschrift*, volume 299 of *Progress in Mathematics*. Springer, New York, 2012.
- [MP17] Thibault Manneville and Vincent Pilaud. Compatibility fans for graphical nested complexes. *J. Combin. Theory Ser. A*, 150:36–107, 2017.
- [Pil13] Vincent Pilaud. Signed tree associahedra. Preprint, [arXiv:1309.5222](https://arxiv.org/abs/1309.5222), 2013.
- [Pos09] Alexander Postnikov. Permutohedra, associahedra, and beyond. *Int. Math. Res. Not. IMRN*, (6):1026–1106, 2009.
- [PPP17] Yann Palu, Vincent Pilaud, and Pierre-Guy Plamondon. Non-kissing complexes and  $\tau$ -tilting for gentle algebras. Preprint, [arXiv:1707.07574](https://arxiv.org/abs/1707.07574), 2017.
- [PPS17] Vincent Pilaud, Pierre-Guy Plamondon, and Salvatore Stella. A  $\tau$ -tilting approach to dissections of polygons. Preprint, [arXiv:1710.02119](https://arxiv.org/abs/1710.02119), 2017.

- [PS12] Vincent Pilaud and Francisco Santos. The brick polytope of a sorting network. *European J. Combin.*, 33(4):632–662, 2012.
- [PS15] Vincent Pilaud and Christian Stump. Brick polytopes of spherical subword complexes and generalized associahedra. *Adv. Math.*, 276:1–61, 2015.
- [Rea06] Nathan Reading. Cambrian lattices. *Adv. Math.*, 205(2):313–353, 2006.
- [Rea07] Nathan Reading. Sortable elements and Cambrian lattices. *Algebra Universalis*, 56(3-4):411–437, 2007.
- [RS09] Nathan Reading and David E. Speyer. Cambrian fans. *J. Eur. Math. Soc.*, 11(2):407–447, 2009.
- [SS93] Steve Shnider and Shlomo Sternberg. *Quantum groups: From coalgebras to Drinfeld algebras*. Series in Mathematical Physics. International Press, Cambridge, MA, 1993.
- [Sta63] Jim Stasheff. Homotopy associativity of H-spaces I, II. *Trans. Amer. Math. Soc.*, 108(2):293–312, 1963.
- [Ste13] Salvatore Stella. Polyhedral models for generalized associahedra via Coxeter elements. *J. Algebraic Combin.*, 38(1):121–158, 2013.
- [Tam51] Dov Tamari. *Monoides préordonnés et chaînes de Malcev*. PhD thesis, Université Paris Sorbonne, 1951.
- [Zel06] Andrei Zelevinsky. Nested complexes and their polyhedral realizations. *Pure Appl. Math. Q.*, 2(3):655–671, 2006.
- [Zie95] Günter M. Ziegler. *Lectures on Polytopes*, volume 152 of *Graduate texts in Mathematics*. Springer-Verlag, New York, 1995.

(Thibault Manneville) LIX, ÉCOLE POLYTECHNIQUE  
E-mail address: [thibault.manneville@lix.polytechnique.fr](mailto:thibault.manneville@lix.polytechnique.fr)  
URL: <http://www.lix.polytechnique.fr/~manneville/>

(Vincent Pilaud) CNRS & LIX, ÉCOLE POLYTECHNIQUE, PALAISEAU  
E-mail address: [vincent.pilaud@lix.polytechnique.fr](mailto:vincent.pilaud@lix.polytechnique.fr)  
URL: <http://www.lix.polytechnique.fr/~pilaud/>