

## AGENT-BASED GENETIC MULTICELLULAR 3D MODELLING OF STEM-CELL SPHEROIDS(GENESPHERE)

Fabrizio Cleri, Chann Lagadec, Ralf Blossey, Pier Luca Palla, Stefano

Giordano

### ▶ To cite this version:

Fabrizio Cleri, Chann Lagadec, Ralf Blossey, Pier Luca Palla, Stefano Giordano. AGENT-BASED GENETIC MULTICELLULAR 3D MODELLING OF STEM-CELL SPHEROIDS(GENESPHERE). 4thInternational Conference on Physics & Biological Systems, 2018 October 22-24, CNRS campus, Oct 2018, Gif-sur-Yvette, France. hal-02344423

### HAL Id: hal-02344423 https://hal.science/hal-02344423

Submitted on 4 Nov 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# AGENT-BASED GENETIC MULTICELLULAR 3D MODELLING OF STEM-CELL SPHEROIDS (GENESPHERE)

 Fabrizio CLERI, Department of Physics and IEMN CNRS UMR8520, University of Lille Chann LAGADEC, Cell Plasticity and Cancer, INSERM U908, Lille
Ralf BLOSSEY, Structural and Functional Glycobiology, CNRS UMR8576, Lille
Pier Luca PALLA, Department of Physics and IEMN CNRS UMR8520, University of Lille
Stefano GIORDANO, Ecole Centrale de Lille and IEMN CNRS UMR8520, Lille


#### **COULD X-ray IRRADIATED BREAST-CANCER CELLS REPROGRAM INTO CANCER-STEM-CELLS ?**


The main objective of GENESPHERE is to produce A a model capable of testing the hypothesis of radiation-induced breast cancer stem cells. In recent experiments, we observed CSC-specific markers to appear in irradiated non-tumorigenic cells (T47D, MCF-7, MDA-MB-231 and SUM159-PT), either isolated, or in presence of preexisting breast-CSCs. We characterized the upregulation of about 10 different genes, among which key elements of the Notch pathway; it was shown, e.g., that simultaneous downregulation of Sox2 and Nanog significantly reduced the generation of new breast-CSCs, contrary to down-regulation of either one alone; also, down-regulation of Notch receptors reduced the ability to form spheroids and radiation-induced reprogramming.

### **GENESPHERE: AN AGENT-BASED MODEL**


The **Agent-Based Model** describes motile cells in a 3D network of Voronoi polyhedra. Each cell-agent has: a local phase clock; duplication with genetic inheritance; normal, stem, or various cancer-cell phenotype; diffusion of oxygen, nutrients, or other chemicals; a basic version of the glycolysis cycle, with the ATP demand setting the level of glucose/oxygen consumption. Cells can absorb radiation/chemical damage, described by a Markov chain, with individual cell-repair mechanisms.

The next level of enrichment of the model is the inclusion of a *cell systemic network*, with a whole new sets of cell markers describing activation/repression of different factors that could characterize the CSC and radiation-induced CSC populations, by means of PDE-based kinetic modeling.









#### **BIBLIOGRAPHY**

Lagadec C, Vlashi E, Della Donna L, Pajonk F, Stem Cells 30, 833 (2012)
Tomezak M, Abbadie C, Lartigau E, Cleri F, J Theor Biol 389, 146 (2015)
Bailleul-Dubois J, Bidan N, LeBourhis X, Lagadec C, Oncologie, 19, 77 (2017)
Cleri F, Eur Phys J E, submitted - biorkiv: <a href="https://doi.org/10.1101/431163">https://doi.org/10.1101/431163</a>

ISEN 0

LILLE

( 🔴

ontralolille

RENATECH

Université Polytechnique HAUTS-DE-FRANCE

Université

de Lille