



HAL
open science

Biophysical Perspective: The Latest Twists in Chromatin Remodeling

R. Blossey, H. Schiessel

► **To cite this version:**

R. Blossey, H. Schiessel. Biophysical Perspective: The Latest Twists in Chromatin Remodeling. Biophysical Journal, 2018. hal-02344693

HAL Id: hal-02344693

<https://hal.science/hal-02344693v1>

Submitted on 4 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Biophysical Perspective: The Latest Twists in Chromatin Remodeling

R. Blossey

University of Lille 1, Unité de Glycobiologie Structurale et Fonctionnelle, CNRS UMR8576, 59000 Lille, France.

H. Schiessel

Institute Lorentz for Theoretical Physics,
Leiden University, Niels Bohrweg 2,
2333 CA Leiden, The Netherlands.

(Dated: February 2, 2018)

In its most restrictive interpretation, the notion of chromatin remodeling refers to the action of chromatin remodeling enzymes on nucleosomes with the aim to displace and remove them from the chromatin fiber (the effective polymer formed by a DNA molecule and proteins). This local modification of the fiber structure can have consequences for the initiation and repression of the transcription process and, when the remodeling processes spreads along the fiber, also results in long-range effects essential for fiber condensation. There are three regulatory levels of relevance that can be distinguished for this process: the first is the intrinsic sequence preference of the histone octamer which rules the positioning of the nucleosome along the DNA, notably in relation to the genetic information coded in DNA, the second is the recognition or selection of nucleosomal substrates by remodeling complexes, and the final one the motor action on the nucleosome exerted by the chromatin remodeler. On each of these three levels recent work has been able to provide crucial insights which add new twists to this exciting and unfinished story, which we highlight in this perspective.

I. INTRODUCTION

The nucleosome, the basic structural unit of the chromatin fiber, remains a key subject of interest for chromatin biologists and physicists. A key basic insight of recent years was the (indirect) sequence-dependence of its positioning along DNA [1]. Already much earlier it was noticed that the nucleosome structure is a flexible, dynamic structure with an intrinsic, thermally driven mobility [2]. In the context of many chromatin-related processes, the nucleosome needs to be actively displaced or removed. The molecular machines that organize this process are called chromatin remodelers [3], commonly grouped in four distinct families, see Figure 1 [4, 5]. Chromatin remodelers are increasingly understood as key factors also in the development of diseases due to the deregulation of gene expression such as in cancer, see, *e.g.* [6]. The definition of the chromatin remodeler families relies on the basic motor unit, the ATPase, which derives from helicases, and the positioning of additional regulatory subunits around the ATPase.

How can we imagine a chromatin remodeler to work on a nucleosome? Being researchers in the low countries (The Netherlands and French Flanders), we allow ourselves to resort to an analogy drawn from a favorite transportation means in our regions, the bicycle. Man-powered bikes may still be the most frequently used type of bikes, but additional power is built in in the ever more popular eBikes. For those who want to have the best of both worlds, there are the ‘clip-on’ motors: if one wants to go faster, one fixes a small add-on motor to power the bike. An example is shown in Figure 2. Chromatin remodelers, within this analogy, are nothing but ‘clip-on’




FIG. 1. Definition of the remodeler families according to [4]. DExx and HELICc (shown in red) are ATPase domains, *Bromo*, SANT and SLIDE, and *Tandem chromo* (shown in blue) are accessory domains that recognize specific histone tail modifications, or DNA. Adapted from [4].

motors to nucleosomes. However, they do not propel the nucleosome in a linear fashion, as many molecular motors do that simply transport cargo. Chromatin remodeling motors displace the DNA wrapped around the histone octamer, and this obviously is a much more complex task than linear transport. Our simplistic analogy, however, also makes us ask simple questions such as: where are these motors actually fixed on the nucleosome? How do they pull on the DNA? How is the DNA transported around the nucleosome? Recent research has given at



FIG. 2. Top: a ‘clip-on’ motor that turns a bike into an eBike; the bottom figure shows how the motor is fixed on the bike. Courtesy: add-e

least partial answers to these basic questions.

Before turning to the topic of the motor action, we will address the two additional levels of chromatin regulation concerning the nucleosome we mentioned at the beginning. The first and most fundamental one is the positioning of the nucleosome along the fiber, which is dictated in a complex way by the DNA sequence - or rather, the physical properties associated with the stacking of the base pairs. The second level is the problem of how the machinery recognizes the right nucleosome at the right time. Some time ago we have postulated a ‘kinetic proofreading’ scenario for this process, for which some experimental justification has become available shortly afterwards. New high-throughput data that have been published only very recently lend new experimental support to our idea. Finally, on level three, recent progress in the understanding of the action of remodeling motors has become available through new experimental techniques that allow to trace the dynamics of the remodeler-nucleosome complexes, which we highlight as we go through some new results. Key in this field have been the substantial advances in cryo-electron microscopy, freshly honoured with the Nobel prize in Chemistry in 2017 for the pioneers of this technique. In the following, we will step through the recent insights on these three levels.

II. NUCLEOSOME POSITIONING

In a nucleosome about one persistence length of DNA is wrapped one and three quarter turns around an octamer of histone proteins, which amounts to a substantial amount of bending energy (about $60 k_B T$ [7]). On the other hand, the elasticity and geometry of the DNA double helix depends on the underlying sequence of base pairs [8, 9]. It is thus to be expected that the affinity of a given stretch of DNA to be complexed in a nucleosome varies strongly with sequence. Moreover, the binding of the DNA to the octamer occurs mainly at 14 patches where the minor groove of the DNA touches the surface of the octamer [10]. As the sugar-phosphate backbones do not change with the underlying base pair sequence, the pure binding energy (mainly based on charge-charge interactions between the DNA phosphates and histones [11]) is expected to be only weakly sequence dependent. Taken together this suggests that the nucleosome performs an indirect readout of the basepair sequence, with the affinity reflecting the overall ability of 147 base pairs to wrap around the octamer.

Nucleosomes show indeed characteristic sequence preferences [1, 12], e.g. high affinity sequences feature an increased occurrence of GC dinucleotides (a G followed by a C) at locations where the minor groove faces outwards, whereas AA, TT and TA dinucleotides peak in between, where the minor groove faces inwards towards the histone octamer (i.e. at the binding sites), see Figure 3(a). These preferences have been proposed to constitute a “genomic code for nucleosome positioning” [1] (for earlier versions of this idea see e.g. Refs. [12, 13]). In fact, the sequence preferences of the nucleosome give rise to two types of positional preferences: rotational and translational positioning. Rotational positioning results from the fact that DNA is attached at its minor groove to the octamer. Therefore when a nucleosome is repositioned one base pair step at a time the DNA needs to perform a corkscrew-like motion. As a given wrapped stretch of DNA has typically a preferred bending direction (even for a completely random base pair sequence), there is every 10 base pairs a minimum in the bending energy, causing the rotational positioning of the nucleosome. The second type of preference, translational positioning, reflects an overall preference of nucleosomes to sit on DNA stretches with high GC content and to avoid certain motifs like e.g. poly(dA:dT) tracts.

Nucleosomes have been mapped genome-wide *in vivo* and *in vitro* (in the latter case nucleosomes are reconstituted on DNA from their pure components) using a variety of methods, e.g. through digestion by nuclease [1] and by a chemical cleavage technique [14]. Such nucleosome maps have taught us that indeed non-random mechanical cues can be found written along genomes. This becomes especially clear by focusing on particular locations, e.g. at transcription start sites (TSSs), and then by averaging over all such locations (e.g. over the TSSs of all genes of an organism). Such studies revealed that yeast shows on




FIG. 3. (a) Nucleosome positioning rules [1]: Key dinucleotides are shown relative to one-half of the symmetric nucleosome. (b) A coarse grained computational model of the nucleosome as employed in Refs. [18–20].

average a depletion of nucleosomes just in front of genes [1, 14]. But is this depletion reflecting nucleosome sequence preferences or is it caused by other mechanisms, e.g. the action of chromatin remodellers [15]? In general this is a complex question and the answer clearly depends on the organism at hand.

To answer this question it is necessary to develop methods that allow to estimate nucleosome sequence preferences along DNA molecules. Bioinformatics models trained on experimental nucleosome maps provide a common approach [1]. Deeper insight into the relation between DNA mechanics and nucleosome positioning can be gained from coarse-grained computer models of nucleosomes that give fairly reliable estimates of sequence preferences [16–18], see e.g. Figure 3(b). These models, however, typically tend to be too slow to perform e.g. genome wide averages of the DNA mechanics around TSSs. The recently developed Mutation Monte Carlo method [18] overcomes this problem as it allows to determine the sequence preferences of a given model nucleosome that in turn can be used to build probabilistic models [19, 20]. Such models are similar to the above-mentioned bioinformatics approaches but ultimately are based on specific microscopic models with well-known ingredients. Starting from an already fairly efficient coarse-grained nucleosome model, this approach leads to a speedup of about 10^5 [19]. Applied to TSSs of yeast [21] we found excellent agreement with experimental data. As the approach is based on a purely mechanical model, this suggests that nucleosome depletion at TSSs in yeast is indeed caused by sequence-dependent DNA mechanics.

An exciting question to ask is to what extent genomes position nucleosomes. This turns out to be a tricky question. For instance, in Ref. [1] it was claimed that about 50% of the nucleosomes on the yeast genomes can be “explained solely by sequence preferences.” But this statement has to be taken with a grain of salt as it does not automatically mean that there are dedicated locations for 50% of the nucleosomes directly encoded into the DNA sequence. When one looks at the *in vivo* nucleosome

maps of yeast it seems that nucleosomes are well-positioned along the beginning of each gene such that one can count them, one after the other (one speaks indeed of the +1 nucleosome, the +2 nucleosome and so on [14]). But when one calculates the average positional preferences of the nucleosomes around the TSSs [21], one finds mainly an “anti-positioning” element just in front of genes, an AT-rich piece of DNA that repels nucleosomes. The appearance of clearly distinguishable “positioned” nucleosomes along the beginning of genes in yeast is then in fact just a statistical effect, caused by the relatively large density of nucleosomes and the presence of a boundary constraint as suggested in the classical work of Kornberg and Stryer [22]. For the much lower density of nucleosomes obtained via *in vitro* reconstitution, these seemingly well-defined peaks vanish completely [23] (the situation is in fact a bit more complex; for details we refer the reader to the excellent review by Struhl and Segal [15]).

Even though nucleosomes seem not to be positioned individually in yeast, the statistical ordering of nucleosomes caused by the presence of a boundary can still be said to be “explained” by nucleosomal sequence preferences. But the more exciting question remains whether there are also examples of nucleosomes that are specifically positioned by sequence at certain genomic locations. There are indeed numerous examples. In fact, the genomes of many organisms attract nucleosomes to TSSs. A rule that seems to be generally true (at least for the 35 eukaryotic genomes we analysed [21]) is that genomes of unicellular organisms repel nucleosomes from TSSs whereas multicellular life forms show a region that attracts nucleosomes. For instance, nematodes like *C. elegans* have (on average) a strongly positioned nucleosome just at the entrance to its genes [21]. Mammals (e.g. mouse, chimpanzee, human) show a broad region with high overall nucleosome affinity [21].

What is the biological function of those various motifs? At this point one can only speculate. For yeast the encoded nucleosome depletion “may indicate that eukaryotic genomes direct the transcriptional machinery to functional sites by encoding unstable nucleosomes over these elements, thereby enhancing their accessibility” [1]; the same would be true for all unicellular lifeforms [21]. In *C. elegans* one finds a single well-positioned nucleosome at the beginning of many genes *in vitro* and *in vivo*, as predicted by our mechanical model [21], and this might function as a mechanism for TSS selection for RNA polymerase [24]. In humans the situation is complex. The high nucleosome occupancy encoded by the sequence at promoters [21] has been speculated “to restrict access to regulatory information that will ultimately be utilized in only a subset of differentiated cells” [25]. However, the *in vivo* nucleosome occupancy does not correlate well with that intrinsic nucleosome affinity, possibly reflecting transcription related processes [26]. A possible explanation is that the mechanical cues serve here a different purpose, namely to determine which nu-

cleosomes are retained in sperm cells (where most nucleosomes are replaced by protamines), allowing the transmission of paternal epigenetic information. In fact, the mechanical cues correlate well with the nucleosome retention score in sperm cells [21, 26]. This does, however, not mean that there are no mechanical cues for nucleosomes in somatic human cells. A spectacular example are 6×10^6 nucleosomes mechanically positioned around nucleosome inhibiting barriers, but the biological function has yet to be determined [27].

We finally mention two more important points that are not well understood yet. The first one is that the nucleosome density varies with cell type or developmental state even though all cells of an organism have the same DNA sequences. This is not inconsistent with the idea of mechanical cues on DNA molecules as regions of nucleosomes might act as “nucleosomal switches” [28]. Another complication is the presence of higher order chromatin structures that might influence nucleosome positioning, a subject that to the best of our knowledge has not yet been addressed.

Sequence-dependent DNA mechanics is thus one of several players that determine nucleosome positions along genomes. We turn now to another key mechanism: the action of chromatin remodelers.

III. NUCLEOSOME RECOGNITION BY REMODELERS

Chromatin remodelers have a very high affinity for DNA so that they will ‘remodel’ even a naked DNA molecule, and this even without ATP [29]. This basic background mechanism therefore needs to be tuned such that the remodelers act on the right objects (the nucleosomes) at the right place (e.g. near a TSS), and at the right time.

A mechanism by which the remodelers identify their targets has been proposed by us in 2008 [30]. It is a variant of the famous kinetic proofreading mechanism originally proposed by John Hopfield in the context of mRNA translation [31]. In the context of chromatin remodeling, histone tail modifications placed by ‘histone writers’ can be read by accessory subunits of the chromatin remodelers which are sensitive to these states. The essentially same idea was put forth shortly after by Geeta Narlikar for the ISWI/ACF system, based on her very detailed experiments [32, 33].

Very recently, a high-throughput analysis confronting different remodelers with the different possible histone tail modifications lends very strong support for the correlation between histone tails states and the outcomes of remodeler actions [34]. While the results have not been interpreted within the proofreading scenario, the observations themselves are clearly in support of it. Apart from the recognition of histone modifications, a key element of the kinetic proofreading scenario, there

are also additional levels of recognition encoded which provide alternative means of regulation. Since ISWI recognizes the H4-tail only without modifications, there is an additional level embedded in this system as a ‘basic patch’ on the tail enters in competition with a like patch on the remodeler itself: the remodeling motor is inhibited by some of its own domains (called AutoN and NegC), an inhibition that is only relieved in the presence of the histone tail [35]. Recently, the multivalency of bromodomains of the BAF chromatin remodeling complex has been elucidated, as they bind to both DNA and the histone tail modification H3K14ac [36]. These insights show that even within the interaction of a specific remodeler type with a nucleosome a rich network of recognition mechanisms has evolved for the fine tuning of remodeler action. These developments are clearly important for more general predictive models of remodeler recruitment on similar large-scale levels as for sequence-directed nucleosome positioning. Statistical physics-based models in this direction were already put forward in [37, 38].

IV. CHROMATIN REMODELER ACTION

A. Motor engagement

Three recent studies have provided new key insights into the engagement of chromatin remodeling motors on the nucleosome; as they consider remodelers from different families, the insights they yield nicely complement each other.

In a study by Liu *et al.* [39], summarized by Wigley and Bowman [40], the interaction between the ATPase unit of Snf2 (also called SWI-SNF) and the nucleosome was studied by cryo-EM. The study reveals that the ATPase is bound to an internal site on superhelical location 2 (SHL2). As the structure is nucleotide-free, the ATP-binding cleft is in an open conformation, which by comparison with closed conformations allows to conclude that the closure of the cleft would push 1 bp of DNA towards the nucleosome dyad. An important insight from this study is that in addition to the binding site of the ATPase at SHL2, an acidic patch on the first lobe of the ATPase contacts the DNA 90 bp away. Upon ATP binding and closure of the cleft, the second lobe can move the DNA duplex toward the dyad axis across the histone octamer. After ATP-hydrolysis, lobe 2 would need to reset without sliding the DNA back with it. This study therefore provides an answer to the question we asked in the introduction. On the eBike, the ‘clip-on’ motor on the nucleosome needs to be fixed on a position at the frame, and then contacts the rolling wheel of the bike to transmit the power. In the case of the ATPase, the remodeler remains attached with its lobe 1 at SHL2, while lobe 2 dynamically resets upon translocation, see the schematic illustration in Figure 4. Very




FIG. 4. Schematic drawing of the remodeler engagement for the Snf2-construct, following Figure 3 a) from [39] and Figure 2 from [40]. Grey: histone octamer; black: nucleosomal DNA; red: ATPase units; blue: accessory remodeler domains.

similar findings on the workings of the remodeler ATPase were obtained in work by Nodelman *et al.* on the Chd1 remodeler [41], which has also been studied by the Bowman group in much detail [42, 43]. Chd1 is a (smaller) remodeling complex that plays a key role in nucleosome positioning over coding regions [44].

The third study we refer to is the work by Sundaramoorthy *et al.* [45], which also considers Chd1. Like other remodelers, it belongs to the Snf2-family and contains two helicase-like domains for which previous structural information was available for the ATPase domain associated to N-terminal chromodomains, which serve to recognize methylation states of the histone tails. Chd1 resembles ISWI in one important aspect, as it also contains a C-terminal DNA-binding domain composed of the SANT and SLIDE domains. The structural information available on all these structures, however, was limited to these subdomains, not reaching up to the level of the full structure. In the experiments described in [45], SAXS data obtained on full Chd1 complexes (without the nucleosome) showed that Chd1 is a monomeric complex. To obtain higher resolution data, the complex was studied with pulsed EPR after cysteine-serine replacements. The mutant protein showed unmodified remodeler activity. Reintroduction of specifically chosen labelled cysteine pairs allowed to determine distances in the complex, which can be related to the crystal structure. The resulting distance distributions reveal the possible dynamics of the structure through their difference to residue distances in the crystal structure. Together with docking poses these data thus allow to develop an idea of the dynamic contacts the complex can undergo. These results also corroborated by a very recent structural study on the Chd1-nucleosome complex published by the Cramer group [46].

Further insights into the role of the N-terminal tail in the regulation of remodeler activity was studied by performing mutations in several regions, and assessing their

effect on the repositioning of remodelers. Finally, also the engagement of the remodeler to the nucleosome was studied by cryo-EM. These highly detailed studies allow substantial insights into the different contacts the remodeler domains make with nucleosomes during remodeling.

B. A dynamical role for the histone octamer

In most studies of chromatin remodeling, the histone octamer is considered as a rather static bystander of remodeling events: it is essentially the spool around which the DNA is wrapped and hence unspooled or transported around this protein structure. A recent study by Sinha, Gross and Narlikar [47] challenges this purely ‘passive’ role of the histone octamer. The findings by Sinha *et al.* have been summarized by Flaus and Owen-Hughes [48]. The key idea of this work has been to first introduce artificial modifications in the histone octamer structure, and to correlate the direct observations of the effect of these modifications on NMR spectra of nucleosomes bound to a remodeler with positioning assays of nucleosomes under the action of remodelers.

Specifically, the authors studied ^{13}C -isotope modifications in the methyl groups of buried H4 side chains, and detected the structural changes these modifications induce via nuclear magnetic resonance spectroscopy. The nucleosomes were bound to an Snf2-remodeler. The observed changes appeared prominently in the H3-H4 dimer near the superhelical location 0 (SHL0), close to the center of the nucleosome, which is distant from the location SHL2 where the remodeler is expected to bind the nucleosome. In a separate set of experiments, the authors engineered disulfide linkages between H3 and H4 to restrict their flexibility. In accord with the NMR data, such crosslinks in the vicinity of SHL0 led to a statistically significant number of disrupted nucleosomes, while crosslinks near SHL2 showed no significant effect. By studying remodelers from different families, ISWI, INO80 and RSC, differences between the remodelers became apparent: the crosslinking near SHL2 affects both ISWI and RSC-remodeled nucleosomes, but shows no relevant effect on INO80.

These results clearly show that the full structure of a nucleosome-remodeler complex features dynamical behaviour in all of its subunits. Remodeling has become yet more dynamic than previously thought.

C. Nucleosomes and CRISPR-Cas

As a final point of this perspective article we address the relevance of chromatin remodeling for the currently hottest topic in DNA biology, CRISPR-Cas9. In a recent study involving one of the co-inventors of the CRISPR technology, the role of nucleosomes on the chromatin fiber undergoing the genome editing process has been

investigated, under conditions of the presence or absence of chromatin remodelers [49]. In biochemical assays in which the preferred positioning of nucleosomes on the 601-sequence was used in specifically constructed sequences, the authors could establish that the efficiency of the Cas9 enzyme to cut the DNA is consistent with the well-known nucleosome breathing behaviour [50]. According to this mechanism nucleosomal DNA unpeels temporarily from the ends favouring the interaction of enzymes with sites that are close to the nucleosomal flanks. This has been shown first for restriction enzymes [50, 51] and can now be observed more directly via fluorescence resonance energy transfer [52, 53]. In addition, Cas9 efficiency is promoted by the presence of chromatin remodelers which displace the nucleosomes from the cut sites. Therefore, due to these two mechanisms, the CRISPR-Cas9 system can reach efficiencies in eukaryotes that equal those in bacterial systems, in which the DNA is more readily accessible.

The authors of Ref. [49] also point out that the design of CRISPR-Cas9 systems can be optimized by taking into account the intrinsic nucleosome positions. But to what extent can such a prediction be made? If nucleosome positions are known, coarse-grained models can predict the sequence-dependent accessibility for proteins to DNA target sites inside nucleosomes [50, 51] as recently demonstrated [54]. However, as mentioned earlier, not all nucleosomes are positioned by mechanical signals alone (see e.g. our discussion of nucleosome positioning at human TSSs). We therefore expect that all three levels of chromatin regulation concerning the nucleosome need to be taken into account to arrive at reliable predictions of Cas9 efficiency.

V. CONCLUSIONS

We have provided an overview about recent developments in understanding how nucleosomes give limited

access to their DNA. One important aspect is the sequence-dependent DNA elasticity which can be tuned over evolutionary time scales to create regions of high or low nucleosome stability. We discussed how in the context of transcription start sites this might give access to regulatory information or, depending on the organisms, might determine nucleosome retention in sperm cells. We then turned to chromatin remodellers and first discussed new detailed experimental evidence how remodellers might recognise the “right” nucleosomes for their action. Finally, we discussed how very recent work begins to provide detailed insight how the motor activity of remodellers translates to nucleosome repositioning. The high level of understanding of the microscopic details of these three mechanisms that influence nucleosome positioning and dynamics makes us hopeful that a comprehensive picture of how nucleosomes influence gene expression might be within reach in the near future. In this perspective we have focused on the very recent developments which in our view significantly advance the field. Obviously, we have not been able to cover all aspects here in which progress has been made in the last years. For an extended description covering earlier work we refer the reader to the recent review by Lai and Pugh [55].

Acknowledgements. This perspective article is dedicated to the memory of Jörg Langowski, a longtime friend and collaborator. As many colleagues could witness, Jörg was a unique individual with a limitless source of friendliness, support and criticism. We deeply miss him.

-
- [1] Segal, E., Y. Fondufe-Mittendorf, ..., J. Widom. 2006. A genomic code for nucleosome positioning. *Nature* 442: 772-778.
 - [2] Eslami-Mossallam, B., H. Schiessel, and J. van Noort. 2016. Nucleosome dynamics: Sequence matters. *Adv. Colloid Interface Sci.* 232: 101-113.
 - [3] Smith, C. L., and C. L. Peterson. 2004. ATP-dependent chromatin remodeling. *Current Topics in Developmental Biology* 65: 115-148.
 - [4] Clapier, C. R. and B. R. Cairns. 2009. The biology of chromatin remodeling complexes. *Annu. Rev. Biochem.* 78: 273-304.
 - [5] Blossey R. 2017. *Chromatin: Structure, Dynamics, Regulation*. CRC Press Boca Raton, USA.
 - [6] Kumar, R. D.-Q. Li, ..., S. Knapp. 2016. Epigenomic regulation of oncogenesis by chromatin remodeling. *Oncogene* 35: 4423-4436.
 - [7] Langowski, J. and H. Schiessel. 2006. Chromatin simulations. In *Computational studies of RNA and DNA*. F. Lankaš and J. Šponer, editors. Springer, Berlin, pp 605-634.
 - [8] Olson, W. K., A. A. Gorin, ..., V. B. Zhurkin. 1998. DNA sequence-dependent deformability deduced from protein-DNA crystal complexes. *Proc. Natl. Acad. Sci. USA* 95: 11163-11168.
 - [9] Lankaš, F., J. Šponer, ..., J. Langowski. 2000. Sequence-dependent elastic properties of DNA. *J. Mol. Biol.* 299: 695-709.

- [10] Davey, C. A., D. F. Sargeny, ..., T. J. Richmond. 2002. Solvent Mediated Interactions in the Structure of the Nucleosome Core Particle at 1.9 Å Resolution. *J. Mol. Biol.* 319: 1097-1113.
- [11] Fenley, A. T., D. A. Adams and A. V. Onufriev. 2010. Charge state of the globular histone core controls stability of the nucleosome. *Biophys. J.* 99: 1577-1585.
- [12] Satchwell, S. C., H. R. Drew, and A. A. Travers. 1986. Sequence periodicities in chicken nucleosome core DNA. *J. Mol. Biol.* 191: 659-675.
- [13] Trifonov, E. N. and J. L. Sussmann. 1980. The pitch of chromatin DNA is reflected in its nucleotide sequence. *Proc. Natl. Acad. Sci. USA* 77: 3816-3820.
- [14] Brogaard, K., L. Xi, ..., J. Widom. 2012. A map of nucleosome positions in yeast at base-pair resolution. *Nature* 486: 496-501.
- [15] Struhl, K. and E. Segal. 2013. Determinants of nucleosome positioning. *Nature Struct. Mol. Biol.* 20: 267-273.
- [16] Morozov, A. V., K. Fortney, ..., E. D. Siggia. 2009. Using DNA mechanics to predict *in vitro* nucleosome positions and formation energies. *Nucl. Acids Res.* 37: 4707-4722.
- [17] Freeman, G. S., J. P. Lequieu, ..., J. J. de Pablo. 2014. DNA Shape Dominates Sequence Affinity in Nucleosome Formation. *Phys. Rev. Lett.* 113: 168101.
- [18] Eslami-Mossallam, B., R. D. Schram, ..., H. Schiessel. 2016. Multiplexing genetic and nucleosome positioning codes: a computational approach. *PLoS ONE* 11: e0156905.
- [19] Tompitak, M., G. T. Barkema, and H. Schiessel. 2017. Benchmarking and refining probability-based models for nucleosome-DNA interaction. *BMC Bioinformatics* 18: 157.
- [20] Wondergem, J. A. J., H. Schiessel, and M. Tompitak. 2017. Performing SELEX experiments in silico *J. Chem. Phys.* 147: 174101.
- [21] Tompitak, M., C. Vaillant, and H. Schiessel. 2017. Genomes of multicellular organisms have evolved to attract nucleosomes to promoter regions. *Biophys. J.* 112: 505-511.
- [22] Kornberg, R. D., and L. Stryer. 1988. Statistical distributions of nucleosomes: nonrandom locations by a stochastic mechanism. *Nucl. Acids Res.* 16: 6677-6690.
- [23] Kaplan, N., I. K. Moore, ..., E. Segal. 2009. The DNA-encoded nucleosome organization of a eukaryotic genome. *Nature* 458: 362-366.
- [24] Dreos, R., G. Ambrosini, and P. Buchner. 2016. Influence of rotational nucleosome positioning on transcription start site selection in animal promoters. *PLoS Comput. Biol.* 12: e1005144.
- [25] Tillo, D., N. Kaplan, ..., T. R. Hughes. 2010. High nucleosome occupancy is encoded at human regulatory sequences. *PLoS ONE* 5: e9129.
- [26] Vavouri, T., and B. Lehner. 2011. Chromatin organization in sperm may be the major functional consequence of base composition variation in the human genome. *PLoS Genet.* 7: e1002036.
- [27] Drillon G., B. Audit, ..., A. Arneodo. 2016. Evidence of selection for an accessible nucleosomal array in human. *BMC Genomics* 17: 526.
- [28] Schwab D. J., R. F. Bruinsma, ..., J. Widom. 2008. Nucleosome switches. *Phys. Rev. Lett.* 100: 228105.
- [29] Lia, G., E. Praly, ..., T. Owen-Hughes. 2006. Direct observation of DNA distortion by the RSC chromatin remodelling complex. *Mol. Cell* 21: 417-425.
- [30] Blossey, R., and H. Schiessel. 2008. Kinetic proofreading of gene activation by chromatin remodeling. *HFSP Journal* 2:167-170.
- [31] Hopfield, J. J. 1974. Kinetic proofreading: a new mechanism for reducing errors in biosynthetic processes requiring high specificity. *Proc. Natl. Acad. Sci. USA* 71: 4135-4139.
- [32] Narlikar, G. J. 2010. A proposal for kinetic proof reading by ISWI family chromatin remodeling motors. *Curr. Op. Chem. Biol.* 14:660-665.
- [33] Blossey, R., and H. Schiessel. 2011. Kinetic proofreading in chromatin remodeling: the case of ISWI/ACF *Biophys. J.* 101:L30-L32
- [34] Dann, G. P, G. Liszczak, ..., T. W. Muir. 2017. ISWI chromatin remodellers sense nucleosome modifications to determine substrate preference. *Nature* 548: 607-611.
- [35] Clapier, C. R. and B. R. Cairns. 2012. Regulation of ISWI involves inhibitory modules antagonized by nucleosomal epitopes. *Nature* 492: 280-284.
- [36] Morrison, E. A., J. C. Sanchez, ..., C. A. Musselman. 2017. DNA binding drives the association of BRG1/hBRM bromodomains with nucleosomes. *Nature Communications.* 8:16080.
- [37] Teif, V. B., and K. Rippe. 2009. Predicting nucleosome positions on the DNA: combining intrinsic sequence preferences and remodeler activities. *Nucl. Acids Res.* 7: 5641-5655.
- [38] Padintheeri, R., and J.F. Marko. 2011. Nucleosome positioning in a model of active remodeling enzymes. *Proc. Natl. Acad. Sci. (USA)* 108: 7799-7803.
- [39] Liu, X., M. Li, ..., Z. Chen. 2017. Mechanism of chromatin remodelling revealed by the Snf2-nucleosome structure. *Nature* 544: 440-445.
- [40] Wigley, D. A., and G. D. Bowman. 2017. A glimpse into chromatin remodeling. *Nature Struct. Mol. Biology* 24: 498-500.
- [41] Nodelman, I. M., F. Bleichert, ..., G. D. Bowman. 2017. Interdomain communication of the Chd1 chromatin remodeler across the DNA gyres of the nucleosome. *Mol. Cell.* 65: 447-459.
- [42] McKnight, J.N., T. Tsukiyama, and G. D. Bowman. 2016. Sequence-targeted nucleosome sliding in vivo by a hybrid Chd1 chromatin remodeler. *Genome Res.* 26: 693-704.
- [43] Winger, J., and G. D. Bowman. 2017. The sequence of nucleosomal DNA modulates sliding by the Chd1 remodeler. *J. Mol. Bio.* 429: 808-822.
- [44] Gkikopoulos, T., P. Schofield, ..., T. Owen-Hughes. 2017. A role for Snf2-related nucleosome-spacing enzymes in genome-wide nucleosome organization. *Science* 333, 1758-1760.
- [45] Sundaramoorthy, R., A. L. Hughes, ..., T. Owen-Hughes. 2017. Structural reorganization of the chromatin remodeling enzyme Chd1 upon engagement with nucleosomes. *eLife* 6: e22510.
- [46] Farnung, L., S. M. Vos, ..., P. Cramer. 2017. Nucleosome-CHD1 structure and implications for chromatin remodeling. *Nature* 550: 539-542.
- [47] Sinha, K. K., J. D. Gross, and G. J. Narlikar. 2017. Distortion of histone octamer core promotes nucleosome mobilization by a chromatin remodeler. *Science* 355: eaaa3761.
- [48] Flaus, A., and T. Owen-Hughes. 2017. Unlocking the nucleosome. *Nature* 355: 245-246.

- [49] Isaac, R. S., F. Jiang, ..., R. Almeida. 2016. Nucleosome breathing and remodeling constrain CRISPR-Cas9 function. *eLife* 5:e13450.
- [50] Polach, K. J. and J. Widom. 1995. Mechanism of protein access to specific DNA sequences in chromatin: a dynamic equilibrium model for gene regulation. *J. Mol. Biol.* 254:130-149.
- [51] Anderson, J. D. and J. Widom. 1995. Sequence and position-dependence of the equilibrium accessibility of nucleosomal DNA target sites. *J. Mol. Biol.* 296:979-987.
- [52] Buning, R. and J. van Noort. 2010. Single-pair FRET experiments on nucleosome conformational dynamics. *Biochimie* 92:1729-1740.
- [53] Gansen, A., A. R. Hieb, ..., J. Langowski. 2013. Closing the gap between single molecule and bulk FRET analysis of nucleosomes. *PLoS ONE* 8:e57018.
- [54] Culkin, J., L. de Bruin, M. Tompitak, R. Phillips and H. Schiessel. 2017. The role of DNA sequence in nucleosome breathing. *Eur. Phys. J. E* 40:106.
- [55] Lai, W. K. M., and B.F. Pugh. 2017. Understanding nucleosome dynamics and their links to gene expression and DNA replication. *Nature Rev. Mol. Cell. Biol.* 18:548-562.