



HAL
open science

Bottom-up effects on biomechanical properties of the skeletal plates of the sea urchin *Paracentrotus lividus* (Lamarck, 1816) in an acidified ocean scenario

Valentina Asnaghi, Marie Collard, Luisa Mangialajo, Jean-Pierre Gattuso, Philippe Dubois

► To cite this version:

Valentina Asnaghi, Marie Collard, Luisa Mangialajo, Jean-Pierre Gattuso, Philippe Dubois. Bottom-up effects on biomechanical properties of the skeletal plates of the sea urchin *Paracentrotus lividus* (Lamarck, 1816) in an acidified ocean scenario. *Marine Environmental Research*, 2019, 144, pp.56-61. 10.1016/j.marenvres.2018.12.002 . hal-02345719

HAL Id: hal-02345719

<https://hal.science/hal-02345719>

Submitted on 18 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Title: Bottom-up effects on biomechanical properties of the skeletal plates of the sea**
2 **urchin *Paracentrotus lividus* (Lamarck, 1816) in an acidified ocean scenario**

3 Valentina Asnaghi ^{1*}, Marie Collard ², Luisa Mangialajo ³, Jean-Pierre Gattuso^{4,5}, Philippe Dubois²

4
5 ¹ Department of Earth, Environment and Life Science, DiSTAV, University of Genoa, Italy

6 ² Laboratoire de Biologie marine CP160/15, Université Libre de Bruxelles, av F.D. Roosevelt, 50, B-1050 Bruxelles,
7 Belgium

8 ³ Université de Nice-Sophia Antipolis, EA 4228 ECOMERS, Nice, France,

9 ⁴ Sorbonne Université, CNRS, Laboratoire d'Océanographie de Villefranche, 181 chemin du Lazaret, F 06230,
10 Villefranche-sur-mer, France

11 ⁵ Institute for Sustainable Development and International Relations, Sciences Po, 27 rue Saint Guillaume, F75007, Paris,
12 France

13 *corresponding author: valentina.asnaghi@unige.it

14

15

16

17

18

19

20 **Keywords**

21 Ocean Acidification; Temperate Reefs; Sea Urchins; Macroalgae; *Paracentrotus lividus*; Trophic Cascade

22

23

24

25

26 **Authors contribution**

27 VA, LM, JPG, PD conceived of or designed study; VA, MC performed research; VA analyzed data, VA, MC,

28 PD wrote the paper.

29 **Abstract**

30 Sea urchins, ecologically important herbivores of shallow subtidal temperate reefs, are considered particularly
31 threatened in a future ocean acidification scenario, since their carbonate structures (skeleton and grazing
32 apparatus) are made up of the very soluble high-magnesium calcite, particularly sensitive to a decrease in pH.
33 The biomechanical properties of their skeletal structures are of great importance for their individual fitness,
34 because the skeleton provides the means for locomotion, grazing and protection from predators. Sea urchin
35 skeleton is composed of discrete calcite plates attached to each other at sutures by organic ligaments. The
36 present study addressed the fate of the sea urchin *Paracentrotus lividus* (Lamarck, 1816) skeleton in acidified
37 oceans, taking into account the combined effect of reduced pH and macroalgal diet, with potential cascading
38 consequences at the ecosystem level. A breaking test on individual plates of juvenile specimens fed different
39 macroalgal diets has been performed, teasing apart plate strength and stiffness from general robustness,.
40 Results showed no direct short-term effect of a decrease in seawater pH nor of the macroalgal diet on single
41 plate mechanical properties. Nevertheless, results from apical plates, the ones presumably formed during the
42 experimental period, provided an indication of a possible diet-mediated response, with sea urchins fed the
43 more calcified macroalga sustaining higher forces before breakage than the one fed the non-calcified algae.
44 This supports the need of longer term experiments to observe substantial differences on skeletal plate structure.

45

46

47

48 **Introduction**

49 Sea urchins are important calcifiers in shallow subtidal areas of temperate regions and play a key
50 ecological role in these ecosystems being generally the most effective benthic herbivores and
51 controlling, through their grazing activity, the dynamic, structure and composition of macroalgal
52 assemblages (Jangoux and Lawrence 1982; Ruitton et al., 2000; Bulleri et al., 2002; Privitera et al.,
53 2008; Bonaviri et al., 2011). Their skeleton, spines and grazing apparatus are made of high-
54 magnesium calcite, a form of calcium carbonate that is particularly vulnerable to dissolution under
55 low pH conditions (Andersson et al., 2008; Hermans et al., 2010). For this reason, sea urchins have
56 long been regarded as particularly threatened by the ongoing decrease of pH and calcium carbonate
57 saturation states of the oceans, referred to as ocean acidification (Kurihara and Shirayama 2004;
58 Dupont et al., 2010; Byrne et al., 2011).

59 Echinoid skeleton is made up of discrete ossicles located in the dermis. Each ossicle consists of a
60 three-dimensional network of mineralized trabeculae, the stereom, delimiting an internal and
61 complementary network filled by connective tissue, the stroma (Dubois and Chen 1989). The
62 perforated calcite plates are attached to each other at sutures by ligaments that wrap around calcite
63 rods, thus sewing together adjacent plates. Trabeculae project from one plate into holes in the adjacent
64 plates, thus interlocking the plates (Moss and Meehan 1967). These processes ensure a relative
65 rigidity of the test. The stereom consists of high-magnesium calcite and of 0.1% (w/w) organic
66 material (the intrastereomic organic matrix; e.g. Weiner 1985). Sutural ligaments among plates
67 strengthen sea urchin skeleton (Kidwell and Baumiller 1990) and, on the basis of histological and
68 morphological evidence, these ligaments may be interpreted as “stress-breakers” that evenly
69 distribute stresses and thus contribute to the structural integrity of echinoid skeletons (Moss and
70 Meehan 1967). The strengthening role of sutural ligaments is different according to size, age, diet
71 and *taxa* (Ellers et al., 1998). Sutural ligaments are known to reinforce urchin skeletons under natural
72 loads such as the action of crab claws, apical or lateral forces from waves and forces generated when
73 an urchin wedges itself in a crack (Ellers et al., 1998).

74 The biomechanical properties of skeletal structures have a great importance for individual fitness
75 (Currey 1989; Meyers et al., 2008) because skeletons provide the means for locomotion, grazing and
76 protection from predators. Few studies have investigated the skeletal biomechanical properties of
77 echinoderms with respect to ocean acidification (reviewed in Dubois 2014; Collard et al., 2016; Dery
78 et al., 2017). Sea urchin spines and test plates are differently affected by seawater acidification.
79 Spines, more vulnerable, showed reduced fracture forces at reduced pH also in short term studies
80 (Holtmann et al., 2013; Dery et al., 2017; Emerson et al., 2017) while plates were not impacted by
81 acidified conditions in short and long term experiments (Holtmann et al., 2013; Moulin et al., 2015;
82 Collard et al., 2016).

83 Only few studies focused on the mechanical resistance of the whole test under low pH conditions.
84 Byrne et al. (2014) reported a reduced crushing force in live *Tripneustes gratilla* juveniles grown
85 from metamorphosis at pH_{NBS} 7.6 (0.5 pH units below control) but this was attributed to differences
86 in urchin size. A decrease in test robustness, also related to test size, was observed on *Paracentrotus*
87 *lividus* and *Diadema africanum* juveniles kept under reduced pH (7.6 pH_{NBS}) for 100 days, compared
88 to control conditions (8.0 pH_{NBS} ; Rodriguez et al., 2017). Similarly, in juvenile *Paracentrotus lividus*
89 maintained for one month at pH_{T} 7.7 (0.4 pH_{T} units below control), the test was less robust (in terms
90 of resistance to an increasing crushing load, tested on dried skeletons) than at higher pH_{T} (7.84, 7.89,
91 8.09; Asnaghi et al., 2013). These differences in test robustness were mirrored by diet-related
92 differences (calcified vs. non-calcified macroalgae) in skeletal composition (particularly Mg/Ca ratio;
93 Asnaghi et al., 2013, 2014), suggesting that diet is another potentially relevant source for bicarbonate
94 uptake (in addition to possible uptake from the seawater). In Asnaghi et al. (2013), the crushing test
95 was performed on entire dry preserved juvenile sea urchins, where organic material and ligaments
96 were still present. This dried organic material is flexurally stiff relative to the calcite plates and might
97 provide tensile reinforcement to the skeleton (Ellers et al., 1998), even if performing the test on dried
98 ligaments could have resulted in biased resistance compared to fresh organisms (Ellers et al., 1998).

99 In the present study, the fate of sea urchin skeletons in acidified oceans has been addressed, teasing
100 apart plate strength and stiffness from general robustness of the test, due to calcium carbonate plates
101 and organic material associated. We performed a breaking test, using a motorized load frame (Instron
102 5543 tensile tester), on individual plates detached from the skeleton and cleaned of organic material
103 in juvenile sea urchins fed different diets. The individuals were part of the experiment presented in
104 Asnaghi et al. (2013, 2014). The individuals tested in this study were not previously tested for whole
105 test strength to avoid any confounding factors such as cracks or breaks.

106 Juvenile sea urchins were treated for one month under four pH levels and fed one of three species of
107 macroalgae with variable carbonate content (i.e. calcified vs. not-calcified). Since sea urchin
108 skeletons grow both by the accretion of calcite at the edges and faces of the plates and by the addition
109 of new plates at the apex resulting in gradual migration of initially apical plates adorally during
110 growth (Deutler 1926; Märkel 1975), we can assume that in our juvenile sea urchins the largest part
111 of the most apical plates were formed during the experimental period, while ambital ones were
112 already formed before the experiment (as proposed by Collard et al., 2015). Consequently, we
113 expected different responses to lowered pH, in terms of individual plate strength and stiffness, for
114 plates under formation (apical) and for already formed ones (ambital), and according to the different
115 macroalgal diets.

116

117 **Materials and methods**

118 **Experimental set-up**

119 The experimental set-up was thoroughly described in Asnaghi et al. (2013). Briefly, a total of 144 4-
120 month old juveniles of *Paracentrotus lividus* (Lamarck, 1816), provided by a sea urchin hatchery in
121 Camogli (NW Mediterranean Sea, Italy), with an average test diameter of 5.8 mm (\pm 0.1 standard
122 error, SE), were randomly selected and transferred to the Laboratoire d'Océanographie de
123 Villefranche (NW Mediterranean Sea, France) where the experiment was performed in July 2011.

124 Four pH_T levels, corresponding to future pCO_2 conditions chosen according to best practices (Barry
125 et al., 2010) and three scenarios were used: (1) present day, $pCO_2 = 390 \mu atm$ ($pH_T \approx 8.1$, control);
126 (2) optimistic scenario (SRES scenario B1), $pCO_2 = 550 \mu atm$ ($pH_T \approx 8.0$); (3) realistic scenario
127 (midway between SRES scenario AB1 and A2), $pCO_2 = 750 \mu atm$ ($pH_T \approx 7.8$) and (4) pessimistic
128 scenario (A1F1), $pCO_2 = 1000 \mu atm$ ($pH_T \approx 7.7$). pCO_2 was controlled by bubbling pure- CO_2 ,
129 according to best practices for ocean acidification experiments (Riebesell et al., 2010).

130 Unfiltered seawater, pumped from a depth of 10 m, was continuously supplied to four 200 l header
131 tanks and bubbled with air. pH was continuously monitored by a pH-stat system (IKS, Karlsbad,
132 Aquastar) and small amounts of pure CO_2 were added to keep pH at the desired level. Manipulated
133 seawater from the four header tanks was delivered to experimental units at a rate of about $6 l h^{-1}$ in
134 an open system.

135 The experiment was performed within a thermostatic chamber kept at $22^\circ C$. Irradiance values in the
136 aquaria were maintained at about $215 \mu mol photons m^{-2} s^{-1}$, with a 12:12 h L:D photoperiod. Juvenile
137 sea urchins were fed three different macroalgal species: a calcified species, *Ellisolandia elongata*
138 (J.Ellis & Solander) K.R.Hind & G.W.Saunders, 2013 (previously known as *Corallina elongata*) and
139 two non-calcified species *Dictyota dichotoma* (Hudson) J.V.Lamouroux, 1809 and *Cystoseira*
140 *amentacea* (C.Agardh) Bory de Saint-Vincent, 1832. Macroalgae were collected prior to the start of
141 the experiment and kept for at least one week in distinct aquaria at the same pH levels as the sea
142 urchins. Feeding was *ad libitum* with macroalgae from the corresponding pH level.

143 Six juvenile sea urchins were placed in each of the 24 experimental units (2 independent replicates
144 for each combination of pH and diet - see details of experimental set-up and sea water parameters in
145 Asnaghi et al., 2013). The experiment lasted one month. At the end of the experiment, all specimens
146 were measured, air-dried and stored pending future analyses.

147 **Mechanical test**

148 For the mechanical test, a subset of 24 sea urchins (one from each experimental unit, representing an
149 independent replicate) was selected. First of all, a cleaning protocol was set up in order to remove the

150 soft tissues surrounding the skeleton without damaging the stereom and allow separation of the plates.

151 Different exposure times were tested in order to select the best cleaning protocol and effectiveness of

152 the different methods was checked under a scanning electron microscope (SEM). The best procedure

153 for cleaning plates without damaging their structure consisted in placing entire dried urchins in 2.5%

154 (v:v) sodium hypochlorite for 1 h to detach the spines and Aristotle's lantern, rinsing 3 times in MilliQ

155 water and air drying for 1 h. Tests devoid of spines and lantern were further cleaned for 30 min in

156 1% (v:v) sodium hypochlorite, rinsed 3 times in MilliQ water and air-dried overnight.

157 Single interambulacral plates were separated under a stereo-microscope; 3 apical (presumably formed

158 during the experimental period) and 3 ambital plates (already formed before the beginning of the

159 experiment) for each specimen were used for biomechanical measurements (Figure 1a).

160 Top and lateral view photographs of each sea urchin plate were obtained with a Nikon Coolpix 995

161 3Mpi under a binocular lens. Photographs were analyzed, using the software ImageJ, to measure

162 tubercle areas (A), which is the area over which the force was actually applied (Figure 1b) and

163 effective length (L_e), which is the height to the edge of the tubercles (Figure 1c). These parameters

164 are necessary for Strain and Stress calculations (see equations below).

165 Due to the small size and flatness of both apical and ambital plates of juvenile sea urchins, an *ad hoc*

166 system, halfway between compression test and three-point bending test, was created to measure the

167 fracture force. Individual cleaned and detached sea urchin plates were placed on a metal stand with a

168 groove in the middle and the mechanical test was carried out using a second metal block fixed on the

169 load frame of the force stand, Instron 5543 tensile tester, at a speed of 0.1 mm min^{-1} , applying constant

170 compression till breakage (Figure 1d). Displacement and force were recorded at a frequency of 10

171 Hz. For each plate, the Bluehill software (Instron) provided information about force at fracture (F_{\max})

172 and displacement (ΔL).

173 Stiffness was measured through Young's modulus (E), a quantity used to characterize materials and

174 defined as the ratio of the stress along an axis over the strain along that same axis.

175

176 Strain and Stress were calculated for each plate, using the following equations:

177 Strain: $\sigma = F/A$

178 Stress: $\epsilon = \Delta L/L_e$


179 where F: force at fracture, A: tubercle area, ΔL : displacement, L_e : effective length.

180 The Young's modulus (E) was calculated as the slope between two points of the final linear part of
181 the curve, in this case the maximum force and the 100th point before that, using the following
182 formula:

183
$$E = \frac{\sigma_{max} - \sigma_{100th\ point}}{\epsilon_{max} - \epsilon_{100th\ point}} \text{ (Pa)}$$

184

185



186

187 Figure 1: Schematic representation of a) interambulacral plates, apical and ambital, and suture geometry with
188 corresponding scale bar at the bottom (modified from Ellers et al., 1998; b) upper and c) lateral view of one plate, in order
189 to perform tubercle area (A) and effective length (L_e) measurements, with corresponding scale bar at the bottom; d) simple
190 *ad hoc* compression device composed by two metal blocks, the lower, where the plate should be placed, with a groove
191 and the upper fixed on the load frame (modified from Collard et al., 2016)

192

193

194 **Data analyses**

195 In order to assess the effect of pH level, algal diet and their interaction on apical and ambital plates
196 strain (F_{\max}/A) and stiffness (Young's modulus, square root transformed data), a crossed ANOVA
197 design was applied, using 'pH' and 'diet' as fixed crossed factors. Since multiple observations were
198 performed on each specimen, a random effect *specimen* nested in the interaction (pH * diet) was
199 added in order to account for the dependency structure in the data.

200 Normality and homogeneity of variance were verified for both the considered response variables
201 (F_{\max}/A and \sqrt{E}) using Shapiro-Wilk test and Bartlett tests, respectively. All statistical analyses were
202 performed using the software R (R Core Team 2014).

203

204 **Results**

205 No significant effect of the interaction between the two fixed factors, pH and diet, was observed in
206 strain (fracture force/surface on which the force is applied) or in stiffness neither for ambital nor for
207 apical plates. Moreover, no significant difference according to factors seawater pH or diet singularly
208 were detected in ambital plates (Tables 1, 2). Similarly, no effect of seawater pH or diet was
209 evidenced in apical plates strain and stiffness (Tables 1, 2).

210 Both strain and stiffness of ambital plates (Fig. 2 a, c) showed more homogeneous values among
211 treatments compared to apical ones (Fig. 2 b, d). Apical plates stiffness slightly decreased according
212 to the pH decrease for *E. elongata* and *C. amentacea*, but not in *D. dichotoma*, where the trend seems
213 to be the opposite (Fig. 2b), even if differences are not statistically significant. A weak gradient of
214 apical plates strain values according to the algal diet can be observed, *i.e.* gradual decrease of the
215 strain value for sea urchins fed *E. elongata*, *C. amentacea* and *D. dichotoma*, respectively (Figure
216 2d).

217

218 Table 1: ANOVA results table for strain data (F_{\max}/A) of ambital and apical plates under factors pH and Diet and their
 219 interaction. The random effect *specimen* nested in the interaction (pH x* diet) is considered in order to account for the
 220 dependency structure in the data


	AMBITAL PLATES					APICAL PLATES				
	Df	SS	MS	F	signif.	Df	SS	MS	F	signif.
pH	3	2.8	0.95	0.01	0.998	3	271	90.2	0.16	0.918
Diet	2	51.1	25.55	0.38	0.694	2	3798	1898.9	3.47	0.065
pH * diet	6	641	106.84	1.58	0.236	6	529	88.1	0.16	0.983
<i>specimen</i> (pH * diet)	12	776.5	67.71	2.91	0.004	12	6567	547.3	3.32	0.001
Residuals	48	1116	23.24			48	7906	164.7		

221
 222
 223 Table 2: ANOVA results table on square root transformed Young's modulus data (\sqrt{E}) of ambital and apical plates under
 224 factors pH and Diet and their interaction. The random effect *specimen* nested in the interaction (pH x* diet) is considered
 225 in order to account for the dependency structure in the data

	AMBITAL PLATES					APICAL PLATES				
	df	SS	MS	F	signif.	Df	SS	MS	F	signif.
pH	3	0.22	0.07	0.88	0.481	3	0.23	0.08	0.61	0.618
Diet	2	0.03	0.01	0.13	0.884	2	0.38	0.19	1.46	0.270
pH * diet	6	0.30	0.05	0.63	0.708	6	1.74	0.29	2.23	0.112
<i>specimen</i> (pH * diet)	12	0.94	0.08	2.00	0.045	12	1.52	0.13	2.20	0.029
Residuals	48	2.09	0.04			48	3.11	0.06		

226

227



228

229 Figure 2: Barplots of Young's modulus (E), upper panels, and strain data (F_{max}/A), lower panels, measured for ambital (a,
 230 c) and apical (b, d) plates under different sea water pH (x axis) and different macroalgal diets (light gray= *E. elongata*,
 231 gray= *C. amentacea*, dark gray= *D. dichotoma*). Error bars: standard error.

232

233 Discussion

234 Ocean acidification caused by anthropogenic carbon dioxide emissions is known to pose major threats
 235 for marine organisms, particularly calcifying ones (e.g. Kroeker et al., 2010; Riebesell et al., 2010),
 236 since their calcium carbonate structures are potentially susceptible to dissolution in acidic waters (Orr
 237 et al. 2005). Sea urchins are important marine calcifiers, playing a relevant ecological role in
 238 temperate ecosystems (Jangoux and Lawrence 1982; Ruitton et al., 2000; Bulleri et al., 2002; Privitera
 239 et al., 2008; Bonaviri et al., 2011). Among calcifying organisms, highly calcified sea urchins are
 240 expected to be the echinoderms more affected by reduced pH, with differences at species, population
 241 and stage level (reviewed in Dupon et al., 2010).

242 Early life stages are acknowledged to be the most sensitive to ocean acidification (Dupont et al.,
 243 2010; Moulin et al., 2011; Byrne et al., 2013; Stump et al., 2013), but also post-metamorphic phases
 244 (juveniles and adults) showed to be affected, mainly in terms of survival and growth, both from
 245 laboratory experiments (e.g. Shirayama and Thornton 2005; Byrne et al., 2011; Albright et al., 2012;

246 Asnaghi et al., 2013, 2014; Moulin et al., 2015; Collard et al., 2016), and from *in situ* records in
247 naturally acidified areas (only on adults, e.g. Hall-Spencer et al., 2008; Bray et al., 2014; Collard et
248 al., 2016). Moreover, different studies investigated coelomic fluid regulatory capacities in adult sea
249 urchins exposed to hypercapnic conditions, reporting contrasting results about their ability to partially
250 or fully compensate extracellular pH (Stumpp et al., 2012; Collard et al., 2013, 2014; Calosi et al.,
251 2013; Kurihara et al., 2013; Moulin et al., 2014, 2015) and prevent skeletal dissolution at low pH
252 (reviewed by Dery et al., 2017).

253 Only few studies addressed the combined effect of reduced pH and macroalgal diet on sea urchins,
254 highlighting potential cascading consequences at the ecosystem level (Johnson and Carpenter, 2012;
255 Asnaghi et al., 2013; 2014). Macroalgae exhibit a broad range of responses to ocean acidification
256 (Hall-Spencer et al., 2008; Nelson, 2009; Martin and Gattuso, 2009; Connell and Russell, 2010;
257 Cornwall et al. 2011, 2017; Poore et al., 2013), mainly linked to their calcium carbonate content
258 (Cornwall et al., 2014; James et al, 2014) and inorganic carbon physiology (Cornwall et al., 2017).
259 The loss of calcium carbonate and tissue modification in macroalgal thalli caused by reduced pH may
260 enhance their palatability to grazers (Duarte et al., 2016; Rich et al., 2018, Rodriguez et al., 2018),
261 leading to potential shifts in the ecosystem equilibrium under an acidified scenario.

262 Different algal feedings, quantity and species of macroalgae available as food are known to directly
263 affect somatic and gonadal production in sea urchins, both from field and laboratory studies (e.g.
264 Ebert 1968; Lawrence 1975; Lilly 1975; Vadas 1977; Larson et al., 1980; Lawrence and Lane 1982;
265 Thompson 1983, 1984; Privitera et al., 2008). Food shortage (i.e. the lack of macroalgal biomass for
266 feeding) have been reported to cause modifications in plastic resource allocation (Ebert 1980; Haag
267 et al., 2016), differences in mechanical properties of sea urchin spines (Moureaux and Dubois 2012)
268 and, in an acidified scenario, reduction of the buffer capacity of the coelomic fluid to compensate the
269 decrease of external pH (Collard et al., 2013).

270 In the present study, the combined effect of pH level and macroalgal diet was investigated in terms
271 of sea urchin plates biomechanical properties, showing no direct short-term effect of seawater pH nor

272 diet on single plate strain and stiffness, in agreement with previous laboratory studies (Holtmann et
273 al., 2013; Moulin et al., 2015; Collard et al., 2016).

274 A field study, instead, highlighted a role of the diet in mediating sea urchins biomechanical properties,
275 showing that test plates from *P. lividus* living in tide pools mainly covered by encrusting calcareous
276 algae exhibit a higher fracture force than test plates of sea urchins living in pools containing erected
277 non-calcifying algae (Collard et al., 2016). Similar role of calcifying macroalgae in strengthen *P.*
278 *lividus* juvenile tests has been shown by Asnaghi et al. (2013) on specimens exposed to the same
279 experimental conditions of the present study, but it is not easy to disentangle the role of plates,
280 ligaments and calcified locking structures in providing test robustness.

281 Results from the present study showed more homogeneous responses of the ambital plates (the ones
282 already formed before the start of the experiment) in terms of strain and stiffness (Fig. 2 a, c), while
283 apical plates, the one presumably formed during the one-month experimental period, showed more
284 variable values, with strain data suggesting a possible diet-mediated response, maybe visible on a
285 longer term (Fig. 2 b, d): sea urchins fed the more calcified macroalga (*E. elongata*) sustained higher
286 fracture force than the one fed the non-calcified algae (*C. amentacea* and *D. dichotoma*). The high
287 variability at the level of the specimen (Table 1 and 2) may have masked patterns on the short term.

288 Mechanical properties can be affected by changes in the growth rate (e.g. Moureaux et al., 2010),
289 which may affect the three-dimensional morphology or density of the stereom (Smith 1980).
290 Alternatively, the structural properties of the material itself may be affected by the formation of
291 imperfections during the CaCO₃ precipitation (e.g. Moureaux et al., 2011).

292 Sea urchins from the present experiment showed different growth rates according to their diet.
293 *Ellisolandia elongata* allowed for a faster growth rate of the sea urchins compared to that of the ones
294 fed *Cystoseira amentacea* and *Dictyota dichotoma*. (Asnaghi et al., 2014). This could be linked to the
295 supply of calcium by calcified algae (Powell et al., 2010) rather than to the carbonate ions, which are
296 not transported through cell membranes, or bicarbonate ions which are available in high concentration
297 in seawater from which they are readily absorbed (from this source; Collard et al., 2014).

298 In the present study, the choice to use juveniles specimens, characterized by higher growth rates
299 compared to adults, was driven by the possibility to observe potential modifications linked to pH and
300 diet treatments on a short time scales (one month). Indeed, a diet-mediated decrease in growth and
301 test robustness in these *P. lividus* juveniles under acidified conditions was proven (Asnaghi et al.,
302 2013), even if not mirrored by substantial modification of test thickness and single plates
303 biomechanical properties. Rodriguez et al. (2017) reported significantly thinner test plates in juvenile
304 *Paracentrotus lividus* and *Diadema africanum* kept for 100 days in one tank at low pH conditions
305 compared to the ones in the control tank, that may have led to less robust test, even if several other
306 sources of variability were present in the experimental design.

307 Those considerations suggest that, in order to observe substantial differences on skeletal plate
308 structure, it is necessary to perform longer term experiments (Dupont et al., 2010; Hendricks et al.,
309 2010), taking into particular account feeding conditions, that are frequently neglected in ocean
310 acidification studies on post-metamorphic individuals (Dubois et al., 2014).

311 The impact of algal diet on sea urchin test resistance to breakage is particularly relevant in the context
312 of global change. Coralline algae are renowned impacted by ocean acidification, while non-calcified
313 algae will be most likely favored in future oceans (Hall-Spencer et al., 2008; Porzio et al., 2011; Koch
314 et al., 2013; Sunday et al., 2017). As a consequence, some sea urchin species might be affected by
315 the expected change in macroalgal diet availability, growing slower and producing a less robust
316 skeleton, with potential relevant cascading consequences on their ability to graze, move, search for
317 shelters and defend from predation and hydrodynamics.

318

319

320

321

322

323

324 **Acknowledgements**

325 Philippe Dubois is a Research Director of the National Fund for Scientific Research (NFSR, Belgium), Marie
326 Collard was holder of a FRIA grant (Belgium). Valentina Asnaghi scientific stay in France was funded by the
327 “Bando VINCI 2010” of the Italian-French University and in Belgium by a grant of the National Fund for
328 Scientific Research (NFSR, Belgium). Work supported by NFSR project 26112046 and J.0219.16 SOFTECHI.
329 This work is a contribution to the “European Project on Ocean Acidification” (EPOCA) funded by European
330 Community’s Seventh Framework Programme (FP7/2007-2013; grant agreement n 211384). We thank Saloua
331 M’Zoudi for technical help, Marco Asnaghi for helping in plate measurements and Mariachiara Chiantore for
332 the scientific support. We thank Andy Davis and the other two anonymous reviewers for their precious
333 comments.

334

335

336 **References**

- 337 1. Albright R, Bland C, Gillette P, Serafy JE, Langdon C, Capo TR 2012. Juvenile growth of the
338 tropical sea urchin *Lytechinus variegatus* exposed to near-future ocean acidification scenarios.
339 Journal of Experimental Marine Biology and Ecology 426–427: 12–17.
- 340 2. Andersson AJ, Mackenzie FT, Bates NB, 2008. Life on the margin: implications of ocean
341 acidification on Mg-calcite, high latitude and cold-water marine calcifiers. Marine Ecology
342 Progress Series 373: 265-273.
- 343 3. Asnaghi V, Chiantore M, Mangialajo L, Gazeau F, Francour P, Alliouane S, Gattuso J-P 2013.
344 Cascading effects of ocean acidification in a rocky subtidal community. PLOS ONE 8 (4):
345 e61978.
- 346 4. Asnaghi V, Mangialajo L, Gattuso JP, Francour P, Privitera D, Chiantore M 2014. Effects of
347 ocean acidification and diet on thickness and carbonate elemental composition of the test of
348 juvenile sea urchins. Marine Environmental Research 93: 78-84.
- 349 5. Barry JP, Tyrrell T, Hansson L, Gattuso J-P 2010. CO₂ targets for ocean acidification
350 perturbation experiments. Riebesell U, Fabry VJ, Hansson L, Gattuso J-P, editors. Guide to

- 351 best practices for ocean acidification research and data reporting. Luxembourg: Publications
352 Office of the European Union, p53-66.
- 353 6. Bonaviri C, Vega Fernández T, Fanelli G, Badalamenti F, Gianguzza P 2011. Leading role of
354 the sea urchin *Arbacia lixula* in maintaining the barren state in southwestern Mediterranean.
355 Marine Biology 158: 2505–2513.
- 356 7. Bray L, Pancucci-Papadopoulou MA, Hall-Spencer JM 2014. Sea urchin response to rising
357 pCO₂ shows ocean acidification may fundamentally alter the chemistry of marine skeletons.
358 Mediterranean Marine Science 15(3): 510–519.
- 359 8. Bulleri F, Bertocci I, Micheli F 2002. Interplay of encrusting coralline algae and sea urchins
360 in maintaining alternative habitats. Marine Ecology Progress Series 243:101–109.
- 361 9. Byrne M, Ho M, Wong E, Soars NA, Selvakumaraswamy P, Shepard-Brennan H,
362 Dworjanyn SA, Davis AR 2011. Unshelled abalone and corrupted urchins: Development of
363 marine calcifiers in a Changing Ocean. Proceedings of the Royal Society B-BiolSci
364 278(1716): 2376-2383.
- 365 10. Byrne M, Lamare M, Winter D, Dworjanyn SA, Uthicke S 2013. The stunting effect of a high
366 CO₂ ocean on calcification and development in sea urchin larvae, a synthesis from the tropics
367 to the poles. Philosophical Transaction of the Royal Society B 368: 20120439.
- 368 11. Byrne M, Smith AM, West S, Collard M, Dubois P, Graba-landry A, Dworjanyn SA 2014.
369 Warming influences Mg²⁺ content, while warming and acidification influence calcification
370 and test strength of a sea urchin. Environmental science & technology 48(21): 12620-12627.
- 371 12. Calosi P, Rastrick SPS, Graziano M, Thomas SC, Baggini C, Carter HA, Hall-Spencer JM,
372 Milazzo M, Spicer JI 2013. Distribution of sea urchins living near shallow water CO₂ vents
373 is dependent upon species acid–base and ion-regulatory abilities. Marine Pollution Bulletin
374 73: 470–484.

- 375 13. Collard M, Laitat K, Moulin L, Catarino A, Grosjean P, Dubois P 2013. Buffer capacity of
376 the coelomic fluid in echinoderms. *Comparative Biochemistry and Physiology Part A*, 166:
377 199 – 206.
- 378 14. Collard M, Dery A, Dehairs F, Dubois P. 2014. Euechinoidea and Cidaroidea respond
379 differently to ocean acidification. *Comparative Biochemistry and Physiology Part A*, 174: 45–
380 55.
- 381 15. Collard M, Rastrick SPS, Calosi P, Demolder Y, Dille J, Findlay HS, Hall-Spencer JM,
382 Milazzo M, Moulin L, Widdicombe S, Dehairs F, Dubois P 2016. The impact of ocean
383 acidification and warming on the skeletal mechanical properties of the sea urchin
384 *Paracentrotus lividus* from laboratory and field observations. *ICES Journal of Marine Science*
385 73(3): 727-738.
- 386 16. Connell SD, Russell BD, 2010. The direct effects of increasing CO₂ and temperature on non-
387 calcifying organisms: increasing the potential for phase shifts in kelp forests. *Proceedings of*
388 *the Royal Society of London B: Biological Sciences*, rspb20092069.
- 389 17. Cornwall CE, Hepburn CD, Pritchard D, Currie KI, McGraw CM, Hunter KA, Hurd CL,
390 2011. Carbon-use strategies in macroalgae: differential responses to lowered pH and
391 implications for ocean acidification. *J. Phycol.* 48: 137–144
- 392 18. Cornwall CE, Boyd PW, McGraw CM, Hepburn CD, Pilditch CA, Morris JN, ... Hurd CL,
393 2014. Diffusion boundary layers ameliorate the negative effects of ocean acidification on the
394 temperate coralline macroalga *Arthrocardia corymbosa*. *PLoS. One* 9, e97235.
- 395 19. Cornwall CE, Revill AT, Hall-Spencer JM, Milazzo M, Raven JA, Hurd CL, 2017.
396 Inorganic carbon physiology underpins macroalgal responses to elevated CO₂.
397 *Scientific reports*, 7, 46297
- 398 20. Currey JD 1989. Biomechanics of mineralized skeletons. *Short Courses in Geology* 5: 11-25

- 399 21. Dery A, Collard M, Dubois P 2017. Ocean acidification reduces spine mechanical strength in
400 euechinoid but not in cidaroid sea urchins. *Environmental Science & Technology* 51: 3640-
401 3648.
- 402 22. Deutler F 1926. Über die Wachstum des Seeigelskeletts. *Zoologische Jahrbücher. Abteilung*
403 für Anatomie und Ontogenie der Tiere Abteilung für Anatomie und Ontogenie der Tiere 48:
404 119-200.
- 405 23. Duarte C, López J, Benítez S, Manríquez PH, Navarro JM, Bonta CC, Torres R, Quijón P,
406 2016. Ocean acidification induces changes in algal palatability and herbivore feeding behavior
407 and performance. *Oecologia* 180: 453-462.
- 408 24. Dubois P, Chen CP 1989. Calcification in echinoderms. Jangoux M, Lawrence JM, editors.
409 Balkema, Rotterdam, The Netherlands. *Echinoderm Studies* 3, p55–136.
- 410 25. Dubois P 2014. The skeleton of postmetamorphic echinoderms in a changing world.
411 *Biological Bulletin* 226: 223–236.
- 412 26. Dupont S, Ortega-Martínez O, Thorndyke MC 2010. Impact of near-future ocean acidification
413 on echinoderms. *Ecotoxicology* 19: 449–462.
- 414 27. Ebert TA 1968. Growth rates of the sea urchin *Strongylocentrotus purpuratus* related to food
415 availability and spine abrasion. *Ecology* 49: 1075:1091.
- 416 28. Ebert TA 1980. Relative growth of sea urchin jaws: an example of plastic resource allocation.
417 *Bulletin of Marine Science* 30: 467–474.
- 418 29. Ellers O, Johnson AS, Moberg PE 1998. Structural strengthening of urchin skeletons by
419 collagenous sutural ligaments. *Biological Bulletin* 195: 136- 144.
- 420 30. Emerson CE, Reinardy HC, Bates NR, Bodnar AG 2017. Ocean acidification impacts spine
421 integrity but not regenerative capacity of spines and tube feet in adult sea urchins. *Royal*
422 *Society open science* 4(5): 170140.

- 423 31. Haag N, Russell MP, Hernandez JC 2016. Effects of spine damage and microhabitat on
424 resource allocation of the purple sea urchin *Strongylocentrotus purpuratus* (Stimpson 1857).
425 Journal of Experimental Marine Biology and Ecology 482: 106-117.
- 426 32. Hall-Spencer JM, Rodolfo-Metalpa R, Martin S, Ransome E, Fine M, Turner SM, Rowley SJ,
427 Tedesco D, Buia M-C 2008. Volcanic carbon dioxide vents show ecosystem effects of ocean
428 acidification. Nature 454(7200) : 96-99.
- 429 33. Hermans J, Borremans C, Willenz P, André L, Dubois P 2010. Temperature, salinity and
430 growth rate dependences of Mg/Ca and Sr/Ca ratios of the skeleton of the sea urchin
431 *Paracentrotus lividus* (Lamarck): an experimental approach. Marine Biology 157: 1293–
432 1300.
- 433 34. Holtmann WC, Stumpp M, Gutowska MA, Syre S, Himmerkus N, Melzner F, Bleich M 2013.
434 Maintenance of coelomic fluid pH in sea urchins exposed to elevated CO₂: the role of body
435 cavity epithelia and stereo dissolution. Marine Biology 160(10):2631-2645.
- 436 35. James RK, Hepburn CD, Cornwall CE, McGraw CM, Hurd CL, 2014. Growth response of an
437 early successional assemblage of coralline algae and benthic diatoms to ocean acidification.
438 Marine Biology 161: 1687–1696.
- 439 36. Jangoux M, Lawrence JM 1982. Echinoderm Nutrition. Balkema, Rotterdam.
- 440 37. Johnson MD, Carpenter RC, 2012. Ocean acidification and warming decrease calcification in
441 the crustose coralline alga *Hydrolithon onkodes* and increase susceptibility to grazing. Journal
442 of Experimental Marine Biology and Ecology, 434 : 94-101.
- 443 38. Kidwell SM, Baumiller T 1990. Experimental disintegration of regular echinoids: roles of
444 temperature, oxygen, and decay thresholds. Paleobiology 16: 247-271.
- 445 39. Koch M, Bowes G, Ross C, Zhang XH 2013. Climate change and ocean acidification effects
446 on seagrasses and marine macroalgae. Global change biology 19(1): 103-132.

- 447 40. Kroeker KJ, Kordas RL, Crim RN, Singh GG, 2010. Meta-analysis reveals negative yet
448 variable effects of ocean acidification on marine organisms. *Ecology letters*, 13(11): 1419-
449 1434.
- 450 41. Kurihara H, Shirayama Y 2004. Effects of increased atmospheric CO₂ on sea urchin early
451 development. *Marine Ecology Progress Series* 274: 161–169.
- 452 42. Kurihara H, Yin R, Nishihara GN, Soyano K, Ishimatsu A 2013. Effect of ocean acidification
453 on growth, gonad development and physiology of the sea urchin *Hemicentrotus pulcherrimus*.
454 *Aquatic Biology* 18: 281–292.
- 455 43. Larson BR, Vadas RL, Keser M 1980. Feeding and nutritional ecology of the sea urchin
456 *Strongylocentrotus droebachiensis*. *Marine Biology* 59: 49-62.
- 457 44. Lawrence JM 1975. On the relationship between marine plants and sea urchins. *Oceanography*
458 *and Marine Biology, An Annual Review* 13: 213-286.
- 459 45. Lawrence JM, Lane JM 1982. The utilization of nutrients by post-metamorphic echinoderms.
460 *Echinoderm nutrition*: 331-371
- 461 46. Lilly GR 1975. The influence of diet on the growth and bioenergetics of the tropical sea
462 urchin, *Tripneustes ventricosus* (Lamarck). Thesis. Vancouver: Univ. of British Columbia.
- 463 47. Martin S, Gattuso JP, 2009. Response of Mediterranean coralline algae to ocean acidification
464 and elevated temperature. *Global Change Biol.* 15: 2089–2100.
- 465 48. Martin S, Richier S, Pedrotti ML, Dupont S, Castejon C, Gerakis Y, Kerros ME, Oberhänsli
466 F, Teyssié, J.L., Jeffree, R., Gattuso, J.P., 2011. Early development and molecular plasticity
467 in the Mediterranean sea urchin *Paracentrotus lividus* exposed to CO₂-driven acidification.
468 *Journal of Experimental Biology* 214(8): 1357-1368.
- 469 49. Märkel K 1975. Wachstum des Coronarskeletes von *Paracentrotus lividus*
470 Lmk.(Echinodermata, Echinoidea). *Zoomorphologie*, 82(2-3): 259-280.
- 471 50. Meyers MA, Chen P-Y, Lin AY-M, Seki Y 2008. Biological materials: structure and
472 mechanical properties. *Progress in Material Science* 53: 1–206.

- 473 51. Moss ML, Meehan MM 1967. Sutural connective tissues in the test of an echinoid *Arbacia*
474 *punctulata*. Acta Anatomica 66: 279-304.
- 475 52. Moulin L, Catarino AI, Claessens T, Dubois P 2011. Effects of seawater acidification on early
476 development of the intertidal sea urchin *Paracentrotus lividus* (Lamarck 1816). Marine
477 Pollution Bulletin 62(1): 48-54.
- 478 53. Moulin L, Grosjean P, Leblud J, Batigny A, Dubois P 2014. Impact of elevated pCO₂ on acid-
479 base regulation of the sea urchin *Echinometra mathaei* and its relation to resistance to ocean
480 acidification: a study in mesocosms. Journal of Experimental Marine Biology and Ecology
481 457: 97–104.
- 482 54. Moulin L, Grosjean P, Leblud J, Batigny A, Collard M, Dubois P 2015. Long-term
483 mesocosms study of the effects of ocean acidification on growth and physiology of the sea
484 urchin *Echinometra mathaei*. Marine environmental research 103: 103-114.
- 485 55. Moureaux C, Pérez-Huerta A, Compère P, Zhu W, Leloup T, Cusack M, Dubois P 2010.
486 Structure, composition and mechanical relations to function in sea urchin spine. Journal of
487 Structural Biology 170: 41– 49.
- 488 56. Moureaux C, Simon J, Mannaerts G, Catarino AI, Pernet P, Dubois P 2011. Effects of field
489 contamination by metals (Cd, Cu, Pb, Zn) on biometry and mechanics of echinoderm ossicles.
490 Aquatic Toxicology 105: 698–707.
- 491 57. Moureaux C, Dubois P 2012. Plasticity of biometrical and mechanical properties of
492 *Echinocardium cordatum* spines according to environment. Marine Biology 159: 471-
493 479.
- 494 58. Nelson WA (2009) Calcified macroalgae critical to coastal ecosystems and vulnerable
495 to change: A review. Mar Freshw Res 60(8):787–801.

- 496 59. O'Donnell MJ, Todgham AE, Sewell MA, Hammond LM, Ruggiero K, Fangué NA, Zippay
497 ML, Hofmann GE 2010. Ocean acidification alters skeletogenesis and gene expression in
498 larval sea urchins. *Marine Ecology Progress Series* 398: 157-171.
- 499 60. Orr JC, Fabry VJ, ..., Yool A, 2005. Anthropogenic ocean acidification over the twenty-first
500 century and its impact on calcifying organisms. *Nature*, 437(7059): 681.
- 501 61. Poore AGB, Graba-Landry A, Favret M, Sheppard Brennan H, Byrne M, Dworjanyn SA,
502 2013. Direct and indirect effects of ocean acidification and warming on a marine plant–
503 herbivore interaction. *Oecologia* 173: 1113–1124.
- 504 62. Porzio L, Buia MC, Hall-Spencer JM, 2011. Effects of ocean acidification on macro-algal
505 communities. *Journal of Experimental Marine Biology and Ecology* 400: 278–287.
- 506 63. Privitera D, Chiantore M, Mangialajo L, Glavic N, Kozul W, Cattaneo-Vietti R 2008. Inter-
507 and intra-specific competition between *Paracentrotus lividus* and *Arbacia lixula* in resource-
508 limited barren areas. *Journal of Sea Research* 60(3): 184-192.
- 509 64. Rich WA, Schubert N, Schläpfer N, Carvalho VF, Horta AC, Horta PA, 2018. Physiological
510 and biochemical responses of a coralline alga and a sea urchin to climate change: Implications
511 for herbivory. *Marine environmental research*, 142, 100-107.
- 512 65. Riebesell U, Fabry VJ, Hansson L, Gattuso J-P (Eds.), 2010. Guide to best practices for ocean
513 acidification research and data reporting. Luxembourg: Publications Office of the European
514 Union, pp. 53–66.
- 515 66. Rodríguez A, Hernández JC, Brito A, Clemente S, 2017. Effects of ocean acidification on
516 juvenile sea urchins: predator-prey interactions. *J. Exp. Mar. Biol. Ecol.* 493: 31–40.
- 517 67. Rodríguez A, Clemente S, Brito A, Hernández JC, 2018. Effects of ocean acidification on
518 algae growth and feeding rates of juvenile sea urchins. *Marine environmental research*, 140,
519 382-389.

- 520 68. Ruitton S, Francour P, Boudouresque C 2000. Relationships between algae, benthic
521 herbivorous invertebrates and fishes in rocky sublittoral communities of a temperate sea
522 (Mediterranean). *Estuarine, Coastal and Shelf Science* 50(2), 217-230.
- 523 69. Shirayama Y, Thornton H 2005. Effect of increased atmospheric CO₂ on shallow water
524 marine benthos. *Journal of Geophysical Research* 110(C9): C09S08.
- 525 70. Smith AB 1980. Stereom microstructure of the echinoid test. *Special Papers in Palaeontology*
526 25: 1-81.
- 527 71. Stumpp M, Dupont S, Thorndyke MC, Melzner F 2011. CO₂ induced seawater acidification
528 impacts sea urchin larval development II: Gene expression patterns in pluteus larvae.
529 *Comparative Biochemistry and Physiology - A Molecular and Integrative Physiology* 160(3):
530 320-330.
- 531 72. Stumpp M, Trubenbach K, Brennecke D, Hu MY, Melzner F 2012. Resource allocation and
532 extracellular acid-base status in the sea urchin *Strongylocentrotus droebachiensis*. *Aquatic*
533 *Toxicology* 110 – 111: 194–207.
- 534 73. Stumpp M, Hu M, Casties I, Saborowski R, Bleich M, Melzner F, Dupont S 2013. Digestion
535 in sea urchin larvae impaired under ocean acidification. *Nature climate change* 3(12): 1044.
- 536 74. Sunday JM, Fabricius KE, Kroeker KJ, Anderson KM, Brown NE, Barry JP, Connell SD,
537 Dupont S, Gaylord B, Hall-Spencer JM, Klinger T, Milazzo M, Munday PL, Russell BD,
538 Sanford E, Thiagarajan V, Vaughan MLH, Widdicombe S, Harley CDG 2017. Ocean
539 acidification can mediate biodiversity shifts by changing biogenic habitat. *Nature Climate*
540 *Change* 7(1): 81.
- 541 75. Thompson RJ 1983. The relationship between food ration and reproductive effort in the green
542 sea urchin, *Strongylocentrotus droebachiensis*. *Oecologia* 56: 50:57.
- 543 76. Thompson RJ 1984. Partitioning of energy between growth and reproduction in three
544 populations of the sea urchin *Strongylocentrotus droebachiensis*. In: Engels W, editor.

- 545 Advances in invertebrate reproduction. 3. Elsevier Science Publishers BV, Amsterdam, p425-
546 431.
- 547 77. Todgham AE, Hofmann GE 2009. Transcriptomic response of sea urchin larvae
548 *Strongylocentrotus purpuratus* to CO₂-driven seawater acidification. Journal of Experimental
549 Biology 212: 2579-2594.
- 550 78. Vadas RL 1977. Preferential feeding: an optimization strategy in sea urchins. Ecological
551 Monographs 47: 337-371.
- 552 79. Weiner S.1985. Organic matrix-like macromolecules associated with the mineral phase of sea
553 urchin skeletal plates and teeth. Journal of Experimental Zoology 234: 7-15.
- 554