



Assessing the ecological status of an estuarine ecosystem: linking biodiversity and food-web indicators

Camille Le Guen, Samuele Tecchio, Jean-Claude Dauvin, Gwenola de Roton, Jérémy Lobry, Mario Lepage, Jocelyne Morin, Géraldine Lassalle, Aurore Raoux, Nathalie Niquil

► To cite this version:

Camille Le Guen, Samuele Tecchio, Jean-Claude Dauvin, Gwenola de Roton, Jérémy Lobry, et al.. Assessing the ecological status of an estuarine ecosystem: linking biodiversity and food-web indicators. Estuarine, Coastal and Shelf Science, 2019, 228, pp.106339. 10.1016/j.ecss.2019.106339 . hal-02346068

HAL Id: hal-02346068

<https://hal.science/hal-02346068>

Submitted on 16 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Assessing the ecological status of an estuarine ecosystem:

linking biodiversity and food-web indicators

Camille Le Guen^{1,2}, Samuele Tecchio^{1,3}, Jean-Claude Dauvin⁴, Gwenola De Roton⁵, Jérémy Lobry⁶, Mario Lepage⁶, Jocelyne Morin⁷, Géraldine Lassalle⁶, Aurore Raoux^{1, 4}, Nathalie Niquil^{1*}

¹ Unité Biologie des Organismes et Ecosystèmes Aquatiques (BOREA), MNHN, CNRS, IRD, Sorbonne Université, Université de Caen Normandie, CS 14032, 14000 CAEN, France

² Pelagic Ecology Research Group, University of St-Andrews, KY16 9TS, St-Andrews, UK.

³ Sinay - Maritime Data Solution, 117 cours Caffarelli, 14000 Caen

⁴ Normandie Univ, UNICAEN, UNIROUEN, Laboratoire Morphodynamique Continentale et Côtière M2C, UMR 6143 M2C, 24 rue des Tilleuls, 14000 Caen, France

⁵ Agence Française pour la Biodiversité, Antenne Manche – Mer du Nord – DDTM-DIRM BP 34, 4 rue du Colonel Fabien – 76 083 LE HAVRE Cedex, France

⁶ IRSTEA, UR EABX (Aquatic Ecosystems and Global Changes Unit), 50 avenue de Verdun, 33612 Cestas CEDEX, France

⁷ IFREMER, Laboratoire Ressources Halieutiques, Avenue du Général de Gaulle, 14520 Port-en-Bessin, France

* Corresponding author: cam.leguen@gmail.com

Pelagic Ecology Research Group, University of St-Andrews, KY16 9TS, St-Andrews, UK

Highlights

The Seine estuary ecosystem was described before and after the Port2000 extension.


A toolbox of various indices was used, describing complementary characteristics.

Port2000 effects were complex to separate from other human-induced and natural effects.

Ecological Network Analysis was useful in defining functional indicators.

The need to continue long-term monitoring and to use multiple tools is emphasized.

Graphical abstract



Abstract

During the last decades, the highly-anthropized Seine estuary has been impacted by modification of its habitats (building of a major extension of Le Havre harbour, i.e. Port2000) and a significant natural decrease in freshwater discharge. A Before/After analysis, using a toolbox of indicators, was applied to characterize the effects of both events on the estuarine ecosystem status. We selected from existing tool boxes several indicators derived from food web modelling or community composition data, such as biodiversity indicators, a guild-based index (*i.e.* Estuarine and Lagoon Fish Index ELFI) and ecological network analysis (ENA) indices. ENA and biodiversity indicators were applied on six spatial boxes describing the Seine estuary and its outlet. Results showed an increase in taxonomic and functional richness over time, mainly due to marinisation, and significant changes in food-web properties in relation to Port2000. ENA indices appeared as a promising method in ecological status assessment, especially for estuaries considered as inherently disturbed.

Keywords: ecology, biodiversity, food web, ecosystem functioning, ecosystem health indicators

1. Introduction

Areas at the land-sea interface are often strongly impacted by human activities (e.g. Borja *et al.*, 2012). Because estuaries are of great economic importance, they are commonly subject to physical modifications such as port development and channelization to enhance harbour and industrial activities (Marmin *et al.*, 2014). In addition, pollution and other anthropogenic disturbances may cause modifications in aquatic communities (*i.e.* group of species or taxa

present in a given area that can be characterised by the distribution of individuals or biomasses among those species (Borja *et al.*, 2012)), that could in turn influence the whole ecosystem functioning. A stressed ecosystem is here defined as an ecosystem undergoing changes in its structure (e.g. productivity of its species) or functioning (e.g. food-web dynamics) following physical, chemical or biological constraints or perturbations, either of a natural or an anthropogenic origin. A stressed ecosystem is therefore expected to show modifications in both its structure (e.g. species assemblages) and functioning (e.g. food-web properties), compared to a defined baseline. Estuarine ecosystems are constituted of a mosaic of habitats linked to natural (e.g. tide, freshwater input) and anthropogenic factors (McLusky and Elliott, 2004; Elliott and Whitfield, 2011). However, it is difficult to distinguish between natural and human-induced disturbances in estuaries due to their naturally high stress level and inherent variability: a concept known as estuarine quality paradox (Dauvin, 2007; Elliott and Quintino, 2007; Dauvin and Ruellet, 2009).

Multiple tools exist to derive various indicators describing and reflecting the structure and functioning of ecosystems (Rombouts *et al.*, 2003; de la Vega *et al.*, 2018). Biodiversity and guild-based indices are the most commonly used and were largely developed for application to coastal and transition water masses in Europe under the EU Water Framework Directive (WFD) (EU, 2000; Hering *et al.*, 2010). Biodiversity indices aim to describe communities' characteristics and enable comparisons at different spatial scales (e.g. species richness, *alpha*, *beta* and *gamma*, diversities). Guild-based indicators are generally developed to assess the response of ecosystems to specific or more general disturbances (*i.e.* the comparison between the Ecology Quality Status of an estuary or a part of an estuary to a reference ecosystem with or without moderate disturbances) (Perez-Dominguez *et al.*, 2012; Lepage *et al.*, 2016). Delpech *et al.* (2010) suggested using a fish-based indicator called Estuarine and Lagoon Fish Index (ELFI) to assess the ecological status of French estuaries. ELFI was based on pressure-

impact models and the purpose was to select relevant metrics that were known to be sensitive to anthropogenic pressures (Tableau *et al.*, 2013; Pasquaud *et al.*, 2013). However, this ecological quality indicator is mainly based on structural or taxonomic elements and do not sufficiently reflect neither the functioning of ecosystems (de Jonge *et al.*, 2006; Rombouts *et al.*, 2013) nor the biodiversity patterns between and within ecosystems (e.g. Vasconcelos *et al.*, 2013).

Ecological Network Analysis (ENA) corresponds to a collection of mathematical indices describing the structure and functioning of a trophic network and characterise the overall structural properties of food webs (Ulanowicz, 1997; Niquil *et al.*, 2014). Mass-balanced food-web models such as Linear Inverse Modelling (Vézina and Platt, 1988; Niquil *et al.*, 2011; Tecchio *et al.*, 2016) and Ecopath with Ecosim (Polovina, 1984) compute values of all carbon flows in a given food web and then calculate ENA indices to assess the properties emerging from the food-web architecture.

A common feature of both biodiversity indices and holistic ENA indicators is that they can be related to concepts of stress, stability, resistance and resilience (Fath *et al.*, 2019; Safi *et al.*, 2019). The stability of an ecosystem corresponds to its ability to maintain a comparable functioning in the presence of perturbations that drive it away from its original state (Vasconcellos *et al.*, 1997; Lobry *et al.*, 2008; Saint-Béat *et al.*, 2015). Grimm & Wissel (1997) defined resistance as the property of populations, communities or ecosystems to remain “essentially unchanged”. Resilience is the ability or the time taken by a system to recover from a change due to perturbations and is linked to stability (Pimm, 1984; Mathevet and Bousquet, 2014). Although ENA indices have great potential to inform management decisions in estuaries and large tidal areas (de la Vega *et al.*, 2018), their implementation in ecosystem health assessment is not sufficiently established yet.

The objective of the present study was to investigate the contribution of ENA indices to the assessment of the Seine estuarine ecosystem quality by confronting them to some more commonly used indicators (i.e. biodiversity and guild-based indices). The Seine estuary is a good example of a highly anthropogenic transitional area (Dauvin and Desroy, 2005; Dauvin *et al.*, 2006). For more than 150 years, the Seine estuary has undergone many transformations, such as shores linearization and calibration of the navigation channel from Le Havre (i.e. entrance of the estuary) to port of Rouen (120 km upstream). Port2000 is an extension of the existing commercial port of Le Havre, located in the Northern Channel of the Seine estuary, that was constructed between 2002 and 2005 and that modified the morphological and hydrological characteristics of the estuary (Dauvin *et al.*, 2006; Tecchio *et al.*, 2016). This extension aimed to optimise the access of long-distance trade ships and to expand the available area for those vessels in the harbour. Moreover, the navigation channel is also continuously dredged to ensure maritime access of ships up to the port of Rouen (Marmin *et al.*, 2014). A recent study using Ecopath models has shown that before the construction of Port2000, the northern and navigation channels of the Seine estuary already presented signs of disturbance (i.e. an effect, either biotic or abiotic, inducing a perturbation such as stress in an ecosystem; relative to a specified reference state, (Rykiel JR, 1985)) due to maritime traffic, dredging, and building works leading to a lack of connectivity between estuarine areas (Tecchio *et al.*, 2015). Later, Tecchio *et al.* (2016) used a statistically more-advanced modelling framework (LIM-MCMC) to estimate flows in the Seine estuary food web and compute ENA indices describing its functional properties before and after Port2000. They showed that the northern channel evolved towards a shorter food web and increased its resistance property. This was considered as a potential impact of the Port2000 construction. Results from ENA indices also evidenced that the food web in the navigation channel was constantly changing, probably due to

continuous dredging activities, and that the southern channel could be seen as the least disturbed area for both periods (Tecchio *et al.*, 2016).

In addition to these anthropogenic disturbances, the Seine Estuary is also subject to an increase of salinity in the lower part of the estuary, called marinisation. Marinisation is due to a decrease in freshwater input coming from the river, associated with intrusion of marine water in the lower part of the estuary. Marinisation is a common evolution in estuaries receiving water from rivers and influenced by human activities (e.g. irrigation, dams) (David *et al.*, 2007; Goberville *et al.*, 2010; Chaalali *et al.*, 2013).

We used here an original combination of methods to assess the ecosystem health of the Seine estuary before and after the construction of Port2000. These analyses were performed on six spatial boxes previously defined to describe the estuary (Tecchio *et al.*, 2016).

We first addressed the methodological question “Do the different indicators used in this study (species-based, guild-based and ecosystem-based indices) bring similar, complementary or different information on the ecological status of this estuarine ecosystem?” We then addressed two ecological questions using this set of indicators: (i) How did physical distinct areas of the estuary (spatial boxes) react to perturbations? (ii) Can we distinguish the effects due to Port2000 from global change effects in the context of the estuary’s natural marinisation?

2. Materials and Methods

2.1. Study site and periods

Six spatial boxes, based on habitat characteristics and sediment types of the Seine estuary and the adjacent eastern part of the Seine Bay, previously defined in Tecchio *et al.* (2015) were considered: (a) three estuarine boxes: the navigation channel (CH), the northern channel (FN),

and the southern channel (FS); (b) an intermediate box located at the mouth of the estuary (E4), and (c) two marine spatial boxes (E2 and E14) (Fig. 1). Hydro-sedimentary and salinity conditions were considered homogenous within each spatial box (Tecchio *et al.*, 2015). For each zone, two sets of data were gathered corresponding to two different periods: 1996–2002, representing the situation before the Port2000 construction, and 2005–2012, corresponding to the recent situation, after the Port 2000 construction.




Fig. 1. Map showing the geographical position of the Seine estuary in France. The spatial boxes are composed of the navigation channel (CH), the northern channel (FN), and the southern channel (FS) as estuary boxes; an intermediate box located at the mouth of the estuary (E4), and two marine boxes (E2 and E14). Source: GIP Seine-Aval and SHOM.

2.2. Indicators based on taxonomic diversity

2.2.1. Data

For the subsequent indices' calculations, both fish data (from beam trawl) and macrozoobenthos data (from 0.1 m² Smith McIntyre or Van Veen grabs) were collected following identical sampling protocols between both periods, as results were standardised to take into account differences in sampling effort. This allows quantitative and qualitative comparisons to be made.

2.2.2. Indicators

The taxonomic richness is defined as the total number of taxa sampled at a given scale. It was computed by pooling all sampled taxa by spatial box and period (Before Port2000 = 1996-2002 vs After Port2000 = 2005-2012). To limit bias due to sampling design, taxonomic richness was standardized with relation to the sampling effort by dividing it by the log-transformed total number of samples realised for a given space-time unit (Nicolas *et al.*, 2010).

Alpha (α), beta (β) and gamma (γ) diversity indices give information on changes in community composition and the processes driving them, between spatial units or periods (Barros *et al.*, 2014). α -diversity describes the taxonomic diversity on a local scale (calculated for each spatial box), γ -diversity corresponds to the taxonomic diversity of the entire region of interest (all six spatial boxes considered in the study) and β -diversity represents the variation in taxonomic composition among the spatial boxes of the studied area (Barros *et al.*, 2014). The mean α -diversity ($\bar{\alpha}$) of a region or area comprising several spatial boxes was calculated as the average of α -diversity in each spatial box (Eq. 1).

$$\bar{\alpha} = \sum_{i=1}^N \alpha_i / N \quad (1); \text{ with } \alpha_i \text{ the taxonomic richness in the spatial box } i, \text{ and } N \text{ the number of spatial boxes.}$$

β -diversity is the ratio between γ and $\bar{\alpha}$ (Eq. 2) (Anderson *et al.*, 2011) and represents the number of times by which the taxonomic richness in the whole region considered (*i.e.* the entire estuary) is greater than the average taxonomic richness in the different spatial boxes. In other words, it corresponds to the ratio between regional and local species diversity and is related to compositional heterogeneity.

$$\beta = \gamma / \bar{\alpha} \quad (2)$$

with γ -diversity = taxonomic diversity of the entire region of interest and $\bar{\alpha}$ = mean α -diversity.

Biodiversity analyses were performed using the software package PRIMER-E v.6 (Clarke & Gorley, 2006). A two-dimensional non-metric Multi-Dimensional Scaling (nMDS) ordination was implemented for the six spatial boxes for each of the two periods, based on square-root-transformed biomasses to reduce the asymmetry in species distributions, and a Bray-Curtis similarity index computed on a species composition matrix. The nMDS plot gives a visual representation of the proximity in species composition between samples: the closer they are on the plot, the more similar they are in their species composition (Clarke and Warwick, 2001). The ‘stress’ index calculated by the routine in PRIMER reflects how well the ordination is capable of preserving the rank of similarities in a bi-dimensional space, providing information on the quality of the representation.

2.3. Multimetric fish-based indicator ELFI

2.3.1. Data

Fish data ‘After Port2000’ were standardized data, collected from surveys following a normalized protocol especially designed for fish monitoring in the French estuaries for the WFD (Appendix A). Fish data used for multimetric fish-based indicator ELFI (Delpech *et al.*, 2010) calculation ‘Before Port2000’ were collected from surveys designed for various other monitoring purposes but were obtained using similar sampling gear (*i.e.* beam trawls with

similar net features). For standardization and comparison, we accounted for sampling effort (trawled surface area and time spent trawling) and considered that the spatial design was relevant regarding WFD fish sampling protocol (> 6 samples by haline zone, defined by salinity level) following the Venice estuarine system: oligohaline (0-5); mesohaline (5-18), polyhaline (18-30) and euhaline (<30) zones (Venice System, 1958; Elliott & McLusky, 2002). This allowed for comparison of ELFI values between both periods.

2.3.2. Indicator calculation

ELFI has already been used to assess the ecological status of French estuaries (Tableau *et al.*, 2013; Teichert *et al.*, 2018) (ecological status categories described in Appendix B). The ELFI indicator is composed of seven metrics: density of (1) benthic fish (DB), (2) diadromous migrant species (DDIA), (3) marine juveniles migrants (DMJ), (4) freshwater species (DFW), (5) resident species (DER), (6) total density of fish (DT) and (7) standardised taxonomic richness (RT/lnS) (Pasquaud *et al.*, 2013). Indeed, taxonomic richness (RT) was divided by the log-transformed total number of samples in each spatial box (lnS) to limit the bias due to sampling design, defining RT/lnS as the standardised taxonomic richness. These metrics are proxies that reflect the physical quality of the habitat (DB, DER), the chemical quality of water and sediment (DB, DMJ, DFW), but also the habitat connectivity (i.e. the degree to which separate patches of habitat are connected; DDIA) (Hall *et al.*, 2011) and the general degradation level, which reflects a general dysfunction in ecosystem health due to accumulation of negative impacts (DT, RT/lnS). Contrary to the diversity indices calculated at the spatial box level, ELFI represents the estuary level due to the high number of samples required to calculate this index. ELFI values were computed using R (R Development Core Team, 2015). This fish index was inter-calibrated at the European level with seven other fish indices in use in the North East Atlantic region of Europe (Lepage *et al.*, 2016) in order to ensure that assessments provided by

these indices are consistent. Data were provided by ‘*Agence de l’Eau Seine Normandie*’ for the ‘After Port2000’ period whereas values were computed from Ifremer and CSLN (*Cellule de Suivi du Littoral Normand*) fish surveys for the ‘Before Port2000’ period.

2.4. Indicators based on food-web models: Ecological Network Analysis (ENA)

2.4.1. Data sources and model construction

We used the LIM model from Tecchio *et al.* (2016) for the three estuarine spatial boxes (FN, FS and CH; Fig. 1) for both periods, using 13 functional groups. A summary of data sources is given in Appendices C and D. Biomasses are expressed in gC.m^{-2} . We constructed three additional LIM models representing the spatial boxes E2, E4 and E14 based on the same functional groups (Appendix C) and the same types of input data, including production, food conversion efficiency (i.e. ratio of the biomass produced by one compartment compared to its food intake (Christensen & Walters, 2004)), respiration, and excretion ratios. Data sources and references used for the construction of the LIM-MCMC models are presented in detail in Tecchio *et al.* (2015, 2016). Briefly, a system of linear equations was set up, linking species metabolic ratios (such as production/biomass, respiration/biomass, excretion/consumption ratios), species biomass, and diet compositions. In these equations, the unknowns were the flows connecting the compartments. The flow constraints were established using data from literature or other modelling works (Appendix C). Known equalities such as those including the biomass stocks were included to constrain the model into a finite multi-dimensional space. Metabolic parameters as well as dietary constraints (i.e. proportion of prey in predator diet) were implemented as "inequalities": minimum and maximum values were included in the linear equations to partially constrain the related flows. This approach produces solutions that are robust to the effect of parameterisation uncertainty (Hines *et al.*, 2015; Hines *et al.*, 2018).

Then, this space was sampled by using a Markov-Chain Monte Carlo routine to obtain 200,000 solutions of the set of flow values. Ecological Network Analysis was applied to these 200,000 solutions, using the R software packages *NetIndices* (Soetaert *et al.*, 2015) and *enaR* (Borrett & Lau, 2014), along with scripts developed for the present study [see Tecchio *et al.* (2016) for additional methodological details].

To assess statistical differences in ENA indices between both periods, the non-parametric effect size statistic introduced by Cliff (1993) was applied [see Tecchio *et al.* (2016) for methodological details] as we were dealing with a situation of large amount of data (each ENA index was calculated 200,000 times before Port2000 and the same number after Port2000) which renders the estimation of classical parametric test statistics unfeasible (Mulholland & Jones, 1968).

2.4.2. Indices

We calculated six relevant ENA indices: the Total System Throughput (T..), the System Omnivory Index (SOI), the Finn's Cycling Index (FCI), the Ascendency (A), the relative Redundancy (R/DC) and the Detritivory/Herbivory ratio (D/H). This set of indices was used by Tecchio *et al.* (2016) to characterize the inner part of the Seine estuary and most of them have been proposed as management tools in the OSPAR Convention context (Safi *et al.*, 2019). The Total system throughput (T..) corresponds to the sum of all flows occurring in the system, and acts as an indicator of system size and activity (Rutledge *et al.*, 1976; Aoki, 1988; Latham, 2006). The Finn's Cycling Index (FCI) represents the fraction of the flows in the system that is generated by recycling and was calculated as the proportion of T.. generated by cycling (Finn, 1976; Finn, 1980). Although other FCI definitions exist, this particular one was chosen to remain consistent with previous work and another common modelling approach, *Ecopath with Ecosim*. The Ascendency was calculated as the product of the Total system throughput (T..)

and the Average Mutual Information (AMI), where AMI expresses the degree of organisation of exchanges between the different functional groups (increasing AMI indicates higher dietary specialisation; Hirata & Ulanowicz, 1984). High ascendancy values mean that the system is more actively channelling flows along more specific pathways, whereas low values have been linked to system immaturity (Ortiz & Wolff, 2002). Relative redundancy (R/DC) was calculated as the ratio between the internal flows overhead and the total development capacity of the ecosystem (Ulanowicz, 1986; Ulanowicz, 2001) and measures the amount of parallel trophic pathways connecting the trophic compartments. R/DC acts as an indicator of inefficiency of the network (Hirata & Ulanowicz, 1984; Bondavalli *et al.*, 2000) but it is also a way to increase ecosystem resilience because parallel pathways can replace each other (Ulanowicz, 1997). The System Omnivory Index (SOI) was calculated as the mean of the omnivory indices of each consumer compartment, weighted by the logarithm of their consumption (Christensen & Walters, 1993; Libralato, 2008). Omnivory is defined as the variability of prey trophic levels where high SOI values correspond to a food web with a web-like structure (i.e. with several pathways between compartments) whereas low SOI values reflect a chain-like structure with fewer pathways (Dimitrios *et al.*, 2018). Therefore, SOI is an indicator of the overall dietary adaptation of the consumers, and an increase in SOI generally corresponds to a stabilizing response to an external disturbance (Fagan, 1997; Libralato, 2008). The Detritivory/Herbivory ratio (D/H) was calculated as the ratio between the sum of all predation flows on the detritus compartment (detritivory, flows from detritus to consumers) and the sum of all predation flows on primary producers (herbivory, flows from phytoplankton and microphytobenthos to consumers). An increase in D/H would indicate that the ecosystem shifts towards a more detritus-based food web, depending less on plant material for trophic interactions (Ulanowicz, 1992; Luong *et al.*, 2014; Niquil *et al.*, 2014).

3. Results

Results concern “species-based” (species diversity and biomasses), “community-based” (fish index ELFI) and “ecosystem-based” (Ecological Network Analysis) indicators, which provide information at different levels.

3.1. Species-based indices related to diversity and biomasses

3.1.1. Biomass trend

In the estuary boxes CH and FN, fish and macrozoobenthos biomasses increased between periods (Fig. 2, Table 1). For the southern channel (FS), while macrozoobenthos biomass increased, fish biomass remained stable, and it appeared that this was the case for all fish groups (Fig. 2). For instance, the biomass of piscivorous fish in FS averaged 0.057 g C.m^{-2} in the first period and approached 0.055 g C.m^{-2} in the second one. In CH, while total fish biomass increased, the biomass of planktivorous fish decreased (Fig. 2). This was mainly related to two pelagic species: *Clupea harengus* and *Sprattus sprattus*. In the same way, for FN, while the total fish biomass increased, the biomass of piscivorous and planktivorous fishes slightly decreased (Fig. 2). This was quantitatively compensated by a strong increase in benthos feeders' biomass. The decrease in planktivorous fish biomass in FN was due to four species *Clupea harengus*, *Gasterosteus aculeatus*, *Osmerus eperlanus* and *Sprattus sprattus*. The overall biomass (including fish, macrozoobenthos and the suprabenthos) increased for the three estuarine spatial boxes. Among the three estuarine boxes, the ratio of fish biomass on benthic invertebrate biomass (BFish/BMacrozoobenthos) was the highest in CH (0.09 in the first period and 0.03 in the second one). Nonetheless, this ratio was divided by three in CH congruently to the sharp increase in benthic macrofauna biomass between periods that was multiplied by 7 (Fig. 2, Table 1).




Fig. 2. Barplots representing fish biomass, macrozoobenthos biomass, the sum of fish and macrozoobenthos biomasses, and the ratio BFish/BMacrozoobenthos, for all spatial boxes before the construction of Port2000 (first bars) and after (second bars). The spatial boxes are composed of the navigation channel (CH), the northern channel (FN), and the southern channel (FS) as estuary boxes; an intermediate box located at the mouth of the estuary (E4), and two marine boxes (E2 and E14).

Table 1. Summary of comparisons between periods (Before and After Port2000 construction) for biomass values, biodiversity indices, and the ELFI index. See text for details on the indices. < indicates that the value of the index is lower Before than After Port2000; > indicates that the value of the index is greater Before than After Port2000.

		Spatial boxes					
		CH	FN	FS	E4	E14	E2
Biomass	Fish	<	<	<	>	<	>
	Benthos	<	<	<	<	<	<
Diversity	TR	Fish	<	<	<	<	>
		Benthos	<	<	<	<	<
	α	Fish	<	<	<	<	>
		Benthos	<	<	<	<	<
	\square^*	Fish			<		
		Benthos			<		
	β^*	Fish			<		
		Benthos			>		
	γ^*	Fish			<		
		Benthos			<		
	ELFI*				<		

* \square , β and γ diversity indices are computed for the whole estuary

3.1.2. Taxonomic richness and species composition

The taxonomic richness increased between periods, from 16 to 47 fish taxa and from 5 to 29 macrozoobenthos taxa in the navigation channel (CH) (Fig. 3). The same trend was observed in the Northern and Southern channels (FN and FS; Fig. 3). The estuary box CH showed particularly low values of macrozoobenthos taxonomic richness compared to the other spatial boxes (Fig. 3). The intermediate box E4 was the richest spatial box in terms of fish (with 51 taxa during the first period and 59 taxa during the second one) and benthic macrofauna (with 177 taxa during the first period and 307 during the second one). No strong changes were observed for the fish taxonomic richness in the marine spatial boxes E14 and E2. All taxa that were gained between periods, both for fish and macrozoobenthos, were exclusively of marine origin. Results on standardised taxonomic richness are given in Figure 3.




Fig. 3. Barplots representing taxonomic richness divided by the log-transformed total number of samples realised within the six spatial boxes before (first bars) and after (second bars) the construction of Port2000 for fish and macrozoobenthos. The spatial boxes are composed of the navigation channel (CH), the northern channel (FN), and the southern channel (FS) as estuary boxes; an intermediate box located at the mouth of the estuary (E4), and two marine boxes (E2 and E14).

3.1.3. Diversity indices

The α , β and γ taxonomic diversity analyses between periods showed that the average number of fish species α increased from 32.8 to 41 species, and from 65.7 to 177.2 for macrozoobenthos. γ -diversity increased from 54 to 71 taxa for fish and from 195 to 353 taxa for macrozoobenthos, and β -diversity increased from 1.65 to 1.73 for fish and decreased from 2.97 to 1.99 for macrozoobenthos (Table 1).

The MDS ordination based on fish composition data showed a remarkable spatial segregation according to the salinity gradient between the estuarine and the marine spatial boxes (Fig. 4 top). The same pattern was observed with the MDS ordination based on macrozoobenthos species (Fig. 4 bottom). The ‘stress’ value of 0.05 was very low, underlying the good quality of the representation.




Fig. 4. MDS ordination based on Bray-Curtis similarity matrices from biomass data (square-root transformed) for fish (on the upper part) and for macrozoobenthos (on the lower part), for all spatial boxes during the first (first bars), before Port2000, and the recent (second bars) periods.

CH appeared as the spatial box that has experienced the greatest change in fish composition between periods, with less than 20% similarity in the community composition before and after Port2000. In addition, the benthic macrofauna composition in this estuary box demonstrated few similarities with the other spatial boxes, and this for both periods. For fish in the 'Before' period, marine boxes E2, E4 and E14 formed a group well separated from the inner estuary,

where FN and FS were grouped and CH isolated from these two other estuarine boxes. Similar results were obtained for macrozoobenthos. But in the ‘After’ period, the marine box E4 was grouped, for fish, with not only the estuary boxes FN and FS, but also CH (Fig. 4). This similarity in biodiversity composition was only limited to fish compartment; the benthos remained structured in a very similar way, in both periods.

3.2. A community-based index for fish: *ELFI*

During the period before Port2000, the Seine estuary showed lower values of ELFI (from 0.1 to 0.14) compared to the period after Port2000, the highest value recorded being 0.68 in 2006 (Fig. 5). The metric DDIA, indicator of habitat connectivity related to diadromous species, showed strong changes in the recent period (after Port2000), with a value of 0.88 in 2006, 0.70 in 2010 and 0.71 in 2012 (missing value in 2011), meaning that this metric gradually decreased after the Port2000 construction.




Fig. 5. ELFI values for the whole estuary for 10 years, between 1995 and 2012, with the class boundaries of the Water Framework Directive.

3.3. Ecosystem-based indicators: food webs and ENA

ENA indices showed that all indices in the two marine zones (E2 and E14) remained similar between the two periods (Table 2). R/DC and SOI decreased between before and after “Port 2000” in the estuary box (FS). All indices, except SOI, increased between the two periods in the intermediate box E4 and in the estuary boxes CH and FN.

Total System Throughput (T.), representing the size and activity of the system, significantly increased in the estuary boxes CH and FN as well as in the intermediate box E4 and remained stable in all the other spatial boxes (Table 2); Cliff’s delta; Fig. 6).

The SOI, reflecting the omnivory in the system, significantly decreased in the estuary box FS between the two periods. (Fig. 6). The D/H ratio significantly increased in the estuary boxes CH, FN and in the intermediate box E4, while remaining stable in the marine boxes E14, E2 and in FS (Table 2, Fig. 6).

The FCI reflecting the recycling in the system, significantly increased between the two periods in the estuary boxes CH, FN and in the intermediate box E4 (Table 2, Fig. 6).

The ascendancy (A) and relative redundancy (R/DC) indices significantly increased in CH between the two periods. The ascendancy also significantly increased in the intermediary box E4 between the two periods (Table 2, Fig. 6).

Table 2. ENA means and standard deviations, derived from LIM-MCMC models, for all spatial boxes before (1996-2002) and after (2005-2012) Port2000 construction. < indicates that the value of the index is lower Before than After Port2000; > indicates that the value of the index is greater Before than After Port2000. Only statistically significant results based on Cliff's delta are listed. ns: non-significant.

ENA	Period	CH	FN	FS	E4	E14	E2
T..	Before	1935 ± 337	2244 ± 251	2600 ± 240	2424 ± 347	2475 ± 296	2447 ± 364
		2332 ± 361	2463 ± 273	2507 ± 227	2687 ± 352	2494 ± 294	2564 ± 328
Evolution T..		<	<	ns	<	ns	ns
FCI	Before	0.087 ± 0.03	0.124 ± 0.03	0.163 ± 0.03	0.128 ± 0.03	0.135 ± 0.02	0.149 ± 0.04
		0.119 ± 0.03	0.175 ± 0.04	0.178 ± 0.04	0.153 ± 0.03	0.131 ± 0.02	0.145 ± 0.03
FCI	After	0.03	0.04	0.04	0.03	0.02	0.03
Evolution FCI		<	<	Ns	<	ns	ns
Ascendency	Before	3542 ± 675	4066 ± 465	4648 ± 461	4382 ± 692	4480 ± 589	4408 ± 721
		4285 ± 742	4453 ± 534	4532 ± 450	4942 ± 737	4590 ± 588	4649 ± 667
Ascendency	After						
Evolution Ascendency		<	<	Ns	<	ns	ns
R/DC (%)	Before	32.6 ± 2	40.6 ± 2	43.7 ± 2	40.6 ± 2	40.9 ± 2	41.5 ± 2
R/DC (%)	After	39.2 ± 2	41.8 ± 1	41.7 ± 2	42.4 ± 2	41.6 ± 2	42.6 ± 2
Evolution R/DC(%)		<	<	>	<	ns	ns
SOI	Before	0.145 ± 0.02	0.133 ± 0.02	0.136 ± 0.01	0.146 ± 0.01	0.143 ± 0.01	0.130 ± 0.01
		0.146 ± 0.01	0.125 ± 0.01	0.119 ± 0.01	0.148 ± 0.01	0.142 ± 0.01	0.137 ± 0.01
SOI	After	0.01	0.01	0.01	0.01	0.01	0.01
Evolution SOI		ns	ns	>	ns	ns	ns
D/H	Before	0.317 ± 0.12	0.589 ± 0.12	0.796 ± 0.13	0.600 ± 0.11	0.637 ± 0.11	0.692 ± 0.16
		0.553 ± 0.12	0.776 ± 0.16	0.758 ± 0.15	0.760 ± 0.15	0.658 ± 0.11	0.718 ± 0.12
D/H	After						
Evolution D/H		<	<	ns	<	ns	ns




Fig. 6. Ecological network analysis (ENA) indices for all spatial boxes for the period before (first bars) and after (second bars) Port2000. The spatial boxes are composed of the navigation channel (CH), the northern channel (FN), and the southern channel (FS) as estuary boxes; an intermediate box located at the mouth of the estuary (E4), and two marine boxes (E2 and E14). Means and standard deviations are represented. Asterisks represent significant changes assessed from Cliff's delta statistics (level 0.05; between before and after construction).

4. Discussion

4.1. Limitations

Fish species were sampled using beam trawls, which target the first 50cm above the seafloor. The fish assemblage was therefore biased, the pelagic species likely being underestimated. In addition, the edges of each habitat were not sampled. Therefore, a few fish species might not have been sampled. However, the sampling gears and methods were similar between periods

and among spatial boxes, allowing comparison between time periods and spatial boxes. ELFI values, for both periods, need to be considered with some caution, given the differences in sampling protocol between periods that may slightly affect the comparability.

4.2. Signs of estuarine marinisation

The biomasses of marine species increased between both periods in the Seine estuary. For example, *Pomatoschistus minutus* biomass increased in two estuarine boxes (FN, FS), and in the intermediate box (E4), and *P. microps* was newly observed in these areas after Port2000. These two fish species, especially *P. minutus* are mostly marine species (Leitão *et al.*, 2006), suggesting a tendency of marinisation. Previously termed by David *et al.* (2007) and later used in several papers (Pasquaud *et al.*, 2012; Chevillot *et al.*, 2016, 2017), « marinisation » describes the process whereby marine waters flow far upstream into the basin along with a higher abundance of marine species in the estuarine transitional zone. The Seine estuary, as many north-eastern Atlantic Ocean estuarine ecosystems (Goberville *et al.*, 2010), is going through a process of ‘marinisation’. The time period 2005-2012 was characterised by a significantly lower discharge of the Seine river (Dauvin and Pezy, 2013), resulting in an increase of salinity up to 5 in the estuarine boxes (Bacq *et al.*, 2013). The salinity in the marine boxes remained stable between periods. The arrival of marine species finding newly suitable conditions in most spatial boxes might explain the increase in taxonomic richness. This marinisation tendency was supported by the MDS ordination showing a clear spatial (along the salinity gradient) and temporal (before and after Port2000) segregation, indicating that marinisation modified the ecological structure of communities. This is also supported by the higher ELFI values in the Seine estuary after Port2000 which is likely a consequence of the marinisation enhancing the taxonomic richness and biomass of fish in the estuarine spatial boxes. The same trend has already been observed in the Gironde estuary where changes in fish

assemblages and an increase of the ELFI index were attributed to an increase in temperature and salinity indicating marinisation (Pasquaud *et al.* 2012; Chevillot *et al.*, 2016, 2017).

4.3. Signs of anthropogenic impacts

4.3.1. The global estuary

In the Seine estuary, the ELFI indicator increases with ecosystem quality (Le Pape *et al.*, 2015). The ecological status of the Seine estuary therefore improved from a ‘bad’ status before Port2000 to a ‘poor’ status after Port2000. Diadromous species abundance was associated to habitat connectivity, as opposed to habitat fragmentation (barriers to migratory fish) (Pasquaud *et al.*, 2013). After Port2000, the highest DDIA value recorded was 0.88 in 2006 and then the general pattern of this metric for that period was characterized by an overall decrease of DDIA, with values dropping down to 0.71 in 2012, likely due to a lower connectivity for migrant species such as eel *Anguilla anguilla* or grey mullet *Liza ramada*, associated with the extension of the breakwaters between FN and CH during the Port2000 construction. As an example, smelt *O. eperlanus* is a diadromous species which was formerly abundant in the upper part of the northern channel for reproduction (Morin *et al.*, 2015). Extending the breakwaters between FN and CH during the Port2000 construction might have impacted the connectivity between these two zones, and *O. eperlanus* has been negatively affected by this rupture of connectivity. Moreover, it seemed that the differences in fish assemblages between the spatial boxes widened (β -diversity increased from 1.65 to 1.73), which might be related to availability of new habitats for marine species inside the estuary. Concerning benthic species, the diversity trend between spatial boxes showed the reverse trend, showing signs of standardisation (β -diversity decreased from 2.97 to 1.99).

4.3.2. Estuarine disturbed boxes

Comparatively to the rest of the estuary, strong effects of perturbations were expected in the navigation channel (CH) because of the continuous dredging and the high hydrodynamics. The estuarine box FN was the spatial box the most directly affected by the construction of Port2000 and by the mitigation measures such as the artificial formation of meanders, which might have modified the hydrodynamics (Cuvilliez *et al.*, 2015). In addition, an increase in water turbidity was noted in this box (S. Lesourd, personal communication).

The increase in detritivory/herbivory ratio (D/H) observed in both CH and FN between the two periods suggests an ecosystem more stressed after the Port2000 construction (Niquil *et al.*, 2014), probably reacting to perturbations by enhancing opportunistic behaviours such as detritivory. The higher carbon recycling (FCI) observed after Port2000 in these two estuarine spatial boxes can also be a sign of increasing stress (Vasconcellos *et al.*, 1997). However, increasing FCI and D/H values could also be interpreted by a sign of increasing maturity (Odum, 1969; Christensen, 1995; Allesina and Ulanowicz, 2004; Vasconcelos *et al.*, 2007) but other ENA indices support the statement of increasing stress. Indeed, the relative redundancy (R/DC) significantly increased across periods in CH and FN, highlighting the fact that the system was enduring a higher level of stress (Hirata & Ulanowicz, 1984; Bondavalli *et al.*, 2000) but was also more able to resist to disturbances as the energy transfers could be maintained through the trophic network via other pathways (Ulanowicz, 1997; Rybarczyk & Elkaïm, 2003).

4.3.3. Estuarine reference box

The FS, which was initially considered as the least varying spatial box (Tecchio *et al.*, 2015) and thus a reference, showed a reduction in redundancy (R/DC), omnivory (SOI) and detritivory (D/H), while maintaining a stable system activity (T..) and carbon recycling (FCI). The trophic specialisation was evidenced by the congruent reduction in omnivory (leading to a more chain-

like structure of the food web) and redundancy (reduction of parallel pathways), which can be a sign of maturity. Results for FS therefore suggested that the reduced disturbance conditions here favoured an expected ecological succession.

4.3.4. Intermediate box

E4 is the transitional area between the inner estuary and the marine environment. Results from ENA indices show an increase in system activity (T..), redundancy (R/DC), carbon cycling (FCI), omnivory (SOI) and detritivory (D/H). Therefore, as for CH and FN, following the same interpretation framework, we could assume that the system in E4 was increasing its stabilizing mechanisms when facing stronger disturbances.

4.3.5. Marine boxes

Results from ENA indices show no significant differences between the two periods for both spatial boxes E14 and E2. Therefore, we can assume that the construction of Port2000 has no negative effect on the food web dynamics of these two marine boxes. However, the spatial box E2 contains the dumping site of Octeville for dredged sediments from the Le Havre harbour (Marmin *et al.*, 2014). For these two spatial boxes, we noticed that the fish taxonomic richness decreased whereas the benthic macrofauna richness increased. The relative importance of benthic communities had intensified in E2, as shown by the decrease in the BFish/BMacrozoobenthos ratio between both periods, contrary to what was noticed in E14. This might be explained by a potential increasing turbidity in E2 due to the dumping of sediments that could affect fish more than the benthic communities. However, the effect of the Port2000 construction on these two marine boxes cannot be evidenced.

4.4. Relative effects of marinisation and port construction

4.4.1. The global estuary

The higher taxonomic diversity and ELFI index values observed after Port2000 in a context of marinisation suggest a decrease of ecosystem stress after Port2000, as taxonomic richness is considered a key factor in ecosystem stability in response to anthropogenic pressures (e.g. McCann, 2000; Chapin *et al.*, 2000). However, investigations in the different spatial boxes were necessary to detect potential effects of the Port2000 construction.

4.4.1. Estuarine disturbed boxes

The food web in CH is continuously disturbed due to the dredging activity, greatly limiting the colonisation of the seafloor by benthic organisms, as seen in other locations (e.g. in the Venice lagoon: Pranovi *et al.*, 2003). In CH, the community was characterized by an extremely low benthos richness and biomass compared to the other spatial boxes. Nonetheless, communities subjected to regular disturbances are also more adapted, less sensitive, and recover more quickly than undisturbed communities (Dernie *et al.*, 2003; Marmin *et al.*, 2014). Results from ENA indices in the two boxes CH and FN suggest an ecosystem more stressed after the Port2000 construction. Species-based indices were very useful to detect the effects of dredging activities which directly affect the settlement of benthic communities. This effect was also reflected in the ENA indices.

4.4.2. Estuarine reference box

In FS, both fish and macrozoobenthos taxonomic richness increased. However, the ratio BFish/BMacrozoobenthos remained stable. The MDS analysis showed that the fish composition in this spatial box did not deeply change between periods. ENA indices suggest a mature system for FS for both periods. Therefore, this estuarine spatial box could be confirmed

as the least stressed area of the Seine estuary, as previously assessed by Tecchio *et al.* (2016), and that species-based indicators, ELFI and ENA indices bring similar information for this spatial box.

4.4.3. Intermediate box

We assumed that E4 should exhibit the highest taxonomic richness and biomasses given its intermediate position between marine and estuarine waters (Dauvin & Desroy, 2005). It had the highest fish and benthic macrofauna taxonomic richness of all spatial boxes. Similar results were found by Martino and Able (2003), who showed that in the Mullica river-Great Bay (USA), fish species richness increased from the marine stations to the ones at the mouth of the estuary, and then strongly decreased towards the estuarine sampling stations. The MDS analysis put forward that E4 could be seen as an intermediate spatial box for fish, but as a marine box for the macrobenthos community. Indeed, this confirmed that E4 itself can be considered as an ‘ecocline’, i.e. a boundary of progressive changes (spatial and ecological) between two different systems, the ocean and the estuary (Attrill and Rundle, 2002; Basset *et al.*, 2013). Finally, results from ENA indices demonstrated an increase in system activity, redundancy, cycling, omnivory and detritivory between the two periods. Therefore, as for CH and FN, ENA indices enabled us to identify that the system was increasing its stabilizing mechanisms, probably to face stronger disturbances.

4.4.4. Marine boxes

Finally, in the bay, E14 and E2 marine boxes appeared similar in terms of fish and macrozoobenthos taxonomic composition between the two periods. Results from ENA indices show no significant differences between the two periods for both spatial boxes. Indices based

on diversity and biomasses showed the importance of benthic communities has intensified in E2 but the origin of this change remains unknown.

4.4.5. Summary

The estuarine marinisation effects are potentially stronger than the effects of the Port2000 construction. Indeed, on the one hand, the marinisation of the Seine estuary mostly changed the species assemblages, therefore affecting the ecosystem structure (i.e. biological and physical architecture of an ecosystem), due to the arrival of numerous marine species. On the other hand, the anthropogenic pressures affected habitat connectivity and food-web dynamics (i.e. transfer of energy from one part of the food web to another, Lindeman, 1942), assessed with ENA indicators such as system activity, omnivory, recycling or detritivory. The main information brought by the combination of indices was summarized in Table 3.

Table 3. Main information about the functioning and the status of the six spatial boxes.

Method	Main results	Conclusions
Biotic indices based on diversity and biomasses => give information about the structure of communities	Increase of the taxonomic richness between periods for the estuarine spatial boxes (CH, FN, FS) and the biomass of fish and macrozoobenthos increased	Arrival of marine species such as <i>P. minutus</i> in relation to the marinisation of the estuary (lower river discharge in the second period)
	CH: low values of macrozoobenthos richness	Low benthos biomass due to dredging and high hydrodynamics in this part of the estuary
	E4: the richest area for both periods	This corresponds to the rich <i>Abra alba-Lagis koreni</i> macrobenthic community and the feeding zone for fish (mainly flatfish).
	The MDS ordination showed a spatial segregation between CH and the other zones, and a temporal change between periods, both for fish and macrozoobenthos	Evolution in salinity plays a determinant role in the spatial gradient of the fauna; both periods were clearly distinguished
Biotic indices based on the classification of species in trophic groups => give information about the functioning of the food web	FN: higher carbon recycling and detritivory (increase in detritus input) + higher redundancy (more species)	Stress increased in the estuary mainly for CH and FN zones. The other zones were less disturbed.
	CH: increase in system activity, carbon cycling, ascendancy, redundancy and detritivory	
Biotic index based on the classification of species in ecological groups => give information about the health status of ecosystems	FS: still in 'high ecological status'	Reference box
	Increase in ELFI values between periods	The ecological status was supposed to have improved but it is likely due to the arrival of marine species only
	Decrease of the metric DDIA (diadromous species) in the second period, indicator of a lower habitat connectivity	Lower connectivity due to the extension of the breakwaters between FN and CH during the Port2000 construction

ELFI was sensitive to the arrival of marine species and therefore was highly relevant in a context of marinisation of the estuary. ENA indices appeared as good 'surveillance indicators', that Shephard *et al.* (2015) proposed as indicators that would be a good complement to operational indicators, for example in the context of the Marine Strategy Framework Directive application. ENA indices are also useful to take ecosystem-based management measures for

such ecosystems (de la Vega *et al.*, 2018; Safi *et al.*, 2019). In addition, their association with biodiversity indicators made the analysis more robust as suggested by Rombouts *et al.* (2013). For instance, the arrival of new species and the higher biomasses observed in some spatial boxes between periods can also explain increases in some ENA indicators such as system activity, omnivory or redundancy but this cannot be evidenced and requires further investigations.

5. Conclusions

The effects of the Port2000 construction were not as evident as expected due to the presence of other natural (e.g. marinisation) and anthropogenic (e.g. continuous harbour activities) perturbations, which makes the interpretation of ecological indicators difficult. ELFI was particularly relevant in a context of marinisation, as this index is sensitive to the arrival of new marine species. But the different metrics used to calculate ELFI are also indicative of anthropogenic impacts (e.g. DDIA and habitat connectivity affected by the construction of Port2000), making ELFI a very powerful tool to assess the health status of this estuary. The use of ENA indices alone was not sufficient to describe changes in ecosystem functioning between the two periods, distinguishing the effects of the Port2000 construction and the process of marinisation, but they appeared as an interesting complementary set of tools for characterising the evolution of food-web dynamics. Indeed, ENA indices allowed us to observe changes that occurred in specific ecosystem processes and to identify how an ecosystem becomes more stable (e.g. by increasing omnivory) under increasing stress. We showed that detecting changes related to anthropogenic disturbances alone in the Seine requires using ecological indicators at different levels (species, communities and ecosystem) and at multiple scales of observations. This work emphasizes the need to use several metrics to assess ecosystem health status and to continue long-term monitoring of such complex and reactive ecosystems.

Acknowledgements

The authors would like to acknowledge Valérie Girardin and Justine Lequesne (UMR CNRS LMNO) for their help in the statistical analysis, Anik Brind’Amour (IFREMER) for her help on fish data integration and statistics, Armonie Tous Rius for creating the AntropoSeine database, Alexandrine Baffreau (UMR CNRS M2C) for her help in checking this database, Barbara Leroy (Agence de l’Eau Seine-Normandie), Philippe Riou (Ifremer Port-en-Bessin), Nicolas Bacq, Stéphanie Moussard and Guillaume Daan (GIP-Seine Aval) for providing advice, maps and data from their Seine estuary sampling programs. We also thank Camille de la Vega and three anonymous referees for their helpful comments on the manuscript. This study was funded by the ANTROPOSEINE project (Seine Aval Public Interest Group, GIPSA, Rouen, France). The authors declare no competing financial interests. Nathalie Niquil is also funded by the EU DG Environment project EcApRHA (‘Addressing gaps in biodiversity indicator development for the OSPAR Region from data to ecosystem assessment: Applying an ecosystem approach to (sub) regional habitat assessments’; Grant No. 11.0661/2015/712630/SUB/ENVC.2 OSPAR).

References

- Allesina S, Ulanowicz RE (2004) Cycling in ecological networks: Finn's index revisited. *Computational Biology and Chemistry*, **28**, 227–233.
- Anderson MJ, Crist TO, Chase JM et al. (2011) Navigating the multiple meanings of β diversity: A roadmap for the practicing ecologist. *Ecology Letters*, **14**, 19–28.
- Aoki I (1988) Entropy laws in ecological networks at steady state. *Ecological Modelling*, **42**, 289–303.
- Attrill M., Rundle S. (2002) Ecotone or Ecocline: Ecological Boundaries in Estuaries. *Estuarine, Coastal and Shelf Science*, **55**, 929–936.
- Bacq N, Lemoine J-P, Moussard S, Le Hir P, Lobry J (2013) *Projet MESSCENE: modélisation des fonctions écologiques de l'estuaire de la Seine et Scénarios d'évolution*. Rapport final du GIP Seine-Aval pour la Fondation de France, 69 pp.
- Barros F, Blanchet H, Hammerstrom K, Sauriau P, Oliver J (2014) A framework for investigating general patterns of benthic β -diversity along estuaries. *Estuarine, Coastal and Shelf Science*, **149**, 223–231.
- Basset A, Barbone E, Elliott M et al. (2013) A unifying approach to understanding transitional waters: Fundamental properties emerging from ecotone ecosystems. *Estuarine, Coastal and Shelf Science*, **132**, 5–16.
- Bondavalli C, Ulanowicz R, Bodini A (2000) Insights into the processing of carbon in the South Florida Cypress Wetlands: a whole-ecosystem approach using network analysis. *Journal of Biogeography*, **27**, 697–710.
- Borja Á, Basset A, Bricker S et al. (2012) Classifying ecological quality and integrity of estuaries. In: *Treatise on Estuarine and Coastal Science* (ed McLusky EW& D), pp. 125–162. Waltham: Academic Press.
- Borrett SR, Lau MK (2014) enaR: An r package for Ecosystem Network Analysis. *Methods in*

- 2007
2008
2009 708 *Ecology and Evolution*, **5**, 1206–1213.
- 2010
2011 709 Chaalali A, Chevillot X, Beaugrand G et al. (2013) Changes in the distribution of copepods in
2012
2013 710 the Gironde estuary: A warming and marinisation consequence? *Estuarine, Coastal and*
2014
2015 711 *Shelf Science*, **134**, 150–161.
- 2016
2017 712 Chapin F, Zavaleta E, Eviner V et al. (2000) Consequences of changing biodiversity. *Nature*,
2018
2019 713 **405**, 234–242.
- 2020
2021 714 Chardy P (1987) Modèle de simulation du système benthique des sédiments grossiers du golfe
2022
2023 715 normand breton (Manche). *Oceanologica acta*, **10**, 421–434.
- 2024
2025 716 Chardy P, Dauvin J-C (1992) Carbon flows in a subtidal fine sand community from the
2026
2027 717 western English 703 Channel: a simulation analysis. *Marine Ecology Progress Series*,
2028
2029 718 **81**, 147–181.
- 2030
2031 719 Chevillot X, Pierre M, Rigaud A, Drouineau H, Chaalali A, Sautour B, Lobry J (2016) Abrupt
2032
2033 720 shifts in the Gironde fish community: an indicator of ecological changes in an estuarine
2034
2035 721 ecosystem. *Marine Ecology Progress Series*, **549**, 137–151.
- 2036
2037 722 Chevillot X, Drouineau H, Lambert P, Carassou L, Sautour B, Lobry J (2017) Toward a
2038
2039 723 phenological mismatch in estuarine pelagic food web? *PLoS ONE*, **12**, e0173752.
- 2040
2041 724 Christensen V (1995) Ecosystem maturity—towards quantification. *Ecological Modelling*, **77**,
2042
2043 725 3–32.
- 2044
2045 726 Christensen V, Walters CJ (1993) Trophic models of aquatic ecosystems. In: *ICLARM*
2046
2047 727 *Conference Proceedings No. 26*, p. 390.
- 2048
2049 728 Christensen V, Walters CJ (2004) Ecopath with Ecosim: Methods, capabilities and
2050
2051 729 limitations. *Ecological Modelling*, **172**, 109–139.
- 2052
2053 730 Clarke K, Gorley R (2006) Primer v6: user manual/tutorial. PRIMER-E, Plymouth.
- 2054
2055 731 Clarke KR, Warwick RM (2001) *Change in marine communities. An approach to statistical*
2056
2057 732 *analysis and interpretation.*, 2nd edn. Plymouth Marine Laboratory, UK, 1-172 pp.
- 2058
2059
2060
2061
2062
2063
2064
2065

- 733 Cliff N (1993) Dominance statistics: Ordinal analyses to answer ordinal questions.
734 *Psychological Bulletin*, **114**, 494–509.
- 735 Cuvilliez A, Lafite R, Deloffre J, Lemoine M, Langlois E, Sakho I (2015) River flow control
736 on intertidal mudflat sedimentation in the mouth of a macrotidal estuary.
737 *Geomorphology*, **239**, 174–181.
- 738 Dauvin J (2007) Paradox of estuarine quality: benthic indicators and indices, consensus or
739 debate for the future. *Marine Pollution Bulletin*, **55**, 271–281.
- 740 Dauvin JC, Desroy N (2005) The food web in the lower part of the Seine estuary: A synthesis
741 of existing knowledge. *Hydrobiologia*, **540**, 13–27.
- 742 Dauvin J, Pezy J (2013) Long-term changes of the Seine estuary suprabenthos (1996-2012).
743 *Journal of Experimental Marine Biology and Ecology*, **448**, 93–103.
- 744 Dauvin JC, Ruellet T (2009) The estuarine quality paradox: Is it possible to define an
745 ecological quality status for specific modified and naturally stressed estuarine
746 ecosystems? *Marine Pollution Bulletin*, **59**, 38–47.
- 747 Dauvin JC, Desroy N, Janson AL, Vallet C, Duhamel S (2006) Recent changes in estuarine
748 benthic and suprabenthic communities resulting from the development of harbour
749 infrastructure. *Marine Pollution Bulletin*, **53**, 80–90.
- 750 David V, Sautour B, Chardy P (2007) Successful colonization of the calanoid copepod
751 *Acartia tonsa* in the oligo-mesohaline area of the Gironde estuary (SW France) - Natural
752 or anthropogenic forcing? *Estuarine, Coastal and Shelf Science*, **71**, 429–442.
- 753 de Jonge VN, Elliott M, Brauer V (2006) Marine monitoring: Its shortcomings and mismatch
754 with the EU water framework directive's objectives. *Marine Pollution Bulletin*, **53**, 5–
755 19.
- 756 de la Vega C, Schückel U, Horn S, Kröncke I, Asmus R, Asmus H (2018) How to include
757 ecological network analysis results in management? A case study of three tidal basins of

- the Wadden Sea, south-eastern North Sea. *Ocean & Coastal Management*, **163**, 401–416.
- Delpech C, Courrat A, Pasquaud S et al. (2010) Development of a fish-based index to assess the ecological quality of transitional waters: The case of French estuaries. *Marine Pollution Bulletin*, **60**, 908–918.
- Dernie KM, Kaiser MJ, Warwick RM (2003) Recovery rates of benthic communities following physical disturbance. *Journal of Animal Ecology*, **72**, 1043–1056.
- Dimitrios K, Konstantinos T, Athanassios M (2018) Assessing ecological and fisheries implications of the EU landing obligation in Eastern Mediterranean. *Journal of Sea Research*, **141**, 99–111.
- Elliott M, McLusky DS (2002) The Need for Definitions in Understanding Estuaries. *Estuarine, Coastal and Shelf Science*, **55**, 815–827.
- Elliott M, Quintino V (2007) The Estuarine Quality Paradox, Environmental Homeostasis and the difficulty of detecting anthropogenic stress in naturally stressed areas. *Marine Pollution Bulletin*, **54**, 640–645.
- Elliott M, Whitfield A (2011) Challenging paradigms in estuarine ecology and management. *Estuarine, Coastal and Shelf Science*, **94**, 306–314.
- EU (2000) Parliament and Council Directive 2000/60/EC of 23rd October 2000. Establishing a Framework for Community Action in the field of Water Policy. Official Journal PE-CONS 3639/1/00 REV 1, Brussels.
- Fagan W (1997) Omnivory as a stabilizing feature of natural communities. *American Naturalist*, **150**, 554–567.
- Fath BD, Asmus H, Asmus R et al. (2019) Ecological network analysis metrics: The need for an entire ecosystem approach in management and policy. *Ocean & Coastal Management*, **174**, 1–14.

- 783 Finn J (1976) Measures of ecosystem structure and function derived from analysis of flows.
784 *Journal of Theoretical Biology*, **56**, 363–380.
- 785 Finn J (1980) Flow analysis of models of the Hubbard Brook ecosystem. *Ecology*, **61**, 562–
786 571.
- 787 Goberville E, Beaugrand G, Sautour B, Tréguer P (2010) Climate-driven changes in coastal
788 marine systems of western Europe. *Marine Ecology Progress Series*, **408**, 129–147.
- 789 Grimm V, Wissel C (1997) Babel, or the ecological stability discussions: an inventory and
790 analysis of terminology and a guide for avoiding confusion. *Oecologia*, **109**, 323–334.
- 791 Le Guellec C, Bodin P (1992) Meiobenthos of the Bay of Saint-Brieuc (North Brittany,
792 France). I: quantitative distribution in subtidal and intertidal zones. *Oceanologica acta*,
793 **15**, 661–671.
- 794 Hall CJ, Jordaan A, Frisk MG (2011) The historical influence of dams on diadromous fish
795 habitat with a focus on river herring and hydrologic longitudinal connectivity. *Landscape
796 Ecology*, **26**, 95–107.
- 797 Hering D, Borja A, Carstensen J et al. (2010) The European Water Framework Directive at
798 the age of 10: A critical review of the achievements with recommendations for the
799 future. *Science of the Total Environment*, **408**, 4007–4019.
- 800 Hines DE, Lisa JA, Song B, Tobias CR, Borrett SR (2015) Estimating the effects of seawater
801 intrusion on an estuarine nitrogen cycle by comparative network analysis. *Marine
802 Ecology Progress Series*, **524**, 137–154.
- 803 Hines DE, Ray S, Borrett SR (2018) Uncertainty analyses for Ecological Network Analysis
804 enable stronger inferences. *Environmental Modelling and Software*, **101**, 117–127.
- 805 Hirata H, Ulanowicz R (1984) Information theoretical analysis of ecological networks.
806 *International Journal of Systems Science*, **15**, 261–270.
- 807 Latham LG (2006) Network flow analysis algorithms. *Ecological Modelling*, **192**, 586–600.

- 808 Le Pape O, Lepage M, Féra P (2015) La démarche de développement d'indicateurs basés sur
809 l'ichtyofaune pour évaluer la qualité écologique des eaux de transition françaises dans le
810 cadre de la DCE : une marche forcée pour des résultats positifs. *Noréis*, **235**, 37–49.
- 811 Leitão R, Martinho F, Neto JM, Cabral H, Marques JC, Pardal M a. (2006) Feeding ecology,
812 population structure and distribution of *Pomatoschistus microps* (Krøyer, 1838) and
813 *Pomatoschistus minutus* (Pallas, 1770) in a temperate estuary, Portugal. *Estuarine,
814 Coastal and Shelf Science*, **66**, 231–239.
- 815 Lepage M, Harrison T, Breine J et al. (2016) An approach to intercalibrate ecological
816 classification tools using fish in transitional water of the North East Atlantic. *Ecological
817 Indicators*, **67**, 318–327.
- 818 Libralato S (2008) System Omnivory Index. *Ecological Indicators*, **4**, 3472–3477.
- 819 Lindeman RL (1942) The Trophic-Dynamic Aspect of Ecology. *Ecology*, **23**, 399–417.
- 820 Lobry J, David V, Pasquaud S, Lepage M, Sautour B, Rochard E (2008) Diversity and
821 stability of an estuarine trophic network. *Marine Ecology Progress Series*, **358**, 13–25.
- 822 Luong A, De Laender F, Olsen Y, Vadstein O, Dewulf J, Janssen C (2014) Inferring time-
823 variable effects of nutrient enrichment on marine ecosystems using inverse modelling
824 and ecological network analysis. *Science of the Total Environment*, **493**, 708–718.
- 825 Marmin S, Dauvin J-C, Lesueur P (2014) Collaborative approach for the management of
826 harbour-dredged sediment in the Bay of Seine (France). *Ocean & Coastal Management*,
827 **102**, 328–339.
- 828 Martino EJ, Able KW (2003) Fish assemblages across the marine to low salinity transition
829 zone of a temperate estuary. *Estuarine, Coastal and Shelf Science*, **56**, 969–987.
- 830 Mathevet R, Bousquet F (2014) *Résilience & environnement - Penser les changements socio-
831 écologiques*. Buchet Chastel, Paris, 170 pp., 170 pp.
- 832 McCann KS (2000) The diversity-stability debate. *Nature*, **405**, 228–233.

- 833 McLusky DS, Elliott M (2004) *The estuarine ecosystem: ecology, threats, and management.*,
834 Third edn. Oxford University Press, 214p pp.
- 835 Morin J, Brind'Amour A, Fisson C et al. (2015) *Peuplements halieutiques en estuaire de*
836 *Seine et baie de Seine orientale - Distribution et évolution des abondances des*
837 *principales espèces capturées au chalut à perche lors des campagnes scientifiques*
838 *réalisées par l'Ifremer entre 1995 et 2010.* Rapport d'étude, 70 pp.
- 839 Mulholland H, Jones C (1968) *Fundamentals of Statistics.* Butterworth-Heinemann, London,
840 301 pp.
- 841 Napoleon C, Raimbault V, Fiant L, Riou P, Lefebvre S, Lampert L, Claquin P (2012)
842 Spatiotemporal dynamics of physicochemical and photosynthetic parameters in the
843 central English Channel. *Journal of Sea Research*, **69**, 43–52.
- 844 Nicolas D, Lobry J, Lepage M et al. (2010) Fish under influence: A macroecological analysis
845 of relations between fish species richness and environmental gradients among European
846 tidal estuaries. *Estuarine, Coastal and Shelf Science*, **86**, 137–147.
- 847 Niquil N, Saint-Béat B, Johnson G a et al. (2011) *Inverse Modeling in Modern Ecology and*
848 *Application to Coastal Ecosystems*, Vol. 9. Elsevier Inc., 115-133 pp.
- 849 Niquil N, Baeta a, Marques J, Chaalali a, Lobry J, Patrício J (2014) Reaction of an estuarine
850 food web to disturbance: Lindeman's perspective. *Marine Ecology Progress Series*, **512**,
851 141–154.
- 852 Odum EP (1969) The Strategy of Ecosystem Development. *Science*, **164**, 262–270.
- 853 Ortiz M, Wolff M (2002) Trophic models of four benthic communities in Tongoy Bay
854 (Chile): comparative analysis and preliminary assessment of management strategies.
855 *Journal of Experimental Marine Biology and Ecology*, **268**, 205–235.
- 856 Pasquaud S, Béguer M, Larsen MH, Chaalali A, Cabral H, Lobry J (2012) Increase of marine
857 juvenile fish abundances in the middle Gironde estuary related to warmer and more

- saline waters, due to global changes. *Estuarine, Coastal and Shelf Science*, **104–105**, 46–53.
- Pasquaud S, Courrat A, Fonseca VF et al. (2013) Strength and time lag of relationships between human pressures and fish-based metrics used to assess ecological quality of estuarine systems. *Estuarine, Coastal and Shelf Science*, **134**, 119–127.
- Perez-Dominguez R, MacI S, Courrat A et al. (2012) Current developments on fish-based indices to assess ecological-quality status of estuaries and lagoons. *Ecological Indicators*, **23**, 34–45.
- Pimm SL (1984) The complexity and stability of ecosystems. *Nature*, **307**, 321–326.
- Polovina J (1984) Model of a coral reef ecosystem. The Ecopath model and its application to French Frigate Shoals. *Coral Reefs*, **3**, 1–11.
- Pranovi F, Libralato S, Raicevich S, Granzotto a., Pastres R, Giovanardi O (2003) Mechanical clam dredging in Venice lagoon: Ecosystem effects evaluated with a trophic mass-balance model. *Marine Biology*, **143**, 393–403.
- R Development Core Team (2015) R: A Language and Environment for Statistical Computing.
- Ratsimbazafy RA (1998) *Spectres de taille des peuplements benthiques de sables grossiers de la Manche. Thèse de doctorat*. Paris, Muséum National d'Histoire Naturelle, 523 pp.
- Rombouts I, Beaugrand G, Artigas L et al. (2013) Evaluating marine ecosystem health : Case studies of indicators using direct observations and modelling methods. *Ecological Indicators*, **24**, 353–365.
- Rutledge R, Basorre B, Mulholland R (1976) Ecological stability: an information theory viewpoint. *Journal of Theoretical Biology*, **57**, 355.
- Rybarczyk H, Elkaïm B (2003) An analysis of the trophic network of a macrotidal estuary: The Seine Estuary (Eastern Channel, Normandy, France). *Estuarine, Coastal and Shelf*

- 883 *Science*, **58**, 775–791.
- 884 Rykiel JR EJ (1985) Towards a definition of ecological disturbance. *Australian Journal of*
- 885 *Ecology*, **10**, 361–365.
- 886 Safi G, Giebels D, Larissa Arroyo N et al. (2019) Vitamine ENA : A framework for the
- 887 development of ecosystem-based indicators for decision makers. *Ocean and Coastal*
- 888 *Management*, **174**, 116–130.
- 889 Saint-Béat B, Baird D, Asmus H et al. (2015) Trophic networks: How do theories link
- 890 ecosystem structure and functioning to stability properties? A review. *Ecological*
- 891 *Indicators*, **52**, 458–471.
- 892 Shephard S, Greenstreet S, Piet G, Rindorf A, Dickey-Collas M (2015) Surveillance
- 893 indicators and their use in implementation of the Marine Strategy Framework Directive.
- 894 *ICES Journal of Marine Science*, **72**, 2269–2277.
- 895 Soetaert K, Kones JK, van Oevelen D (2015) Package ‘NetIndices’. Estimating network
- 896 indices, including trophic structure of foodwebs in R.
- 897 Spilmont N, Migné A, Lefebvre A, Artigas L, Rauch M, Davoult D (2005) Potential
- 898 contribution of sporadic pelagic deposits on the benthic metabolism of an exposed sandy
- 899 beach during emersion (Wimereux, eastern English Channel, France). *Journal of Sea*
- 900 *Research*, **53**, 161–167.
- 901 Tableau A, Drouineau H, Delpech C et al. (2013) A fish-based index of estuarine ecological
- 902 quality incorporating information from both scientific fish survey and experts
- 903 knowledge. *Ecological Indicators*, **32**, 147–156.
- 904 Tecchio S, Rius AT, Dauvin J-C et al. (2015) The mosaic of habitats of the Seine estuary:
- 905 Insights from food-web modelling and network analysis. *Ecological Modelling*, **312**, 91–
- 906 101.
- 907 Tecchio S, Chaalali A, Raoux A et al. (2016) Evaluating ecosystem-level anthropogenic

- impacts in a stressed transitional environment: The case of the Seine estuary. *Ecological Indicators*, **612**, 833–845.
- Teichert N, Lepage M, Lobry J (2018) Beyond classic ecological assessment: The use of functional indices to indicate fish assemblages sensitivity to human disturbance in estuaries. *Science of the Total Environment*, **639**, 465–475.
- Ulanowicz RE (1986) *Growth and development: Ecosystems phenomenology*. Springer-Verlag, New York, USA.
- Ulanowicz R (1992) Ecosystem health and trophic flow networks. In: *Ecosystem Health: New Goals for Environmental Management* (eds Costanza R, Norton B, Haskell B). Island Press, Washington, DC & Covelo, CA, USA.
- Ulanowicz RE (1997) *Ecology, the Ascendent Perspective*. Columbia University Press, New York, 201 pp.
- Ulanowicz RE (2001) Information theory in ecology. *Journal of Computational Chemistry*, **25**, 393–399.
- Vasconcellos M, Mackinson S, Sloman K, Pauly D (1997) The stability of trophic mass-balanced models of marine ecosystems: a comparative analysis. *Ecological modelling*, **100**, 125–134.
- Vasconcelos RP, Reis-Santos P, Fonseca V et al. (2007) Assessing anthropogenic pressures on estuarine fish nurseries along the Portuguese coast: A multi-metric index and conceptual approach. *Science of the Total Environment*, **374**, 199–215.
- Vasconcelos RP, Le Pape O, Costa MJ, Cabral HN (2013) Predicting estuarine use patterns of juvenile fish with Generalized Linear Models. *Estuarine, Coastal and Shelf Science*, **120**, 64–74.
- Venice System (1958) Symposium on the Classification of Brackish Waters, Venice, April 8–14, 1958. *Archives Oceanography and Limnology*, **11**, 1–248.

2538
2539
2540 933 Vézina AF, Platt T (1988) Food web dynamics in the ocean. I. Best-estimates of flow
2541
2542 934 networks using inverse methods. *Marine Ecology Progress Series*, **42**, 269–287.
2543
2544 935
2545
2546 936
2547
2548
2549
2550
2551
2552
2553
2554
2555
2556
2557
2558
2559
2560
2561
2562
2563
2564
2565
2566
2567
2568
2569
2570
2571
2572
2573
2574
2575
2576
2577
2578
2579
2580
2581
2582
2583
2584
2585
2586
2587
2588
2589
2590
2591
2592
2593
2594
2595
2596

Figure legends

Fig. 1. Map showing the geographical position of the Seine estuary in France. The spatial boxes are composed of the navigation channel (CH), the northern channel (FN), and the southern channel (FS) as estuary boxes; an intermediate box located at the mouth of the estuary (E4), and two marine boxes (E2 and E14). Source: GIP Seine-Aval and SHOM.

Fig. 2. Barplots representing fish biomass, macrozoobenthos biomass, the sum of fish and macrozoobenthos biomasses, and the ratio B_{Fish}/B_{Macrozoobenthos}, for all spatial boxes before the construction of Port2000 (first bars) and after (second bars). The spatial boxes are composed of the navigation channel (CH), the northern channel (FN), and the southern channel (FS) as estuary boxes; an intermediate box located at the mouth of the estuary (E4), and two marine boxes (E2 and E14).

Fig. 3. Barplots representing taxonomic richness divided by the log-transformed total number of samples realised within the six spatial boxes before (first bars) and after (second bars) the construction of Port2000 for fish and macrozoobenthos. The spatial boxes are composed of the navigation channel (CH), the northern channel (FN), and the southern channel (FS) as estuary boxes; an intermediate box located at the mouth of the estuary (E4), and two marine boxes (E2 and E14).

Fig. 4. MDS ordination based on Bray-Curtis similarity matrices from biomass data (square-root transformed) for fish (on the upper part) and for macrozoobenthos (on the lower part), for all spatial boxes during the first (first bars), before Port2000, and the recent (second bars) periods.

Fig. 5. ELFI values for the whole estuary for 10 years, between 1995 and 2012, with the class boundaries of the Water Framework Directive.

Fig. 6. Ecological network analysis (ENA) indices for all spatial boxes for the period before (first bars) and after (second bars) Port2000. The spatial boxes are composed of the navigation channel (CH), the northern channel (FN), and the southern channel (FS) as estuary boxes; an intermediate box located at the mouth of the estuary (E4), and two marine boxes (E2 and E14). Means and standard deviations are represented. Asterisks represent significant changes assessed from Cliff's delta statistics (level 0.05; between before and after construction).

Appendices

Appendix A. Data sources used in the calculation of ELFI index

Indicator	Before Port2000	After Port2000	Boundary classes
ELFI index	Biomasses from		
	Ifremer and CSLN.	Index values from	
	Use of an R script	Agence de l'Eau Seine-	Delpech et al., 2010
	provided by Mario	Normandie	
	Lepage (Irstea, UR EABX)		

Fish data ‘After Port2000’ are standardized data from surveys designed for ELFI calculation following normalized protocol. Fish data used for ELFI calculation ‘Before Port2000’ were obtained using similar sampling gear (beam trawls with similar net features), accounting for sampling effort (trawled surface area) and considering a relevant spatial design regarding WFD fish sampling protocol (> 6 tows by haline zone)

Appendix B. Range values between the five ‘Ecological Quality Status’ levels considered for the ELFI indicator (Delpech et al., 2010). Five levels of EQS classified as ‘high’, ‘good’, ‘moderate’, ‘poor’, and ‘bad’ were used. ‘High status’ means very low human pressure. ‘Good status’ means a ‘slight’ deviation from this condition, ‘moderate status’ means moderate deviation, ‘poor status’ means high deviation and ‘bad status’ means very high human pressure.

Range values for	
EQS level	ELFI
High	$0.90 \leq \text{ELFI} \leq 1.00$
Good	$0.68 \leq \text{ELFI} < 0.90$
Moderate	$0.45 \leq \text{ELFI} < 0.68$
Poor	$0.23 \leq \text{ELFI} < 0.45$
Bad	$0.00 \leq \text{ELFI} < 0.23$

Appendix C. Data sources used in the calculation of the different indicators

Group	Compartment	Biomass data	Biodiversity indicators	ENA indices	ELFI
1	Seabirds	-		X	
2	Fish, piscivores	Ifremer, CSLN, Le Havre port authority sampling programs (Liteau, Port2000, Grand Canal, COLMATAGE 2008-2009-2010), and GIP Seine-Aval	X	X	X
3	Fish, planktivores		X	X	X
4	Fish, benthos feeders		X	X	X
5	Benthic predators		X	X	
6	Benthic filter feeders	PECTOW, BENTHOSEINE, and COLMATAGE projects, Le Havre port authority, and GIP Seine-Aval	X	X	
7	Benthic deposit feeders		X	X	
8	Suprabenthos	National Natural History Museum Paris, Wimereux marine station, Caen University, Le Havre port authority, and GIP Seine-Aval sampling programs		X	
9	Zooplankton	From Rybarczyk and Elkaïm (2003)		X	
10	Meiofauna	Le Guellec and Bodin (1992); Ratsimbazafy, (1998); Spilmont <i>et al.</i> (2005)		X	
11	Bacteria	Chardy (1987); Chardy and Dauvin (1992)		X	
12	Microphytobenthos	Spilmont <i>et al.</i> (2005), Napoleon <i>et al.</i> (2012)		X	
13	Phytoplankton	SURVAL database (http://envlit.ifremer.fr/resultats/surval)		X	
14	Detritus	Seine-Aval sampling program		X	

Appendix D. Biomasses entered in LIM-MCMC models for all trophic compartments and for both periods, before (1996-2002) and after (2005-2012) Port2000 construction. The values of biomass are mean over the years. Data sources are presented in Appendix A. More information can be found in Tecchio et al. 2016. For the spatial boxes, see Fig.1.

Trophic compartments			Period	Biomass (gC m ⁻² y ⁻¹)					
				CH	FN	FS	E4	E14	E2
1	OIS	Seabirds	1996-2002						
			2005-2012						
2	FPI	Fish, piscivorous	1996-2002	0.019	0.060	0.057	0.143	0.103	0.264
			2005-2012	0.044	0.048	0.055	0.061	0.543	0.004
3	FPV	Fish, planktivorous	1996-2002	0.040	0.037	0.025	0.043	0.026	0.016
			2005-2012	0.012	0.027	0.028	0.047	0.013	0.004
4	FBF	Fish, benthos feeders	1996-2002	0.003	0.005	0.230	0.560	0.244	0.554
			2005-2012	0.088	0.327	0.252	0.307	0.387	0.358
5	IPR	Invertebrates, predators	1996-2002	0.163	17.15	9.830	1.000	2.590	0.981
			2005-2012	1.118	26.88	41.65	1.109	14.69	2.817
6	IFF	Invertebrates, filter feeders	1996-2002	0.442	1.897	22.32	3.08	3.81	12.71
			2005-2012	1.963	4.340	5.262	9.425	3.123	10.82
7	IDF	Invertebrates, deposit feeders	1996-2002	0.052	1.587	3.140	6.134	6.134	4.255
			2005-2012	1.420	1.466	0.865	7.200	3.705	5.923
8	SUP	Suprabenthos	1996-2002	2.208	2.030	2.010	0.101	0.101	0.101
			2005-2012	0.542	0.960	0.930	0.020	0.020	0.020
9	ZOO	Zooplankton	1996-2002						
			2005-2012						
10	NEM	Meiofauna	1996-2002	0.427	0.119	0.119	0.377	0.402	0.167
			2005-2012	0.427	0.119	0.119	0.377	0.402	0.167
11	BAC	Bacteria	1996-2002						
			2005-2012						
12	MPB	Microphytobenthos	1996-2002	0.050	1.725	1.720	0.300	0.350	0.380
			2005-2012	0.050	1.725	1.720	0.300	0.350	0.380
13	PHY	Phytoplankton	1996-2002						
			2005-2012						