



Cainotheriidae (Mammalia, Artiodactyla) from Dams (Quercy, SW France): phylogenetic relationships and evolution around the Eocene–Oligocene transition (MP19–MP21)

Romain Weppe, Cécile Blondel, Monique Vianey-Liaud, Gilles Escarguel,
Thierry Pelissie, Pierre-Olivier Antoine, Maeva J. Orliac

► To cite this version:

Romain Weppe, Cécile Blondel, Monique Vianey-Liaud, Gilles Escarguel, Thierry Pelissie, et al.. Cainotheriidae (Mammalia, Artiodactyla) from Dams (Quercy, SW France): phylogenetic relationships and evolution around the Eocene–Oligocene transition (MP19–MP21). *Journal of Systematic Palaeontology*, 2020, 18 (7), pp.541-572. 10.1080/14772019.2019.1645754 . hal-02349546


HAL Id: hal-02349546

<https://hal.science/hal-02349546>

Submitted on 23 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Cainotheriidae (Mammalia, Artiodactyla) from Dams (Quercy, SW France); phylogenetic relationships and evolution around the Eocene-Oligocene transition (MP19-MP21).

Journal:	<i>Journal of Systematic Palaeontology</i>
Manuscript ID	TJSP-2019-0012.R2
Manuscript Type:	Original Article
Keywords:	Grande Coupure, Quercy karstic infillings, Western Europe, decidual dentition, morpho-anatomical phylogeny, faunal turnover
Note: The following files were submitted by the author for peer review, but cannot be converted to PDF. You must view these files (e.g. movies) online.	
nexus_matrix_Cainotheriidae.nex	

SCHOLARONE™
Manuscripts

**Title: Cainotheriidae (Mammalia, Artiodactyla) from Dams (Quercy, SW France);
phylogenetic relationships and evolution around the Eocene-Oligocene transition
(MP19-MP21).**

Authors: Weppe Romain^{a*}, Blondel Cécile^b, Vianey-Liaud Monique^a, Escarguel Gilles^c, Pélissié Thierry^d, Antoine Pierre-Olivier^a, Orliac Maëva Judith^a

^a*Institut des Sciences de l'Évolution de Montpellier, Université de Montpellier, CNRS, IRD, EPHE, Place*

Eugène Bataillon, 34095 Montpellier Cedex 05, France; bLaboratoire Paléontologie Évolution

Paléoécosystèmes Paléoprimateologie: UMR 7262, Bât. B35 TSA 51106, 6 rue M. Brunet, 86073 Poitiers

Cedex9, France ; ^cLaboratoire d'Ecologie des Hydrosystèmes Naturels et Anthropisés, Université de Lyon 1,

CNRS, UMR 5023 – LEHNA , Bat. Forel, F-69622 Villeurbanne Cedex, France; ^dRéserve naturelle nationale

géologique du Lot; Parc régional et Géoparc mondial UNESCO Causses du Quercy.

* Corresponding authors. Email: romain.weppe@umontpellier.fr

Running title: Cainotheriidae from Dams, Quercy

Abstract:

Cainotheriids are a family of small artiodactyls, known in the fossil record from the late Eocene to the middle Miocene in Western Europe. Contrary to several European endemic lineages that became extinct at the end of the Eocene or close to the Eocene-Oligocene transition (*Grande Coupure*), cainotheriids crossed this boundary and diversified during the Miocene. The evolutionary history of cainotheriids around the Grande Coupure remains poorly documented and only a few works deal with the modalities of their evolution, notably because of the scarcity of available Eocene remains. A new fossiliferous karstic network named *Le Chêne* was discovered during field prospections in the Quercy area (Tarn-et-Garonne, France).

1
2 1 It notably displays two infillings that yielded a great abundance of cainotheriid remains, namely
3 2 DAM1 (MP19, late Eocene) and DAM3 (MP22, early Oligocene), bracketing the Eocene–
4 3 Oligocene transition. A detailed study of cainotheriid mandibular and dental remains from these
5 4 infillings reveals that only *Paroxacron valdense* occurs at DAM1, while five species are found
6 5 in DAM3. The karstic network at Dams seemingly records a local taxonomic diversification of
7 6 cainotheriids after the Grande Coupure, with Cainotheriinae being particularly successful. Our
8 7 phylogenetic analysis, including cainotheriids from Dams, constitutes the first formal
9 8 phylogeny of Cainotheriidae. Our results, based on mandibular and dental characters, allow for
10 9 (i) clarifying relationships within Cainotheriidae (ii) erecting a new family Robiacinidae, as
11 10 being the sister taxon to Cainotheriidae among Cainotherioidea and, more broadly, (iii)
12 11 discussing the controversial position of Cainotheriidae within Artiodactyla, as being more
13 12 closely related to Ruminantia than to Tylopoda.

13
32 14 **Keywords:** Grande Coupure, Quercy karstic infillings, Western Europe, deciduous
33 15 dentition, morpho-anatomical phylogeny, faunal turnover.

1 1 Introduction

2
3 Cainotheriids are a family of small artiodactyls, known from the late Eocene to the
4 middle Miocene in Western Europe (Blondel 2005). This family is composed of two sub-
5 families and at least five genera (Blondel 2005): Oxacroninae Hürzeler, 1936, with *Oxacron*
6 Fihol, 1884 and *Paroxacron* Hürzeler, 1936; and Cainotheriinae Camp & VanDerHoof, 1940,
7 with *Plesiomeryx* Gervais, 1873, *Caenomeryx* Hürzeler, 1936, and *Cainotherium* Bravard,
8 1828. Their size range is variable, from small rabbit-sized species (e.g., *Oxacron*, *Paroxacron*)
9 to a size slightly smaller than the extant tragulid ruminant *Tragulus javanicus* (Theodor 2010)
10 (e.g., *Cainotherium*). Cainotheriids are characterised by brachydont selenodont molars. The
11 upper molars bear five cusps arranged in a unique '*Cainotherium plan*' (Stehlin 1906), resulting
12 from a posterior migration of the protocone (Erfurt & Métais 2007). Upper molars therefore
13 have two anterior cusps (paracone and paraconule), and three posterior ones (metacone,
14 metaconule, protocone).

15 Like many mammalian species, several European endemic ungulates became extinct at
16 the end of the Eocene (Sudre & Legendre 1992; Blondel 2001), or close to the Eocene-
17 Oligocene transition. This major mammalian faunal turnover, called the Grande Coupure
18 (Stehlin 1910), is linked to climatic, geographic and oceanic circulation changes (Legendre
19 1987; Berggren & Prothero 1992; Akhmetiev & Beniamovski 2009). Unlike most other
20 endemic artiodactyl taxa, cainotheriids crossed the Eocene-Oligocene boundary and diversified
21 during the Oligocene (Blondel 2005). However, the evolutionary history of cainotheriids
22 around this transition remains poorly documented and understood. Indeed, only few studies
23 (Legendre 1980; Hooker & Weidmann 2000) are dedicated to Eocene cainotheriids and deal
24 with the modalities of their evolution. This is primarily due to the apparent monotony of their

1 dental characters and to the scarcity of cainotheriid remains documented in pre-Grande Coupure
2 localities.

3 Knowledge of most cainotheriid genera (e.g. *Oxacron*, *Plesiomeryx*, or *Caenomeryx*)
4 principally relies on material collected from the Phosphorites karstic fillings from Quercy
5 (southwestern France) during the last century (Blondel 2005). This fossiliferous area yielded
6 an unparalleled fossil record with over 170 localities recording local faunal assemblages with a
7 very short time interval of ~1 million years or less (Aguilar *et al.* 1997, Escarguel *et al.* 2008),
8 and a continuous record spanning the 38–26 Ma interval (late Eocene to late early Oligocene).

9 In 2016, a new fossiliferous karstic network named “Dams” was discovered during field
10 prospecting in the Quercy area (Tarn-et-Garonne, France). This karstic network was partly
11 curated during the extensive phosphate exploitation in Quercy during the late 19th century. The
12 Dams network however still houses a great quantity of clay infillings and preserves several
13 fossiliferous channels distinct in age. Specifically, two fossiliferous infillings, named DAM1
14 and DAM3, yielded a great amount of cainotheriid remains. Thanks to a low-energy transport,
15 osseous and dental material is particularly well preserved, and the smallest and most delicate
16 dental elements are documented, including incisors and deciduous dentition. Biochronological
17 dating of Dams fossiliferous infillings builds on their rodent assemblages; morphological
18 analyses of their dental and cranial remains from Quercy localities have allowed for
19 identification of several chronospecies (lineages), characterised by evolutionary changes in the
20 dental features of related species (e.g. Schmidt-Kittler & Vianey-Liaud 1987; Vianey-Liaud
21 1974, 1976, 1979, 1998, 2015; Vianey-Liaud *et al.* 1995; Vianey-Liaud & Ringeade, 1993;
22 Vianey-Liaud & Schmid 2009). They have proved to be reliable biochronological markers (e.g.
23 Aguilar *et al.* 1997). Rodents, especially theridomorphs, are abundant in the Dams material.
24 The assemblages found in the different Dams infillings indicate that they bracket the Eocene-
25 Oligocene transition (see Material and Methods section for more details). Dams therefore

1 provides an unprecedented window on cainotheriid evolution, before and after the Grande
2 Coupure, i.e. a time interval experiencing a dramatic faunal turnover under drastic
3 environmental changes.

4 This work builds principally on the exceptionally abundant cainotheriid material from
5 DAM1 (MP19, pre-Grande Coupure) assigned here to the single species *Paroxacron valdense*.
6 This material allows to revise and complete the diagnosis of this species and to describe for the
7 first time the complete deciduous and permanent dentition of Cainotheriidae. Study of the less
8 abundant, but more diverse post-Grande Coupure material of DAM3, allows for discussion the
9 diversity dynamics of cainotheriids across the Eocene-Oligocene transition.

10 The phylogenetic position of Cainotheriidae within Artiodactyla remains an open
11 question. They present a mosaic of plesiomorphic characters (unfused tarsals, complete digits
12 II and V, complete dentition) and autapomorphies (selenodont molars with unique
13 ‘Cainotherium’ plan) (Theodor 2010), making their phylogenetic affinities difficult to assess.
14 According to Romer (1966) Cainotheriidae would be placed within Tylopoda, together with
15 Anoplotheriidae, Xiphodontidae, and Amphimerycidae. Webb & Taylor (1980) subsequently
16 removed them from Tylopoda; while Gentry & Hooker (1988) supported placement of
17 cainotheriids within Tylopoda in their ‘preferred analysis’ (manual, Gentry & Hooker 1988,
18 fig. 9.8). However, the results of their formal analysis (using PAUP (Swofford 1984), Gentry
19 & Hooker 1988, fig. 9.7) retrieved them in a much more basal position, outside of the
20 “Selenodontia” clade (i.e., Tylopoda + Ruminantia). More recent phylogenetic analyses placed
21 them either close to Tylopoda (Geisler & Uhen 2003, Geisler *et al.* 2007, Thewissen *et al.* 2007)
22 or as an early offshoot of a wide clade including Ruminantia (Geisler & Uhen 2005; O’Leary
23 & Gatesy 2007, Lihoreau *et al.* 2015). The revision of the dental nomenclature of cainotheriids
24 performed in this work sets a sound basis for their inclusion in phylogenetic studies at the
25 Artiodactyla level and identifies for the first time their closest relatives. Based on the abundant

1
2 Dams cainotheriid material, we here perform the first formal phylogenetic analysis of the
3 family, in order to support the systematic framework and to discuss the phylogenetic affinities
4 of Cainotheriidae.
5
6

7 **Material and Methods**

8

9 **Material**

10

11 In order to document and compare the exceptionally abundant cainotheriid dental
12 remains from DAM1, we selected and measured 270 specimens. In addition to this material
13 from DAM1, 152 specimens were also selected from the less abundant material of DAM3. The
14 assignment of different species in DAM3 relies principally on morphological features of the
15 premolars and size criterion. The attribution of several isolated specimens (mainly molars)
16 remains uncertain. In this work we compare cainotheriid remains from Dams [DAM1 (MP19),
17 DAM3 (MP22)] with material from other late Eocene and Oligocene fossiliferous localities of
18 Quercy. All specimens from Dams are housed in the collection of Montpellier University, as
19 other artiodactyl species used in this work, except for *Cainotherium laticurvatum* from the
20 Museum d'Histoire naturelle de Toulouse. These taxa were scored through direct observation
21 by RW. Only *Anoplotherium latipes*, *Parvitragulus priscus*, *Paratoceras coatesi*, and
22 *Eotylopus reedi* were scored from the literature (see Phylogenetic analysis section).

23 The age of the material described here was determined using rodent species associations
24 in the two localities DAM1 and DAM3. The assemblage documented in DAM1 consists in the
25 co-occurrence of *Patriotheridomys altus* Vianey-Liaud, 1974 and *Suevosciurus cf. minimus*
26 (Major 1873), pointing to a late Priabonian age. It corresponds to the Mammalian Paleogene
27 (MP) reference level MP19. The specific diversity retrieved in DAM3 is higher with five
28 associated taxa: *Sciuromys cayluxi* Schlosser, 1884; *Pseudolitomys major* Vianey-Liaud,
29

1
2 1 1976; *Issiodoromys medius* Vianey-Liaud, 1976; *Blainvillimys gregarius* Schlosser, 1884;
3 2 *Toeniodus curvistriatus* Pomel, 1853. This association points to MP22, coinciding with an early
4
5 3 Rupelian age.
6
7
8
9
10 4
11
12 5 **Methods**
13
14

15 6 **Measurements.** Measurements (millimetres) were taken using a mesuroscope Nikon 10. The
16 7 global surface of the tooth is expressed through: L, the maximal mesio-distal length; l, the
17 8 maximal bucco-lingual dimension. The bucco-lingual aspect of the tooth is also expressed
18 9 through two measurements: lp, maximal width of posterior lobe of the teeth; la, maximal width
19 10 of anterior lobe. The height (h) corresponds to the maximal height of the crown. L3 and l3
20 11 respectively correspond to maximal length and width of the third lobe (distal) of the lower M3.
21 12 Lam, corresponds to the maximal length of the third lobe of the lower DP4. Finally, some
22 13 diastemata lengths or premolar rows have been estimated based on alveoli when one of the teeth
23 14 was missing. The variability of dental measures is estimated by the variation coefficient (V).
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

15 6 **Phylogenetic analysis.** In order to discuss cainotheriid systematics and the position of
16 7 Cainotheriidae within Artiodactyla, we performed a phylogenetic analysis including
17 8 representatives of all cainotheriid genera, as well as representatives of Tylopoda, Ruminantia,
18 9 and European endemic artiodactyl taxa and retrieved closely related to cainotheriids in previous
19 10 studies. 23 artiodactyl species were included as operational terminal units among which eight
20 11 constitute the ingroup Cainotheriidae: *Paroxacron valdense*, *Paroxacron bergeri*, *Oxacron*
21 12 *courtoisii*, *Plesiomeryx huerzeleri*, *Plesiomeryx cadurcensis*, *Caenomeryx procommunis*,
22 13 *Caenomeryx filholi*, *Cainotherium laticurvatum*; thirteen species stand for the branching group
23 14 (Antoine 2002, Orliac *et al.* 2010) which consists in members of Ruminantia (*Parvitragulus*
24 15 *priscus*, *Lophiomeryx chalaniati*, *Archaeomeryx optatus*), Tylopoda (*Paratoceras coatesi*,
25 16 *Eotylopus reedi*) and taxa of disputed phylogenetic position (*Robiacina minuta*, *Robiacina*

1
2 1 *quercyi*, *Robiacina lavergnensis*, *Dacrytherium ovinum*, *Anoplotherium latipes*, *Mixtotherium*
3 2 *cuspidatum*, *Amphimeryx murinus*, *Xiphodon castrense*); two species form the outgroup:
4 3 *Eurodexis russelli* and *Dichobune leporina*. Dental characters (Dataset. S1) and taxa/characters
5 matrix (Dataset. S2-3) provided in the Supplementary Online Material (S). 44 are original and
6 29 are based on or modified from previous analyses. Character coding is based on direct
7 observations of specimens and on illustrations and descriptions from the literature: Stehlin
8 (1908); Matthew & Reed (1910); Hürzeler (1936); Berger (1959); Emry (1978); Sudre (1978);
9 Gentry & Hooker (1988); Erfurt & Métais (2007); Hooker (2007); Hooker & Weidmann
10 (2000); Rincon *et al.* (2015). The matrix of characters was established with the software NDE
11 1.0 (Page 2001). Phylogenetic analysis was performed with PAUP* logical version 4.0a161
12 (Swofford 2002), using a heuristic search (>12 taxa) with TBR algorithm, at 1000 repetitions
13 (10 trees retained by repetition).

14 13 **Dental nomenclature** (Fig. 1-2). The nomenclature of Boissarie *et al.* (2010) was used in this
15 work to name crests/cristids. That of VanDerMade (1996, fig. 6) and Boivin *et al.* (2018) is
16 used for deciduous dentition. The nomenclature of incisors was inspired from VanDerMade
17 (1996, fig. 13). The justification of cusp/cuspids names is provided in the Discussion section.

18 18 Abbreviations

19 MX/: upper molars; M/X: lower molars; PX/: upper premolars; P/X: lower molars; DX:
20 deciduous teeth; BSP: Bayerische Staatssammlung für Paläontologien, München, Germany;
21 DAM1, DAM3: localities of Dams; MNHN: Muséum national d'histoire naturelle, Paris,
22 France; NMB: Naturhistorisches Museum Basel, Switzerland.

Systematic palaeontology

Class **Mammalia** Linnaeus, 1758

Order **Artiodactyla** Owen, 1848

Superfamily **Cainotherioidea** Camp & VanDerHoof, 1940

Emended Diagnosis. Small artiodactyls with large auditory bullae and large orbits. Paracone and metacone sub-equal on P3/; hypoconid small on P2-P3; postprotoconulid present. Mesoconid small to large on the lower molars; entoconid slightly more to more distal than the hypoconid. Protocone sub-central to distal on the upper molars and lingual with respect to the paraconule position on M3/.

Remarks. The justification for using the super-family Cainotherioidea is provided in the following section and ensues the results of phylogenetic analysis.

Family **Cainotheriidae** Camp & VanDerHoof, 1940

Emended Diagnosis. Cainotherioidea with a size ranging from small rabbit-sized species (e.g., *Oxacron*, *Paroxacron*) to slightly smaller than the extant tragulid ruminant *Tragulus javanicus* (e.g., *Cainotherium*). Complete postorbital bar; enlarged bullae and orbits; morphology of the lower jaw variable within the family but tends to become massive and deepen posteriorly in some genera. Presence of diastemata within the premolar row in the Eocene species and tend to be reduced or lost (*Cainotherium*) in the Oligocene and Miocene forms. Lower incisors with two endocristulids; lower canine incisiform. Mesial premolars narrow; P4/ tends to be molarized; P1/ and P2/ without mesial lobe; P3/ triangular and labially concave, with a lingual cusp; P4/ subrectangular. Crowns of cheek teeth moderately high. Lower molars with a large medio-lingual metaconulid; entoconid very distal and close to hypoconulid; neotrigonid (Sudre 1977) present; absence of hypoconulid on M1/ and M2/. Upper molars sub-triangular to

1 quadrangular with a W-shaped ectoloph and crescent-shaped lingual cusps; protocone disto-
2 lingual and paraconule mesiolingual; postprotocrista long and distally orientated;
3 postparacristula present; lack of preprotocrista and paraconule junction. Limbs tetradactyl with
4 a reduction of lateral digits (II and V); forelimb shorter than hindlimb; tail long.

5 Subfamily **Oxacroninae** Hürzeler, 1936

6 **Emended Diagnosis.** Small cainotheriids, comprised around 400g (e. g. *Oxacron courtoisii*) –
7 900g (e. g. *Paroxacron bergeri*). Muzzle generally elongated and narrow; orbits large with
8 respect to the small size of the skull; ethmoidal fissures absent. Diastemata present (except *P.*
9 *bergeri*), between P1-P2, P2-P3 and P1-P2/. Premolars narrow; postprotoconulid weak;
10 protocone more mesial than paracone on P4/; P1 biradiculate. Upper molars subtriangular to
11 quadrangular (e. g. *P. bergeri*); styles without crest and mesostyle bounded (except *P.*
12 *bergeri*); postparacristula relatively short.

13 Genus **Paroxacron** Hürzeler, 1936

14 **Type species.** *Paroxacron valdense* Stehlin, 1906 from Mormont Entreroches, Switzerland
15 (MP19). Lectotype: right cranial fragment with P2-/M3/ (NMB Mt.230; Stehlin 1906),
16 **Original Diagnosis.** Oxacroninae with P1/, P1, P2/ and P2 elongated and premolariform, with
17 a short diastema between P1-/P2/, P1-P2 and P2-P3 or no diastemata at all; anterior lobe of
18 P3/ elongated; upper molars with mesostyle notch (Hürzeler 1936).

19 ***Paroxacron valdense* (Stehlin, 1906)**

20 (Fig. 2-7)

21 Synonymy: 1906 *Oxacron valdense* Stehlin; pp. 681–683, pl. 12:20, text-fig. 97.
22 1936 *Paroxacron valdense* (Stehlin); Hürzeler, pp. 105–106, fig. 72
23 1980 *Paroxacron* sp. Legendre, pp. 120–127, tab. 3–4.

1
2 **Lectotype.** A right cranial fragment with P2/-M3/ (NMB Mt.230; Stehlin, 1906), from
3
4 2 Mormont Entreroches, Switzerland (MP19).

5
6 **Emended Diagnosis.** Species of *Paroxacron* with diastemata short or occasionally absent
7
8 4 between P1/-P2/ and P/2-P/3, and very short to absent between P/1-P/2 and C-P/1; mandible
9
10 5 with a gracile and relatively shallow horizontal ramus of constant height between the premolar
11
12 6 area and the last molar; coronoid process of the mandible particularly high above the occlusal
13
14 7 plane; mandibular condyle long medio-laterally and inclined medially in posterior view;
15
16 8 angular apophysis not extended ventrally and stretched distally, with deeply concave dorsal
17
18 9 outlines; shallow *incisura vasorum*.

24
25 **Differential Diagnosis.** *Paroxacron valdense* differs from Cainotheriinae by the presence of
26
27 11 sub-triangular upper molars without crested styles, a shorter postparacristula, narrow premolars,
28
29 12 a weaker postprotoconulid, a P/1 biradiculate, diastemata between P/2-P/3 and P1/-P2/, and a
30
31 13 protocone more mesial than the paracone on P4/. It differs also from *Plesiomeryx* and
32
33 14 *Caenomeryx* species by a smaller size (except *Ple. cadurcensis*), a small protocone and a longer
34
35 15 mesial lobe on P3/, as well as the absence of a protocone on P2/. It diverges from *Cainotherium*
36
37 16 by a smaller size, the absence of cingulum/id on premolars, a triangular P3/, and the presence
38
39 17 of a valley between the entoconid and the mesoconid on lower molars. *Paroxacron valdense*
40
41 18 differs from the other Oxacroninae *Oxacron courtoisii* by the presence of more elongate and
42
43 19 premolariform premolars (P1 and P2), an elongate anterior lobe on the P3/ and a larger
44
45 20 paroconule on the upper molars. Within *Paroxacron*, *Paroxacron valdense* distinguishes from
46
47 21 *P. bergeri* by a smaller size, sub-triangular upper molars, looped mesostyle, and the presence
48
49 22 of diastemata. Finally, *Paroxacron valdense* differs from *Paroxacron* sp. by a larger size, a M1/
50
51 23 more triangular and a shorter diastema between P/1-P/2.

52
53 24 **New material from DAM1.** DAM1: four left maxillary fragments with P2/-M3/ (DAM1 01-
54
55 25 03, DAM1 07); two left maxillary fragments with P2/, P4/-M3/ (DAM1 04, DAM1 09); two

1
2
3 1 left maxillary fragments with P3-/M3/ (DAM1 05-06); one right maxillary fragment with P1/-
4 2 M3/ (DAM1 08); two left maxillary fragments with P4-/M3/ (DAM1 10, DAM1 13); two right
5 3 maxillary fragments withs P4-/M3/ (DAM1 11, DAM1 14); one left maxillary fragments with
6
7 4 P2-/M2/ (DAM1 12); two left maxillary fragments with P2-/P4/ (DAM1 15, DAM1 17); one
8 5 right maxillary fragment with P3-/M2/ (DAM1 16); one left maxillary fragment with P2-/P3/
9 6 (DAM1 18); one left juvenile maxillary fragment with DP3-/M2/ (DAM1 19); one left
10 7 mandibular fragment with P/2(partial)-M/3 (DAM1 20); one right mandibular fragment with
11 8 P/2-M/3 (DAM1 21); two left mandibular fragments with P/1(partielle)-M/3 (DAM1 22-23);
12 9 one right mandibular fragment with P/3-M/3 (DAM1 24); four left mandibular fragments with
13 10 P/3-M/3 (DAM1 25-28); one left mandibular fragment with P/2, P/4-M/3 (DAM1 29); one left
14 11 mandibular fragment with P/4-M/3 (DAM1 30); two right mandibular fragments with P/4M/3
15 12 (DAM1 31-32); one left mandibular fragmentt with P/3, M/1-M/3 (DAM1 33); one left
16 13 mandibular fragment with P/1-M/1 (DAM1 34); one right mandibular fragment with P/2-M/1
17 14 (DAM1 35); one right mandibular fragment with P/2-M/2 (DAM1 36); one left mandibular
18 15 fragment with P/1-P/4 (DAM1 37); one right mandibular fragment with P/2-P/4 (DAM1 38);
19 16 six left P/1 (DAM1 39, DAM1 41-42, DAM1 44, DAM1 46, DAM1 48); seven right P/1
20 17 (DAM1 40, DAM1 43, DAM1 45, DAM1 47, DAM1 49-51); eight left P1/ (DAM1 52-57,
21 18 DAM1 63, DAM1 65); seven right P1/ (DAM1 58-62, DAM1 64, 66); 23 left P3/ (DAM1 67,
22 19 DAM1 69-71, DAM1 73-80, DAM1 85, DAM1 87-88, DAM1 90, DAM1 92-93, DAM1 95-
23 20 96, DAM1 99-101); twelve right P3/ (DAM1 68, DAM1 72, DAM1 81-84, DAM1 86, DAM1
24 21 89, DAM1 91, DAM1 94, DAM1 97-98); eight right DP/4 (DAM1 102, DAM1 107-111,
25 22 DAM1 113-114); seven left DP/4 (DAM1 103-106, DAM1 112, DAM1 115-116); six left
26 23 DP/2-3 (DAM1 117-120, DAM1 122-123); three right DP/2-3 (DAM1 121, DAM1 124-125);
27 24 five right DP/1 (DAM1 126-128, DAM1 130-131); two left DP/1 (DAM1 129, DAM1 132);
28 25 six left DP4/ (DAM1 133-134, DAM1 140-141, DAM1 145, DAM1 147); nine right DP4/

1
2 1 (DAM1 135-139, DAM1 142-144, DAM1 146); six right DP3/ (DAM1 148, DAM1 151,
3 2 DAM1 154-155, DAM1 158, DAM1 160); nine left DP3/ (DAM1 149-150, DAM1 152-153,
4 3 DAM1 156-157, DAM1 159, DAM1 161-162); four right DP1-2/ (DAM1 163-165, DAM1
5 4 167); one left DP1-2/ (DAM1 166); seven right I/1 (DAM1 168, DAM1 170, DAM1 172,
6 5 DAM1 174, 180-182); eight left I/1 (DAM1 169, DAM1 171, DAM1 173, DAM1 175-179);
7 6 nine right I/2-3 (DAM1 183, DAM1 186-187, DAM1 189-191, DAM1 193, DAM1 196-197);
8 7 six left I/2-3 (DAM1 184-185, DAM1 188, DAM1 192, DAM1 194-195); eleven right lower
9 8 canines (DAM1 198-201, DAM1 204-206, DAM1 208-209, DAM1 211-212); four left lower
10 9 canines (DAM1 202-203, DAM1 207, DAM1 210); six right I1/ (DAM1 213-214, DAM1 217-
11 10 219, DAM1 221); four left I1/ (DAM1 215-216, DAM1 220, DAM1 222); ten right I2/ (DAM1
12 11 223-224, DAM1 227-230, DAM1 232, DAM1 234-235, DAM1 237); five left I2/ (DAM1 225-
13 12 226, DAM1 231, DAM1 233, DAM1 236); seven left I3/ (DAM1 238, DAM1 242, DAM1 244-
14 13 247, DAM1 251); eight right I3/ (DAM1 239-241, DAM1 243, DAM1 248-250, DAM1 252);
15 14 eight right upper canines (DAM1 253-254, DAM1 258, DAM1 260, DAM1 262-264, DAM1
16 15 266); seven left upper canines (DAM1 255-257, DAM1 259, DAM1 261, DAM1 265, DAM1
17 16 267); one right mandibular fragment with M/3 (DAM1 268); two left P/3 (DAM1 269-270).

40 17 Description.

41 18 **Lower dentition (Tab. S1; Fig. 2-3, 6).** The lower incisors are characterised by dorso-
42 19 ventrally straight crown and root.

43 20 I/1 (Fig. 2F, 6A) has a dorso-ventrally elongated and mesio-distally short crown. It is
44 21 characterised by two endocristulids related to the pre-conulid and post-conulid, respectively;
45 22 they form two pronounced lobes, surrounded by three fossids (pre, endo and post). The posterior
46 23 lobe tends to be larger and more curved lingually than the anterior one. I/1 present also a small
47 24 protoconid, weakly pronounced apical stylids, a post-fossid deeper than other fossids and wider
48 25 than pre-fossid, the endofossid being the most important. They have an apical crest

1 (mediocristid) connecting the two conulids slightly disto-lingually inclined, and a long and thin
2 root. Morphological variation observed within the DAM1 sample (15 specimens) is negligible.

3 I/2-3 (Fig. 3K, 6B) are very similar morphologically. They differ from I/1 by a dorso-
4 ventrally shorter crown, an anterior part of the tooth more mesially projected, and more
5 pronounced lobes, with a posterior lobe more lingually directed than the anterior lobe. They
6 also present a pre-styliid tending to be more pronounced than the poststyliid, bigger pre- and post-
7 fossid, a more disto-lingually inclined crown, with a mediocristid strongly inclined disto-
8 lingually, and a bulge at the base of mesial lobe. Morphological variation is rather small and
9 concerns mainly the curvature of the crown (30 specimens).

10 Lower canine (Fig. 3L, 6C): The lower canine is incisiform. It is larger than the incisors
11 and it differs from the latter teeth by a posterior part more mesially directed, a well-pronounced
12 distal lobe and bigger than mesial lobe, a more marked prestyliid, a bucco-lingually deep, and
13 wide and mesio-distally short postfossid. It further differs from incisors by a mesio-distally
14 deeper and larger endofossid, a mesio-distally elongated and dorso-ventrally short prefossid, a
15 less disto-lingually inclined mediocristid. The lower canine also presents a bigger bulge at the
16 base of the mesial lobe and a wider and longer root. Morphological variation is negligible and
17 is focused on the post-fossid development (15 specimens).

18 P/1 and P/2 (Fig. 3A-B, 6D-E, 6L) are very similar morphologically and they essentially
19 differ in size (Tab. S1) and root orientation. P/1 is short mesio-distally, with close oblique roots,
20 whereas P/2 is longer with more remote roots. P/1 and P/2 are characterised by two main cuspids
21 with a protoconid larger and higher than paraconid; a bending at the mesial margin of the tooth
22 (postparacristid and preprotocristid junction), the postparacristid being disto-buccally directed
23 and the preprotocristid mesio-buccally directed; a single and straight distal cristid disto-
24 buccally orientated, connecting the protoconid to the distal margin of the tooth; a small
25 accessory cusp at the back of the protoconid, called a postprotoconulid; a weak disto-lingual

1
2 1 cingulid with a small distal stylid (distostyloid); a disto-lingual fossid (endofossid). Inter-
3 2 specimen variability mainly concerns the paraconid size. The bigger it is, the deeper the bending
4 3 between protoconid and paraconid is. Some P/1 do not show this bending. Other variations are
5 4 also observed as the distance between the roots, the size of the endofossid, the presence of an
6 5 endoprotocristid, and the size of the postprotoconulid.

7 6 P/3 (Fig. 3C-D, 6L, 6N) is the longest premolar, and it may occasionally have a small
8 7 root between the two main roots (Fig. 6M). Its morphology is very similar to that of P/1 and
9 8 P/2, but it differs from the latter in its distal part by similarities with P/4. The postprotoconulid
10 9 of P/3 is bigger and the distal cristid lingually curves on the distal margin of the tooth, and bears
11 10 two small accessory cusps, called conulid 1 (end) and conulid 2. Mesially, the paraconid is also
12 11 larger than on P/1-2. Marked morphological variations are observed on P/3s from DAM1: i) the
13 12 distal cristid can be long to very short lingually, joining a small to absent conulid 1, ii) the depth
14 13 and width of the endofossid vary, iii) a small distal and buccal cristid may be present, iv) the
15 14 postprotoconulid is more or less large and may have a small endocristid disto-lingually curved,
16 15 as on P/4.

17 16 P/4 (Fig. 3E-F, 6L, 6N) is shorter and more molariform than P/3 and it differs from it
18 17 by the presence of three main cusps with a large protoconid, a paraconid and a metaconid, the
19 18 protoconid being more mesial than the metaconid. It also shows an anterior part (protoconid
20 19 and paraconid) more lingually inclined, with a preprotocristid mesio-lingually orientated and a
21 20 paracristid more lingually inclined. The entoconid and hypoconid are weakly pronounced, the
22 21 latter having an extended prehypocristid. The postprotoconulid is smaller, with a more extended
23 22 endocristid, and there are two posterior cristids on the protoconid and the metaconid, joining at
24 23 the center of the tooth. P/4 further has disto-buccal and distal cingulids. Morphological variation
25 24 only concerns the size of cristids and cingulids, and paracristid orientation. The endocristid of

1
2 1 the postprotoconulid may in some specimens connect to an accessory cusp or link the
3 2 metaconid.
4
5

6
7 3 M/1-2 (Fig. 3G-H, 6L, 6N) are very difficult to distinguish from each other. They are
8
9 4 characterised by five main cusps with a high metaconid, a metaconulid mesio-lingually located
10 5 and an entoconide disto-lingually. The entoconid has a loop-like shape, notably created by
11 6 respectively distally convex and concave preentocristid and postentocristid. M/1-2 present also
12 7 a small to absent lingual paraconid and an entoconid more distal than the hypoconid. The
13 8 mesiodistal cristid (mediocristid) is short, connecting the premetacristulid and postmetacristid,
14 9 and closing the transverse valley. There is a continuous cristid formed by a long prehypocristid
15 10 and short premetacristulid; an accessory cusp is present on this last cristid, called a
16 11 centroconulid, and bears a cristid (postendocentrocristulid) disto-lingually orientated. This set
17 12 corresponds to the neotrigonid described by Sudre (1977). Finally, M/1-2 bear a major buccal
18 13 cristid formed by the junction of the posthypocristid and postentocristid that is directly
19 14 connected and a large distal cingulid. Morphological variation is negligible.
20
21

22 15 M/3 (Fig. 2C-D, 3I-J, 6L, 6N) exhibits the same pattern as M/1-2, differing from the
23 16 latter in having a hypoconulid. This conulid connects the rest of the talonid by the
24 17 prehypocristulid, which joins the postentocristid (short) and the posthypocristid (long). The
25 18 hypoconulid and the entoconid are very close to one-another; the talonid is smaller than that of
26 19 M/1-2. There are some variations between specimens (posthypocristulid sometimes present;
27 20 roots are either fused or separate).
28
29

30 21 **Upper dentition (Tab. S2; Fig. 2, 4, 7).** The upper incisors are characterised by a
31 22 spatulate-like crown (lingually concave) and an inclined root.
32
33

34 23 The I1/ (Fig. 2E, 7C) is a very simple tooth with a dorso-ventrally elongated crown. It
35 24 is characterised by a mesio-distally symmetrical crown, with the apical portion of the tooth
36 25 mesially convex. It bears two main cusps, with a small paracone and a metacone having
37
38

1
2 1 respectively an endoparacrista and endometacrista, and small apical styles. The lingual surface
3 2 bears a little mesial fossa (pre-fossa), a very tiny and lightly deep distal fossa (post-fossa), a
4 3 mesio-distally large medial fossa (endo-fossa) and a lingual cingulum. The root is thin and
5 4 short. Variation observed within our sample (11 specimens) is rather weak and concerns the
6 5 mesial length of the paracone/parastyle portion of the tooth.

7 6 I2/ (Fig. 4K, 7B) is more complex and mesio-distally longer than I1/. It differs from the
8 7 latter by a less spatulate-like and mesio-distally assymmetrical crown, a more mesially
9 8 projected anterior part, more pronounced metacone and paracone (the latter being bigger), more
10 9 marked styles, with a metastyle with a small bulge at its base. The pre-fossa and the post-fossa
11 10 are mesio-distally wider and deeper, the endo-fossa is mesio-distally smaller. The apical crest
12 11 connecting the paracone and the metastyle (postcrista) is inclined disto-lingually. The lingual
13 12 cingulum is larger, and the root is longer and less arched than that of I1/. Morphological
14 13 variation affects the size of cusps and styles and the degree of crests inclination (15 specimens).

15 14 I3/ (Fig. 4J, 7A) is mesio-distally shorter than I1/-2/. It differs from I2/ by a lower crown,
16 15 a less mesially projected mesial part, a larger paracone giving the tooth a caniniform shape, a
17 16 less dorso-ventrally extended metastyle and placed at the level of its bulge, at the base of the
18 17 crown. The parastyle is more pronounced. The post-fossa is smaller but deeper, while the pre-
19 18 fossa is larger, as wide as the central fossa. The post-fossa is strongly inclined distally. The
20 19 precrista is more mesio-lingually inclined and the apical crest connecting the paracone and
21 20 metacone (mediocrista) is more disto-lingually inclined. The root of I3/ is stockier than that of
22 21 I2/. The morphological variation (15 specimens) is rather weak and similar to I2/. Some teeth
23 22 present a strong mesial projection of their mesial part.

24 23 Upper canine (Fig. 4L, 7D) is taller and stockier than I3/. It differs from it by a more
25 24 mesio-distally directed mesial part, larger paracone and metacone, more pronounced styles, the
26 25 metastyle having a big basal bulge. The post-fossa is deeper, the pre-fossa is mesio-distally

1 wider, and the endo-fossa is more compressed. The mesial fossa is thus more important than
2 the endo-fossa. The mediocrista and postcrista are strongly disto-lingually inclined. The root of
3 the upper canine is longer and very thick. Variation is mainly in metastyle and metacone
4 development. Some teeth present an anterior part more mesially projected (15 specimens).

5 P1/ (Fig. 4A-B, 7I-J) is the smallest upper premolar. It is characterised by two medial
6 main cusps, with a paracone larger and higher than the metacone, mesial (parastyle) and distal
7 (metastyle) styles weaker marked, the former being higher and more salient than the latter.
8 The posterior part of P1/ is almost as wide as the anterior part. The paracristyla is mesio-
9 lingually projected and the preparacrista slightly mesio-buccally orientated. There is a very
10 small posterior fossa, and a weak lingual cingulum. The roots are close and oblique. The
11 morphology of P1/ is highly variable: i) the general shape of the tooth can be elongated to very
12 short, ii) the metacone varies also greatly in size from very small to as large as the paracone,
13 iii) the paracristyla is variously mesio-lingually projected; a ectoprotocrista can be present, iv)
14 the distance between the two roots varies in size.

15 P2/ (Fig. 4C-D, 7M) is mesiodistally longer than P1/, but quite similar in morphology.
16 It differs from the latter by a smaller paracone, a parastyle more salient and individualised, a
17 crest connecting the metacone to metastyle disto-buccally inclined, and the appearance of a
18 lingual loop-like crest (postmetacristula), on the distal part of the tooth. The distal part of the
19 tooth is enlarged, with a slightly bigger lingual cingulum. P2/ is more variable than P1/,
20 especially in the distal part: i) a cingular cusp and an ectometacristyla can occur, ii) the
21 postmetacristyla can be either long or short. Some teeth have a small root between the two main
22 roots.

23 P3/ (Fig. 4E-F, 7M) differs morphologically from P1/-2/. It is subtriangular in shape and
24 bears three roots. It is characterised by well-pronounced paracone, metacone and protocone,
25 salient parastyle and metastyle, a very elongated mesial lobe (paracone and parastyle) and a

1
2 1 crest connecting the metastyle to parastyle that is buccally concave. There are lingual and distal
3 2 cingula. P3/ is highly variable: i) the protocone can be wide to mesio-distally compressed and
4 3 it can be located in a distal to medial position, ii) mesial and disto-lingual (protocone) lobes can
5 4 be variously elongated and wide, iii) the crests connecting the paracone and the parastyle
6 5 (preparacrista and postparacristyla) can be more or less mesio-buccally inclined, iv) the crest
7 6 formed by the junction of the postmetacrista and premetacristyla is variously inclined disto-
8 7 buccally, v) an ectoprotocrista can be occasionally present, oriented towards the metacone, vi)
9 8 cingulum can be variously expanded. The P3/ of this species therefore have a wide
10 9 morphological diversity, ranging from a small, stocky and short tooth to an elongated and
11 10 narrow tooth.

12 11 P4/ (Fig. 4G-H, 7M) is easily recognisable from other premolars by much shorter mesio-
13 12 distal dimensions. It presents a paracone larger and higher than the mesial protocone, a small
14 13 metacone, well individualised and salient parastyle and metastyle, medio-lingually bent. A crest
15 14 connects the parastyle and the metastyle is buccally concave. An accessory cusp is present on
16 15 the postprotocrista (conule 1), and it bears a crest extended buccally. The disto-lingual cingulum
17 16 is large. Variations between specimens are important and mainly related to the cingulum
18 17 development. The P4/ can vary from stocky and quadrangular to narrow and triangular.

19 18 Upper molars (Fig. 2A-B, 4I, 7M) are morphologically similar. They are characterised
20 19 by the '*Cainotherium*' plan, composed of five main cusps (Stehlin, 1906). The paracone is
21 20 mesio-buccal, the paraconule mesio-lingual, while the protocone is disto-lingual, located more
22 21 mesially than the metacone and the metaconule. Upper molars present also well-pronounced
23 22 styles, with a loop-like mesostyle; a short and continuous premetacristula; a preprotocrista
24 23 tending to bind with the postparacristula; a preparacristula connecting the preparacristyla, and
25 24 the postmetacristula joining the postparacristyla; a mesial cingulum; a lingual cingulum
26 25 symbolised by cingular cusps in front of (mediostyle) or behind the protocone (distostyle). The

1 morphology of upper molars is highly variable: i) the crown shape can be triangular to
2 rectangular (bucco-lingually), ii) the inclination of the crown is variously pronounced, with a
3 small mesially and distally convex curvature, accompanied by a parastyle more or less buccal,
4 iii) the preprotocrista can be long buccally in some teeth, iii) the distostyle is variously marked.

5 $M2/$ tends to be wider than the other two molars, whereas $M1/$ is the shortest. On $M1/$,
6 the paraconule is more lingually located and the general outlines of the crown concurrently
7 rectangular. However, because of important morphological variation, the distinction between
8 $M1/$ and $M2/$ of our sample is essentially done by size criterion (Tab. S2).

9 $M3/$ further differs from other molars in having a more triangular shape, with a more
10 buccal paraconule and a less prominent mediostyle. It has a paracone higher than the metacone
11 in buccal view, unlike $M1/$ and $M2/$ where these two cusps are of equal height.

12 **Lower deciduous teeth (Tab. S3; Fig. 5A-F, 6F-K).** The lower deciduous teeth differ
13 from the permanent ones by a lower and more mesio-distally elongated crown, with thinner
14 enamel layer, and more individualised cuspids/stylids.

15 $DP/1$ (Fig. 5A-B, 6F-G) is very similar in morphology and variability to $P/1$. It differs
16 from the latter in having a larger postprotoconulid, a cristid connecting the protoconid and
17 postprotoconulid mesio-distally projected, a cristid joining the postprotoconulid and distal part
18 of the tooth more disto-buccally oriented and curving lingually, a smaller lingual posterior
19 fossid, spaced roots.

20 $DP/2-3$ (Fig. 5C-D, 6H-I) are morphologically similar to $P/3$ but they differ in their
21 posterior part by pronounced hypoconid and entoconid; a larger conulid 2 with a small cristid
22 lingually directed; a cristid descending from the postprotoconulid to the conulid 2 and joining
23 the prehypocristid and postentocristid, strongly disto-buccally inclined; a disto-lingually
24 inclined cristid starting from the postprotoconulid; a mesio-distal cristid connecting the

1
2 1 protoconid and postprotoconulid. The morphological variation observed in our sample is low
3
4 2 (nine specimens).

5
6 3 DP/4 (Fig. 5E-F, 6J-K) is very similar in morphology and variability to lower molars
7
8 4 but it differs from them in the presence of a third lobe in its anterior part, consisting of a lingual
9
10 5 primoconid and a buccal paraconid, the latter being little more mesial than the former. The
11
12 6 talonid (posterior lobe) is larger than the mesial lobe. The postprimocristid is directed disto-
13
14 7 lingually and joins the preprotocristid and premetacristid. There is a bucco-lingually inclined
15
16 8 cristid formed by the junction of the endoprotocristid and ectometacristid, and a bucco-lingually
17
18 9 inclined cristid originating from the hypoconid (endohypocristid). DP/4 bears an open
19
20 10 transverse valley and an absent to weak mesial cingulid.

21
22 11 **Upper deciduous teeth (Tab. S3; Fig. 5G-L, 7E-H, 7K-L).** Compared to the
23
24 12 permanent dentition, the upper deciduous teeth present the same general characteristics as the
25
26 13 lower ones and they differ by a lower, narrower, and more mesio-distally elongated crown.

27
28 14 DP1-2/ (Fig. 5G-H, 7E-F) are premolariform and very close morphologically to P2/.
29
30 15 However, they differ from P2/ in having a shorter postmetacristyla and in lacking a lingual
31
32 16 cingular cusp. Morphological variation is negligible (five specimens).

33
34 17 DP3/ (Fig. 5I-J, 7G-H) is distally molarised and mesially premolarised. It is very similar
35
36 18 to DP4/ and upper molars in its distal part and to DP1-2/ in its mesial part. However, DP3/ is
37
38 19 shorter and wider than DP1/ and DP2/. It is characterised by a distal part with a metacone,
39
40 20 metaconule and protocone, a mesial part with a big paracone, salient styles with posterior
41
42 21 metastyle and mesostyle, as well as a highly mesially prominent anterior parastyle. The crown
43
44 22 shows a significant distally concave curvature owing to a strong mesio-distally compressed
45
46 23 protocone, splitting disto-buccal and disto-lingual parts. The anterior part of the tooth is very
47
48 24 elongated and bucco-lingually compressed (paracone to parastyle). The paracristyla is mesio-
49
50 25 buccally inclined, the ectoprotocrista binds with the premetacristula and the preprotocrista with

1 the mediostyle. The distostyle is marked and the lingual cingulum is big. Morphological
2 variation mainly affects the size of the protocone and the bucco-lingual compression of the
3 distal part of the tooth (15 specimens).

4 DP4/ (Fig. 5K-L, 7K-L) morphologically resembles upper molars in its mesial half and
5 DP3/ in its distal half. However, DP4/ is shorter and wider than the latter and it differs from
6 other molars by its small size, and by a strong inclination of the crown, with a mesially convex
7 and distally concave outline. Styles are also more salient and individualised, the parastyle being
8 more buccal and metastyle strongly disto-buccally projected. The postprotocrista is more
9 inclined disto-lingually. The distostyle is marked at the back of the protocone. DP4/ further
10 differs from DP3/ by lack of a distal cingulum, presence of a lingual cingulum and a less marked
11 mediostyle. Variation concerns size (Tab. S3) and the degree of inclination of the crown (15
12 specimens).

13 **Diastemata/Mandible (Fig. 6L, 6N-R).** Specimens are characterised by diastemata
14 short or occasionally absent between P1/ and P2/, P/2 and P/3, and shorter or absent between
15 P/1 and P/2 (Tab. S1; Fig. 6L, 6O). The posterior tip of the mandibular symphysis is located at
16 the level of the distal margin of P/2 (Fig. 6L, 6N). Two mental foramina are constantly present
17 on the horizontal ramus, one below the mesial part of P/2 and one below the mesial part of P/4
18 (Fig. 6O, 6R). The horizontal ramus is gracile, relatively shallow and its height is rather constant
19 between the premolar area and the last molar. However, specimens show almost no variation.
20 The mesial margin of the ascending ramus is wide at its root (Fig. 6O-P). The coronoid process
21 is particularly high above the occlusal plane and its dorsal tip forms a sharp crest (Fig. 6P-Q).
22 The condyle is narrow and convex mesio-distally (Fig. 6Q). In posterior view, the condyle is
23 medio-laterally long and medially inclined. The masseteric fossa, in lateral view, is deep in its
24 mesial portion. There is a wide mandibular foramen on the medial aspect of the ascending ramus
25 base (Fig. 6P-Q). The mandibles are characterised by a very elongated angular apophysis,

1
2 1 making a weak angle with ascending ramus. In lateral view, the dorsal outline of the angular
3 process is deeply concave and it forms a pointed tip at the distal most part. The angular
4 apophysis does not extend much ventrally beyond the level of the ventral margin of the
5 horizontal ramus. Almost no variation is observed on the ventral extension of the angular
6 process. The angular apophysis is mesially delimited by a shallow *incisura vasorum*.
7
8
9
10
11
12
13

14
15 6 **Remarks.** The cainotheriid material from DAM1 is referred here to *Paroxacron* based on the
16 presence of i) small diastemata between P1-P2, P2-P3 and P1-P2/, ii) premolariform P1
17 and P2, and iii) a P3/ with a long mesial lobe. Mandibular remains from DAM1 differ from
18 those of *Paroxacron bergeri* in having a more elongated angular apophysis and a shorter angle
19 with the ascending ramus, as well as a smaller and less marked *incisura vasorum* and the
20 presence of diastemata. They also differ in the size and shape of their upper molars which have
21 more triangular outlines, and smaller size.
22
23
24
25
26
27
28
29
30

31 13 The morphology of DAM1 specimens is compatible with the diagnosis of *Paroxacron*
32 *valdense* as proposed by Stehlin (1906) and Hürzeler (1936). The maxilla Mt. 230 from
33 Entreroches (Mormont, Switzerland), lectotype of the species, and the mandibles Mt. 232-233
34 from Cinq-Sous (Mormont, Switzerland) described by Stehlin (1906), as well as maxillae QS.
35
36 15 486 and the mandible QU. 32 from old Quercy collections described by Hürzeler (1936), are
37 indeed morphologically and biometrically very similar. However, as mentioned by Hooker &
38 Weidmann (2000) based on the new material from Eclepens C and Cinq-Sous (Mormont,
39 Switzerland), some diagnostic characters like the diastemata length and the anterior premolar
40 shape, present significant intraspecific variation. This variation is indeed also observed in
41 DAM1 material (see description section above). However, specimens from Cinq-Sous and
42 Eclepens C differ from those from Quercy and Entreroches by smaller dimensions of anterior
43 premolars, by a P4/ and upper molars that are more triangular, producing a more primitive shape
44 according to Hooker & Weidmann (2000). These differences may be related to the slightly older
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 age of the first two sites (Hooker & Weidmann 2000). The differences observed between
2 DAM1 material and *Paroxacron bergeri* and the similarities with *P. valdense* from the type
3 locality of Entreroches allow the DAM1 specimens to be identified as *P. valdense*.

4 Legendre (1980) described two Oxacroninae species, attributed to *Oxacron courtoisii*
5 and *Paroxacron* sp. in the late Priabonian locality of Escamps (MP19, Quercy). According to
6 Legendre (1980), *Paroxacron* sp. from Escamps differs from *Paroxacron valdense* from
7 Mormont in being slightly smaller and in having different mandible shape. However, we think
8 that slight differences encountered with Mormont specimens, are attributable to intraspecific
9 variation. Specimens from Escamps are here considered as conspecific with the DAM1 sample,
10 based on biometrical and morphological characters (Tab. S1-S3; Legendre 1980, tab. 3-4), and
11 are thus attributed to *Paroxacron valdense*. The comparison of DAM1 material with specimens
12 referred to *Paroxacron* from Palembert (MP19) (Quercy locality, collection of Montpellier
13 University) indicates that these specimens could also belong to *P. valdense*.

14 ***Paroxacron* sp.**

15 (Fig. 8)

16 **New material from DAM3.** Three left M3/ (DAM3 01-03); one right M2/ (DAM3 04); one
17 left M2/ (DAM3 05); two right M1/ (DAM3 06-07); one left M1/ (DAM3 08); one right P4/
18 (DAM3 09); one right P1/ (DAM3 10); one right M/3 (DAM3 11); two right M/2 (DAM3 12,
19 DAM3 14); one left M/2 (DAM3 13); two left M/1 (DAM3 15-16); one left DP/4 (DAM3 17);
20 one right DP/4 (DAM3 18); one right P/2 (DAM3 19); one left P/1 (DAM3 20); one left
21 mandibular fragment with M/2 (DAM3 21).

22 **Description (Tab. S4-5; Fig. 8).** The double-rooted P/1 and P/2 are premolariform, elongated
23 and narrow (Fig. 8O-R). The paraconid and protoconid are weakly developed, their protoconid
24 being the higher. The DP/4 present well individualised cusps, a mesial lobe smaller than the
25 distal lobe (talonid) and no mesial cingulid (Fig. 8S-T).

1
2 P1/ is relatively elongated, premolariform, and double-rooted (Fig. 8M-N). The
3 paracone is thicker than the weakly-marked parastyle and metastyle. P4/ is relatively narrow
4 and triangular, with a large paracone distally located with respect to the protocone (Fig. 8A-B).
5 Styles are salient and the disto-lingual cingulum is weak. Upper molars are subtriangular with
6 a loop-like mesostyle (Fig. 8C-H).

7
8 **Remarks.** The presence of small teeth and double-rooted premolariform P1 and P2, elongated
9 and narrow with weakly developed cusps, and a P4/ with a mesial protocone, allows to attribute
10 these specimens to an unnamed species of *Paroxacron*, but which is different from either of the
11 named ones. Specimens described by Blondel (2005) from Aubrelong 1, Ravet (MP21) and
12 Plante 2 (MP22) of Quercy are similar to those of DAM3. However, dimensions are close to
13 those of *Plesiomeryx cadurcensis*, and it is sometimes difficult to differentiate the two species
14 on the basis of isolated upper and lower molars.

15 According to the description of Blondel (2005), *Paroxacron* sp. differs from
16 *Paroxacron valdense* by an anterior lobe of the P3/ similar to that observed in *Plesiomeryx*
17 *cadurcensis*, a more quadrangular M1/ and a longer diastema between P1-P2. The
18 dimensions are also smaller, especially for upper premolars. Owing to the small available
19 sample (with respect to DAM 1 material), it is not possible to test intraspecific variation
20 accurately. Yet, morphological and metrical features of these Oligocene specimens are
21 consistently distinct from those of DAM1. It is therefore possible to consider that the Aubrelong
22 1, Ravet, La Plante 2 and DAM3 sites include a younger species of *Paroxacron* which also
23 differs from *Paroxacron bergeri* by its small size, its less quadrangular molars and the presence
24 of a diastema between P2-P3. This species may be new.

25 ***Paroxacron bergeri* Heissig, 1978**

26 (Fig. 9)

27 Synonymy: 1959 cf. *Cainotherium? elongatum* Filhol (?Berger); Berger, p. 45, pl. 4, fig. 5.

1
2 1 1959 *Cainotherium?* n. sp.; Berger, p. 47, pl. 5, fig. 6–10.
3
4
5

6 2 **Holotype.** A right mandibular fragment I/3-M/3 (BSP 1879XV201), locality of Mouillac in
7
8 3 Phosphorites of Quercy. The precise age and type locality are unknown (Old Quercy
9
10 4 collections).

11
12
13 5 **Diagnosis.** Species of *Paroxacron* without a diastema between P/2-P/3. The edge of the
14
15 6 mandible is more strongly bent than in *P. valdense*. The *angulus mandibulae* is short, high and
16
17 7 angular. It is limited by a short but strong *incisura vasorum* anteriorly. P/3 and P/4 have a more
18
19 8 complete talonid and the internal wall of the lower molars is higher than in *P. valdense* (Heissig
20
21 9 1978).

22
23
24 10 **Differential Diagnosis.** *Paroxacron bergeri* differs from Cainotheriinae species by the absence
25
26 11 of crested styles on the upper molars, a shorter postparacristula, the presence of narrow
27
28 12 premolars and a protocone more mesial than the paracone on P4/. It differs also from
29
30 13 *Plesiomeryx* and *Caenomeryx* species by a small protocone and a longer mesial lobe on P3/,
31
32 14 and the absence of diastemata and protocone on P2/. It distinguishes from *Cainotherium* by the
33
34 15 absence of cingulum/id on premolars and a triangular P3/. *P. bergeri* diverges from the
35
36 16 Oxacroninae *Oxacron courtoisii*, by a larger size, the presence of more elongate and
37
38 17 premolariform premolars (P1 and P2), an elongate anterior lobe on the P3/, as well as a larger
39
40 18 paroconule and pinched mesostyle on the upper molars. *P. bergeri* differs from *P. valdense* and
41
42 19 *P. sp.* by a larger size, quadrangular upper molars, a pinched mesostyle and the absence of
43
44 20 diastemata.

45
46
47 21 **New material from DAM3.** Two left M3/ (DAM3 22-23); one right M3/ (DAM3 24); two left
48
49 22 M2/ (DAM3 25, DAM3 27); one right M2/ (DAM3 26); two left M1/ (DAM3 28-29); one left
50
51 23 P4/ (DAM3 30); one right P3/ (DAM3 31); one left P2/ (DAM3 32); one right maxillary
52
53 24 fragment with P1/ (DAM3 33).

1
2 **1 Description (Tab. S6; Fig. 9).** P1/ and P2/ are elongated and premolariform (Fig. 9I-L). The
3 **2 P3/** presents a well elongated anterior lobe, with salient parastyle and metastyle, and a
4 **3 pronounced protocone** (Fig. 9M-N). The crest connecting these two styles is buccally concave,
5 **4 the disto-buccal lobe being very buccally inclined.** The P4/ shows prominent paracone and
6 **5 mesial protocone**, the paracone being higher (Fig. 9A-B). The parastyle and metastyle are well
7 **6 individualised and salient, and conule 1** presents an extended buccal crest. Upper molars are
8 **7 quadrangular with crested styles** (Fig. 9C-H).

19
20 **8 Remarks.** The presence of quadrangular molars, a P3/ with a very elongated anterior lobe, a
21 **9 pronounced protocone and large size allow for referring these specimens to as** *Paroxacron*
22 **10 bergeri.** These teeth are similar in terms of size and morphology to *Paroxacron cf. bergeri* from
23 **11 Mas de Got (MP22), Pech Crabit, Roqueprune 2, and Mounayne (MP23) localities described**
24 **12 by Blondel (2005).** According to this author, some of these specimens are comparable in size
25 **13 to those referred to** *P. huerzeleri* by Berger (1959), but they differ from the latter species by the
26 **14 morphology of the P2/ (smaller median cusp and presence of a lingual cingulum)** and the P3/
27 **15 (greater anterior lobe and smaller protocone).** The same observations are performed in DAM3,
28 **16 and it is sometimes difficult to differentiate the two species on some isolated molars.**

41 **17 Subfamily Cainotheriinae Camp & VanDerHoof, 1940**

42
43 **18 Emended diagnosis.** Small to large cainotheriids, comprised around 500g (e. g. *Plesiomeryx*
44 **19 cadurcensis**) – 2,4 kg (e. g. *Cainotherium laticurvatum*). Muzzle elongated and narrow (e. g.
45 **20 *Plesiomeryx***) to short and wide (e. g. *Caenomeryx*); Orbita large; ethmoidal fissures present.
46
47 **21 Diastemata present (except** *Cainotherium***), between P1-P2 and C-P1/. Premolars wide;**
48 **22 postprotoconulid large; protocone at the same level than the paracone on P4/; P1 uniradiculate.**
49
50 **23 Upper molars quadrangular; styles crested and mesostyle pinched; postparacristula buccally**
51
52 **24 long and joins the preprotocrista.**

1
2 1 Genus *Plesiomyx* Gervais, 1873
3
4 2 *Plesiomyx cadurcensis* Gervais, 1873
5
6 3 (Fig. 10)

7
8 4 **Holotype.** Right mandibular P/2-M/3 (MNHN-QU 1772). Mouillac locality, Phosphorites of
9 5 Quercy. The precise age and type locality are unknown (Old Quercy collections).

10
11 6 **Diagnosis.** Protocone and paraconule present (stage B, Hürzeler, 1936: 8). Long diastema
12 7 between P1/ and C and between P/1 and P/2. P1/, P2/, P/2 and P/3 are short, P3/ with a strong
13 8 lingual cone, P2/ with a weak lingual cone. The premaxilla reaches the frontal. The nasal is
14 9 long, narrow and straight. The ethmoidal slit is falciform. The choanae are large. The horizontal
15 10 portion of the lower jaw is slender (Hürzeler, 1936).

16
17 11 **Differential Diagnosis.** *Plesiomyx cadurcensis* differs from Oxacroninae species by the
18 12 presence of quadrangular upper molars with a pinched mesostyle (except *P. bergeri*) and crested
19 13 styles, a longer postparacristula, larger premolars, a larger postprotoconulid, a P/1
20 14 uniradicate, diastemata between P/1-P/2 and C-P1/, and a protocone at the same level than
21 15 the paracone on P4/. It differs also from *Paroxacron* species by a shorter anterior lobe on the
22 16 P3/ and a larger protocone on the P3/ and P2/. *Ple. cadurcensis* diverges from *Oxacron* by a
23 17 larger protocone on the P3/ and P2/, as well as premolariform mesial premolars. *Ple.*
24 18 *cadurcensis* diverges from *Cainotherium* by a smaller size, a smaller paraconule on the upper
25 19 molars, the absence of cingulum/id on premolars, the presence of a triangular P3/, a larger
26 20 protocone on the P3/ and P2/, a valley between the entoconid and the mesoconid on lower
27 21 molars, and the presence of diastemata. It distinguishes from *Caenomeryx* species by a smaller
28 22 size, a smaller protocone on the P3/ and P2/, a shorter mesial lobe on the P3/ and a smaller
29 23 paraconule. Finally, *Ple. cadurcensis* differs from *Ple. huerzeleri* by a smaller size and a labial
30 24 cone of the P/4 more distal than the lingual cone.

1
2 1 **New material from DAM3.** One left M3/ (DAM3 34); two right M3/ (DAM3 35-36); two left
3 2 M2/ (DAM3 37, DAM3 40); three right M2/ (DAM3 38-39, DAM3 41); two right M1/ (DAM3
4 3 42, DAM3 44); one left M1/ (DAM3 43); one right DP4/ (DAM3 45); one left DP4/ (DAM3
5 4 46); one right P4/ (DAM3 47); one left P3/ (DAM3 48); one right P2/ (DAM3 49); one right
6 5 P1/ (DAM3 50); one right maxillary fragment with P2/ (DAM3 51); two left M3/ (DAM3 52,
7 6 DAM3 54); one right M3/ (DAM3 53); three right M2/ (DAM3 55, DAM3 57-58); two left
8 7 M2/ (DAM3 56, DAM3 59); two left M1/ (DAM3 60, DAM3 62); three right M1/ (DAM3 61,
9 8 DAM3 63-64); three right P4/ (DAM3 65, DAM3 67-68); one left P4/ (DAM3 66); four right
10 9 P3/ (DAM3 69-70, DAM3 72, DAM3 74); two left P3/ (DAM3 71, DAM3 73); two left P2/
11 10 (DAM3 75, DAM3 76); one right P2/ (DAM3 77); one left mandibular fragment with M1/
12 11 (DAM3 78); one right mandibular fragment with P3-P4 (DAM3 79).

13 12 **Description (Tab. S7-8; Fig. 10).** P1/ is uniradicate and probably premolariform (only
14 13 alveoli are preserved). The diastema that separates P1/ from P2/ is short. The double-rooted
15 14 P2/, P3/, and P4/ are short and wide (Fig. 10S-V). P2/ and P3/ present a strong protoconid. P3/
16 15 shows a short anterior lobe, a weak lingual cingulum, as well as an expanded distal cristid
17 16 lingually curved. The protoconid of the P4/ is more mesial than the metaconid. The
18 17 postprotoconulid is more or less large. The mandible bears a mandibular symphysis located on
19 18 the distal margin of P2/ (Fig. 10U-V). A mental foramen is present, below the mesial part of
20 19 P2/. The horizontal ramus is gracile and shallow.

21 20 P1/ is premolariform, short, wide, and biradicular (Fig. 10M-N). P2/ is more elongated
22 21 and wider than P1/ (anterior lobe as wide as posterior lobe). It has a weakly pronounced
23 22 protocone and a large paracone (Fig. 10O-P). P3/ is short with a well-developed protocone (Fig.
24 23 10A-B). The paracone and the protocone of P4/ seem to be median (in the same plan), but the
25 24 only preserved tooth is strongly worn (Fig. 10C-D). Upper molars are quadrangular in occlusal
26 25 view, with a pinched mesostyle, styles being very weakly crested, and a small paraconule (Fig.

1 10E-J). The DP4/ are similar to upper molars, but present an important mesially convex
2 curvature, with a mesially projected parastyle (Fig. 10Q-R).

3 **Remarks.** The presence of a uniradiculate and premolariform P/1, of a short diastema between
4 P/1-P/2, of a strong protoconid on P/2 and P/3 (Blondel, 2005), and of upper molars with a
5 small paraconule point to a representative of *Plesiomeryx*. Its small size and premolar
6 morphology match those of *Plesiomeryx cadurcensis*, as described by Hürzeler (1936), Berger
7 (1959), and Blondel (2005) from Aubrelong 1, Ravet (MP 21), Mas de Got, La Plante 2 (MP
8 22), Pech Crabit, Mounayne et Itardies (MP23, Quercy localities). Even if some isolated upper
9 and lower molars are hardly distinguishable from those of *Paroxacron* sp. from DAM3, a
10 slightly taller size, more quadrangular upper molars and a more pinched mesostyle in *P.*
11 *cadurcensis* may allow to distinguish them.

***Plesiomeryx huerzeleri* Berger, 1959**

(Fig. 11)

14 Synonymy: 1937 *Caenotherium gracile* Pomel; Dehm, p. 352.

15 **Holotype.** Skull BSP 1952 II 1149. Gaimersheim fissure filling (Germany) (MP28).

16 **Diagnosis.** Species of *Plesiomyx* similar to *P. cadurcensis*, but larger. The fronto-nasal suture
17 is short. The lingual cingulum of P/3 is missing. The lingual and labial cones of P/4 reach the
18 same level (Berger 1959).

19 **Differential Diagnosis.** *Plesiomeryx huerzeleri* differs from other Cainotheriidae species in the
20 same way as that is described for *Plesiomeryx cadurcensis*, but present a larger size than
21 *Paroxacron valdense*, *Paroxacron*. sp. and *Oxacron courtoisii*. *Ple. huerzeleri* distinguishes
22 from *Ple. cadurcensis* by a larger size, as well as the labial and lingual cones of the P4/ that
23 reach at the same level.

1
2 1 **New material from DAM3.** Three right M3/ (DAM3 80-82); four right M2/ (DAM3 83-85,
3 2 DAM3 88); two left M2/ (DAM3 86-87); three left M1/ (DAM3 89, DAM3 94-95); six right
4 3 M1/ (DAM3 90-93, DAM3 96-97); two left P2/ (DAM3 98-99); one left P1/ (DAM3 100); one
5 4 left maxillary fragment with P3/-P4/ (DAM3 101); two right M3 (DAM3 102-103); two left
6 5 M3 (DAM3 104-105); two left M2 (DAM3 106, DAM3 109); two right M2 (DAM3 107-
7 6 108); two left M1 (DAM3 110-111); two right M1 (DAM3 112-113); one left DP4
8 7 (DAM3 114); three left DP3 (DAM3 115-117); four right P3 (DAM3 118-119, DAM3 121,
9 8 DAM3 125); four left P3 (DAM3 120, DAM3 122-124); one right P2 (DAM3 126); one right
10 9 P1 (DAM3 127); one left mandibular fragment with P4-P3 (DAM3 128).
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

10 **Description (Tab. S9-S10; Fig. 11).** P1/ is uniradiculate and premolariform (Fig. 11S-T). A
11 short diastema is present between P1/ and P2/ (Fig. 11W-X). The double-rooted P2, P3 and
12 P4 are short and wide (Fig. 11U-X). P2/ and P3/ present a strong protoconid. In lateral view,
13 the protoconid of the P4 is at the same level than the metaconid. The postprotoconulid is more
14 or less pronounced. DP4 and DP3 are elongated, cusps being well individualised and cingulids
15 absent (Fig. 11Y-B'). The mandible bears a mandibular symphysis located on the distal margin
16 of P2/ (Fig. 11W-X). A mental foramen is present, below the mesial part of P2/. The horizontal
17 ramus is relatively gracile on its mesial part.

18 P1/ and P2/ are short and wide with a large paracone (Fig. 11O-R). P2/ bears a weakly
19 developed protocone. The triangular P3/ bears a strong protocone, and the parastyle and
20 metastyle are not very prominent (Fig. 11A-B). The mesial lobe is quite short and the crest
21 connecting the metastyle and parastyle is weakly buccally concave. P4/ is wide and rectangular
22 in occlusal view, the paracone and the protocone being median. Upper molars are quadrangular
23 with a small paraconule and a pinched mesostyle, styles being weakly crested (Fig. 11C-H).

24 **Remarks.** A larger size than in *Plesiomeryx cadurcensis*, a lack of lingual cingulum in P3/ and
25 buccal and lingual cones at the same level in P4/, allow some specimens to be assigned to

1 *Plesiomyx huerzeleri*. The morphology and size are very similar to those of Quercy specimens
2 described by Blondel (2005) from Mas de Got, La Plante 2 (MP 22), Pech Crabit, Roqueprune
3 2 and Mounayne (MP 23). They are also similar to those specimens from Gaimersheim (MP28,
4 Germany; Berger 1959, tab. 3, pl. 3; Bolliger 2013, fig. 3a). Some isolated upper molars are
5 however hardly differentiated from those of *Paroxacron bergeri*. Slightly smaller size, a more
6 quadrangular shape and a more pinched mesostyle in *Ple. huerzeleri* may allow them to be
7 discriminated.

Genus *Caenomeryx* Hürzeler, 1936

Caenomeryx procommunis (Filhol, 1877)

Synonymy: 1877 *Cainotherium procommune* Filhol: 212, fig. 269–270.
1936 *Caenomeryx procommunis* (Filhol), Hürzeler, p. 100.

Holotype. Fragment of mandible bearing P/2-M/3 (Filhol collections, MNHN). Precise age and type locality unknown (Old Quercy collections).

Emended Diagnosis. Upper molars are wider than long. Protocone and protoconule present (stage B, Hürzeler, 1936; p. 8). Diastema between P1/ and C is short. Diastema very short or missing between P1/ and P2. Premolars reduced, P3/ with a strong lingual cone, P2/ with a weaker lingual cone, P1/ very small. The premaxilla does not reach the frontal. The wide nasal bones broaden behind the posterior extremities of the premaxilla. The ethmoidal gap is irregular, the choanae are large and the mandible is high, stocky and massive (Hürzeler, 1936).

Remarks: According to our observations, the C-P1/ and P1/-P2 diastemata are rather short than absent (*contra* the original diagnosis of Hürzeler (1936)).

Differential Diagnosis. *Caenomeryx procommunis* differs from Oxacroninae species by a larger size, the presence of quadrangular upper molars with a pinched mesostyle (except *P. bergeri*) and crested styles, a longer postparacristula, larger premolars, a larger

1 postprotoconulid, a P/1 uniradiculate, diastemata between P/1-P/2 and C-P1/, and a protocone
2 at the same level than the paracone on P4/. It differs also from *Paroxacron* species by a shorter
3 anterior lobe on the P3/ and a larger protocone on the P3/ and P2/. *Cae. procommunis* diverges
4 from *Oxacron* by the presence of a large protocone on the P3/ and P2/, and premolariform
5 mesial premolars. *Cae. procommunis* distinguishes from *Cainotherium* by rectangular upper
6 molars, the absence of important cingulum/id on premolars, the presence of a triangular P3/, a
7 larger protocone on the P3/ and P2/, a valley between the entoconid and the mesoconid on lower
8 molars, and the presence of diastemata. *Cae. procommunis* diverges from *Plesiomeryx* species
9 by a larger size, a larger protocone on the P3/ and P2/, a longer mesial lobe on the P3/, a larger
10 paraconule, smaller diastemata and more rectangular upper molars. Finally, *Cae. procommunis*
11 differs from *Cae. filholi* by a longer mesial lobe on the P3/ and shorter diastemata.

Caenomeryx cf. procommunis (Filhol, 1877)

(Fig. 12)

14 **New material from DAM3.** One right M3/ (DAM3 129); two left M2/ (DAM3 130-131); one
15 right M2/ (DAM3 132); two left M1/ (DAM3 133-134); one right P3/ (DAM3 135); four left
16 M3 (DAM3 136-139); one left M2 (DAM3 140); one right M1 (DAM3 141); one left M1
17 (DAM3 142); one right P4 (DAM3 143); one left P4 (DAM3 144); two right DP3
18 (DAM3 145-146); one right P3 (DAM3 147); one left P2 (DAM3 148); one left P1
19 (DAM3 149); one right mandibular fragment with M2 (DAM3 150); one left mandibular
20 fragment with P4-M1 (DAM3 151); one right mandibular fragment with P4-M3
21 (DAM3 152).

22 **Description (Tab. S11-S12; Fig. 12).** P/1 is premolariform and single-rooted (Fig. 12M-N).
23 P/2, P/3, and P/4 are short and wide (Fig. 12K-L, 12O-R). P/2 and P/3 present a pronounced
24 paraconid; P/3 has a strong protoconid. DP/3 is elongated mesiodistally with a large mesial part
25 (Fig. 12I-J). The mandible bears a mandibular symphysis located on the distal margin of P/2

1 (Fig. 12K-L). A mental foramen is present, below the distal part of P/4. The horizontal ramus
2 is wide, dorso-ventrally extended and its height increase between the premolar area and the last
3 molar.

4 The anterior lobe of the P3/ is relatively elongated mesio-distally and the crest
5 connecting the parastyle and metastyle is buccally concave (Fig. 12A-B). The metastyle is well-
6 marked and the protocone, althought broken away, seems to be strong. Upper molars are
7 quadrangular in occlusal view (Fig. 12C-H). They are rather bucco-lingually elongated and they
8 have a pinched mesostyle, styles being crested.

9 **Remarks.** The presence of large specimens, with a wide and tall corpus mandibulae, as well as
10 short premolars with large cusps and rectangular upper molars (Hürzeler, 1936), suggests the
11 occurrence of *Caenomeryx* within the DAM3 material. The morphology is similar to the
12 undated specimens of the old Quercy collections attributed to *Caenomeryx procommunis* by
13 Hürzeler (1936). It is also similar to the specimens later attributed to *Cae. cf. procommunis* by
14 Blondel (2005) from Mas de Got (MP22), Pech Crabit, Itardies, Mounayne and Roqueprune 2
15 (MP23), and by Sudre (1995) from Le Garouillas (MP25). The dimension of DAM3 specimens
16 are however a little smaller than previously described specimens from these localities.
17 Moreover, it may be difficult to discriminate some isolated upper molars from those of *P.*
18 *bergeri* and *Ple. huerzeleri*. In any case, a larger size, a more rectangular shape on the upper
19 molars and a larger protocone on the premolars in *Caenomeryx* may allow to distinguish them.

20 Class **Mammalia** Linnaeus, 1758

21 Order **Artiodactyla** Owen, 1848

22 Superfamily **Cainotherioidea** Camp & VanDerHoof, 1940

23 Family **Robiacinidae**, fam. nov.

24 **Type genus.** *Robiacina* Sudre, 1969

1
2 1 **Included genera.** Type genus only.
3
4

5 2 **Original Diagnosis.** As for its type and only genus *Robiacina* (Sudre 1978).
6
7

8 3 **Emended Diagnosis.** Small Cainotherioidea similar in size to *Oxacron*. Skull globular in shape
9 4 with a sub-spherical cerebral part and muzzle extremely short; very large orbits; lacrymal small;
10 5 in the bottom of the orbital cavity, palatine does not contact the frontal, due to the great posterior
11 6 development of the lacrimal; auditory bullae of large size, sub-spherical; ethmoid fissure absent.
12 7 Lower and upper premolars short mesio-distally; diastemata absent to reduced; P/3 with a small
13 8 medio-lingual cusp; P/1 and P/2 unicuspid. Lower molars with a pinched trigonid and a wide
14 9 talonid; presence of a small to medium-sized metaconulid on the distal aspect of the metaconid;
15 10 entoconid slightly more distal than the hypoconid; presence of a small hypoconulid on M/1-2
16 11 and a posthypocristulid on M/3. Upper molars subtriangular with five cusps and a central or
17 12 slightly mesial protocone; metacone higher than the paracone; paraconule small; preparacrista
18 13 short; preprotocrista linked to the paraconule; buccal wall of the paracone with or without a
19 14 cingulum.
20
21

22 15 **Remarks.** The justification for the erection of the new family Robiacinidae is provided in the
23 16 following section and ensues the results of phylogenetic analysis.
24
25

26 18 **Results and Discussion** 27 28

29 20 **Dental nomenclature** 30 31

32 21 Premolar and molar cusp/id patterns of Cainotheriidae differ from other Artiodactyla,
33 22 which resulted in different dental nomenclatures for premolars and lower molars (Fig. 1A-E,
34 23 I). According to Berthet (2003), the disto-lingual cusp of upper premolars corresponds to an
35 24 hypocone while, according to Hürzeler (1936), Gentry & Hooker (1988), and Boisserie *et al.*

(2010), this cusp would correspond to the protocone instead. For lower premolars, these authors identify the mesial cuspid as the paraconid and the medio-lingual cuspid of P/4 as a metaconid. Berthet (2003) considers the mesial cuspid of lower premolars as a protoconulid, and the medio-lingual cuspid of P/4 an entoconid. Problems of identification of the cusps/ids also occur for the lower molars. According to Hürzeler (1936), the three lingual cusps of lower molars correspond to: a mesial cuspid identified as a metaconid, and two distal ones, that would correspond to the entoconid and to the hypoconulid that would lie in a lingual position. According to Sudre (1977), the medio-lingual cuspid would not be an entoconid but a neoformation called the mesoconid. This mesoconid would originate from a duplication of the metaconid (Sudre 1977). Gentry & Hooker (1988) and Hooker & Weidmann (2000) identify the medio-lingual cuspid as a metastylid, originating from the cingulid. In this framework, the entoconid would also be placed distally.

We based our hypotheses of dental cusp(id) homologies of cainotheriids on the examination of: i) the genus *Robiacina* whose cranial features and dental pattern are very close to those of Eocene cainotheriids, and ii) the earliest documented cainotheriid, *Oxacron courtoisii* (Blondel 2005). According to Sudre (1969, 1977), *Robiacina* would be phylogenetically close to Cainotheriidae. Gentry & Hooker (1988) considered the genus as part of Cainotheriidae.

Concerning the premolars, the disto-lingual cusp of the upper premolars of *Oxacron* and *Robiacina* does not derive from the cingulum, and therefore is identified here as the protocone. For lower premolars, we consider that the medio-lingual cuspid of P/4 represent the metaconid and that the distal cuspid is the entoconid, by homology of position with lower molars. This corresponds to the nomenclature proposed by Gentry & Hooker (1988) and Boissierie *et al.* (2010).

1
2
3 1 For the lower molars, we identify the medio-lingual cuspid as a neoformed structure, in
4
5 agreement with Sudre (1977), based on the presence of a small neoformed accessory cuspid at
6
7 the base of the metaconid, in the lower molars of *Robiacina* (Fig. 1F, 1H). Indeed, the
8
9 hypothesis of a duplication of the metaconid proposed by Sudre (1977) seem to be more
10
11 parsimonious than a medio-lingual position of the entoconid and a concurrent lingual
12
13 displacement of the hypoconulid as hypothesised by Hürzeler (1936). The latter hypothesis
14
15 would imply a fusion/disappearance of the anterior and/or posterior crests of the talonid
16
17 crescent, with a mesial migration of the entoconid and a lingual displacement of the
18
19 hypoconulid. Moreover, according to this hypothesis, the entoconid would be located in front
20
21 of the protocone during the occlusion with upper molars, contrary to most other artiodactyls
22
23 (Sudre 1977). Finally, if Hürzeler's (1936) hypothesis is considered, with the hypoconulid in
24
25 disto-lingual position on M/1 and M/2, what would then be the distalmost cuspid on the M/3?
26
27 As concluded by Sudre (1977), the distalmost cuspid on M/3 is necessarily the hypoconulid. A
28
29 hypoconulid is clearly visible on the M/1 and M/2 of *Robiacina*, positioned just distal to the
30
31 entoconid (Fig. 1F). This disposition of the hypoconulid relative to the entoconid is also found
32
33 on M/3 of Cainotheriidae, which lack a hypoconulid on M/1 and M/2. Finally, the hypothesis
34
35 of a metastyloid as medio-lingual cuspid, proposed by Gentry & Hooker (1988) and Hooker &
36
37 Weidmann (2000), is here rejected as it does not originate from the cingulid, neither in
38
39 *Robiacina*, nor in *Oxacron*. Contrary to Sudre (1977), we use the term 'metaconulid' instead of
40
41 'mesoconid', in order to homogenized with the nomenclatural system of Boisserie *et al.* (2010),
42
43 and because mesoconid is used for another structure in other mammalian group (Klingener
44
45 1964; Skwara 1986; Hooker & Russell 2012).

46
47
48
49
50
51 23 Thus, according to the hypotheses of Sudre (1977; molars) and of Gentry & Hooker
52
53 (1988) and Boisserie *et al.* (2010; premolars), the three lingual cusps of lower molars would
54
55 correspond to the metaconid (mesial), the metaconulid (medial, originates from metaconid
56
57
58
59
60

1 duplication), and the entoconid (distal). The disto-lingual cusp of upper premolars is here
2 identified as the protocone, and the mesial cuspid of lower premolars as the paraconid. The P/4
3 bears a metaconid (medio-lingual) and an entoconid (one distal).

5 Phylogenetic analysis

6 The heuristic search of the parsimony analysis retained a single parsimonious tree (L =
7 170, CI = 0.48, RI = 0.73; CI with only parsimony-informative characters = 0.46) (Fig. 13),
8 reflecting a well-structured signal and relatively weakly homoplastic characters. Cainotheriidae
9 are monophyletic and split into two subgroups, interpreted as subfamilies: Oxacroninae, with
10 *Oxacron* and *Paroxacron*; and Cainotheriinae, with *Plesiomeryx*, *Caenomeryx*, and
11 *Cainotherium*. Cainotheriidae, together with Mixtotheriidae, Anoplotheriidae and
12 representatives of *Robiacina* form a clade, sister group to ruminants. Tylopoda split earlier in
13 the tree, together with *Amphimeryx* and *Xiphodon*.

14 The monophyly of Cainotheriidae is supported by 13 non-ambiguous synapomorphies
15 and a high Bremer index (IB = 7). They are characterised by the presence on their upper molars
16 of a large parastyle similar to other styles (8¹; IR = 0.67), a disto-lingual protocone (11²; IR =
17 0.94), a very strong dilambdodonty (20¹; IR = 0.88), a long (26¹; IR = 0.40) and distally
18 orientated (27¹; IR = 1.00) postprotocrista, a postparacristula (29¹; IR = 0.86) and the lack of
19 preprotocrista and paraconule junction (15¹; IR = 0.67); on their lower molars, they present a
20 large (34²; IR = 1.00) and individualised (35⁰; IR = 1.00) metaconulid, a very distal entoconid
21 and close to the hypoconulid (40²; IR = 1.00), a neotrigonid (47¹; IR = 1.00), the absence of
22 posthypocristulid (50⁰; IR = 0.67) and hypoconulids on M/1 and M/2 (38⁰; IR = 0.75).
23 Cainotheriidae are composed of two clades: the Oxacroninae, characterised by the presence of
24 styles without crests on upper molars (7⁰; IR = 0.44), a protocone more mesial than the paracone
25 on P4/ (56¹; IR= 0.50), as well as a biradiculate P/1 (70¹; IR = 1.00); Cainotheriinae, supported

1 by the presence of a postparacristula buccally long joining the preprotocrista on upper molars
2 (30¹; IR = 1.00)³⁰ and of a large postprotoconulid on lower premolars (69¹; IR = 1.00). They
3 further differ from Oxacroninae by the presence of a uniradiculate P1 (70⁰; IR= 1.00) and large
4 premolars (73¹; IR = 0.67), which represent plesiomorphic states of characters.

5 Within Oxacroninae, *Paroxacron* is paraphyletic. *Paroxacron bergeri* is excluded from
6 the (*Paroxacron valdense*, *Oxacron courtoisii*) clade by the presence of quadrangular molars
7 (31²; IR = 0.60) with a pinched mesostyle (10¹; IR = 1.00) and the absence of diastemata
8 between P1/-P2/ (5⁰; IR = 0.50) and P/2-P/3 (6⁰; IR = 0.33).

9 Within Cainotheriinae, *Cainotherium* is sister taxon to the clade (*Caenomeryx*,
10 *Plesiomeryx*). *Cainotherium laticurvatum* is characterised by numerous autapomorphies, such
11 as the absence of a valley between the entoconid and the metaconulid on lower molars (41⁰; IR
12 = 0.78), the presence of a protocone at the same level (bucco-lingual axis) as the paraconule on
13 the M3/ (17⁰; IR= 0.50), the absence of a triangular P3/ (quadrangular shape) (61⁰; IR = 0.50)
14 and a wide cingulum/id on the premolars (72¹; IR = 0.50)⁷². The clade (*Caenomeryx*,
15 *Plesiomeryx*) is supported by the presence of a protocone on P2/ (64¹; IR = 0.57) and diastemata
16 between C-P1/ (3¹; IR=0.75) and P/1-P/2 (5¹; IR = 0.86).

17 *Robiacina* is sister taxon to the Cainotheriidae clade. This position is supported by the
18 presence of a more distal (11 ;IR = 0.94) and lingual protocone (17¹; IR = 0.50) on upper molars
19 and M3/, respectively, but also by sub-equal paracone and metacone on P3/ (60¹; IR = 1.00),
20 the absence of protocone on P2/ (64⁰; IR = 0.57), the presence of a small hypoconid on P/2-P/3
21 (67⁰; IR = 0.75) and of a postprotoconulid on premolars (68¹; IR = 1.00). This result does not
22 support the hypothesis of Sudre (1977) who placed *Robiacina* within Anoplotheriidae, close to
23 Anoplotheriinae. Our results also slightly contrast with the hypothesis of Gentry & Hooker
24 (1988) who placed *Robiacina* within Cainotheriidae. The clade composed of representatives of
25 *Robiacina* is excluded from the Cainotheriidae clade by 13 unambiguous characters (see

support of Cainotheriidae clade above) and it further differs from it by sharing three synapomorphies: sub-triangular upper molars (31^1 ; IR = 0.60), a small paraconule (14^0 ; IR = 0.43), and a short preparacrista (25^1 ; IR = 1.00). This marked morphological divergence between Cainotheriidae and the *Robiacina* clade leads us to recognise this clade as a new family, here named Robiacinidae. It would therefore constitute the closest family to Cainotheriidae, and forms with it the superfamily Cainotherioidea Camp & VanDerHoof, 1940.

Anoplotheriidae and Cainotherioidea form a clade (B) supported by the presence of a metaconulid (33^1 ; IR = 1.00), an entoconid positioned at the back of the hypoconid (40^1 ; IR = 1.00) and a valley between entoconid and metaconulid on the lower molars (41^1 ; IR = 0.78), as well as a connected paracone and metacone on P4/ (55^0 ; IR = 0.50). The Mixtotheriidae, Anoplotheriidae, and Cainotherioidea together form a clade (A) supported by the presence of a very large parastyle (8^0 ; IR = 0.67) and an absent to reduct postprotocrista (26^0 ; IR=0.40) on the upper molars, a ‘loop-like’ entoconid (39^1 ; IR = 1.00) on the lower molars, and the absence of a diastema between P1-P2 (5^0 ; IR = 0.86).

According to our results, Cainotherioidea, Anoplotheriidae and Mixtotheriidae are closer to Ruminantia than to Tylopoda. They share the presence of a protocone and a metaconule at the same level on M3/ (17^0 ; IR = 0.50), crested styles on P3-/P4/ (58^1 ; IR = 0.71), a large lingual cusp mesio-distally on P3/ (63^1 ; IR = 0.56), as well as a large hypoconid on P3 and P2 (67^1 ; IR = 0.75).

Impact on artiodactyl systematics

Cainotheriidae. According to our results (Fig. 13), *Paroxacron* is paraphyletic within Oxacroninae. Hooker & Weidmann (2000) proposed to synonymise the genera *Paroxacron* and *Oxacron*. However, *Oxacron* exhibits three autapomorphic characters: a short mesial lobe on the P3/ (62^0 ; IR = 0.25), a caniniform P1/ (71^0 ; IR= 0.50), and small paraconule (14^0 ; IR=0.43).

1
2 1 We therefore chose to maintain the genus *Oxacron*. *Paroxacron valdense* is the type species of
3 2 *Paroxacron*. Our results raise the question of the erection of a new genus for the species *P.*
4 3 *bergeri*. However, the latter has no dental autapomorphies based on our results. For the time
5 4 being, based on dental data only, we therefore choose to keep *Paroxacron* as a paraphyletic
6 5 assemblage.
7
8

9
10 6 **Robiacinidae, fam. nov.** Our results retrieve a monophyletic assemblage of *Robiacina* species.
11 7 This clade appears as sister taxon to Cainotheriidae. This contrasts with the hypothesis of
12 8 Hooker & Weidmann (2000), according to which *Robiacina lavergnensis* should be referred to
13 9 *Mixtotherium*. Following these authors, *R. lavergnensis* shares many characters with the species
14 10 of *Mixtotherium*: a short premetacristula, a pre-cingulum and crested styles on the upper molars;
15 11 a prehypocristid fused with the postprotocristid, a basined hypoconulid in basin and a high
16 12 crown on the lower molars, and a caniniform lower canine. However, we disagree on the coding
17 13 of some of these characters for *Mixtotherium* by Hooker & Weidmann (2000). According to
18 14 our observations, there is no basin on the hypoconulid, the premetacristula is long and the
19 15 prehypocristid is unfused. Besides, the results of our phylogenetic analyses place *Robiacina* as
20 16 sister taxon to Cainotheriidae, and far apart from *Mixtotherium*. The presence of crested styles
21 17 in *Mixtotherium* and *R. lavergnensis* as mentioned by Hooker & Weidmann (2000) is here a
22 18 symplesiomorphy. Contrary to Hooker & Weidmann (2000), we retain the binomen *Robiacina*
23 19 *lavergnensis* instead of *Mixtotherium lavergnense*.
24
25

26
27 20 **Cainotherioidea.** We propose here to recognize the clade (Robiacinidae + Cainotheriidae) as
28 defining the Cainotherioidea. The concept of this compound was defined by Simpson (1945) as
29 single-family, then Hooker & Weidmann (2000) extended it by including the genera *Robiacina*
30 and *Mixtotherium*. However, and according to our results, we propose here to exclude
31 *Mixtotherium* from the Cainotherioidea.
32
33

34 25
35
36

1 1 Relationships of Cainotheriidae within Artiodactyla

2 The phylogenetic position of Cainotheriidae within Artiodactyla remains controversial.
3 Gentry & Hooker (1988, fig. 9.8) supported the placement of cainotheriids within Tylopoda in
4 their (informal) “preferred analysis”, but retrieved them in a much more basal position, in their
5 formal analysis (Gentry & Hooker 1988, fig. 9.7), i.e. outside of the “Selenodontia” clade (i.e.,
6 Tylopoda + Ruminantia). More recent phylogenetic analyses placed them either close to
7 Tylopoda (e.g., Geisler & Uhen 2003, Geisler *et al.* 2007, Thewissen *et al.* 2007), or as an early
8 offshoot of a wide clade including Ruminantia (e.g., Geisler & Uhen 2005; O’Leary & Gatesy
9 2007, Lihoreau *et al.* 2015). Our results are congruent with the previous studies that excluded
10 Cainotheriidae from Tylopoda (Webb & Taylor 1980), and placed Cainotheriidae closer to
11 Ruminantia (Geisler & Uhen 2005; O’Leary & Gatesy 2007; Lihoreau *et al.* 2015). Indeed,
12 according to our sample, Cainotherioidea, Anoplotheriidae and Mixtotheriidae are sister group
13 to Ruminantia. As far as direct relationships of Cainotherioidea are concerned, according to our
14 results, Anoplotheriidae are their closest relatives.

15 It is worth noting that the nomenclature chosen here has an impact on the primary
16 homology hypotheses underlaying the definition of phylogenetic characters. Indeed, the
17 presence of a metaconulid (33) is retrieved as a synapomorphy of the Anoplotheriidae and
18 Cainotherioidea clade (clade B in Figure 13, cf. below). If we consider an alternative primary
19 homology hypothesis and identify the medio-lingual cuspid of the lower molars as a metastylid
20 (following Gentry and Hooker, 1988 and Hooker and Weidmann, 2000), as for Mixtotheriidae,
21 coding of four characters (33 – 36) would be modified. However, the topology of the tree is not
22 impacted by this new coding.

23 These relationships rely on dental characters only and it is necessary to extend the data
24 matrix with mandibular, cranial and post-cranial characters, in order to test further the

1 phylogenetic relationship of Cainotheriidae. These characters could bring additional support to
2 clades with relatively low Bremer index.

3

4 **Evolutionary history of Cainotheriidae around the Grande Coupure**

5 Dams karstic network is a new exceptional fossiliferous compound locality
6 documenting the evolutionary history of Cainotheriidae around the Grande Coupure (pre- and
7 -post), through MP19 and MP22 fossiliferous infillings. The first documented representatives
8 of Cainotheriidae are Oxacroninae (Fig. 14). They appear in the fossil record during the late
9 Eocene between MP18-MP19 with two species, *Oxacron courtoisii* and *Paroxacron valdense*
10 (Blondel 2005; Hooker & Weidmann 2000; Legendre 1980). Representatives of Cainotheriinae
11 occur shortly after Oxacroninae, around MP19 with *Cainotherium commune?* (Blondel 2005).
12 This diversity of Cainotheriidae during the late Eocene is not recorded at Dams, where only
13 *Paroxacron valdense* is documented in DAM1 (MP19). However, Dams highlights a second,
14 post-Grande Coupure, phase of diversification of Cainotheriidae, as suggested – based on other
15 localities – by Blondel (2005), that took place at the beginning of the Oligocene. Indeed, at least
16 five species are retrieved in DAM3 (MP22) and this specific diversity is also observed in other
17 fossiliferous localities of Quercy such as Pech Crabit (six species) and Mas de Got (four species;
18 Blondel 2005). The phase of diversification occurs mainly in Cainotheriinae, with the
19 differentiation of five species belonging to *Plesiomeryx* (*Ple. cadurcensis* and *Ple. huerzeleri*),
20 *Caenomeryx* (*Cae. cf. procommunis* and *Cae. filholi*) and *Cainotherium* (*Cainotherium* sp.)
21 (Fig. 14). Among Oxacroninae, two new species also appear in the fossil record after the Grande
22 Coupure: *Paroxacron* sp. and *Paroxacron bergeri*. The two subfamilies of Cainotheriidae thus
23 survive to the Grande Coupure.

24 The Eocene-Oligocene transition further documents a body size increase in
25 Cainotheriidae, as shown in *Plesiomeryx huerzeleri*, *Paroxacron bergeri* but also

1 representatives of *Caenomeryx* and *Cainotherium* (see Tab. S1-S12). However, small species
2 are still present after the Grande Coupure with *Paroxacron* sp. and *Plesiomeryx cadurcensis*.
3 This reflects an increase in the size range of Cainotheriidae after the Grande Coupure, and thus
4 likely an expansion in the occupation of different ecological niches. It is probable that the
5 Cainotheriidae were quite generalist in terms of diet, and that they took the ecological niches
6 released by endemic families of Europe which had became extinct at this transition, such as
7 Xiphodontidae, Dacrytheriinae, or Mixtotheriidae (Blondel 2001).

8 The Grande Coupure coincides with a major mammalian faunal turnover concurrent to
9 climatic, geographic, and oceanic circulation changes (Legendre 1987; Berggren & Prothero
10 1992; Akhmetiev & Beniamovski 2009). The increase of the chewing surface, with the
11 appearance of more quadrangular molars, and the reduction of diastemata as observed in some
12 species of Oligocene Cainotheriidae (notably in Dams) could be linked to changes in their diet
13 (Fig. 14; Blondel 2001). Environmental changes were drastic during the concerned interval,
14 with a shift from a rainforest under tropical/sub-tropical conditions to open environments under
15 a much drier climate (Collinson 1992; Cavagnetto & Anadón 1996; Akhmetiev & Beniamovski
16 2009).

18 Conclusion

19
20 Dams is an exceptional fossil site that yields distinct deposits broadly bracketing the
21 Grande Coupure. Cainotheriid remains are documented by thousands specimens and as such,
22 they allow for in-depth insights into the history and systematics of this family. In the DAM1
23 deposits (MP19, late Eocene), a single species occurs: *Paroxacron valdense*. The DAM3
24 channel (MP22, early Oligocene) yields a higher taxonomic diversity with at least five species:
25 *Paroxacron* sp., *Paroxacron bergeri*, *Plesiomeryx huerzeleri*, *Plesiomeryx cadurcensis*, and

1
2 1 *Caenomeryx* cf. *procommunis*. Dams localities highlight a phase of local diversification of
3 2 cainotheriids by early Oligocene times, thus confirming the hypothesis of Blondel (2005).
4
5 3 Among Cainotheriidae, this diversity increase mainly concerns Cainotheriinae. Yet, both
6
7 4 Cainotheriinae and Oxacroninae survive to major environmental changes related to the Grande
8
9 5 Coupure, contrary to many other European endemic artiodactyls.

10
11
12
13
14 6 Our phylogenetic analysis, including Dams taxa described in this work, constitutes the
15
16 7 first formal phylogeny of Cainotheriidae. Our results, based on dental characters, allow for (i)
17
18 8 clarifying relationships within Cainotheriidae, with Oxacroninae as sister group to
19
20 9 Cainotheriinae, (ii) defining *Robiacina* (and the new family Robiacinidae) as being the sister
21
22 10 taxon to Cainotheriidae among Cainotherioidea and, more broadly, (iii) discussing the
23
24 11 controversial position of Cainotheriidae within Artiodactyla, as being more closely related to
25
26 12 Ruminantia than to Tylopoda.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgments

16 15 We are grateful to all the team of the Cloup d'Aural and to the Quercy research team
17 16 (Palevoprim, Poitiers; M. Godinot, MNHN, Paris; S. Couette, EPHE, Dijon; Margot Bernardi,
18 17 EPHE, Dijon; ISEM, Montpellier; Christian Bousquet, Cloup d'Aural) for their work in the
19 18 field. We thank Christine Bibal for the bibliographic research she has done. We also thank
20 19 Suzanne Jiquel and Bernard Marandat for their availability and help to obtain the specimens
21 20 from the University's collections. We are also grateful to Sylvie Agret and Laurent Marivaux,
22 21 for kindly welcoming in their office for the measurements and photos of the material. Finally,
23 22 we acknowledge the anonymous reviewers for their constructive comments. This work was
24 23 financially supported by the ANR program DEADENDER (ANR-18-CE02-0003-01) - PI M.J.
25 24 Orliac.

1 1 References

- 2 2
- 3 3 **Aguilar, J.-P., Legendre, S. & Michaux, J.** 1997. Biochronologie mammalienne du cénozoïque en Europe et
4 4 domaines reliés. Pp. 1-818 in J. P. Aguilar, S. Legendre & J. Michaux (eds) *Actes Du Congrès*
5 5 *BiochroM'97*. Mémoires et Travaux de l'Institut de Montpellier de l'Ecole Pratique des Hautes Etudes,
6 6 Montpellier.
- 7 7 **Akhmetiev, M. A. & Beniamovski, V. N.** 2009. Paleogene floral assemblages around epicontinental seas and
8 8 straits in Northern Central Eurasia: proxies for climatic and paleogeographic evolution. *Geologica Acta: an international earth science journal*, **7**(1-2), 297-309.
- 9 9
- 10 10 **Antoine, P. A.** 2002. *Phylogénie et évolution des Elasmotheriinae (Mammalia, Rhinocerotidae)*. Mémoires du
11 11 Muséum national d'Histoire naturelle, **188**, 1-359.
- 12 12 **Berger, F. E.** 1959. Untersuchungen an Schädel und Gebissresten von Cainotheriidae. *Palaeontographica*,
13 13 *Abteilung A*, **112**, 1-58.
- 14 14 **Berggren, W. A. & Prothero, D. R.** 1992. Eocene-Oligocene climatic and biotic evolution: an overview. Pp. 1-
15 15 28 in W. A. Berggren & D. R. Prothero (eds) *Eocene-Oligocene climatic and biotic evolution*. Princeton
16 16 University Press, Oxford.
- 17 17 **Berthet, D.** 2003. *Le genre Cainotherium (Mammalia, Artiodactyla): étude morphométrique, révision*
18 18 *systématique, implications évolutives et paléobiogéographiques, extinction*. Unpublished PhD thesis,
19 19 Lyon 1, 204 pp.
- 20 20 **Blondel, C.** 2001. The Eocene-Oligocene ungulates from Western Europe and their environment.
21 21 *Palaeogeography, Palaeoclimatology, Palaeoecology*, **168**(1-2), 125-139.
- 22 22 **Blondel, C.** 2005. New data on the Cainotheriidae (Mammalia, Artiodactyla) from the early Oligocene of south-
23 23 western France. *Zoological Journal of the Linnean Society*, **144**(2), 145-166.
- 24 24 **Boisserie, J. R., Lihoreau, F., Orliac, M. J., Fisher, R. E., Weston, E. M. & Ducrocq, S.** 2010. Morphology
25 25 and phylogenetic relationships of the earliest known hippopotamids (Cetartiodactyla, Hippopotamidae,
26 26 Kenyapotaminae). *Zoological Journal of the Linnean Society*, **158**(2), 325-366.
- 27 27 **Boivin, M., Orliac, M. J., Antunes, M. T., Godinot, M., Laurent, Y., Marandat, B., Vidalenc, D. & Tabuce,**
28 28 **R.** 2018. New material of Diacodexis (Mammalia, Artiodactyla) from the early Eocene of Southern
29 29 Europe. *Geobios*, **51**(4), 285-306.

- 1
2 1 **Bolliger, T.** 2013. Late Oligocene micromammal teeth (MP28/29) from the Lower Freshwater Molasse (USM) of
3 2 Chrummorge (Lägern, N-Switzerland). *Swiss Journal of Geosciences*, **106**(2), 147-159.
4
5 3 **Cavagnetto, C. & Anadón, P.** 1996. Preliminary palynological data on floristic and climatic changes during the
6 4 Middle Eocene-Early Oligocene of the eastern Ebro Basin, northeast Spain. *Review of Palaeobotany and*
7 5 *Palynology*, **92**(3-4), 281-305.
8
9 6 **Collinson, M. E.** 1992. Vegetational and floristic changes around the Eocene/Oligocene boundary in Western and
10 7 Central Europe. Pp. 437-450 in W. A. Berggren & D. R. Prothero (eds) *Eocene-Oligocene climatic and*
11 8 *biotic evolution*. Princeton University Press, Oxford.
12
13 9 **Emry, R. J.** 1978. A new hypertragulid (Mammalia, Ruminantia) from the early Chadronian of Wyoming and
14 10 Texas. *Journal of Paleontology*, **52**(5), 1004-1014.
15
16 11 **Erfurt, J. & Métais, G.** 2007. Endemic European Paleogene Artiodactyls: Cebochoeridae, Choeropotamidae,
17 12 Mixtotheriidae, Cainotheriidae, Anoplotheriidae, Xiphodontidae, and Amphimerycidae. Pp. 59-84 in D.
18 13 R. Prothero & S. E. Foss (eds) *The Evolution of Artiodactyls*. Johns Hopkins University Press, Baltimore,
19 14 Maryland.
20
21 15 **Escarguel, G., Legendre, S. & Sigé, B.** 2008. Unearthing deep-time biodiversity changes: the Palaeogene
22 16 mammalian metacommunity of the Quercy and Limagne area (Massif Central, France). *Comptes Rendus*
23 17 *Geoscience*, **340**(9-10), 602-614.
24
25 18 **Geisler, J. H. & Uhen, M. D.** 2003. Morphological support for a close relationship between hippos and whales.
26 19 *Journal of Vertebrate Paleontology*, **23**(4), 991-996.
27
28 20 **Geisler, J. H. & Uhen, M. D.** 2005. Phylogenetic relationships of extinct cetartiodactyls: results of simultaneous
29 21 analyses of molecular, morphological, and stratigraphic data. *Journal of Mammalian Evolution*, **12**, 145-
30 22 160.
31
32 23 **Geisler, J. H., Theodor, J. M., Uhen, M. D. & Foss, S. E.** 2007. Phylogenetic relationships of cetaceans to
33 24 terrestrial artiodactyls. Pp. 19-31 in D. R. Prothero & S. E. Foss (eds) *The Evolution of Artiodactyls*. Johns
34 25 Hopkins University Press, Baltimore, Maryland.
35
36 26 **Gentry, A.W. & Hooker, J. J.** 1988. The phylogeny of the Artiodactyla. Pp. 235-272 in M. J Benton (eds) *The*
37 27 *phylogeny and classification of the tetrapods: Mammals*. Clarendon Press, Oxford.
38
39 28 **Hooker, J. J.** 2007. Bipedal browsing adaptations of the unusual Late Eocene-earliest Oligocene tylopod
40 29 Anoplotherium (Artiodactyla, Mammalia). *Zoological Journal of the Linnean Society*, **151**(3), 609-659.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2 1 **Hooker, J. J. & Weidmann, M.** 2000. *The Eocene mammal faunas of Mormont, Switzerland: systematic revision*
3 2 and resolutions of dating problems. Schweizerische Paläontologische Abhandlungen, **120**, pp. 141.
4
5 3 **Hooker, J. J. & Russell, D. E.** 2012. Early Palaeogene Louisinidae (Macroscelidea, Mammalia), their
6 4 relationships and north European diversity. *Zoological Journal of the Linnean Society*, **164**(4), 856-936.
7
8 5 **Hürzeler, J.** 1936. *Osteologie und odontologie der Caenotheriden*. Abhandlungen der Schweizerischen
9 6 Paläontologischen Gesellschaft, Birkhäuser, **58–59**, pp. 111.
10
11 7 **Klingener, D.** 1963. Dental evolution of Zapus. *Journal of Mammalogy*, **44**(2), 248-260.
12
13 8 **Legendre, S.** 1980. *Etude du gisement de Port-la-Nouvelle; étude des Cainotheriidae d'Escamps*. Unpublished
14 9 DEA of Paleontology, University of Montpellier, Montpellier.
15
16 10 **Legendre, S.** 1987. Les communautés de mammifères d'Europe occidentale de l'Eocène supérieur et Oligocène:
17 11 structures et milieux. *Münchener geowissenschaftliche Abhandlungen*, **10**(A), 310-312.
18
19 12 **Lihoreau, F., Boisserie, J. R., Manthi, F. K. & Ducrocq, S.** 2015. Hippo stem from the longest sequence of
20 13 terrestrial cetartiodactyl evolution in Africa. *Nat. Commun.*, **6**:6264.
21
22 14 **Matthew, W. D. & Reed, W. H.** 1910. On the skull of Apterodus and the skeleton of a new artiodactyl. *Bulletin*
23 15 *of the American Museum of Natural History*, **28**(5), 33-42.
24
25 16 **Métais, G. & Vislobokova, I.** 2007. Basal ruminants. Pp. 189-212 in D. R. Prothero & S. E. Foss (eds) *The*
26 17 *evolution of artiodactyls*. Johns Hopkins University Press, Baltimore, Maryland.
27
28 18 **O'Leary, M. A. & Gatesy, J.** 2007. Impact of increased character sampling on the phylogeny of Cetartiodactyla
29 19 (Mammalia): combined analysis including fossils. *Cladistics*, **23**, 1-46.
30
31 20 **Orliac, M. J., Pierre-Olivier, A. & Ducrocq, S.** 2010. Phylogenetic relationships of the Suidae (Mammalia,
32 21 Cetartiodactyla): new insights on the relationships within Suoidea. *Zoologica Scripta*, **39**(4), 315-330.
33
34 22 **Page, R. D. M.** 2001. Nexus Data Editor.
35
36 23 **Rincon, A. F., Bloch, J. I., MacFadden, B. J. & Jaramillo, C. A.** 2015. New early Miocene protoceratids
37 24 (Mammalia, Artiodactyla) from Panama. *Journal of Vertebrate Paleontology*, **35**(5), 1-22.
38
39 25 **Romer, A. S.** 1966. *Vertebrate Paleontology*. 3^{ème} édition. University of Chicago Press, Chicago Illinois, 468 pp.
40
41 26 **Schmidt-Kittler, N. & Vianey-Liaud, M.** 1987. Morphometric analysis of the genus *Issiodoromys*
42 27 (Theridomyidae, Rodentia) of the European Oligocene. *Proceedings of the Koninklijke Nederlandse*
43 28 *Akademie van Wetenschappen*, **90**, 281–306.
44
45 29 **Simpson, G. G.** 1945. The principles of classification and a classification of mammals. *Bulletin of the American*
46 30 *Museum of Natural History*, **85**, 1-350.

- 1
2 1 **Skwara, T.** 1986. A new ‘flying squirrel’ (Rodentia: Sciuridae) from the early Miocene of southwestern
3 2 Saskatchewan. *Journal of Vertebrate Paleontology*, **6**(3), 290-294.
4
5 3 **Stehlin, H. G.** 1906. Die Säugetiere des schweizerischen Eocaens. *Abhandlungen des schweizerischen*
6 4 *paläontologischen Gesellschaft*, 1-33.
7
8 5 **Stehlin, H. G.** 1908. Die Säugetiere des Schweizerischen Eocaens, Cristischer Catalog der Materialen, T. 5.
9 6 *Abhandlungen Schweizerischen Paläontologischen Gesellschaft*, **34**, 691-837.
10
11 7 **Stehlin, H. G.** 1910. Remarques sur les faunules de mammifère des couches éocènes et oligocènes du Bassin de
12 8 Paris. *Bull Soc Géol France*, **19**, 488-520.
13
14 9 **Sudre, J.** 1969. Les gisements de Robiac (Eocène supérieur) et leurs faunes de Mammifères. *Palaeovertebrata*, **2**,
15 10 95-156.
16
17 11 **Sudre, J.** 1977. L’évolution du genre Robiacina Sudre 1969, et l’origine des Cainotheriidae; implications
18 12 systématiques. *Geobios*, **10**, 213–231.
19
20 13 **Sudre, J.** 1978. *Les artiodactyles de l'éocène moyen et supérieur d'Europe occidentale: (systématique et*
21 14 *évolution)*. Unpublished PhD thesis, Montpellier, Institut EPHH, 229 pp.
22
23 15 **Sudre, J. & Legendre S.** 1992. Ungulates from Paleogene of Western Europe: relationships between their
24 16 evolution and environmental changes during that period. Pp 15-25 in G. Janeau, F. Spitz, G. Gonzalez &
25 17 S. Aulagnier (eds) *Ongulés/Ungulates*. Toulouse.
26
27 18 **Swofford, D. L.** 1984. PAUP-Phylogenetic Analysis Using Parsimony. *Illinois Natural History Survey*,
28 19 Champaign, Illinois.
29
30 20 **Swofford, D. L.** 2002. PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods). Version 4.0a161.
31 21 *Sinauer Associates*, Sunderland, Massachusetts.
32
33 22 **Theodor, J. M.** 2010. Micro-Computed Tomographic Scanning of the Ear Region of Cainotherium: Character
34 23 Analysis and Implications. *Journal of Vertebrate Paleontology*, **30**(1), 236-243.
35
36 24 **Thewissen, J. G., Cooper, L. N., Clementz, M. T., Bajpai, S. & Tiwari, B. N.** 2007. Whales originated from
37 25 aquatic artiodactyls in the Eocene epoch of India. *Nature*, **450**, 1190-1195.
38
39 26 **Van der Made, J.** 1996. Listriodontinae (Suidae, Mammalia), their evolution, systematics and distribution in time
40 27 and space. *Mededelingen van de Werkgroep voor Tertiaire en Kwartaire Geologie*, **33**(1/4), 3-254.
41
42 28 **Vianey-Liaud, M.** 1974. Les Rongeurs de l’Oligocène inférieur d’Escamps. *Palaeovertebrata*, **6**, 197–241.
43
44 29 **Vianey-Liaud, M.** 1976. Les Issiodoromyinae (Rodentia, Theridomyidae) de l’Eocène supérieur à l’Oligocène
45 30 supérieur en Europe Occidentale. *Palaeovertebrata*, **7**, 1–115.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2 **1** Vianey-Liaud, M. 1979. Evolution des rongeurs à l'Oligocène en Europe Occidentale. *Palaeontographica, A* **166**,
3 **2** 136–236.
- 4
5 **3** Vianey-Liaud, M. 1998. La radiation des Theridomyidae (Rodentia) à l'Oligocène inférieur: modalités et
6 **4** implications biochronologiques. *Geologica et Palaeontologica*, Marburg, **32**, 253–285.
- 7
8 **5** Vianey-Liaud, M. 2015. Parallelism in the evolution of dental pattern and systematic implications: The case of
9 **6** *Protechimys major* Schlosser, 1884 (Theridomyinae, Rodentia, Mammalia) and its associated rodents.
10 **7** *Comptes Rendus – Palevol*, **14**, 451–470.
- 11
12 **8** Vianey-Liaud, M. & Ringeade, M. 1993. La radiation des Theridomyidae (Rodentia) hypsodontes à l'Eocène
13 **9** supérieur. *Géobios*, Lyon, **26**, 455–495.
- 14
15 **10** Vianey-Liaud, M., Comte, B. & Lévêque, F. 1995. Le Garouillas et les sites contemporains (Oligocène MP25)
16 **11** des phosphorites du Quercy (Lot, Tarn et Garonne, France) et leurs faunes de vertébrés: Rongeurs.
17 **12** *Palaeontographica, A* **236**, 257–326.
- 18
19 **13** Vianey-Liaud, M. & Schmid, B. 2009. Diversité, datation et paléoenvironnement de la faune de mammifères
20 **14** oligocène de Cavalé (Quercy, SO France): contribution de l'analyse morphométrique des Theridomyinae
21 **15** (Mammalia, Rodentia). *Geodiversitas*, **31**, 909–941.
- 22
23 **16** Vianey-Liaud, M. & Schmidt-Kittler, N. 1987. Biostratigraphie de l'Oligocène d'Europe: importance des lignées
24 **17** guides de rongeurs Theridomyidae, et particulièrement des Issiodoromyinae. *Münchner
25 **18** Geowissenschaftliche Abhandlungen, A* **10**, 211–216.
- 26
27 **19** Webb, S. D. & Taylor, B. E. 1980. The phylogeny of hornless ruminants and a description of the cranium of
28 **20** Archaeomeryx. *Bulletin of the American Museum of Natural History*, **167**, 120–157.

22 **Figure captions**

24 **Figure 1.** Dental diagrams in occlusal view. P/4 of Cainotheriidae: **A**, Nomenclature of Berthet (2003);
25 **B**, Nomenclature of Boisserie *et al.* (2010). M/1-M/2 of Cainotheriidae: **C**, Nomenclature of Hürzeler
26 (1936); **D**, Nomenclature of Sudre (1977); **E**, Nomenclature of Gentry & Hooker (1988). M/1-M/2: **F**,
27 *Robiacina*; **G**, *Dacrytherium*. M/3: **H**, *Robiacina*; **I**, Cainotheriidae. **Proto^{ulid}.**, protoconulid. For the rest
28 of the legend, see Fig. 2.

1
2 **Figure 2.** Dental diagrams of the *Paroxacron valdense* dentition according to dental nomenclature of
3 Boisserie *et al.* (2010). **A, B**, M1/-M2/; **C, D**, M/3; **E, F**, I1/. In occlusal view: **A, B, C, D**. In lingual
4 view: **E, F**. Abbreviations: dist., distostyle; med., mediostyle; mes., mesostyle; met., metastyle; par.,
5 parastyle; **Meta.**, metacone; **Meta^{ul}.**, metaconule; **Para.**, paracone; **Para^{ul}.**, paraconule; **Proto.**,
6 protocone; post^{ylid}., poststylid; pre^{ylid}., prestylid; **Centro^{ulid}.**, centroconulid; **Ento^d.**, entoconid; **Hypo^d.**,
7 hypoconid; **Hypo^{ulid}.**, hypoconulid; **Meta^d.**, metaconid; **Meta^{ulid}.**, metaconulid; **Para^d.**, paraconid;
8 **Post^{ulid}.**, postconulid; **Pre^{ulid}.**, preconulid; **Proto^d.**, protoconid; -c., -crista; -c^{ula}., -cristula; -c^{yla}., -
9 cristyla ; -c^{id}., -cristid ; -c^{ulid}., -cristulid; -c^{ylid}., -cristylid; -f., fossa; -f^{ld}., fossid. Full circles:
10 cusp/style/conule/conulid; full lines: crest; dashed lines: valley/fosse; dotted lines: cingulum/cingulid;
11 bold dashed lines: lobe contour; bold full lines: crown contour.
12
13

14 **Figure 3.** Dental diagrams of the permanent lower dentition of *Paroxacron valdense*. **A, B**, P/1-2; **C, D**,
15 P/3; **E, F**, P/4; **G, H**, M/1-2; **I, J**, M/3; **K**, I/2-3; **L**, lower canine. In occlusal view: **A, C, E, G, I**. In
16 lingual view: **B, D, F, K, L**. In buccal view: **H, J**. Scale 1mm. Abbreviations: **Postproto^{ulid}.**,
17 postprotoconulid; disto^{ylid}., distostylid; co^{ulid} x., conulid x.
18
19

20 **Figure 4.** Dental diagrams of the permanent upper dentition of *Paroxacron valdense*. **A, B**, P1/; **C, D**,
21 P2/; **E, F**, P3/; **G, H**, P4/; **I**, M1-2/; **J**, I3/; **K**, I2/; **L**, upper canine. In occlusal view: **A, C, E, G**. In
22 lingual view: **B, D, F, H, I, J, K, L**. Scale 1mm. Abbreviations: co^{ul} x., conule x.
23
24

25 **Figure 5.** Dental diagrams of the deciduous dentition of *Paroxacron valdense*. **A, B**, DP/1; **C, D**, DP/2-
26 3; **E, F**, DP/4; **G, H**, DP1-2/; **I, J**, DP3/; **K, L**, DP4/. In occlusal view: **A, C, E, G, I, K**. In lingual view:
27 **B, D, H, J, L**. In buccal view: **F**. Scale 1 mm. Abbreviations: **Primo^{id}.**, primoconid.
28
29

30 **Figure 6.** Material of the lower dentition of *Paroxacron valdense*. **A**, right I/1 (DAM1 174); **B**, left I/2-3
31 (DAM1 194; mirror view); **C**, right lower canine (DAM1 199); **D, E**, right P/1 (DAM1 40); **F, G**, right
32 DP/1 (DAM1 126); **H**, ; **I**, left DP/2-3 (DAM1 117); **J**, ; **K**, left DP/4 (DAM1 103); **L**, left mandibular
33 fragment with P/1-M/3 (DAM1 22); **M**, left P/3 (DAM1 269); **N**, right mandibular fragment with P/2-
34 3 (DAM1 195).
35
36

1 M/3 (DAM1 21); **O**, left mandibular fragment with P/1-M/3 (DAM1 22); **P**, **Q**, right mandibular
2 fragment with M/3 (DAM1 268); **R**, right mandibular fragment with P/2-M/3 (DAM1 21). In lingual
3 view: **A**, **B**, **C**, **E**, **G**, **I**, **Q**. In occlusal view: **D**, **F**, **H**, **J**, **L**, **N**. In buccal view: **K**, **M**, **O**, **P**, **R**. Scale 1
4 mm.
5

6 **Figure 7.** Material of the upper dentition of *Paroxacron valdense*. **A**, right I3/ (DAM1 252) ; **B**, right
7 I2/ (DAM1 228); **C**, right I1/ (DAM1 213); **D**, left upper canine (DAM1 261); **E**, **F**, right DP1-2/
8 (DAM1 164); **G**, **H**, right DP3/ (DAM1 160); **I**, **J**, left P1/ (DAM1 66); **K**, **L**, right DP4/ (DAM1 144);
9 **M**, right maxillary fragment with P1/-M3/ (DAM1 08). In lingual view: **A**, **B**, **C**, **D**, **F**, **H**, **I**, **L**. In
10 occlusal view: **E**, **G**, **J**, **K**, **M**. Scale 1 mm.
11

12 **Figure 8.** Material of the *Paroxacron* sp. dentition. **A**, **B**, right P4/ (DAM3 09; mirror view); **C**, **D**, right
13 M1/ (DAM1 06; mirror view); **E**, **F**, left M2/ (DAM3 04); **G**, **H**, left M3/ (DAM3 03); **I**, **J**, left M2/
14 (DAM3 21; mirror view); **K**, **L**, right M3 (DAM3 11); **M**, **N**, right P1/ (DAM3 10; mirror view); **O**, **P**,
15 right P1 (DAM3 20; mirror view); **Q**, **R**, right P2 (DAM3 19); **S**, **T**, left DP/4 (DAM3 17; mirror view);
16 **U**, **V**, left M/1 (DAM3 16; mirror view). In occlusal view: **A**, **C**, **E**, **G**, **I**, **K**, **M**, **O**, **Q**, **S**, **U**. In lingual
17 view: **B**, **D**, **F**, **H**, **N**, **P**, **R**. In buccal view: **J**, **L**, **T**, **V**.
18

19 **Figure 9.** Material of the *Paroxacron bergeri* dentition. **A**, **B**, left P4/ (DAM3 30); **C**, **D**, left M1/
20 (DAM3 29); **E**, **F**, right M2/ (DAM3 26; mirror view); **G**, **H**, right M3/ (DAM3 24; mirror view); **I**, **J**,
21 left P1/ (DAM3 33); **K**, **L**, left P2/ (DAM3 32); **M**, **N**, right P3/ (DAM3 31; mirror view). In occlusal
22 view: **A**, **C**, **E**, **G**, **I**, **K**, **M**. In lingual view: **B**, **D**, **F**, **H**, **J**, **L**, **N**.
23

24 **Figure 10.** Material of the *Plesiomeryx cadurcensis* dentition. **A**, **B**, right P3/ (DAM3 48; mirror view);
25 **C**, **D**, right P4/ (DAM3 47; mirror view); **E**, **F**, right M1/ (DAM3 44; mirror view); **G**, **H**, left M2/
26 (DAM3 37); **I**, **J**, left M3/ (DAM3 34); **K**, **L**, left M3 (DAM3 52; mirror view); **M**, **N**, right P1/ (DAM3
27 50; mirror view); **O**, **P**, right P2/ (DAM3 51; mirror view); **Q**, **R**, right DP4/ (DAM3 45; mirror view);
28 **S**, **T**, right P2 (DAM3 77); **U**, **V**, right mandibular fragment with P/3-P/4 (DAM3 79); **W**, **X**, right M/1

1
2 1 (DAM3 61); **Y**, **Z**, left M/2 (DAM3 56; mirror view). In occlusal view: **A**, **C**, **E**, **G**, **I**, **K**, **M**, **O**, **Q**, **S**,
3
4 2 **U**, **W**, **Y**. In lingual view: **B**, **D**, **F**, **H**, **J**, **N**, **P**, **R**, **T**, **V**. In buccal view: **L**, **X**, **Z**.
5
6 3
7
8
9
10 4 **Figure 11.** Material of the *Plesiomeryx huerzeleri* dentition. **A**, **B**, left maxillary fragment with P3-/P4/
11
12 5 (DAM3 101); **C**, **D**, right M1/ (DAM3 88; mirror view); **E**, **F**, right M2/ (DAM3 83; mirror view);
13
14 6 **G**, **H**, right M3/ (DAM3 81; mirror view); **I**, **J**, right M1 (DAM3 111); **K**, **L**, right M/2 (DAM3 107);
15
16 7 **M**, **N**, right M/3 (DAM3 103); **O**, **P**, left P1/ (DAM3 100); **Q**, **R**, left P2/ (DAM3 99); **S**, **T**, right P1
17
18 8 (DAM3 127); **U**, **V**, right P/2 (DAM3 126); **W**, **X**, left mandibular fragment with P/4-P/3 (DAM3 128;
19
20 9 mirror view); **Y**, **Z**, right DP/2-3 (DAM3 115); **A'**, **B'**, left DP/4 (DAM3 114; mirror view). In occlusal
21
22 10 view: **A**, **C**, **E**, **G**, **I**, **K**, **M**, **O**, **Q**, **T**, **U**, **W**, **Y**, **A'**. In lingual view: **B**, **D**, **F**, **H**, **P**, **R**, **S**, **V**, **X**, **Z**. In
23
24 11 buccal view: **J**, **L**, **N**, **B'**.
25
26
27 12
28
29 13 **Figure 12.** Material of the *Caenomeryx cf. procommunis* dentition. **A**, **B**, right P3/ (DAM3 135; mirror
30
31 view); **C**, **D**, left M1/ (DAM3 133); **E**, **F**, right M2/ (DAM3 132; mirror view); **G**, **H**, right M3/
32
33 15 (DAM3 129; mirror view); **I**, **J**, right DP/2-3 (DAM3 145); **K**, **L**, right mandibular fragment with P/4-
34
35 M/3 (DAM3 152); **M**, **N**, left P1/ (DAM3 149; mirror view); **O**, **P**, left P2/ (DAM3 148; mirror view);
36
37 17 **Q**, **R**, right P3/ (DAM3 147). In occlusal view: **A**, **C**, **E**, **G**, **I**, **K**, **M**, **O**, **Q**. In lingual view: **B**, **D**, **F**, **H**,
38
39 18 **J**, **N**, **P**, **R**. In buccal view: **L**.
40
41
42 19
43
44 20 **Figure 13.** Phylogeny of Cainotheriidae and their position within artiodactyls. Most parsimonious tree
45
46 at 170 steps, with IC=0.48 and IR=0.73. The Bremer Index (IB) is indicated in bold at each node. **X** ●
47
48 a - b: X, node number; a, number of synapomorphies; b, number of ambiguous characters (Acctran
49
50 optimization).
51
52
53
54 25 **Figure 14.** Trends and stratigraphical extensions of cainotheriid species from Quercy, around the
55
56 Grande Coupure (modified from Blondel (2005; fig. 5) on the stratigraphical position of *Paroxacron*
57
58 27 *valdense* and *Paroxacron* sp., with the attribution of Escamps specimens to *Paroxacron valdense*).
59
60

1
2
3 1 Dotted-line: long diastemata, sub-triangular molars; simple line: diastemata reduction trend,
4
5 2 quadrangular molars.
6
7 3
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only




Figure 1. Dental diagrams in occlusal view. P/4 of Cainotheriidae: A, Nomenclature of Berthet (2003); B, Nomenclature of Boissarie et al. (2010). M/1-M/2 of Cainotheriidae: C, Nomenclature of Hürzeler (1936); D, Nomenclature of Sudre (1977); E, Nomenclature of Gentry & Hooker (1988). M/1-M/2: F, Robiacina; G, Dacrytherium. M/3: H, Robiacina; I, Cainotheriidae. Protoulid., protoconulid. For the rest of the legend, see Fig. 2.

173x232mm (300 x 300 DPI)




Figure 2. Dental diagrams of the *Paroxacron valdense* dentition according to dental nomenclature of Boisserie et al. (2010). A, B, M1/-M2/; C, D, M3/; E, I1/; F, I1/. In occlusal view: A, B, C, D. In lingual view: E, F. Abbreviations: dist., distostyle; med., mediostyle; mes., mesostyle; met., metastyle; par., parastyle; Meta., metacone; Metaul., metaconule; Para., paracone; Paraull., paraconule; Proto., protocone; poststylid., poststylid; preylid., prestylid; Centroulid., centroconulid; Entod., entoconid; Hypod., hypoconid; Hypoulid., hypoconulid; Metad., metaconid; Metaulid., metaconulid; Parad., paraconid; Postulid., postconulid; Preulid., preconulid; Protod., protoconid; -c., -crista; -cula., -cristula; -cyla., -cristyla; -cid., -cristid; -culid., -cristulid; -cylid., -cristylid; -f., fossa; -fid., fossid. Full circles: cusp/style/conule/conulid; full lines: crest; dashed lines: valley/fosse; dotted lines: cingulum/cingulid; bold dashed lines: lobe contour; bold full lines: crown contour.

173x228mm (300 x 300 DPI)




Figure 3. Dental diagrams of the permanent lower dentition of *Paroxacron valdense*. A, B, P/1-2; C, D, P/3; E, F, P/4; G, H, M/1-2; I, J, M/3; K, I/2-3; L, lower canine. In occlusal view: A, C, E, G, I. In lingual view: B, D, F, K, L. In buccal view: H, J. Scale 1mm. Abbreviations: Postprotoulid., postprotoconulid; distostyloid., distostyloid; coulid x., conulid x.

173x232mm (300 x 300 DPI)




Figure 4. Dental diagrams of the permanent upper dentition of *Paroxacron valdense*. A, B, P1/; C, D, P2/; E, F, P3/; G, H, P4/; I, M1-2/; J, I3/; K, I2/; L, upper canine. In occlusal view: A, C, E, G. In lingual view: B, D, F, H, I, J, K, L. Scale 1mm. Abbreviations: coul x., conule x.

173x232mm (300 x 300 DPI)




Figure 5. Dental diagrams of the deciduous dentition of *Paroxacron valdense*. A, B, DP/1; C, D, DP/2-3; E, F, DP/4; G, H, DP1-2/; I, J, DP3/; K, L, DP4/. In occlusal view: A, C, E, G, I, K. In lingual view: B, D, H, J, L. In buccal view: F. Scale 1 mm. Abbreviations: Primoid., primoconid.

173x232mm (300 x 300 DPI)




Figure 6. Material of the lower dentition of *Paroxacron valdense*. A, right I/1 (DAM1 174); B, left I/2-3 (DAM1 194; mirror view); C, right lower canine (DAM1 199); D, E, right P/1 (DAM1 40); F, G, right DP/1 (DAM1 126); H, ; I, left DP/2-3 (DAM1 117); J, ; K, left DP/4 (DAM1 103); L, left mandibular fragment with P/1-M/3 (DAM1 22); M, left P/3 (DAM1 269); N, right mandibular fragment with P/2-M/3 (DAM1 21); O, left mandibular fragment with P/1-M/3 (DAM1 22); P, Q, right mandibular fragment with M/3 (DAM1 268); R, right mandibular fragment with P/2-M/3 (DAM1 21). In lingual view: A, B, C, E, G, I, Q. In occlusal view: D, F, H, J, L, N. In buccal view: K, M, O, P, R. Scale 1 mm.

160x231mm (300 x 300 DPI)




Figure 7. Material of the upper dentition of *Paroxacron valdense*. A, right I₃/ (DAM1 252); B, right I₂/ (DAM1 228); C, right I₁/ (DAM1 213); D, left upper canine (DAM1 261); E, F, right DP₁₋₂/ (DAM1 164); G, H, right DP₃/ (DAM1 160); I, J, left P₁/ (DAM1 66); K, L, right DP₄/ (DAM1 144); M, right maxillary fragment with P₁-M₃/ (DAM1 08). In lingual view: A, B, C, D, F, H, I, L. In occlusal view: E, G, J, K, M. Scale 1 mm.

172x185mm (300 x 300 DPI)




Figure 8. Material of the *Paroxacron* sp. dentition. A, B, right P4/ (DAM3 09; mirror view); C, D, right M1/ (DAM1 06; mirror view); E, F, left M2/ (DAM3 04); G, H, left M3/ (DAM3 03); I, J, left M2/ (DAM3 21; mirror view); K, L, right M3/ (DAM3 11); M, N, right P1/ (DAM3 10; mirror view); O, P, right P1/ (DAM3 20; mirror view); Q, R, right P2/ (DAM3 19); S, T, left DP4/ (DAM3 17; mirror view); U, V, left M1/ (DAM3 16; mirror view). In occlusal view: A, C, E, G, I, K, M, O, Q, S, U. In lingual view: B, D, F, H, N, P, R. In buccal view: J, L, T, V.

170x231mm (300 x 300 DPI)




Figure 9. Material of the *Paroxacron bergeri* dentition. A, B, left P4/ (DAM3 30); C, D, left M1/ (DAM3 29); E, F, right M2/ (DAM3 26; mirror view); G, H, right M3/ (DAM3 24; mirror view); I, J, left P1/ (DAM3 33); K, L, left P2/ (DAM3 32); M, N, right P3/ (DAM3 31; mirror view). In occlusal view: A, C, E, G, I, K, M. In lingual view: B, D, F, H, J, L, N.

170x231mm (300 x 300 DPI)




Figure 10. Material of the *Plesiomeryx cadurcensis* dentition. A, B, right P3/ (DAM3 48; mirror view); C, D, right P4/ (DAM3 47; mirror view); E, F, right M1/ (DAM3 44; mirror view); G, H, left M2/ (DAM3 37); I, J, left M3/ (DAM3 34); K, L, left M3 (DAM3 52; mirror view); M, N, right P1/ (DAM3 50; mirror view); O, P, right P2/ (DAM3 51; mirror view); Q, R, right DP4/ (DAM3 45; mirror view); S, T, right P2 (DAM3 77); U, V, right mandibular fragment with P3-P4 (DAM3 79); W, X, right M1 (DAM3 61); Y, Z, left M2 (DAM3 56; mirror view). In occlusal view: A, C, E, G, I, K, M, O, Q, S, U, W, Y. In lingual view: B, D, F, H, J, N, P, R, T, V. In buccal view: L, X, Z.

170x231mm (300 x 300 DPI)




Figure 11. Material of the *Plesiomeryx huerzeleri* dentition. A, B, left maxillary fragment with P3/-P4/ (DAM3 101); C, D, right M1/ (DAM3 88; mirror view); E, F, right M2/ (DAM3 83; mirror view); G, H, right M3/ (DAM3 81; mirror view); I, J, right M1/ (DAM3 111); K, L, right M2/ (DAM3 107); M, N, right M3/ (DAM3 103); O, P, left P1/ (DAM3 100); Q, R, left P2/ (DAM3 99); S, T, right P1/ (DAM3 127); U, V, right P2/ (DAM3 126); W, X, left mandibular fragment with P4-P3/ (DAM3 128; mirror view); Y, Z, right DP2-3 (DAM3 115); A', B', left DP4/ (DAM3 114; mirror view). In occlusal view: A, C, E, G, I, K, M, O, Q, T, U, W, Y, A'. In lingual view: B, D, F, H, P, R, S, V, X, Z. In buccal view: J, L, N, B'.

170x231mm (300 x 300 DPI)




Figure 12. Material of the *Caenomeryx procommunis* dentition. A, B, right P3/ (DAM3 135; mirror view); C, D, left M1/ (DAM3 133); E, F, right M2/ (DAM3 132; mirror view); G, H, right M3/ (DAM3 129; mirror view); I, J, right DP/2-3 (DAM3 145); K, L, right mandibular fragment with P/4-M/3 (DAM3 152); M, N, left P/1 (DAM3 149; mirror view); O, P, left P/2 (DAM3 148; mirror view); Q, R, right P3/ (DAM3 147). In occlusal view: A, C, E, G, I, K, M, O, Q. In lingual view: B, D, F, H, J, N, P, R. In buccal view: L.

170x231mm (300 x 300 DPI)




Figure 13. Phylogeny of Cainotheriidae and their position within artiodactyls. Most parsimonious tree at 170 steps, with IC=0.48 and IR=0.73. The Bremer Index (IB) is indicated in bold at each node. X - a - b: X, node number; a, number of synapomorphies; b, number of ambiguous characters (Acctran optimization).

172x86mm (300 x 300 DPI)




Figure 14. Trends and stratigraphical extensions of cainotheriid species from Quercy, around the Grande Coupure (modified from Blondel (2005; fig. 5) on the stratigraphical position of *Paroxacron valdense* and *Paroxacron* sp., with the attribution of Escamps specimens to *Paroxacron valdense*). Dotted-line: long diastemata, sub-triangular molars; simple line: diastemata reduction trend, quadrangular molars.

172x113mm (300 x 300 DPI)

Supplementary data: Cainotheriidae (Mammalia, Artiodactyla) from Dams (Quercy, SW France); phylogenetic relationships and evolution around the Eocene-Oligocene transition (MP19-MP21).

Dataset S1. Characters list

- 1.** **Lower incisors.**
0. one endocristulid
1. two endocristulids

2. **Lower canine.** Hooker & Weidmann (2000)
0. caniniform
1. incisiform

3. **Diastema between C-P1/.**
0. absence
1. presence

4. **Diastema between P1-/P2/.**
0. absence
1. presence

5. **Diastema between P1-P2/.**
0. absent to reduced
1. short to long

6. **Diastema between P2/P3/.**
0. absent to reduced
1. short to long

7. **Upper M: styles, crests.** Hooker & Weidmann (2000)
0. absence
1. presence

8. **Upper M: parastyle, size.** Modified from Gentry & Hooker (1988)
0. very large, superior to other styles
1. large, like other styles

9. **Upper M: mesostyle.**
0. absence
1. presence

10. **Upper M: mesostyle, shape.** Gentry & Hooker (1988)
0. looped
1. pinched

11. **Upper M: position of the protocone (A).**
0. mesial
1. subcentral
2. distal

12. **M1/-M2/: hypocone.**
0. absence
1. presence

13. **Upper M: paraconule.** Modified from Gentry & Hooker (1988)
0. absence
1. presence

- 1
2
3 **14. Upper M: paraconule, size.** Modified from Gentry & Hooker (1988)
4 0. small
5 1. large
- 6 **15. Upper M: preprotocrista and paraconule, junction.**
7 0. absence
8 1. presence
- 9
10 **16. Upper M: metaconule, size.** Modified from Gentry & Hooker (1988)
11 0. small
12 1. large
- 13 **17. M3/: position of protocone relative to mesial or distal lingual cusp (bucco-lingual axis).** Modified from Hooker & Weidmann (2000)
14 *Lingual code when protocone is significatively more lingual than the mesial or distal lingual cusp.*
15 0. at the same level
16 1. protocone more lingual
- 17
18 **18. Upper M: cingulum surrounding the protocone (A).** Modified from Hooker & Weidmann (2000)
19 0. absent to very weak
20 1. weak
21 2. wide
22 3. very wide
- 23
24 **19. Upper M: dilambodonty.**
25 0. absence
26 1. presence
- 27 **20. Upper M: dilambodonty, type.** Modified from Hooker & Weidmann (2000)
28 0. strong, with crests weakly oblique and pronounced
29 1. very strong, with crests very oblique and pronounced
- 30
31 **21. Upper M: premetacristula.**
32 0. absence
33 1. presence
- 34 **22. Upper M: premetacristula, length.** Modified from Hooker & Weidmann (2000)
35 0. short
36 1. long, extend into the central valley
- 37
38 **23. Upper M: premetacristula, shape.** Modified from Hooker & Weidmann (2000)
39 0. continuous
40 1. discontinuous
- 41
42 **24. Upper M: postmetacristula and metastyle, junction.**
43 0. absence
44 1. presence
- 45
46 **25. Upper M: preparacrista, length.**
47 0. as long as the postparacrista
48 1. shorter than postparacrista
- 49
50 **26. Upper M: postprotocrista.**
51 0. absent to reduced
52 1. long
- 53
54 **27. Upper M: postprotocrista, direction.**
55 0. disto-buccally directed
56 1. distally directed
- 57
58 **28. Upper M: postprotocrista, bifurcation.**
59 0. absence
60 1. presence
- 61
62 **29. Upper M: postparacristula.**
63 0. absence
64 1. presence

- 1
2
3
4 **30.** **Upper M: postparacristula long and joins preprotocrista in the center of the tooth.** Modified from Erfurt &
5 Métais (2007)
6 0. absence
7 1. presence
- 8 **31.** **Upper M: shape (A).** Modified from Erfurt & Métais (2007)
9 0. triangular
10 1. subtriangular
11 2. quadrangular
- 12 **32.** **Upper M: elongated bucco-lingually.**
13 0. absence
14 1. presence
- 15 **33.** **Lower M: metaconulid.**
16 0. absence
17 1. presence
- 18 **34.** **Lower M: metaconulid, size (A).**
19 0. small
20 1. medium
21 2. large (like other cusps)
- 22 **35.** **Lower M: metaconulid and metaconid, linked.**
23 0. absence
24 1. presence
- 25 **36.** **Lower M: metastyliid.** Gentry & Hooker (1988)
26 0. absence
27 1. presence
- 28 **37.** **M/3: hypoconulid in basin-shape.** Hooker & Weidmann (2000)
29 0. absence
30 1. presence
- 31 **38.** **M/1-M/2: hypoconulid.**
32 0. absence
33 1. presence
- 34 **39.** **Lower M: entoconid in ‘loop-like’.**
35 *In ‘loop’: postentocristid convex and preentocristid concave distally*
36 0. absence
37 1. presence
- 38 **40.** **Lower M: entoconid and hypoconid, relative position (mesio-distal axis) (A).** Modified from Gentry & Hooker
39 (1988)
40 0. opposed, to entoconid more mesial
41 1. entoconid slightly more distal
42 2. entoconid very distal
- 43 **41.** **Lower M: valley between entoconid and metaconid or metaconulid.**
44 0. absence
45 1. presence
- 46 **42.** **Lower M: paraconid.** Modified from Gentry & Hooker (1988)
47 0. absent to small
48 1. large
- 49 **43.** **Lower M: prehypocristid, fused (cusp or crest).**
50 0. absence
51 1. presence
- 52 **44.** **Lower M: prehypocristid (oblique cristid), junction with (NA).** Modified from Hooker & Weidmann (2000)
53 0. postprotocristid
54 1. metaconulid or metastyliid
55 2. preentocristid

- 1
2
3 3. postprotocristid then preentocristid
4
5 45. **Lower M: prehypocristid.**
6 0. straight to convex buccally
7 1. concave buccally
8
9 46. **Lower M: postprotocristid and postmetacristid, fused.**
10 0. absence
11 1. presence
12
13 47. **Lower M: neotrigonid (Centroconulid + Postendocentrocristulid).** Sudre (1977)
14 0. absence
15 1. presence
16
17 48. **M/3: prehypocristulid, fused (cusp or crest).**
18 0. absence
19 1. presence
20
21 49. **M/3: prehypocristulid, junction with.**
22 0. posthypocristid
23 1. posthypocristid and postentocristid
24
25 50. **M/3: posthypocristulid.**
26 0. absence
27 1. presence
28
29 51. **M/3: posthypocristulid, inclined.**
30 0. mesio-buccally
31 1. mesio-distally
32
33 52. **Lower M: hypolophid.** Gentry & Hooker (1988)
34 0. absence
35 1. presence
36
37 53. **Lower M: straight buccal wall.**
38 *Straight buccal wall: buccal crests are aligned mesio-distally*
39 0. absence
40 1. presence
41
42 54. **P4/: metacone.** Modified from Hooker & Weidmann (2000)
43 0. absence
44 1. presence
45
46 55. **P4/: paracone et metacone, individualized.** Modified from Hooker & Weidmann (2000)
47 0. absence
48 1. presence
49
50 56. **P4/: paracone and protocone, level.**
51 0. at the same level
52 1. protocone more mesial
53 2. paracone more mesial
54
55 57. **P4/: postprotocrista.** Modified from Gentry & Hooker (1988)
56 0. absent to reduced
57 1. long
58
59 58. **P3-/P4/: styles, crests.** Hooker & Weidmann (2000)
60 0. absence
61 1. presence
62
63 59. **P3/: metacone.**
64 0. absence
65 1. presence
66
67 60. **P3/: metacone and paracone, relative size.** Hooker & Weidmann (2000)
68 0. metacone smaller than paracone
69 1. metacone and paracone sub-equal

- 1
2
3
4 **61. P3/: triangular.**
5 0. absence
6 1. presence
7 **62. P3/: mesial lobe, very elongated.**
8 0. absence
9 1. presence
10 **63. P3/: protocone, mesio-distal width.** Modified from Hooker & Weidmann (2000)
11 0. small
12 1. large
13
14 **64. P2/: protocone.**
15 0. absence
16 1. presence
17
18 **65. P/3: metaconid (visible groove):** Modified from Gentry & Hooker (1988)
19 0. absence
20 1. presence
21
22 **66. P/3: entoconid.**
23 0. absence
24 1. presence
25
26 **67. P/2-P/3: hypoconid.**
27 0. absent to small
28 1. large
29
30 **68. Lower P : postprotoconulid.**
31 0. absence
32 1. presence
33
34 **69. Lower P : postprotoconulid, size.**
35 0. very small
36 1. small to large
37
38 **70. P/1: roots.**
39 0. uniradiculate
40 1. biradiculate
41
42 **71. P/1: shape.** Modified from Gentry & Hooker (1988)
43 0. caniniform
44 1. premolariform
45
46 **72. P: cingulum/id pronounced.**
47 0. absence
48 1. presence
49
50 **73. Lower P: large premolars with strong main cusp.**
51 0. absence
52 1. presence
53
54
55 **(A) Ordered characters; (NA) Unordered characters**
56
57
58
59
60

Dataset S2. Taxa/characters matrix from dental material (provided as .nex format)

#NEXUS

[File saved by NDE version 0.5.0, Sun Jan 13 10:07:54 2019]

1
2
3 BEGIN TAXA;
4 DIMENSIONS NTAX=23;
5 TAXLABELS
6 'Paroxacron valdense'
7 'Paroxacron bergeri'
8 'Oxacron courtoisii'
9 'Plesiomeryx huerzeleri'
10 'Plesiomeryx cadurcensis'
11 'Caenomeryx procommunis'
12 'Caenomeryx filholi'
13 'Cainotherium laticurvatum'
14 'Robiacina minuta'
15 'Robiacina quercyi'
16 'Robiacina lavergnensis'
17 'Dacrytherium ovinum'
18 'Anoplotherium latipes'
19 'Mixtotherium cuspidatum'
20 'Parvitragulus priscus'
21 'Lophiomeryx chalaniati'
22 'Archaeomeryx optatus'
23 'Amphimeryx murinus'
24 'Xiphodon castrense'
25 'Paratoceras coatesi'
26 'Eotylopus reedi'
27 'Dichobune leporina'
28 'Eurodexis russelli'
29 ;
30 ENDBLOCK;
31
32 BEGIN CHARACTERS;
33 DIMENSIONS NCHAR=73;
34 FORMAT DATATYPE=STANDARD MISSING=? GAP=- SYMBOLS="0123";
35 CHARLABELS
36 [1] 'Lower incisors'
37 [2] 'Lower canine'
38 [3] 'Diastema between C-P1/'
39 [4] 'Diastema between P1-/P2/'
40 [5] 'Diastema between P/1-P/2'
41 [6] 'Diastema between P/2-P/3'
42 [7] 'Upper M: styles, crests'
43 [8] 'Upper M: parastyle, size'
44 [9] 'Upper M: mesostyle'
45 [10] 'Upper M: mesostyle, shape'
46 [11] 'Upper M: position of the protocone (A)'
47 [12] 'M1/-M2/: hypocone'
48 [13] 'Upper M: paraconule'
49 [14] 'Upper M: paraconule, size'
50 [15] 'Upper M: preprotocrista and paraconule, junction'
51 [16] 'Upper M: metaconule, size'
52 [17] 'M3/: position of protocone relative to mesial or distal lingual cusp (bucco-lingual axis)'
53 [18] 'Upper M: cingulum surrounding the protocone (A)'
54 [19] 'Upper M: dilambdodonty'
55 [20] 'Upper M: dilambdodonty, type'
56 [21] 'Upper M: premetacristula'
57 [22] 'Upper M: premetacristula, length'
58 [23] 'Upper M: premetacristula, shape'
59 [24] 'Upper M: postmetacristula and metastyle, junction'
60 [25] 'Upper M: preparacrista, length'
61 [26] 'Upper M: postprotocrista'
62 [27] 'Upper M: postprotocrista, direction'
63 [28] 'Upper M: postprotocrista, bifurcation'
64 [29] 'Upper M: postparacristula'
65 [30] 'Upper M: postparacristula long and joins preprotocrista in the center of the tooth'
66 [31] 'Upper M: shape (A)'
67 [32] 'Upper M: elongated bucco-lingually'
68 [33] 'Lower M: metaconulid'
69 [34] 'Lower M: metaconulid, size (A)'

- 1
2
3 [35] 'Lower M: metaconulid and metaconid, linked'
4 [36] 'Lower M: metastylid'
5 [37] 'M/3: hypoconulid in basin-shape'
6 [38] 'M/1-M/2: hypoconulid'
7 [39] 'Lower M: entoconid in 'loop-like''
8 [40] 'Lower M: entoconid and hypoconid, relative position (mesio-distal axis) (A)'
9 [41] 'Lower M: valley between entoconid and metaconid or metaconulid'
10 [42] 'Lower M: paraconid'
11 [43] 'Lower M: prehypocristid, fused (cusp or crest)'
12 [44] 'Lower M: prehypocristid (oblique cristid), junction with (NA)'
13 [45] 'Lower M: prehypocristid'
14 [46] 'Lower M: postprotocristid and postmetacristid, fused'
15 [47] 'Lower M: neotrigonid (Centroconulid + Postendocentrocristulid)'
16 [48] 'M/3: prehypocristulid, fused (cusp or crest)'
17 [49] 'M/3: prehypocristulid, junction with'
18 [50] 'M/3: posthypocristulid'
19 [51] 'M/3: posthypocristulid, inclined'
20 [52] 'Lower M: hypolophid'
21 [53] 'Lower M: straight buccal wall'
22 [54] 'P4/: metacone'
23 [55] 'P4/: paracone et metacone, individualized'
24 [56] 'P4/: paracone and protocone, level'
25 [57] 'P4/: postprotocrista'
26 [58] 'P3/-P4/: styles, crests'
27 [59] 'P3/: metacone'
28 [60] 'P3/: metacone and paracone, relative size'
29 [61] 'P3/: triangular'
30 [62] 'P3/: mesial lobe, very elongated'
31 [63] 'P3/: protocone, mesio-distal width'
32 [64] 'P2/: protocone'
33 [65] 'P/3: metaconid (visible groove)'
34 [66] 'P/3: entoconid'
35 [67] 'P/2-P/3: hypoconid'
36 [68] 'Lower P : postprotoconulid'
37 [69] 'Lower P : postprotoconulid, size'
38 ;
39 STATELABELS
40 1 'one endocristulid'
41 'two endocristulids',
42 2 caniniform
43 incisiform,
44 3 absence
45 presence,
46 4 absence
47 presence,
48 5 'absent to reduced'
49 'short to long',
50 6 'absent to reduced'
51 'short to long',
52 7 absence
53 presence,
54 8 'very large, superior to other styles'
55 'large, like other styles',
56 9 absence

1		
2		
3		presence,
4	10	looped
5		pinched,
6	11	mesial
7		subcentral
8		distal,
9	12	absence
10		presence,
11	13	absence
12		presence,
13	14	small
14		large,
15	15	absence
16		presence,
17	16	small
18		large,
19	17	'at the same level'
20		'protocone more lingual',
21	18	'absent to very weak'
22		'weak'
23		'wide'
24		'very wide',
25	19	absence
26		presence,
27	20	'strong, with crests weakly oblique and pronounced'
28		'very strong, with crests very oblique and pronounced',
29	21	absence
30		presence,
31	22	short
32		'long, extend into the central valley',
33	23	continuous
34		discontinuous,
35	24	absence
36		presence,
37	25	'as long as the postparacrista'
38		'shorter than postparacrista',
39	26	'absent to reduced'
40		long,
41	27	'disto-buccally directed'
42		'distally directed',
43	28	absence
44		presence,
45	29	absence
46		presence,
47	30	absence
48		presence,
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

1		
2		
3	31	triangular subtriangular quadrangular,
4		
5	32	absence presence,
6		
7	33	absence presence,
8		
9	34	small medium 'large (like other cuspids)',
10		
11	35	absence presence,
12		
13	36	absence presence,
14		
15	37	absence presence,
16		
17	38	absence presence,
18		
19	39	absence presence,
20		
21	40	'opposed, to entoconid more mesial' 'entoconid slightly more distal' 'entoconid very distal',
22		
23	41	absence presence,
24		
25	42	'absent to small' large,
26		
27	43	absence presence,
28		
29	44	postprotocristid 'metaconulid or metastylid' preentocristid 'postprotocristid then preentocristid',
30		
31	45	'straight to convex buccally' 'concave buccally',
32		
33	46	absence presence,
34		
35	47	absence presence,
36		
37	48	absence presence,
38		
39	49	absence presence,
40		
41	50	'posthypocristid ' 'posthypocristid and postentocristid',
42		
43	51	absence presence,
44		
45	51	'mesio-buccally'
46		
47		
48		
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

1		
2		
3		'mesio-distally',
4	52	absence
5		presence,
6	53	absence
7		presence,
8	54	absence
9		presence,
10	55	absence
11		presence,
12	56	absence
13		presence,
14	57	'at the same level'
15		'protocone more mesial'
16		'paracone more mesial',
17	58	'absent to reduced'
18		long,
19	59	absence
20		presence,
21	60	absence
22		presence,
23	61	'metacone smaller than paracone'
24		'metacone and paracone sub-equal',
25	62	absence
26		presence,
27	63	absence
28		presence,
29	64	small
30		large,
31	65	absence
32		presence,
33	66	absence
34		presence,
35	67	'absent to small'
36		'large',
37	68	absence
38		presence,
39	69	'very small'
40		'small to large',
41	70	uniradiculated
42		biradiculated,
43	71	caniniform
44		premolariform,
45	72	absence
46		presence,
47	73	absence
48		
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

```

1
2
3           presence,
4
5           ;
6           MATRIX
7           'Paroxacron valdense'      1101010110 2011011111 1001011010 1012000012 1011011110 -001011111
8           'Paroxacron bergeri'     ?100000111 2011011111 1001011010 201200???2 20????????? ??01011111
9           'Oxacron courtoisii'    1101010110 2010011111 1001011010 1012000012 1011011110 -001011111
10          '1000000101 000'
11          'Plesiomeryx huerzeleri' 1110100111 2010011111 1001011011 2012000012 1011011110 -001001111
12          '1011000110 101'
13          'Plesiomeryx cadurcensis' ?110100111 2010011111 1001011011 2012000012 1011011110 -001001111
14          '1011000110 101'
15          'Caenomeryx procommunis' 1110101111 2011011111 1001011011 2112000012 1011011110 -001001111
16          '1011000110 101'
17          'Caenomeryx filholi'     ??10101111 2011011111 1001011011 2112000012 1011011110 -001001111
18          '1011000110 101'
19          'Cainotherium laticurvatum' 1100001111 2011010111 1001011011 2012000012 0011011110 -001001111
20          '0000000110 111'
21          'Robiacina minuta'       ?1??010011 1010111010 111111000- 1010101111 1011010111 0001000011
22          '100?100100 100'
23          'Robiacina querocyi'    ?100??1011 1010111010 111110-00- 1010101111 1011010111 0001000011
24          '1000??0??? ?00'
25          'Robiacina lavergnensis' ?0?????1011 1010111210 100110-00- 1011011111 1010010111 0001011111
26          '101?100??0 ?00'
27          'Dacrytherium ovinum'     0100001011 0011110210 100100-00- 2011101111 1011010111 0001001110
28          '00111010?-? 111'
29          'Anoplotherium latipes'   ??00001011 001111?210 100100-00- 201110??11 11110101?? ?001001110
30          '00111010?-? 111'
31          'Mixtotherium cuspidatum' ?000001011 0010110210 110100-00- 200--10110 000-110111 0001100110
32          '10110010?-? 101'
33          'Parvitragulus priscus'   ???10110- 000--1000- 11010100-- 200--00000 0013010101 1011101110 101110?0-0
34          '000'
35          'Lophiomeryx chalaniati'  ???101111 000--10310 11010100-- 200--00100 0012010100 -000-01010 01111010-0
36          '010'
37          'Archaeomeryx optatus'    ?1--101111 000--10310 11010100-- 200--00?00 0012010101 0000-11110 1110?2010-0
38          '000'
39          'Amphimeryx murinus'      ??????0111 0011011211 1001010010 200--00100 000-110110 -00100101?
40          '110????0-? ???'
41          'Xiphodon castrense'     ?????01011 0011010210 1001010010 200--00000 00120100-1 001100101? 000?0100-
42          '? ?10'
43          'Paratoceras coatesi '   ?????101111 000--1120- 10000100-- 200--00000 000-0000-1 0000-01010 00000100-0
44          '011'
45          'Eotylopus reedi'        0111100111 000--1110- 10000101-- 200--00000 000-0000-1 0000-01010 00100100-0
46          '111'
47          'Dichobune leporina'     ?????11010- 011101130- 0--100-00- 200--?100 1?0-010100 -101121??? ??????000-0
48          '??0'
49          'Eurodexis russelli'    ??????010- 101010130- 0--101000- 000--00100 010-010101 0000-0000- 100?????-?
50          '??'
51          ;
52          ENDBLOCK;
53
54          BEGIN ASSUMPTIONS;
55          OPTIONS DEFTYPE=UNORD POLYTCOUNT=MINSTEPS;
56          TYPESET * default = ORD: 11 18 31 34 40;
57          ENDBLOCK;
58
59
60

```

Dataset S3. Apomorphy lists

Branch	Character	Steps	CI	Change
<hr/>				

1			
2			
3	node_36 --> node_37	7 (Upper M: styles, crests) 9 (Upper M: mesostyle) 18 (Upper M: cingulum surrounding the protocone) 19 (Upper M: dilambdodonty) 21 (Upper M: premetacristula) 49 (M/3: prehypocristulid, junction with) 59 (P3/: metacone)	1 0.167 0 - -> 1 1 0.500 0 ==> 1 1 0.375 3 ==> 2 1 0.333 0 ==> 1 1 1.000 0 ==> 1 1 0.500 0 - -> 1 1 1.000 0 - -> 1
4			
5			
6			
7			
8			
9			
10			
11			
12			
13			
14	node_37 --> node_38	15 (Upper M: preprotocrista and paraconule, j) 17 (M3/: position of protocone relative to me) 22 (Upper M: premetacristula, length) 43 (Lower M: prehypocristid, fused (cusp or c)) 58 (P3/-P4/: styles, crests) 63 (P3/: protocone, mesio-distal width) 64 (P2/: protocone) 65 (P/3: metaconid (visible groove)) 67 (P/2-P/3: hypoconid)	1 0.333 0 - -> 1 1 0.250 1 ==> 0 1 0.333 0 - -> 1 1 0.333 0 - -> 1 1 0.333 0 ==> 1 1 0.200 0 ==> 1 1 0.250 0 - -> 1 1 0.333 0 - -> 1 1 0.500 0 ==> 1
15			
16			
17			
18			
19			
20			
21			
22			
23			
24			
25			
26			
27			
28			
29	node_38 --> node_39	5 (Diastema between P/1-P/2) 8 (Upper M: parastyle, size) 26 (Upper M: postprotocrista) 39 (Lower M: entoconid in ?loop-like?) 44 (Lower M: prehypocristid (oblique cristid)) 71 (P/1: shape) 73 (Lower P: large premolars with strong mai)	1 0.500 1 ==> 0 1 0.333 1 ==> 0 1 0.250 1 ==> 0 1 1.000 0 ==> 1 1 1.000 2 - -> 1 1 0.333 0 - -> 1 1 0.250 0 - -> 1
30			
31			
32			
33			
34			
35			
36			
37			
38			
39			
40	node_39 --> node_40	22 (Upper M: premetacristula, length) 33 (Lower M: metaconulid)	1 0.333 1 - -> 0 1 1.000 0 ==>
41			
42	1		
43			
44		37 (M/3: hypoconulid in basin-shape)	1 0.500 0 - -> 1
45		40 (Lower M: entoconid and hypoconid, relativ)	1 1.000 0 ==> 1
46		41 (Lower M: valley between entoconid and met)	1 0.333 0 ==> 1
47		55 (P4/: paracone et metacone, individualized)	1 0.500 1 ==> 0
48			
49			
50	node_40 --> node_41	1 (Lower incisors) 11 (Upper M: position of the protocone (A)) 17 (M3/: position of protocone relative to me) 18 (Upper M: cingulum surrounding the protocone) 60 (P3/: metacone and paracone, relative size) 63 (P3/: protocone, mesio-distal width) 64 (P2/: protocone)	1 1.000 0 - -> 1 1 0.667 0 ==> 1 1 0.250 0 ==> 1 1 0.375 2 - -> 1 1 1.000 0 ==> 1 1 0.200 1 - -> 0 1 0.250 1 ==> 0
51			
52			
53			
54			
55			
56			
57			
58			
59			
60			

1			
2			
3		67 (P/2-P/3: hypoconid)	1 0.500 1 ==> 0
4		68 (Lower P : postprotoconulid)	1 1.000 0 ==> 1
5		73 (Lower P: large premolars with strong mai)	1 0.250 1 - -> 0
6			
7	node_41 --> node_42	8 (Upper M: parastyle, size)	1 0.333 0 ==> 1
8		11 (Upper M: position of the protocone (A))	1 0.667 1 ==> 2
9		15 (Upper M: preprotocrista and paraconule, i)	1 0.333 1 ==> 0
10		20 (Upper M: dilambdodonty, type)	1 0.500 0 ==> 1
11		26 (Upper M: postprotocrista)	1 0.250 0 ==> 1
12		27 (Upper M: postprotocrista, direction)	1 1.000 0 ==> 1
13		29 (Upper M: postparacristula)	1 0.500 0 ==> 1
14		34 (Lower M: metaconulid, size (A))	1 1.000 1 ==> 2
15		35 (Lower M: metaconulid and metaconid, linked)	1 1.000 1 ==> 0
16		37 (M/3: hypoconulid in basin-shape)	1 0.500 1 - -> 0
17		38 (M/1-M/2: hypoconulid)	1 0.333 1 ==> 0
18		40 (Lower M: entoconid and hypoconid, relativ)	1 1.000 1 ==> 2
19		47 (Lower M: neotrigonid (Centroconulid + Pos))	1 1.000 0 ==> 1
20		50 (M/3: posthypocristulid)	1 0.250 1 ==> 0
21		65 (P/3: metaconid (visible groove))	1 0.333 1 - -> 0
22			
23	node_42 --> node_43	7 (Upper M: styles, crests)	1 0.167 1 ==> 0
24		56 (P4/: paracone and protocone, level)	1 0.500 0 ==> 1
25		62 (P3/: mesial lobe, very elongated)	1 0.250 0 - -> 1
26		70 (P1/: roots)	1 1.000 0 ==> 1
27			
28	node_43 --> node_44	4 (Diastema between P1-/P2/)	1 0.500 0 ==> 1
29		6 (Diastema between P/2-P/3)	1 0.333 0 ==> 1
30		10 (Upper M: mesostyle, shape)	1 1.000 1 ==> 0
31		31 (Upper M: shape (A))	1 0.500 2 ==> 1
32			
33	node_44 --> O. courtoisii	14 (Upper M: paraconule, size)	1 0.200 1 ==> 0
34		62 (P3/: mesial lobe, very elongated)	1 0.250 1 - -> 0
35		71 (P1/: shape)	1 0.333 1 ==> 0
36			
37	node_42 --> node_27	30 (Upper M: postparacristula long and joins)	1 1.000 0 ==> 1
38		69 (Lower P : postprotoconulid, size)	1 1.000 0 ==> 1
39		73 (Lower P: large premolars with strong mai)	1 0.250 0 - -> 1
40			
41	node_27 --> node_26	3 (Diastema between C-P1/)	1 0.500 0 ==> 1
42		5 (Diastema between P/1-P/2)	1 0.500 0 ==> 1
43		63 (P3/: protocone, mesio-distal width)	1 0.200 0 - -> 1
44		64 (P2/: protocone)	1 0.250 0 ==> 1
45			
46			
47			
48			
49			
50			
51			
52			
53			
54			
55			
56			
57			
58			
59			
60			

1			
2			
3	node_26 --> node_24	7 (Upper M: styles, crests)	1 0.167 1 ==> 0
4		14 (Upper M: paraconule, size)	1 0.200 1 ==> 0
5			
6	node_26 --> node_25	32 (Upper M: elongated bucco-lingually)	1 1.000 0 ==> 1
7			
8	node_27 --> Cai. laticurvatum	17 (M3/: position of protocone relative to me)	1 0.250 1 ==> 0
9		41 (Lower M: valley between entoconid and met)	1 0.333 1 ==> 0
10		61 (P3/: triangular)	1 0.250 1 ==> 0
11		72 (P: cingulum/id pronounced)	1 0.250 0 ==> 1
12			
13	node_41 --> node_29	6 (Diastema between P2-P3)	1 0.333 0 -> 1
14		14 (Upper M: paraconule, size)	1 0.200 1 ==> 0
15		25 (Upper M: preparacrista, length)	1 1.000 0 ==> 1
16		31 (Upper M: shape (A))	1 0.500 2 ==> 1
17			
18	node_29 --> node_28	18 (Upper M: cingulum surrounding the proto)	1 0.375 1 ==> 0
19		22 (Upper M: premetacristula, length)	1 0.333 0 ==> 1
20		23 (Upper M: premetacristula, shape)	1 1.000 0 ==> 1
21		34 (Lower M: metaconulid, size (A))	1 1.000 1 ==> 0
22		57 (P4/: postprotocrista)	1 0.333 1 ==> 0
23		58 (P3-/P4/: styles, crests)	1 0.333 1 ==> 0
24			
25	node_28 --> R. minuta	7 (Upper M: styles, crests)	1 0.167 1 ==> 0
26		26 (Upper M: postprotocrista)	1 0.250 0 ==> 1
27			
28	node_29 --> R. lavergnensis	2 (Lower canine)	1 0.500 1 ==> 0
29		18 (Upper M: cingulum surrounding the proto)	1 0.375 1 -> 2
30		44 (Lower M: prehypocristid (oblique cristid))	1 1.000 1 ==> 0
31		56 (P4/: paracone and protocone, level)	1 0.500 0 ==> 1
32		63 (P3/: protocone, mesio-distal width)	1 0.200 0 -> 1
33			
34	node_40 --> node_30	61 (P3/: triangular)	1 0.250 1 ==> 0
35		72 (P: cingulum/id pronounced)	1 0.250 0 ==> 1
36			
37	node_30 --> A. latipes	42 (Lower M: paraconid)	1 0.500 0 ==> 1
38			
39	node_39 --> M. cuspidatum	2 (Lower canine)	1 0.500 1 ==> 0
40		14 (Upper M: paraconule, size)	1 0.200 1 ==> 0
41		36 (Lower M: metastyloid)	1 1.000 0 ==> 1
42		43 (Lower M: prehypocristid, fused (cusp or c))	1 0.333 1 -> 0
43		45 (Lower M: prehypocristid)	1 0.500 0 ==> 1
44		57 (P4/: postprotocrista)	1 0.333 1 ==> 0
45		65 (P3/: metaconid (visible groove))	1 0.333 1 -> 0
46			
47	node_38 --> node_32	13 (Upper M: paraconule)	1 0.500 1 ==> 0
48		49 (M3/: prehypocristulid, junction with)	1 0.500 1 -> 0
49			
50			
51			
52			
53			
54			
55			
56			
57			
58			
59			
60			

1			
2			
3	node_32 --> P. priscus	9 (Upper M: mesostyle) 18 (Upper M: cingulum surrounding the protocone) 19 (Upper M: dilambdodonty) 38 (M1-M2: hypoconulid) 44 (Lower M: prehypocristid (oblique cristid)) 51 (M3: posthypocristulid, inclined) 53 (Lower M: straight buccal wall)	1 0.500 1 ==> 0 2 0.375 2 ==> 0 1 0.333 1 ==> 0 1 0.333 1 ==> 0 1 1.000 2 ==> 3 1 1.000 0 ==> 1 1 0.500 0 ==> 1
4			
5			
6			
7			
8			
9			
10			
11			
12			
13			
14	node_32 --> node_31	18 (Upper M: cingulum surrounding the protocone) 54 (P4/: metacone) 62 (P3/: mesial lobe, very elongated)	1 0.375 2 ==> 3 1 0.333 1 ==> 0 1 0.250 0 ==> 1
15			
16			
17			
18			
19	node_31 --> L. chalaniati	50 (M3: posthypocristulid) 58 (P3-P4/: styles, crests) 61 (P3/: triangular) 72 (P: cingulum/id pronounced)	1 0.250 1 ==> 0 1 0.333 1 ==> 0 1 0.250 1 ==> 0 1 0.250 0 ==> 1
20			
21			
22			
23			
24			
25	node_31 --> Arc. optatus	56 (P4/: paracone and protocone, level) 64 (P2/: protocone)	1 0.500 0 ==> 1 1 0.250 1 ==> 0
26			
27			
28			
29	node_37 --> node_35	3 (Diastema between C-P1/) 4 (Diastema between P1-P2/) 29 (Upper M: postparacristula) 55 (P4/: paracone et metacone, individualized)	1 0.500 0 -> 1 1 0.500 0 -> 1 1 0.500 0 ==> 1 1 0.500 1 ==> 0
30			
31			
32			
33			
34			
35			
36			
37			
38	node_35 --> Amp. murinus	66 (P3/: entoconid) 72 (P: cingulum/id pronounced)	1 1.000 0 -> 1 1 0.250 0 -> 1
39			
40			
41			
42			
43			
44			
45			
46	node_35 --> node_34	7 (Upper M: styles, crests) 20 (Upper M: dilambdodonty, type) 45 (Lower M: prehypocristid) 50 (M3: posthypocristulid) 62 (P3/: mesial lobe, very elongated)	1 0.167 1 -> 0 1 0.500 0 ==> 1 1 0.500 0 ==> 1 1 0.250 1 ==> 0 1 0.250 0 ==> 1
47			
48			
49			
50			
51	node_34 --> X. castrense	38 (M1-M2: hypoconulid) 48 (M3: prehypocristulid, fused (cusp or crest)) 61 (P3/: triangular)	1 0.333 1 ==> 0 1 1.000 1 ==> 0 1 0.250 1 ==> 0
52			
53			
54			
55			
56			
57			
58			
59	node_34 --> node_33	8 (Upper M: parastyle, size) 17 (M3/: position of protocone relative to me) 43 (Lower M: prehypocristid, fused (cusp or crest)) 53 (Lower M: straight buccal wall)	1 0.333 1 ==> 0 1 0.250 1 ==> 0 1 0.333 0 ==> 1 1 0.500 0 ==> 1
60			

1						
2						
3		24 (Upper M: postmetacristula and metastyle,)			1 1.000 1 ==> 0	
4		46 (Lower M: postprotocristid and postmetaci)			1 1.000 1 ==> 0	
5		54 (P4/: metacone)			1 0.333 1 ==> 0	
6		73 (Lower P: large premolars with strong mai)			1 0.250 0 ==> 1	
7						
8	node_33 --> E. reedi	7 (Upper M: styles, crests)			1 0.167 1 - -> 0	
9		18 (Upper M: cingulum surrounding the protoco)			1 0.375 2 ==> 1	
10		28 (Upper M: postprotocrista, bifurcation)			1 1.000 0 ==> 1	
11		63 (P3/: protocone, mesio-distal width)			1 0.200 0 ==> 1	
12		71 (P1/: shape)			1 0.333 0 - -> 1	
13	node_36 --> D. leporina	6 (Diastema between P2-P3)			1 0.333 0 - -> 1	
14		12 (M1/-M2/: hypocone)			1 1.000 0 ==> 1	
15		26 (Upper M: postprotocrista)			1 0.250 1 ==> 0	
16		41 (Lower M: valley between entoconid and met)			1 0.333 0 ==> 1	
17		50 (M/3: posthypocristulid)			1 0.250 1 ==> 0	
18		52 (Lower M: hypolophid)			1 1.000 0 ==> 1	
19		56 (P4/: paracone and protocone, level)			1 0.500 0 ==> 2	
20	node_36 --> E. russelli	11 (Upper M: position of the protocone (A))			1 0.667 0 ==> 1	
21		14 (Upper M: paraconule, size)			1 0.200 1 ==> 0	
22		15 (Upper M: preprotocrista and paraconule, j)			1 0.333 0 - -> 1	
23		16 (Upper M: metaconule, size)			1 1.000 1 ==> 0	
24		31 (Upper M: shape (A))			2 0.500 2 ==> 0	
25		42 (Lower M: paraconid)			1 0.500 0 - -> 1	
26		54 (P4/: metacone)			1 0.333 1 ==> 0	
27		57 (P4/: postprotocrista)			1 0.333 1 ==> 0	
28						
29						
30						
31						
32						
33						
34						
35						
36						
37						
38						
39						
40						
41						
42						
43						
44						
45						

Table S1. Measurements and statistical data of the permanent lower teeth of *Paroxacron valdense* of DAM1. N, number of specimens; L/lp/la/h/L3 and l3, see Methods; LV, limit values; M ± SE, Mean ± Standard Error; σ, standard deviation; V, coefficient of variation.

	N	L/Lp/la/h	LV	M ± SE	Σ	V
I/1	15	L	1.01-1.28	1.17 ± 0.02	0.08	6.93
		lp	0.70-0.95	0.82 ± 0.05	0.19	6.91
		h	1.43-2.07	1.84 ± 0.04	0.17	9.01
I/2-3	15	L	0.97-1.24	1.12 ± 0.02	0.07	6.54
		lp	0.72-0.90	0.81 ± 0.02	0.06	7.40

			N	L/lp/la/h	LV	M ± SE	σ	V
1								
2								
3								
4								
5	Lower	15		h	1.73-2.00	1.84 ± 0.02	0.08	4.17
6	canine			L	1.26-1.98	1.46 ± 0.04	0.16	11.1
7				lp	0.79-0.97	0.87 ± 0.01	0.04	5.08
8	P/1	15		h	1.76-2.13	1.91 ± 0.03	0.11	5.96
9		16		L	2.35-3.17	2.83 ± 0.07	0.26	9.34
10				lp	0.86-1.10	0.97 ± 0.02	0.08	7.90
11				la	0.76-0.96	0.87 ± 0.02	0.06	7.01
12	P/2	10		h	1.43-1.83	1.63 ± 0.03	0.12	7.09
13				L	3.10-3.58	3.32 ± 0.05	0.17	5.16
14				lp	1.02-1.18	1.08 ± 0.01	0.04	4.00
15				la	0.86-0.95	0.91 ± 0.01	0.04	4.23
16	P/3	17		h	1.45-1.97	1.69 ± 0.05	0.16	9.29
17				L	3.48-4.05	3.76 ± 0.04	0.16	4.24
18				lp	1.38-1.71	1.56 ± 0.02	0.10	6.50
19				la	1.05-1.29	1.16 ± 0.02	0.07	5.69
20				h	1.41-2.06	1.82 ± 0.04	0.18	9.80
21	P/4	18		L	2.92-3.42	3.14 ± 0.03	0.14	4.51
22				lp	1.83-2.50	2.12 ± 0.04	0.18	8.42
23				la	1.44-1.79	1.62 ± 0.03	0.11	6.64
24				h	1.34-2.35	1.79 ± 0.07	0.29	16.20
25	M/1	17		L	2.96-3.40	3.15 ± 0.03	0.12	3.92
26				lp	2.33-2.81	2.58 ± 0.03	0.14	5.53
27				la	2.12-2.73	2.37 ± 0.04	0.15	6.50
28				h	1.33-2.15	1.73 ± 0.05	0.22	12.49
29	M/2	15		L	3.19-3.51	3.34 ± 0.02	0.09	2.82
30				lp	2.54-3.01	2.72 ± 0.04	0.14	5.18
31				la	2.39-3.01	2.60 ± 0.04	0.17	6.68
32				h	1.48-2.38	1.99 ± 0.07	0.26	13.01
33	M/3	15		L	4.35-4.77	4.51 ± 0.03	0.12	2.69
34				lp	2.25-2.63	2.42 ± 0.03	0.12	5.06
35				la	2.17-2.67	2.39 ± 0.04	0.16	6.56
36				h	1.73-2.50	2.18 ± 0.06	0.24	11.16
37				L3	1.02-1.43	1.22 ± 0.03	0.12	9.45
38				I3	1.05-1.52	1.23 ± 0.03	0.12	9.87
39	M/1-M/3	14		L	10.62-11.39	10.97 ± 0.07	0.26	2.36
40	P/3-P/4	19		L	6.26-7.29	6.82 ± 0.06	0.26	3.80
41	P/2-P/4	10		L	10.32-11.61	11.06 ± 0.13	0.40	3.59
42	P/1-P/4	2		L	13.39-13.44	13.41		
43	Dia P/2-P/3	16		L	0.45-1.58	1.11 ± 0.08	0.33	30.00
44	Dia P/1-P/2	8		L	0.05-0.85	0.49 ± 0.06	0.21	42.81

Table S2. Measurements and statistical data of the permanent upper teeth of *Paroxacron valdense* of DAM1. N, number of specimens; L/lp/la/h, see Methods; LV, limit values; M ± SE, Mean ± Standard Error; σ , standard deviation; V, coefficient of variation.

	I1/	10	L	1.73-2.03	1.92 ± 0.03	0.11	5.56
			lp	0.80-0.93	0.87 ± 0.01	0.04	4.97
			h	1.55-1.93	1.72 ± 0.05	0.15	8.53
	I2/	15	L	1.53-1.90	1.74 ± 0.03	0.1	5.73
			lp	0.70-0.95	0.81 ± 0.02	0.07	8.71
			h	1.63-2.14	1.88 ± 0.03	0.13	6.86
	I3/	15	L	1.45-1.75	1.56 ± 0.02	0.09	5.83
			lp	0.65-0.94	0.78 ± 0.02	0.07	8.65
			h	1.49-1.94	1.77 ± 0.04	0.14	8.21
	Upper canine	15	L	1.56-2.00	1.75 ± 0.03	0.13	7.31
			lp	0.90-1.12	1.02 ± 0.02	0.06	6.23
			h	1.99-2.23	2.08 ± 0.02	0.07	3.56
	P1/	16	L	2.67-3.54	3.08 ± 0.06	0.23	7.59
			lp	0.82-1.05	0.94 ± 0.01	0.07	5.98
			la	0.87-1.05	0.96 ± 0.01	0.05	5.26
			h	1.49-1.96	1.76 ± 0.03	0.13	7.56
	P2/	11	L	3.46-4.15	3.75 ± 0.06	0.21	5.67
			lp	1.15-1.73	1.34 ± 0.05	0.18	13.2
			la	1.04-1.20	1.11 ± 0.02	0.06	5.49
			h	1.49-1.87	1.72 ± 0.04	0.14	8.03
	P3/	12	L	3.21-3.86	3.53 ± 0.06	0.22	6.31
			lp	2.55-3.18	2.83 ± 0.05	0.17	6.17
			la	1.39-1.86	1.68 ± 0.04	0.14	8.09
			h	1.39-2.18	1.85 ± 0.07	0.23	12.2
	P4/	17	L	2.39-2.78	2.55 ± 0.03	0.12	4.64
			lp	3.07-3.66	3.42 ± 0.04	0.15	4.46
			h	1.65-2.58	2.03 ± 0.06	0.24	11.92
	M1/	15	L	2.94-3.46	3.24 ± 0.04	0.15	4.54
			lp	3.71-4.33	3.99 ± 0.05	0.20	4.92
			la	3.26-3.75	3.52 ± 0.04	0.14	4.11
			h	1.40-1.90	1.58 ± 0.03	0.13	8.53
	M2/	15	L	3.34-3.74	3.55 ± 0.03	0.12	3.35
			lp	4.13-4.71	4.37 ± 0.04	0.16	3.60
			la	3.81-4.36	4.07 ± 0.05	0.18	4.40
			h	1.62-2.09	1.81 ± 0.04	0.16	8.78
	M3/	13	L	3.13-3.65	3.37 ± 0.04	0.15	4.55
			lp	3.92-4.53	4.13 ± 0.04	0.16	3.76
			la	3.79-4.16	3.97 ± 0.03	0.12	3.05
			h	1.59-1.97	1.79 ± 0.04	0.13	7.14
	M1-M3/	13	L	9.08-10.57	9.89 ± 0.1	0.36	3.64
	P3/-P4/	13	L	5.38-6.51	5.95 ± 0.09	0.33	5.48
	P2/-P4/	10	L	8.95-10.44	9.62 ± 0.16	0.51	5.33
	P1/-P4/	1	L		13.229		
	Dia P1/-P2/	10	L	0-1.46	0.94 ± 0.13	0.41	43.17

Table S3. Measurements and statistical data of the deciduous upper and lower teeth of *Paroxacron valdense* of DAM1. N, number of specimens; L/lp/la/h and lam, see Methods; LV, limit values; M \pm SE, Mean \pm Standard Error; σ , standard deviation; V, coefficient of variation.

	N	L/lp/la/h	LV	M ± SE	σ	V
DP1-2/	5	L	3.56-3.80	3.66 ± 0.04	0.09	2.38
		lp	1.13-1.55	1.26 ± 0.08	0.17	13.61
		la	0.80-1.00	0.90 ± 0.04	0.08	8.90
		h	1.20-1.44	1.29 ± 0.04	0.09	6.90
DP3/	15	L	3.24-3.71	3.54 ± 0.04	0.15	4.16
		lp	2.14-2.73	2.49 ± 0.04	0.17	6.70
		la	1.23-1.54	1.40 ± 0.03	0.10	6.90
		h	1.10-1.43	1.28 ± 0.03	0.10	7.74
DP4/	15	L	2.87-3.14	3.02 ± 0.02	0.08	2.74
		lp	3.10-3.85	3.40 ± 0.05	0.20	5.84
		la	2.91-3.29	3.03 ± 0.03	0.13	4.22
		h	1.10-1.79	1.51 ± 0.05	0.18	11.77
DP/1	7	L	2.92-3.28	3.10 ± 0.05	0.13	4.33
		lp	0.78-0.94	0.87 ± 0.02	0.06	6.77
		la	0.71-0.94	0.84 ± 0.03	0.08	9.45
DP/2-3	9	L	1.07-1.35	1.24 ± 0.04	0.10	7.81
		lp	3.11-3.87	3.66 ± 0.08	0.24	6.52
		la	1.22-1.70	1.47 ± 0.05	0.14	9.17
		h	0.92-1.21	1.06 ± 0.03	0.10	9.61
DP/4	15	L	1.40-1.95	1.68 ± 0.07	0.20	11.88
		lp	3.61-4.05	3.80 ± 0.03	0.12	3.19
		la	1.90-2.22	2.08 ± 0.03	0.11	5.35
		h	1.57-1.76	1.62 ± 0.01	0.05	2.90
		lam	1.42-2.15	1.73 ± 0.05	0.21	12.08

Table S4. Measurements and statistical data of the upper teeth of *Paroxacron* sp. of DAM3. N, number of specimens; L/lp/la/h, see Methods; LV, limit values; M ± SE, Mean ± Standard Error; σ , standard deviation; V, coefficient of variation.

	N	L/lp/la/h	LV	M ± SE	σ	V
P1/	1	L		2.73		
		lp		1.17		
		la		1.01		
		h		1.56		
P4/	1	L		2.25		
		lp		2.85		
		h		1.93		
M1/	3	L	2.59-2.71	2.64 ± 0.04	0.06	2.42
		lp	3.47-3.70	3.61 ± 0.07	0.13	3.50
		la	3.05-3.58	3.39 ± 0.17	0.29	8.69
		h	1.08-1.36	1.23 ± 0.08	0.14	11.50
M2/	2	L	2.94-3.04	2.99 ± 0.05	0.07	2.48
		lp	3.94-3.95	3.94 ± 0.01	0.01	0.20
		la	3.26-3.67	3.46 ± 0.20	0.29	8.35

			h	1.62-1.68	1.65 ± 0.03	0.04	2.53
4	M3/	3	L	2.88-3.09	3.00 ± 0.06	0.10	3.44
5			lp	3.58-3.92	3.76 ± 0.10	0.17	4.59
6			la	3.49-3.61	3.53 ± 0.04	0.07	1.96
7			h	1.30-1.75	1.52 ± 0.13	0.23	15.08

Table S5. Measurements and statistical data of the lower teeth of *Paroxacron* sp. of DAM3. N, number of specimens; L/lp/la/h/L3/l3 and lam, see Methods; LV, limit values; M \pm SE, Mean \pm Standard Error; σ , standard deviation; V, coefficient of variation.

		N	L/Lp/la/h	LV	$M \pm SE$	σ	V
21	P/1	1	L		2.53		
22			lp		1.00		
23			la		0.79		
24			h		1.54		
25	P/2	1	L		3.02		
26			lp		1.15		
27			la		1.00		
28			h		1.39		
29	DP/4	2	L	3.16-3.25	3.21 ± 0.05	0.07	2.14
30			lp	1.81-1.92	1.87 ± 0.05	0.08	4.13
31			la	1.40-1.67	1.53 ± 0.13	0.19	12.26
32			h	1.13-1.32	1.23 ± 0.09	0.13	10.74
33			lam	1.41-1.65	1.53 ± 0.12	0.17	11.09
34	M/1	2	L	2.55-2.94	2.75 ± 0.20	0.28	10.06
35			lp	1.86-2.09	1.98 ± 0.11	0.16	8.09
36			la	1.71-1.95	1.83 ± 0.12	0.17	9.23
37			h	1.03-1.37	1.20 ± 0.17	0.24	19.85
38	M/2	4	L	2.87-3.18	2.97 ± 0.07	0.14	4.83
39			lp	2.01-2.25	2.17 ± 0.05	0.11	4.91
40			la	1.88-2.11	1.99 ± 0.05	0.09	4.61
41			h	1.36-1.92	1.68 ± 0.01	0.23	13.93
42	M/3	1	L		3.89		
43			lp		2.04		
44			la		1.96		
45			h		2.22		
46			L3		1.04		
47			l3		1.34		

Table S6. Measurements and statistical data of the upper teeth of *Paroxacron bergeri* of DAM3. N, number of specimens; L/lp/la/h, see Methods; LV, limit values; M \pm SE, Mean \pm Standard Error; σ , standard deviation; V, coefficient of variation.

	N	L/lp/la/h	LV	M ± SE	σ	V
P1/	1	L		3.04		
		lp		1.25		
		la		1.33		
		h		1.06		
P2/	1	L		4.22		
		lp		1.63		
		la		1.52		
		h		2.48		
P3/	1	L		3.79		
		lp		2.84		
		la		2.01		
		h		2.08		
P4/	1	L		3.09		
		lp		3.98		
		la				
		h		2.36		
M1/	2	L	4.08-4.48	4.28 ± 0.20	0.28	6.64
		lp	4.85-5.11	4.98 ± 0.13	0.18	3.66
		la	4.55-4.81	4.68 ± 0.13	0.19	4.05
		h	1.68-2.45	2.07 ± 0.38	0.54	26.27
M2/	3	L	4.70-5.05	4.83 ± 0.11	0.19	4.00
		lp	5.14-5.69	5.50 ± 0.18	0.31	5.70
		la	4.67-5.44	5.13 ± 0.24	0.41	7.96
		h	2.25-2.78	2.59 ± 0.17	0.30	11.47
M3/	13	L	4.52-4.61	4.56 ± 0.03	0.05	1.06
		lp	5.06-5.63	5.40 ± 0.17	0.30	5.54
		la	4.98-5.32	5.17 ± 0.10	0.17	3.34
		h	1.95-2.75	2.38 ± 0.24	0.41	17.23

Table S7. Measurements and statistical data of the upper teeth of *Plesiomeryx cadurcensis* of DAM3.

N, number of specimens; L/lp/la/h, see Methods; LV, limit values; M ± SE, Mean ± Standard Error; σ , standard deviation; V, coefficient of variation.

	N	L/lp/la/h	LV	M ± SE	σ	V
P1/	1	L		3.05		
		lp		1.28		
		la		1.07		
		h		1.25		
P2/	2	L	3.42-3.48	3.45 ± 0.03	0.04	1.15
		lp	1.48-1.69	1.59 ± 0.11	0.15	9.55
		la	1.26-1.33	1.29 ± 0.04	0.05	3.84
		h	1.52-1.69	1.60 ± 0.08	0.11	7.10
P3/	1	L		3.44		
		lp		2.94		
		la		1.81		
		h		1.54		
P4/	1	L		2.50		
		lp		3.25		

			h		1.44	
DP4/	2	L	3.01-3.07	3.04 ± 0.03	0.04	1.37
		lp	3.62-3.91	3.77 ± 0.15	0.21	5.51
		la	3.08-3.58	3.33 ± 0.25	0.35	10.64
		h	1.20-1.24	1.22 ± 0.02	0.03	2.66
M1/	3	L	3.08-3.34	3.22 ± 0.08	0.13	4.07
		lp	4.15-4.28	4.22 ± 0.04	0.06	1.50
		la	3.57-3.90	3.76 ± 0.10	0.17	4.62
		h	1.40-1.90	1.72 ± 0.10	0.17	9.89
M2/	5	L	3.51-3.78	3.61 ± 0.05	0.10	2.84
		lp	4.50-4.85	4.65 ± 0.06	0.14	3.04
		la	3.83-4.32	4.07 ± 0.10	0.22	5.52
		h	1.75-2.01	1.86 ± 0.05	0.10	5.49
M3/	3	L	3.14-3.69	3.45 ± 0.17	0.31	8.78
		lp	4.17-4.60	4.45 ± 0.14	0.25	5.57
		la	3.86-4.12	4.01 ± 0.08	0.14	3.50
		h	1.74-2.00	1.84 ± 0.08	0.14	7.62

Table S8. Measurements and statistical data of the lower teeth of *Plesiomeryx cadurcensis* of DAM3.

N, number of specimens; L/lp/la/h/L3 and l3, see Methods; LV, limit values; M ± SE, Mean ± Standard Error; σ, standard deviation; V, coefficient of variation.

	N	L/Lp/la/h	LV	M ± SE	Σ	V
P/2	3	L	2.52-2.81	2.70 ± 0.09	0.16	5.80
		lp	1.07-1.29	1.16 ± 0.07	0.11	9.75
		la	0.93-1.03	0.97 ± 0.03	0.05	5.57
		h	1.46-1.77	1.61 ± 0.09	0.16	9.66
P/3	7	L	2.89-3.27	3.03 ± 0.05	0.14	4.50
		lp	1.35-1.61	1.49 ± 0.04	0.11	7.22
		la	1.00-1.35	1.21 ± 0.04	0.11	9.50
		h	1.60-2.13	1.79 ± 0.08	0.20	11.35
P/4	5	L	2.99-3.51	3.16 ± 0.09	0.20	6.46
		lp	1.85-2.03	1.96 ± 0.04	0.08	4.09
		la	1.51-1.76	1.60 ± 0.04	0.10	6.04
		h	1.50-2.58	1.88 ± 0.19	0.42	22.38
M/1	6	L	3.05-3.20	3.09 ± 0.02	0.06	1.85
		lp	2.24-2.57	2.37 ± 0.05	0.13	5.67
		la	1.89-2.41	2.13 ± 0.08	0.19	9.01
		h	1.40-2.08	1.79 ± 0.12	0.30	17.01
M/2	5	L	3.25-3.67	3.42 ± 0.08	0.17	5.08
		lp	2.47-2.89	2.64 ± 0.08	0.18	6.69
		la	2.14-2.59	2.42 ± 0.08	0.17	7.17
		h	1.79-2.29	1.98 ± 0.09	0.19	9.73
M/3	3	L	4.18-4.39	4.30 ± 0.06	0.11	2.52
		lp	2.14-2.28	2.21 ± 0.04	0.07	3.31
		la	2.13-2.25	2.19 ± 0.04	0.06	2.84
		h	1.89-2.29	2.14 ± 0.13	0.22	10.32

		L3	1.12-1.16	1.14 ± 0.01	0.02	2.05
		I3	1.26-1.65	1.44 ± 0.11	0.20	13.70
P/3-P/4	1	L		5.97		

Table S9. Measurements and statistical data of the upper teeth of *Plesiomeryx huerzeleri* of DAM3. N, number of specimens; L/lp/la/h, see Methods; LV, limit values; M \pm SE, Mean \pm Standard Error; σ , standard deviation; V, coefficient of variation.

	N	L/lp/la/h	LV	M \pm SE	σ	V
P1/	1	L		3.23		
		lp		1.23		
		la		1.36		
		h		1.81		
P2/	2	L	4.24-4.46	4.35 ± 0.11	0.16	3.64
		lp	2.00-2.08	2.04 ± 0.04	0.06	2.85
		la	1.62-1.86	1.74 ± 0.12	0.17	9.60
		h	2.08-2.87	2.47 ± 0.39	0.55	22.42
P3/	1	L		4.08		
		lp		3.52		
		la		2.04		
		h		1.39		
P4/	1	L		3.25		
		lp		4.37		
		h		1.46		
M1/	9	L	3.75-4.06	3.92 ± 0.03	0.08	2.16
		lp	4.68-5.29	5.00 ± 0.07	0.20	4.00
		la	4.11-4.80	4.48 ± 0.07	0.22	4.99
		h	1.36-2.26	1.85 ± 0.09	0.27	14.75
M2/	6	L	3.95-4.49	4.23 ± 0.10	0.25	5.88
		lp	5.17-5.79	5.46 ± 0.10	0.24	4.39
		la	4.49-5.33	4.92 ± 0.12	0.30	6.01
		h	1.58-2.97	2.08 ± 0.21	0.52	24.87
M3/	13	L	4.15-4.25	4.20 ± 0.03	0.05	1.24
		lp	5.00-5.25	5.12 ± 0.07	0.13	2.48
		la	4.14-4.95	4.56 ± 0.24	0.41	8.95
		h	1.55-2.25	2.00 ± 0.22	0.39	19.46
P3/-P4/	1	L		7.14		

Table S10. Measurements and statistical data of the lower teeth of *Plesiomeryx huerzeleri* of DAM3. N, number of specimens; L/lp/la/h/L3/I3 and lam, see Methods; LV, limit values; M \pm SE, Mean \pm Standard Error; σ , standard deviation; V, coefficient of variation.

	N	L/Lp/la/h	LV	M ± SE	σ	V
P/1	1	L		2.77		
		lp		1.37		
		la		1.25		
		h		1.65		
P/2	1	L		3.41		
		lp		1.50		
		la		1.39		
		h		1.85		
P/3	9	L	3.51-4.17	3.84 ± 0.07	0.20	5.18
		lp	1.61-2.01	1.81 ± 0.05	0.14	7.61
		la	1.40-1.64	1.52 ± 0.03	0.09	6.18
		h	2.05-2.59	2.37 ± 0.06	0.17	7.06
DP/3	3	L	3.83-4.08	3.92 ± 0.08	0.14	3.53
		lp	1.77-1.91	1.85 ± 0.04	0.07	3.78
		la	1.45-1.66	1.53 ± 0.07	0.11	7.47
		h	1.29-2.01	1.66 ± 0.21	0.36	21.72
P/4	1	L		4.00		
		lp		2.34		
		la		2.12		
		h		2.31		
DP/4	1	L		4.92		
		lp		2.76		
		la		2.33		
		h		1.75		
		lam		2.35		
M/1	4	L	3.70-4.02	3.85 ± 0.07	0.13	3.48
		lp	2.73-3.09	2.89 ± 0.09	0.18	6.19
		la	2.49-2.93	2.66 ± 0.10	0.19	7.17
		h	1.60-2.19	1.96 ± 0.14	0.28	14.34
M/2	4	L	4.17-4.57	4.36 ± 0.08	0.17	3.87
		lp	3.12-3.20	3.16 ± 0.02	0.03	1.06
		la	2.90-3.00	2.95 ± 0.03	0.05	1.81
		h	2.16-2.84	2.35 ± 0.16	0.33	13.91
M/3	4	L	4.94-4.51	5.25 ± 0.12	0.23	4.46
		lp	2.60-2.80	2.70 ± 0.04	0.08	3.14
		la	2.57-2.76	2.67 ± 0.04	0.08	3.01
		h	1.38-2.44	2.02 ± 0.23	0.46	22.78
		L3	1.18-1.44	1.32 ± 0.06	0.12	8.81
		l3	1.62-1.92	1.77 ± 0.07	0.14	7.78
P/3-P/4	1	L		7.64		

Table S11. Measurements and statistical data of the upper teeth of *Caenomeryx cf. procommunis* of DAM3. N, number of specimens; L/lp/la/h, see Methods; LV, limit values; M ± SE, Mean ± Standard Error; σ , standard deviation; V, coefficient of variation.

	N	L/lp/la/h	LV	M ± SE	σ	V
P3/	1	L		3.46		
		lp				
		la		2.36		
		h		2.66		
M1/	2	L	4.80-5.13	4.96 ± 0.16	0.23	4.66
		lp	5.76-6.09	5.92 ± 0.17	0.23	3.94
		la	5.40-5.49	5.44 ± 0.04	0.06	1.16
		h	1.67-2.11	1.89 ± 0.22	0.31	16.59
M2/	3	L	5.07-5.35	5.18 ± 0.09	0.16	3.01
		lp	6.07-6.16	6.11 ± 0.03	0.05	0.73
		la	5.60-5.79	5.67 ± 0.06	0.11	1.87
		h	2.66-3.06	2.85 ± 0.11	0.20	6.96
M3/	1	L		4.83		
		lp		6.07		
		la		5.50		
		h		2.43		

Table S12. Measurements and statistical data of the lower teeth of *Caenomeryx cf. procommunis* of DAM3. N, number of specimens; L/lp/la/h/L3 and l3, see Methods; LV, limit values; M ± SE, Mean ± Standard Error; σ , standard deviation; V, coefficient of variation.

	N	L/lp/la/h	LV	M ± SE	σ	V
P/1	1	L		3.19		
		lp		1.39		
		la		1.22		
		h		1.69		
P/2	1	L		4.21		
		lp		1.89		
		la		1.72		
		h		2.72		
P/3	1	L		4.02		
		lp		2.03		
		la		1.81		
		h		2.64		
DP/3	2	L	4.57-4.66	4.62 ± 0.04	0.06	1.32
		lp	1.82-1.90	1.86 ± 0.04	0.06	3.01
		la	1.64-1.69	1.66 ± 0.02	0.03	1.53
		h	1.82-2.23	2.03 ± 0.21	0.29	14.38
P/4	4	L	4.01-4.40	4.23 ± 0.08	0.17	3.93
		lp	2.79-3.13	2.92 ± 0.07	0.15	5.09
		la	1.95-2.49	2.20 ± 0.11	0.23	10.25
		h	1.66-3.47	2.37 ± 0.39	0.78	33.02
M/1	4	L	4.13-4.39	4.32 ± 0.06	0.13	2.91
		lp	3.23-3.84	3.57 ± 0.14	0.28	7.87
		la	3.15-3.66	3.40 ± 0.12	0.24	6.96

1						
2						
3						
4	M/2	5	h	1.36-2.80	2.05 ± 0.34	0.67
5			L	4.32-4.47	4.39 ± 0.03	0.06
6			lp	3.23-3.68	3.44 ± 0.07	0.16
7			la	3.08-3.66	3.30 ± 0.10	0.22
8			h	2.17-2.80	2.50 ± 0.10	0.23
9	M/3	4	L	6.02-6.77	6.33 ± 0.18	0.35
10			lp	2.79-3.41	3.13 ± 0.11	0.25
11			la	2.90-3.35	3.15 ± 0.11	0.22
12			h	2.34-3.44	3.03 ± 0.25	0.49
13			L3	1.64-2.14	1.99 ± 0.09	0.21
14			l3	2.04-2.27	2.16 ± 0.05	0.11
15	M/1-M/3	1	L		14.73	
16						