



HAL
open science

Modulatory effects of sesame oil and ascorbic acid on abamectin-induced oxidative stress in rat liver and brain tissues

Abeer M. Radi, Eman T. Mohammed, Abdelrahman Ibrahim Abushouk, Lotfi Aleya, Mohamed M. Abdel-Daim

► To cite this version:

Abeer M. Radi, Eman T. Mohammed, Abdelrahman Ibrahim Abushouk, Lotfi Aleya, Mohamed M. Abdel-Daim. Modulatory effects of sesame oil and ascorbic acid on abamectin-induced oxidative stress in rat liver and brain tissues. *Science of the Total Environment*, 2019, pp.134882. 10.1016/j.scitotenv.2019.134882 . hal-02349580

HAL Id: hal-02349580

<https://hal.science/hal-02349580>

Submitted on 21 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

Modulatory effects of sesame oil and ascorbic acid on abamectin-induced oxidative stress in rat liver and brain tissues

Abeer M. Radi¹, Eman T. Mohammed², Abdelrahman Ibrahim Abushouk³, Lotfi Aleya^{*4}

Mohamed M. Abdel-Daim^{5,6}

¹ Pharmacology Department, Faculty of Veterinary Medicine, Beni-Suef University, Beni-Suef 62515, Egypt

² Biochemistry Department, Faculty of Veterinary Medicine, Beni-Suef University, Beni-Suef 62515, Egypt

³ Faculty of Medicine, Ain Shams University, Cairo, 11566, Egypt

^{*4}Chrono-Environnement Laboratory, UMR CNRS 6249, Bourgogne Franche-Comté University, F-25030, Besançon Cedex, France. (* **Corresponding author: lotfi.aleya@univ-fcomte.fr**)

⁵Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia,

⁶ Pharmacology Department, Faculty of Veterinary Medicine, Suez Canal University, Ismailia 41522, Egypt

29 **Abstract**

30
31 The present work was designed to assess the modulatory effects of sesame oil (SO) and
32 ascorbic acid (AA) on abamectin (ABM)-induced oxidative stress and altered gene expression of
33 hepatic cytochrome P450 2E1 (CYP-2E1), p38 MAPK, and caspase-3 and cerebral P-
34 glycoprotein (Abcb1a receptor). Male rats were distributed into five groups (6 rats/group),
35 receiving distilled water, ABM 2 mg/kg bwt 1/5 LD₅₀ orally for 5 days, ABM+AA 100 mg/kg
36 bwt orally, ABM+SO 5 mL/kg bwt orally, or ABM+SO+AA at the aforementioned doses.
37 Nineteen compounds were identified in the SO sample by GC-MS analysis, including
38 tetradecane,2,6,10-trimethyl, octadecane, 1-hexadecanol,2-methyl, and octadecane,6-methyl.
39 Abamectin significantly upregulated the hepatic CYP-2E1 expression with excess generation of
40 oxidative radicals, as evident by the significant depletion of reduced glutathione and elevation of
41 malondialdehyde concentration ($p \leq 0.05$) in rat liver and brain tissues. Further, ABM
42 significantly increased TNF- α concentration, the expression of caspase-3 and p38 MAPK in the
43 liver, as well as p-glycoprotein and GABA-A receptor in the brain. These results were in line
44 with the observed histopathological changes. Sesame oil and/or AA supplementation alleviated
45 ABM-induced cell damage by modulating all tested parameters. In conclusion, ABM induces
46 oxidative stress and increases the expression of CYP-2E1, caspase-3, and p38 MAPK in the
47 liver, as well as P-gp and GABA-A receptor in the brain. These effects could be ameliorated by
48 SO and AA, alone and in combination, probably due to their anti-oxidant, anti-apoptotic, and
49 gene-regulating activities.

50 **Keywords:** abamectin; oxidative stress; sesame oil; ascorbic acid; liver and brain; rats

51

52 **Introduction**

53 Pesticides are used worldwide to control pests, insects, and disease vectors in veterinary
54 medicine and agriculture (Jones, 2018). Abamectin (ABM) belongs to the avermectins family;
55 16-membered macrocyclic lactones with a disaccharide substituent at the carbon-13 position. It
56 is generated from soil actinomycete fungi (*Streptomyces avermitilis*) and is a combination of 80%
57 avermectin B1a plus 20% avermectin B1b. Abamectin is used as an anthelmintic agent in cattle
58 and sheep to eliminate gastrointestinal nematodes, lung worms and nasal bots (Campbell, 2012).
59 Non-therapeutic exposures to ivermectin and other macrocyclic lactones may lead to toxic
60 effects; only after oral ingestion of a large amount. Although the exact mechanisms remain
61 unclear, macrocyclic lactones, when taken in large doses, may pass through the blood-brain
62 barrier and produce GABA-mimetic toxic effects. In mammals, avermectin intoxication begins
63 with hyperexcitability, incoordination, tremors, hypotension, and later develops into ataxia,
64 coma, respiratory failure, and even death (Yang, 2012).

65 Avermectin is poorly metabolized in mammals; 80–98% of the initially administered dose gets
66 eliminated in stool (Sun et al., 2005). The highest levels of ABM were detected in the liver and
67 fat (owing to its lipophilic nature), while the lowest was found in the brain (Gonzalez Canga et
68 al., 2009). Therefore, the detoxification of ABM may disturb the functions of hepatocytes. The
69 biodistribution of ivermectin in the host depends on the efflux by P-glycoprotein (P-gp) and
70 biotransformation by cytochromes P450 (Alberich et al., 2014). P-gp prevents brain toxicity by
71 limiting the penetration of toxic compounds across the blood-brain barrier (Roulet et al., 2003).
72 Besides, it contributes to the intestinal excretion of ivermectin (Ballent et al., 2006). Hepatic
73 drug metabolism is attained by the microsomal cytochrome P450 enzyme system that facilitates
74 transformation into toxic intermediates, followed by reactive oxygen species (ROS) production,

75 inflammatory cytokines release and lipid peroxidation (Lu et al., 2012; Maioli et al., 2013).
76 These events initiate apoptosis and tissue inflammation through interaction with caspases and
77 mitogen-activated protein kinases (MAPKs) (Bayir and Kagan, 2008).

78 Oxidative stress is a key factor in avermectins-induced cytotoxicity (Zhu et al., 2013). Therefore,
79 the use of antioxidants to ameliorate its toxicity is a logical approach. Ascorbic acid is a water-
80 soluble non-enzymatic antioxidant that defends the cellular compartments against reactive
81 oxygen species (ROS) (Jurczuk et al., 2007). It is an essential element; must be supplied in diet
82 because it cannot be synthesized in mammalian bodies. Ascorbic acid showed the ability to
83 prevent lipid peroxidation and protect lipid membranes, proteins and DNA from oxidative harm
84 (Granger and Eck, 2018). Moreover, it could antagonize the toxic effects of many xenobiotics
85 (Abdel-Daim et al., 2019a; Abdel-Daim et al., 2019b; Özkan et al., 2012) and has been shown to
86 renovate the antioxidant capacity of vitamin E (Serbecic and Beutelspacher, 2005).

87 Sesame oil (SO) is extracted from *Sesamum indicum* seeds. These seeds contain many
88 phytochemicals as flavonoids, phenolic acids, tannins, alkaloids, terpenoids, cephalin and
89 lecithin and some minerals like iron, calcium, magnesium, copper, zinc, manganese and
90 phosphorus (Anilakumar et al., 2010; Sani et al., 2013). Several lignans, such as sesamol,
91 sesamin, and γ -tocopherol are potent polyphenolic antioxidants found in sesame seeds
92 (Rangkadilok et al., 2010). For example, sesamin protects against oxidative stress and enhances
93 hepatic drug detoxification (Shuang et al., 2018; Zhang et al., 2016a). Over decades of research,
94 sesame (oil) exhibited anti-inflammatory, antibacterial, hypolipidemic, antitumor (Anilakumar et
95 al., 2010) and anti-allergic effects (Jung et al., 2018).

96 The goals of this experiment were to explore the impact of acute exposure to the commercial
97 formulation of ABM on the expression of metabolic cytochromes P450 2E1 (CYP-2E1), p38

98 MAPK and caspase-3 enzymes in the liver, as well as membrane p-glycoprotein efflux
99 transporter and GABA-A signaling in rat brain. Further, we tested the preventive capacities of
100 SO and/or AA against acute ABM toxicity.

101

102 **2. Materials and methods**

103 **2.1. Chemicals:** Abamectin (*Vertemic*[®], 1.8% EC) was bought from *Syngenta Agro Services AG*,
104 Egypt. Sesame Oil was supplied by *El-Captain Company* (El-Obour City, Egypt) and AA
105 (*Ascorbin*[®] 100% B.p) was obtained from *Newvetrovit Company*, Egypt. Commercial kits for
106 malondialdehyde (MDA), reduced glutathione (GSH) and catalase (CAT) were purchased from
107 *Biodiagnostic Company*, Egypt. The kits for SGPT, SGOT, ALP enzymes, and direct bilirubin
108 were supplied from *Greiner Diagnostic GmbH-Bahlingen, Germany*. The rest of used chemicals
109 were of analytical grade. The chemical composition of sesame oil was analyzed using Trace GC-
110 ISQ Mass Spectrometer (Thermo Scientific, Austin, TX, USA) as described in [Supplementary](#)
111 [file 1](#).

112 **2.2. Experimental design:** Thirty mature male albino rats (weighing 120 to 150 g), were
113 purchased from the *Egyptian Company for Biological Products and Vaccines*. Animals were
114 kept in stainless steel cages, fed rat chow and water *ad libitum*, and maintained at lab
115 temperature of 25 ± 2 °C. All maintenance and care procedures were approved by the Research
116 Ethical Committee at the Faculty of Veterinary Medicine, Beni-Suef University, Egypt.

117 After two weeks of acclimation, rats were randomly assigned to five groups ($n = 6$ /group). The
118 first group received distilled water (negative control). The second group (ABM) received a daily
119 oral dose of ABM (2 mg/kg bwt, 1/5 LD₅₀) for 5 days (LD₅₀: 10 mg/kg) (Abdel-Daim and
120 Abdellatif, 2018). The third group (ABM+AA) received ABM plus a daily dose of AA (100

121 mg/kg bwt, orally) (Abdel-Daim and El-Ghoneimy, 2015; Seo and Lee, 2002), while the fourth
122 group (ABM+SO) was given ABM plus a daily dose of SO (5 ml/kg bwt, orally) (Saleem et al.,
123 2014). The fifth group (Combination) received SO and AA with the same doses as before.
124 Groups (3 to 5) were given SO and/ or AA for 10 days before ABM exposure and 1 hour before
125 ABM administration for 5 successive days. **Figure 1** summarizes the experimental design and
126 observed findings.

127 **2.3. Blood sampling and tissue processing:** On the day following the last ABM dose, rats were
128 anesthetized using isoflurane for blood withdrawal from the retro-orbital plexus, and then killed
129 via cervical dislocation. The samples were centrifuged (at 3000 rpm for 15 min) to separate sera.
130 The liver and brain were excised, washed with saline, and then blotted over filter paper. The
131 tissue was divided into three parts: the first (0.5 g) was homogenized in phosphate buffer saline
132 (pH 7.4, 5 ml). Homogenates were later centrifuged at 3000 rpm for 15 min at 4 °C using a
133 cooling, high-speed centrifuge; supernatants were collected and preserved at -80 °C until
134 analysis of tissue oxidant and antioxidant indices. The second tissue part was preserved at -80 °C
135 for molecular investigations. The third tissue part was prepared for histopathological
136 examination.

137 **2.4. Biochemical estimations:** The measurements of MDA, GSH concentrations and CAT
138 activity were performed as per the methods described by Mihara and Uchiyama (Mihara and
139 Uchiyama, 1978), Beutler et al. (Beutler et al., 1963) and Aebi (Aebi, 1984), respectively. Serum
140 samples were used to measure the activities of SGPT and SGOT (Reitman and Frankel, 1957),
141 ALP (Tietz et al., 1983), and bilirubin (Tolman and Rej, 1999).

142
143 **2.5. Detection of TNF- α by ELISA:** Tumor necrosis factor- α (TNF- α) levels were determined

144 by ELISA technique as per the methods of Tietz 1995 (Tietz, 1995). ELISA kits were purchased
145 from *R&D system* (MN, USA).

146 **2.6. Quantitative analysis of gene expression of CYP-2E1, Caspase-3, GABA-A receptor and** 147 **p-gp by real time PCR**

148 **Total RNA extraction:** Total RNA was extracted from the brain and liver tissue homogenates
149 according to the manufacturer's instruction, using *SV Total RNA Isolation System (Promega,*
150 *Madison, WI, USA)*. The RNA concentrations and purity were measured with UV
151 spectrophotometer.

152 **Complementary DNA (cDNA) synthesis:** The extracted RNA was reverse-transcribed into
153 cDNA using high capacity cDNA reverse transcription kits (*#K1621, Fermentas, USA*) following
154 the manufacturer's instructions.

155 **Real-time quantitative PCR:** Real-time qPCR amplification and analysis were performed using
156 an Applied Biosystem with software version 3.1 (*StepOne™, USA*) to measure the expression of
157 mRNAs of target genes in the liver and brain, with *B-actin* as an internal reference (house-
158 keeping gene). The isolated cDNA was amplified using SYBR Green Master Mix (*Applied*
159 *Biosystems*) following the manufacturer's protocol. The primers used in the amplification are
160 shown in [Table 1](#) and were designed by Gene Runner Software (*Hasting Software, Inc., Hasting,*
161 *NY*) from RNA sequences in the gene bank (based on published rat sequences). Data from real-
162 time assays were analyzed using the v1.7 sequence detection software from PE Biosystems
163 (*Foster City, CA*). Relative expression of the studied gene mRNA was calculated using the
164 comparative Cycle threshold (Ct) method. All values were normalized to β -actin and reported as
165 fold change over background levels detected in the treated groups. All these steps were
166 performed according to the methods of Livak and Schmittgen (Livak and Schmittgen, 2001).

167 **2.7. Detection of p38 MAPK protein by Western Blot technique:** p38 MAPK protein was
168 detected following the manufacturer's protocol (*V3 Western Workflow™ Complete System, Bio-*
169 *Rad® Hercules, CA*). In brief, ice-cold radio immune precipitation assay (RIPA) buffer, along
170 with phosphatase/protease inhibitors, were used for protein extraction from liver tissue
171 homogenates. To visualize p38 MAPK protein, we used enhanced chemiluminescence (from
172 *ECL plus; Amersham, IL*), followed by Molecular Analyst Software (*Bio-Rad, Richmond, CA*)
173 for quantification. Protein levels were expressed in relation to β -actin.

174 **2.8. Histopathological studies:** Sections from the liver and brain tissues (M1 motor cortex and
175 hippocampus) were fixed in buffered formalin, then stained with Hematoxylin & Eosin to
176 examine pathological findings under a light microscope.

177 **2.9. Statistical analysis:** Data from the five experimental groups were summarized as means \pm
178 standard errors (SEM), and then transferred to a data sheet on the SPSS software (version 22,
179 IBM Co., Armonk, NY). We used the ANOVA followed by Tukey's post-hoc test for
180 experimental group comparison. P value ≤ 0.05 was accepted for significance.

181

182 **3. Results**

183 **3.1. Sesame oil GC-MS analysis:** In the present study, 19 compounds have been identified in
184 the SO sample by GC-MS analysis ([Table 2](#)). The major components were tetradecane,2,6,10-
185 trimethyl- (38.41%), octadecane (20.52%), 1-hexadecanol, 2-methyl- (9.14 %), octadecane, 6-
186 methyl- (5.72%), 1-tetradecanol (3.85%) and 9-octadecenoic acid (Z)- (3.70%). The mass
187 spectra of all major components in the studied sample are shown in [Supplementary file 1](#).

188 **3.2. Serum biochemical analysis:** Oral administration of ABM was associated with significant
189 ($p \leq 0.0001$) elevations in serum levels of hepatocyte injury biomarkers (SGOT and SGPT) and

190 biliary tract injury biomarkers (ALP and direct bilirubin), compared to the control group. Our
191 results revealed the hepatoprotective effects of SO and AA, alone or in combination, as they
192 significantly ameliorated the ABM-induced alterations in the four parameters. The obtained
193 values in the ABM+SO and ABM+AA+SO combination groups were more frequently
194 comparable to the control group, relative to the ABM+AA group (Table 3).

195 **3.3. Hepatic oxidant/antioxidant markers:** Acute ABM exposure was associated with
196 significant ($p \leq 0.05$) increases in the hepatic tissue MDA concentration and CAT enzyme
197 activity, as well as a significant drop in GSH level, compared to the control rats. Ascorbic acid
198 and/or SO supplementation significantly ($p \leq 0.05$) restored MDA, CAT and GSH to the normal
199 control levels. There were no significant variations among the treated groups (ABM+AA,
200 ABM+SO or Combination group) (Table 4).

201 **3.4. Brain oxidant/antioxidant markers:** Our analysis showed a significant decrease ($p \leq 0.05$)
202 in MDA concentration and significant increases in GSH content and CAT activity in the brain
203 tissues of rats, treated with SO and/or AA, in comparison with ABM-exposed rats. Both
204 treatments, either alone or in combination, could restore MDA and GSH to normal levels;
205 however, the effect of combined treatment was more pronounced than that of a single treatment
206 (Table 4).

207 **3.5. Hepatic tumor necrosis factor- α concentration:** Following ABM exposure, we observed
208 significant elevations ($p \leq 0.05$) in liver tissue TNF- α concentrations in comparison to control
209 rats. Treatments of ABM-intoxicated rats with SO, AA, or their combination was associated with
210 significant reductions in hepatic TNF- α in comparison to rats, exposed to ABM alone. There
211 were no significant variations among the treated groups (ABM+SO, ABM+AA or Combination
212 group); Figure 2.

213 **3.6. Expression of CYP-2E1, caspase-3, and p38 MAPK in the liver:** Abamectin
214 administration was associated with a significant boost in the hepatic tissue expression of CYP-
215 2E1, caspase-3 and p38 MAPK proteins. In contrast, SO and/or AA supplementation ameliorated
216 the ABM-induced increases in CYP-2E1, caspase-3, and p38 MAPK expression. There were no
217 significant variations among the treated groups (ABM+SO, ABM+AA, or Combination groups).
218 **Figure 3** show the observed changes in CYP-2E1, caspase-3, and p38 MAPK expression in rats'
219 liver tissues. **Figure 4** shows the western blotting analysis of p38 MAPK hepatic expression in
220 relation to β -actin.

221 **3.7. Expression of ABC efflux transporter (*Abcb1a*) and GABA-A receptor in the brain:** In
222 this study, the expression of genes encoding the major ABC transporter (*Abcb1a*) was
223 significantly increased following ABM exposure. This over-expression was significantly
224 ameliorated in rats, treated with SO and AA, alone or in combination. Similarly, ABM exposure
225 significantly upregulated GABA-A receptor expression in the brain of ABM-intoxicated rats
226 (compared to normal controls), which was alleviated in rats treated with ABM plus SO, AA, or
227 their combination (**Figure 5**).

228 **3.8. Histopathological findings:** Liver tissue sections from the control group show normal
229 hepatic lobules with granulated and radiating hepatocytic cords from the central vein. However,
230 tissue sections from ABM-exposed rats display lymphocytic infiltration and hemorrhage in the
231 portal area, sinusoidal dilatation, as well as cellular hydropic degeneration, nuclear displacement,
232 and focal ballooning of hepatocytes around the central vein. Marked reductions in these
233 pathological alterations were observed after administration of SO and AA or their combination.
234 Hepatic sections were nearly normal in the three treated groups except that those treated with SO
235 and AA alone showed blood vessels congestion (**Figure 6**).

236 Cerebral cortex tissue sections from the control group show normal histological structure. In
237 contrast, sections from ABM-intoxicated rats show dark stained nuclei, neuropil vaculation and
238 hemorrhage. Rat groups, treated with SO or AA alone, showed marked reduction of hemorrhage,
239 while the combination group sections show lightly-stained vesicular nuclei (Figure 7).
240 Hippocampal tissue sections from the control group show normal outer pleomorphic, middle
241 pyramidal, and inner molecular layers. However, sections from ABM-intoxicated rats show
242 pyknosis, dark stained pyramidal neurons and degenerative changes in the middle pyramidal
243 layer. Interestingly, ABM-intoxicated rats treated with SO only showed few pyknotic pyramidal
244 cells, while rats treated with AA and SO+AA combination show nearly normal pyramidal cells
245 with euchromatic nuclei (Figure 8).

246

247 **4. Discussion**

248 Oxidative stress is a key player in ABM-induced toxicity. The heme protein cytochrome P450
249 multi-enzymatic complex act mainly as mono-oxygenases for the metabolism of many
250 compounds. During the metabolism of toxic substrates by CYP2E1, more reactive and toxic
251 products are formed with excess generation of ROS (Danielson, 2002). These ROS further
252 degrade the CYP heme protein to release iron, which catalyzes the Fenton's reaction, potentiating
253 lipid peroxidation (Caro and Cederbaum, 2004). In the present work, the detected overexpression
254 of hepatic CYP2E1 gene and resulting ROS probably played a role in the observed hepatic injury
255 in ABM-intoxicated rats as indicated by the significant increases in the hepatic lipid peroxidation
256 indicator (MDA concentration) and depletion of GSH stores in the liver. In addition, the brain
257 redox markers were altered following ABM exposure.

258 Reduced glutathione is the principal antioxidant that removes ROS, generated by CYP2E1 (Chen
259 and Cederbaum, 1998), thus GSH depletion (as observed in this study) leads to H₂O₂
260 accumulation, lipid peroxidation and cell damage. Catalase is a prominent endogenous
261 antioxidant enzyme. The higher activity of hepatic CAT, noticed in the present study, may have
262 been an adaptive response against H₂O₂ produced by ABM metabolism. In contrast, CAT
263 activity was reduced in the brain of ABM-intoxicated rats. This result is parallel with that
264 reported by Nasr et al. (Nasr et al., 2016), which reflects the failure of the brain antioxidant
265 capacity to overcome ABM-induced oxidative stress, probably due to the high oxidative
266 metabolism in the brain (Gandhi and Abramov, 2012).

267 The link between ROS and TNF- α is complicated; ROS increase TNF- α release and TNF- α
268 increases ROS production (Blaser et al., 2016). Reactive oxygen species play as second
269 messengers in the intracellular signal transduction pathways, including apoptosis. The p38
270 MAPK pathway is a series of serine/threonine kinases in mammalian cells, activated by
271 excessive ROS generation (Di Lisa et al., 2011) and inflammatory cytokines (Segales et al.,
272 2016). It mediates inflammatory response in various cell types by up-regulating TNF- α ,
273 interleukin-1 and interleukin-8 (Cuenda and Rousseau, 2007). In this study, ABM induced
274 hepatocyte apoptosis, as confirmed by the increased caspase-3 expression in rat liver. These
275 results are comparable to those detected in isolated rat hepatocytes (Maioli et al., 2013) and the
276 pigeon liver (Zhu et al., 2013). The increased caspase-3 expression detected in our study may be
277 related to the increased TNF- α level, which is involved in the extrinsic apoptotic pathway (Perez
278 and White, 2000). Therefore, we suggest the possibility of ABM-induced apoptosis based on the
279 extrinsic apoptotic pathway.

280 In addition, our results show that oral administration of ABM 1/5 LD50 (2 mg/kg bwt.) for 5
281 days significantly increased the serum activities of SGPT, SGOT and ALP in ABM-treated rats,
282 compared to the control group. In parallel with prior investigations (Hsu et al., 2001; Khaldoun-
283 Oularbi et al., 2013), these findings indicate that ABM causes permeability alteration and blood
284 leakage of intracellular enzymes from the damaged hepatocytes. Oxidative stress and
285 inflammatory cytokines activate heme-oxygenase-1 that regulates the synthesis of bilirubin
286 (Yamamoto et al., 2007). Direct bilirubin is actively excreted at the canalicular membrane after
287 binding to transporter proteins (Jedlitschky et al., 1997). Therefore, the damaged hepatocytes
288 may be less able to produce the transporter proteins required for active transportation of direct
289 bilirubin into the gall bladder, explaining the observed increase in its serum levels in the present
290 study.

291 Because avermectins act as GABA-A receptor agonists in vertebrates, their safety in animals
292 requires an intact blood brain barrier with integral P-gp. These are efflux transporters from the
293 ATP-Binding Cassette (ABC) transporters superfamily (Jones and George, 2004), expressed
294 within the capillary endothelial cells in the brain, placenta, and intestine (Ballent et al., 2006). P-
295 gp protects animals against the diffusion of avermectins into the brain, avoiding the consequent
296 neurotoxicity (Macdonald and Gledhill, 2007). In rodent genome, P-gp is encoded as two genes
297 known as *MDR1a/Abcb1a* and *MDR1b/Abcb1b*. *Abcb1a* is more similar to the human gene and
298 is the main form at the blood-brain barrier and intestine (Croop et al., 1989). In the current study,
299 the expression of *Abcb1a* gene was significantly increased following ABM exposure. This
300 finding is in accordance with a former study on ivermectin in murine hepatocytes (Ménez et al.,
301 2012). In contrast, earlier studies showed that avermectin induces neurotoxicity via either
302 oxidative damage, apoptosis (Li et al., 2013) or down regulation of P-gp (Sun et al., 2010).

303 The increased P-gp expression following ABM exposure in this study can be explained by
304 several mechanisms. Reactive oxygen species increase the hepatic expression of P-gp (Deng et
305 al., 2001) and brain endothelial cells in rats (Hong et al., 2006). Moreover, p38 MAPK can
306 activate NF- κ B expression (Saha et al., 2007), which activates MDR1 gene transcription
307 (Bentires-Alj et al., 2003). Additionally, previous investigations have shown that up-regulation
308 of P-gp is stimulated by TNF- α in primary hepatocytes (Hirsch-Ernst et al., 1998) and the p38
309 MAPK pathway in rat prostate cancer cells (Sauvant et al., 2008). In the present study, despite
310 the detected overexpression of *Abcb1a* in the brain, the neurotoxic effect of ABM supports that it
311 is an effective inhibitor of p-gp facilitated transport (Lankas et al., 1998). Therefore, ABM could
312 increase *Abcb1a* expression as an adaptive response to oxidative stress. At the same time, ABM
313 inhibits the P-gp efflux transport function, increasing its penetration into the CNS and facilitating
314 the interaction with GABA receptors.

315 Another finding of this study is that ABM could significantly upregulate GABA-A receptor
316 expression in the cerebral tissue of ABM-intoxicated rats. The mechanism of action of
317 macrocyclic lactones-induced neurotoxicity (including ABM) in pests relies on their high affinity
318 for glutamate-gated Cl channels in neuronal and muscular cells. In mammals, they bind to the
319 receptor for the inhibitory neurotransmitter (GABA) and open the ionotropic GABA-A receptor-
320 gated Cl channels that are limited to the CNS (McCavera et al., 2007). Chloride ions then flow
321 into the postsynaptic neuron in excess, causing hyperpolarization of the membrane potential and
322 disrupting nerve signal transmission (Novelli et al., 2012).

323 Ascorbic acid and SO are commonly used as dietary supplements. Oral SO and/or AA
324 supplementation were able to inhibit the upregulation of CYP-2E1 expression in the liver of
325 treated rats. Similarly, sesamin reduced the expression of CYP-2E1 in hepatocytes (Zhang et al.,

2016b) and alleviated oxidative stress in acetaminophen or carbon tetrachloride-treated mouse liver (Ma et al., 2014). Previous studies have demonstrated that AA could directly react with lipid peroxides, increase GSH and antioxidant enzyme levels, and prevent resulting apoptosis (Santos et al., 2009; Wang et al., 2007). Our results showed that antioxidants like SO and AA eliminated ROS and lipid peroxides in the liver and brain tissues. Additionally, the effect of combined treatment was more pronounced in alleviating the oxidative stress in the brain tissue than that of a single treatment. This may be attributed to the augmenting effect of both treatments.

Administration of SO and AA significantly alleviated ABM-induced apoptosis by inhibiting the hepatic expression of caspase-3, as well as p38 MAPK and TNF- α . The anti-apoptotic activities of SO components and AA have been described in earlier studies. Ascorbic acid inhibited p38 MAPK phosphorylation in vitro (Carcamo et al., 2004). Similarly, Hou et al. reported that the inhibitory effect of sesamol on caspase-3, p38 MAPK activation and ROS production could protect microglia against cell injury (Hou et al., 2004). Ma et al. found that CCl₄-induced apoptosis was inhibited in liver by sesamin through reduction of hepatic TNF- α , Bak, Bax, Cyt. C and caspase-3 expression levels (Ma et al., 2014).

Sesame oil, AA, and their combination could ameliorate the significant increases in serum SGPT, SGOT and ALP, caused by ABM. These changes were more pronounced in combination group compared with that of single treated groups. Previous studies have shown that AA ameliorated the elevated serum SGPT and SGOT levels in rat models of malathion hepatotoxicity (Kalender et al., 2010) and organophosphate pesticide toxicity (Ambali et al., 2007). Further, SO ameliorated subacute diazinon toxicity (Abdel-Daim et al., 2016) and protected brain cells against cypermethrin toxicity (Hussien et al., 2013). Sesame oil

349 significantly alleviated ABM-induced elevation serum direct bilirubin, as observed previously in
350 diazinon intoxicated rats (Al-Attar et al., 2017). These effects may be explained in the light of
351 GC-MS analysis findings; several compounds within the analyzed SO sample, such as
352 octadecane and 9-octadecenoic acid have anti-inflammatory activities and have been shown to
353 mitigate oxidative stress and inhibit the formation of arachidonic acid.

354 In this study, oral SO and/or AA supplementation counteracted the ABM-induced up-regulation
355 of *Abcb1a* expression. Previous research has proven the ability of flavonoids to inhibit P-gp
356 transporters via inhibition of ATPase activity (Pulido et al., 2006) or competitive inhibition
357 (Alvarez et al., 2010). Thus, the phenolic sesamin, sesamol, and flavonoids in SO may be
358 responsible for these effects concerning *Abcb1a* gene (Anilakumar et al., 2010). Further, SO and
359 AA alleviated ABM-induced overexpression of GABA-A receptor in rat brain. The brain is
360 susceptible to oxidative injury due to its high oxygen consumption and a high content of
361 polyunsaturated fatty acids (Gandhi and Abramov, 2012). Redox agents can regulate the GABA-
362 A receptors function (Calero and Calvo, 2008). In contrast, AA has been shown able to control
363 the activity of glutamate and GABA receptors to protect neurons against glutamate excitotoxicity
364 (Calero et al., 2011). Therefore, in the present study, the simultaneous administration of SO and
365 AA could significantly modulate GABA-A receptor mRNA levels owing to their synergistic
366 ROS-scavenging effect.

367 In conclusion, ABM induced oxidative stress and increases the expression of TNF- α , caspase-3
368 and p38 MAPK. Moreover, it upregulates the expression of drug detoxifying genes; the brain
369 *Abcb1a* efflux transporter and the hepatic CYP-2E1 enzyme. However, pretreatments with SO
370 and AA effectively ameliorated ABM-induced oxidative stress and apoptosis. Simultaneous

371 administration of SO and AA was more efficient in protecting the rat liver and brain than single
372 agent use.

373 **Conflicts of Interest:** All authors declare no conflicts of interest

374 **Acknowledgement:** This project was supported King Saud University, Deanship of Scientific
375 Research, College of Science Research Center.

376

377 **Abbreviation:** AA: Ascorbic acid, ABM: Abamectin, ALP: Alkaline phosphatase, CAT:
378 Catalase, CYP-2E1: Cytochrome P-450 2E1, GABA: Gamma aminobutyric acid, GSH: Reduced
379 glutathione, MDA: Malondialdehyde, MAPK: Mitogen-activated protein kinases, P-gp: P-
380 glycoprotein, SGOT: Serum glutamic oxaloacetic transaminase, SGPT: Serum glutamic pyruvic
381 transaminase, SO: Sesame oil, TNF- α : Tumor necrosis factor- α

382

383 **References**

- 384 Abdel-Daim MM, Abdellatief SA. Attenuating effects of caffeic acid phenethyl ester and betaine
385 on abamectin-induced hepatotoxicity and nephrotoxicity. *Environmental Science and*
386 *Pollution Research* 2018: 1-9.
- 387 Abdel-Daim MM, Abushouk AI, Donia T, Alarifi S ,Alkahtani S, Aleya L, et al. The
388 nephroprotective effects of allicin and ascorbic acid against cisplatin-induced toxicity in
389 rats. *Environmental Science and Pollution Research* 2019a: 1-8.
- 390 Abdel-Daim MM, Ahmed A, Ijaz H, Abushouk AI, Ahmed H, Negida A, et al. Influence of
391 *Spirulina platensis* and ascorbic acid on amikacin-induced nephrotoxicity in rabbits.
392 *Environmental Science and Pollution Research* 2019b: 1-7.
- 393 Abdel-Daim MM, El-Ghoneimy A. Synergistic protective effects of ceftriaxone and ascorbic
394 acid against subacute deltamethrin-induced nephrotoxicity in rats. *Renal failure* 2015; 37:
395 297-304.
- 396 Abdel-Daim MM, Taha R, Ghazy EW, El-Sayed YS. Synergistic ameliorative effects of sesame
397 oil and alpha-lipoic acid against subacute diazinon toxicity in rats: hematological,
398 biochemical, and antioxidant studies. *Can J Physiol Pharmacol* 2016; 94: 81-8.
- 399 Aebi H. Catalase in vitro. *Methods Enzymol* 1984; 105: 121-6.
- 400 Al-Attar AM, Elnaggar MHR, Almalki EA. Protective effect of some plant oils on diazinon
401 induced hepatorenal toxicity in male rats. *Saudi J Biol Sci* 2017; 24: 1162-1171.
- 402 Alberich M, Menez C, Sutra JF, Lespine A. Ivermectin exposure leads to up-regulation of
403 detoxification genes in vitro and in vivo in mice. *Eur J Pharmacol* 2014; 740: 428-35.
- 404 Alvarez AI, Real R, Perez M, Mendoza G, Prieto JG, Merino G. Modulation of the activity of
405 ABC transporters (P-glycoprotein, MRP2, BCRP) by flavonoids and drug response. *J*
406 *Pharm Sci* 2010; 99: 598-617.
- 407 Ambali S, Akanbi D, Igbokwe N, Shittu M, Kawu M, Ayo J. Evaluation of subchronic
408 chlorpyrifos poisoning on hematological and serum biochemical changes in mice and
409 protective effect of vitamin C. *J Toxicol Sci* 2007; 32: 111-20.
- 410 Anilakumar KR, Pal A, Khanum F, Bawa AS. Nutritional, medicinal and industrial uses of
411 sesame (*Sesamum indicum* L.) seeds-an overview. *Agriculturae Conspectus Scientificus*
412 2010; 75: 159-168.
- 413 Ballent M, Lifschitz A, Virkel G, Sallovitz J, Lanusse C. Modulation of the P-glycoprotein-
414 mediated intestinal secretion of ivermectin: in vitro and in vivo assessments. *Drug Metab*
415 *Dispos* 2006; 34: 457-63.
- 416 BayIr H, Kagan VE. Bench-to-bedside review: Mitochondrial injury, oxidative stress and
417 apoptosis—there is nothing more practical than a good theory. *Critical care* 2008; 12: 206.

- 418 Bentires-Alj M, Barbu V, Fillet M, Chariot A, Relic B, Jacobs N, et al. NF-kappaB transcription
419 factor induces drug resistance through MDR1 expression in cancer cells. *Oncogene* 2003;
420 22: 90-7.
- 421 Beutler E, Duron O, Kelly BM. Improved method for the determination of blood glutathione. *J*
422 *Lab Clin Med* 1963; 61: 882-8.
- 423 Blaser H, Dostert C, Mak TW, Brenner D. TNF and ROS crosstalk in inflammation. *Trends in*
424 *cell biology* 2016; 26: 249-261.
- 425 Calero CI, Calvo DJ. Redox modulation of homomeric rho1 GABA receptors. *J Neurochem*
426 2008; 105: 2367-74.
- 427 Calero CI, Vickers E, Moraga Cid G, Aguayo LG, von Gersdorff H, Calvo DJ. Allosteric
428 modulation of retinal GABA receptors by ascorbic acid. *J Neurosci* 2011; 31: 9672-82.
- 429 Campbell WC. Ivermectin and abamectin: Springer Science & Business Media, 2012.
- 430 Carcamo JM, Pedraza A, Borquez-Ojeda O, Zhang B, Sanchez R, Golde DW. Vitamin C is a
431 kinase inhibitor: dehydroascorbic acid inhibits IkappaBalpha kinase beta. *Mol Cell Biol*
432 2004; 24: 6645-52.
- 433 Caro AA, Cederbaum AI. Oxidative stress, toxicology, and pharmacology of CYP2E1. *Annu*
434 *Rev Pharmacol Toxicol* 2004; 44: 27-42.
- 435 Chen Q, Cederbaum AI. Cytotoxicity and apoptosis produced by cytochrome P450 2E1 in Hep
436 G2 cells. *Mol Pharmacol* 1998; 53: 638-48.
- 437 Croop JM, Raymond M, Haber D, Devault A, Arceci RJ, Gros P, et al. The three mouse
438 multidrug resistance (mdr) genes are expressed in a tissue-specific manner in normal
439 mouse tissues. *Mol Cell Biol* 1989; 9: 1346-50.
- 440 Cuenda A, Rousseau S. p38 MAP-kinases pathway regulation, function and role in human
441 diseases. *Biochim Biophys Acta* 2007; 1773: 1358-75.
- 442 Danielson PB. The cytochrome P450 superfamily: biochemistry, evolution and drug metabolism
443 in humans. *Curr Drug Metab* 2002; 3: 561-97.
- 444 Deng L, Lin-Lee YC, Claret FX, Kuo MT. 2-acetylaminofluorene up-regulates rat mdr1b
445 expression through generating reactive oxygen species that activate NF-kappa B
446 pathway. *J Biol Chem* 2001; 276: 413-20.
- 447 Di Lisa F, Kaludercic N, Paolocci N. beta(2)-Adrenoceptors, NADPH oxidase, ROS and p38
448 MAPK: another 'radical' road to heart failure? *Br J Pharmacol* 2011; 162: 1009-11.
- 449 Gandhi S, Abramov AY. Mechanism of oxidative stress in neurodegeneration. *Oxid Med Cell*
450 *Longev* 2012; 2012: 428010.
- 451 Gonzalez Canga A, Sahagun Prieto AM, Jose Diez Liebana M, Martinez NF, Vega MS, Vieitez
452 JJ. The pharmacokinetics and metabolism of ivermectin in domestic animal species.
453 *Veterinary Journal* 2009; 179: 25-37.

- 454 Granger M, Eck P. Dietary vitamin C in human health. *Advances in food and nutrition research.*
455 83. Elsevier, 2018, pp. 281-310.
- 456 Hirsch-Ernst KI, Ziemann C, Foth H, Kozián D, Schmitz-Salue C, Kahl GF. Induction of *mdr1b*
457 mRNA and P-glycoprotein expression by tumor necrosis factor alpha in primary rat
458 hepatocyte cultures. *J Cell Physiol* 1998; 176: 506-15.
- 459 Hong H, Lu Y, Ji ZN, Liu GQ. Up-regulation of P-glycoprotein expression by glutathione
460 depletion-induced oxidative stress in rat brain microvessel endothelial cells. *J Neurochem*
461 2006; 98: 1465-73.
- 462 Hou RC, Wu CC, Yang CH, Jeng KC. Protective effects of sesamin and sesamol on murine
463 BV-2 microglia cell line under hypoxia. *Neurosci Lett* 2004; 367: 10-3.
- 464 Hsu DZ, Hsu CH, Huang BM, Liu MY. Abamectin effects on aspartate aminotransferase and
465 nitric oxide in rats. *Toxicology* 2001; 165: 189-93.
- 466 Hussien HM, Abdou HM, Yousef MI. Cypermethrin induced damage in genomic DNA and
467 histopathological changes in brain and haematotoxicity in rats: the protective effect of
468 sesame oil. *Brain Res Bull* 2013; 92: 76-83.
- 469 Jedlitschky G, Leier I, Buchholz U, Hummel-Eisenbeiss J, Burchell B, Keppler D. ATP-
470 dependent transport of bilirubin glucuronides by the multidrug resistance protein MRP1
471 and its hepatocyte canalicular isoform MRP2. *Biochem J* 1997; 327 (Pt 1): 305-10.
- 472 Jones B. *Pesticides and Insecticides: Development and Use* Syrawood Publishing House, 2018.
- 473 Jones PM, George AM. The ABC transporter structure and mechanism: perspectives on recent
474 research. *Cell Mol Life Sci* 2004; 61: 682-99.
- 475 Jung T-D, Choi S-I, Choi S-H, Cho B-Y, Sim W-S, Lee S, et al. Changes in the Anti-Allergic
476 Activities of Sesame by Bioconversion. *Nutrients* 2018; 10: 210.
- 477 Jurczuk M, Brzoska MM, Moniuszko-Jakoniuk J. Hepatic and renal concentrations of vitamins E
478 and C in lead- and ethanol-exposed rats. An assessment of their involvement in the
479 mechanisms of peroxidative damage. *Food Chem Toxicol* 2007; 45: 1478-86.
- 480 Kalender S, Uzun FG, Durak D, Demir F, Kalender Y. Malathion-induced hepatotoxicity in rats:
481 the effects of vitamins C and E. *Food Chem Toxicol* 2010; 48: 633-8.
- 482 Khaldoun-Oularbi H, Richeval C, Djenas N, Lhermitte M, Humbert L, Baz A. Effect of sub-
483 acute exposure to abamectin “insecticide” on liver rats (*Rattus norvegicus*). *Annales de*
484 *Toxicologie Analytique*. 25. EDP Sciences, 2013, pp. 63-70.
- 485 Lankas GR, Wise LD, Cartwright ME, Pippert T, Umbenhauer DR. Placental P-glycoprotein
486 deficiency enhances susceptibility to chemically induced birth defects in mice. *Reprod*
487 *Toxicol* 1998; 12: 457-63.

- 488 Li M, You TZ, Zhu WJ, Qu JP, Liu C, Zhao B, et al. Antioxidant response and histopathological
489 changes in brain tissue of pigeon exposed to avermectin. *Ecotoxicology* 2013; 22: 1241-
490 5.4
- 491 Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative
492 PCR and the 2(-Delta Delta C(T)) Method. *Methods* 2001; 25: 402-8.
- 493 Lu D, Ma Y, Zhang W, Bao D, Dong W, Lian H, et al. Knockdown of cytochrome P450 2E1
494 inhibits oxidative stress and apoptosis in the cTnTR141W dilated cardiomyopathy
495 transgenic mice. *Hypertension* 2012; 60: 81-89.
- 496 Ma JQ, Ding J, Zhang L, Liu CM. Hepatoprotective properties of sesamin against CCl4 induced
497 oxidative stress-mediated apoptosis in mice via JNK pathway. *Food Chem Toxicol* 2014;
498 64: 41-8.
- 499 Macdonald N, Gledhill A. Potential impact of ABCB1 (p-glycoprotein) polymorphisms on
500 avermectin toxicity in humans. *Arch Toxicol* 2007; 81: 553-63.
- 501 Maioli MA, de Medeiros HC, Guelfi M, Trinca V, Pereira FT, Mingatto FE. The role of
502 mitochondria and biotransformation in abamectin-induced cytotoxicity in isolated rat
503 hepatocytes. *Toxicol In Vitro* 2013; 27: 570-9.
- 504 McCavera S, Walsh TK, Wolstenholme AJ. Nematode ligand-gated chloride channels: an
505 appraisal of their involvement in macrocyclic lactone resistance and prospects for
506 developing molecular markers. *Parasitology* 2007; 134: 1111-21.
- 507 Ménez C, Mselli-Lakhal L, Foucaud-Vignault M, Balaguer P, Alvinerie M, Lespine A.
508 Ivermectin induces P-glycoprotein expression and function through mRNA stabilization
509 in murine hepatocyte cell line. *Biochemical pharmacology* 2012; 83: 269-278.
- 510 Mihara M, Uchiyama M. Determination of malonaldehyde precursor in tissues by thiobarbituric
511 acid test. *Anal Biochem* 1978; 86: 271-8.
- 512 Nasr HM, El-Demerdash FM, El-Nagar WA. Neuro and renal toxicity induced by chlorpyrifos
513 and abamectin in rats. *Environmental Science and Pollution Research* 2016; 23: 1852-
514 1859.
- 515 Novelli A, Vieira BH, Cordeiro D, Cappelini LT, Vieira EM, Espindola EL. Lethal effects of
516 abamectin on the aquatic organisms *Daphnia similis*, *Chironomus xanthus* and *Danio*
517 *rerio*. *Chemosphere* 2012; 86: 36-40.
- 518 Özkan F, Gündüz SG, Berköz M, Hunt AÖ, Yalın S. The protective role of ascorbic acid
519 (vitamin C) against chlorpyrifos-induced oxidative stress in *Oreochromis niloticus*. *Fish*
520 *physiology and biochemistry* 2012; 38: 635-643.
- 521 Perez D, White E. TNF-alpha signals apoptosis through a bid-dependent conformational change
522 in Bax that is inhibited by E1B 19K. *Mol Cell* 2000; 6: 53-63.

- 523 Pulido MM, Molina AJ, Merino G, Mendoza G, Prieto JG, Alvarez AI. Interaction of
524 enrofloxacin with breast cancer resistance protein (BCRP/ABCG2): influence of
525 flavonoids and role in milk secretion in sheep. *J Vet Pharmacol Ther* 2006; 29: 279-87.
- 526 Rangkadilok N, Pholphana N, Mahidol C, Wongyai W, Saengsooksree K, Nookabkaew S, et al.
527 Variation of sesamin, sesamolin and tocopherols in sesame (*Sesamum indicum* L.) seeds
528 and oil products in Thailand. *Food Chemistry* 2010; 122: 724-730.
- 529 Reitman S, Frankel S. A colorimetric method for the determination of serum glutamic oxalacetic
530 and glutamic pyruvic transaminases. *American journal of clinical pathology* 1957; 28:
531 56-63.
- 532 Roulet A, Puel O, Gesta S, Lepage JF, Drag M, Soll M, et al. MDR1-deficient genotype in Collie
533 dogs hypersensitive to the P-glycoprotein substrate ivermectin. *Eur J Pharmacol* 2003;
534 460: 85-91.
- 535 Saha RN, Jana M, Pahan K. MAPK p38 regulates transcriptional activity of NF-kappaB in
536 primary human astrocytes via acetylation of p65. *J Immunol* 2007; 179: 7101.9-
- 537 Saleem MT, Chetty MC, Kavimani S. Antioxidants and tumor necrosis factor alpha-inhibiting
538 activity of sesame oil against doxorubicin-induced cardiotoxicity. *Ther Adv Cardiovasc*
539 *Dis* 2014; 8: 4-11.
- 540 Sani I, Sule FA, Warra AA, Bello F, Fakai IM, Abdulhamid A. Phytochemicals and mineral
541 elements composition of white *Sesamum indicum* L. seed oil. *International Journal of*
542 *Traditional and Natural Medicines* 2013; 2: 118-130.
- 543 Santos IM, Tome Ada R, Saldanha GB, Ferreira PM, Militao GC, Freitas RM. Oxidative stress
544 in the hippocampus during experimental seizures can be ameliorated with the antioxidant
545 ascorbic acid. *Oxid Med Cell Longev* 2009; 2: 214-21.
- 546 Sauvant C, Nowak M, Wirth C, Schneider B, Riemann A, Gekle M, et al. Acidosis induces
547 multi-drug resistance in rat prostate cancer cells (AT1) in vitro and in vivo by increasing
548 the activity of the p-glycoprotein via activation of p38. *Int J Cancer* 2008; 123: 2532-42.
- 549 Segales J, Perdiguero E, Munoz-Canoves P. Regulation of Muscle Stem Cell Functions: A Focus
550 on the p38 MAPK Signaling Pathway. *Front Cell Dev Biol* 2016; 4: 91.
- 551 Seo M-Y, Lee S-M. Protective effect of low dose of ascorbic acid on hepatobiliary function in
552 hepatic ischemia/reperfusion in rats. *Journal of hepatology* 2002; 36: 72-77.
- 553 Serbecic N, Beutelspacher SC. Anti-oxidative vitamins prevent lipid-peroxidation and apoptosis
554 in corneal endothelial cells. *Cell Tissue Res* 2005; 320: 465-75.
- 555 Shuang D, Zhou J, Fang H, Nie Z, Chen S, Peng H. Sesamin protects the femoral head from
556 osteonecrosis by inhibiting ROS-induced osteoblast apoptosis in rat model. *Frontiers in*
557 *physiology* 2018; 9: 1787.
- 558 Sun Y, Diao X, Zhang Q, Shen J. Bioaccumulation and elimination of avermectin B1a in the
559 earthworms (*Eisenia fetida*). *Chemosphere* 2005; 60: 699-704.

- 560 Sun YJ, Long DX, Li W, Hou WY, Wu YJ, Shen JZ. Effects of avermectins on neurite
561 outgrowth in differentiating mouse neuroblastoma N2a cells. *Toxicol Lett* 2010; 192:
562 206-11.
- 563 Tietz N, Burtis C, Duncan P, Ervin K, Petittlerc C, Rinker A, et al. A reference method for
564 measurement of alkaline phosphatase activity in human serum. *Clinical chemistry* 1983;
565 29: 751-761.
- 566 Tietz NW. *Clinical Guide to Laboratory Tests (ELISA)*. 3rd Edition, W.B.: Saunders, Co.,
567 Philadelphia, 1995.
- 568 Tolman KG, Rej R. Liver function. in: Burtis, C.A., Ashwood, E.R., editors. *Tietz Textbook of*
569 *clinical Chemistry*. Third ed. Philadelphia; W.B.: Saunders company, 1999.
- 570 Wang C, Liang J, Zhang C, Bi Y, Shi X, Shi Q. Effect of ascorbic Acid and thiamine
571 supplementation at different concentrations on lead toxicity in liver. *Ann Occup Hyg*
572 2007; 51: 563-9.
- 573 Yamamoto M, Maeda H, Hirose N, Radhakrishnan G, Katare R, Hayashi Y, et al. Bilirubin
574 oxidation provoked by nitric oxide radicals predicts the progression of acute cardiac
575 allograft rejection. *American journal of transplantation* 2007; 7: 1897-1906.
- 576 Yang CC. Acute human toxicity of macrocyclic lactones. *Curr Pharm Biotechnol* 2012; 13: 999-
577 1003.
- 578 Zhang R, Yu Y, Hu S, Zhang J, Yang H, Han B, et al. Sesamin ameliorates hepatic steatosis and
579 inflammation in rats on a high-fat diet via LXR α and PPAR α . *Nutr Res* 2016a;
580 36: 1022-1030.
- 581 Zhang R, Yu Y, Hu S, Zhang J, Yang H, Han B, et al. Sesamin ameliorates hepatic steatosis and
582 inflammation in rats on a high-fat diet via LXR α and PPAR α . *Nutrition Research* 2016b;
583 36: 10.1030-22
- 584 Zhu WJ, Li M, Liu C, Qu JP, Min YH, Xu SW, et al. Avermectin induced liver injury in pigeon:
585 mechanisms of apoptosis and oxidative stress. *Ecotoxicol Environ Saf* 2013; 98: 74-81.
- 586
- 587

588 Figure Legends

589 **Figure 1:** Synthesis of the experimental design along with assessed molecular mechanisms. AA:
590 Ascorbic acid, ABM: Abamectin, CAT: Catalase, CYP-2E1: Cytochrome P450 2E1, GSH:
591 Reduced glutathione, MDA: Malondialdehyde, ROS: reactive oxygen species, SO: Sesame oil,
592 TNF- α : Tumor necrosis factor- α

593 **Figure 2:** The hepatic concentration of tumor necrosis factor- α (pg/mg) in different experimental
594 groups. AA: Ascorbic acid, SO: Sesame oil. Columns represent means \pm SEM (n=6). Columns
595 with different superscript letters are significantly different at $p \leq 0.05$.

596 **Figure 3:** Expression of cytochrome P450-2E1 (CYP-2E1), caspase-3, and p38 MAPK proteins
597 in the liver of different experimental groups (in relation to β -actin). AA: Ascorbic acid, SO:
598 Sesame oil. Columns represent means \pm SEM (n=6). Columns with different superscript letters
599 are significantly different at $p \leq 0.05$.

600 **Figure 4:** Western blotting analysis of p38 MAPK hepatic expression (in relation to β -actin) in
601 different experimental groups.

602 **Figure 5:** Relative expression of brain P-glycoprotein (*Abcb1a*) in different groups and GABA-
603 A receptor in different groups. AA: Ascorbic acid, SO: Sesame oil. Columns represent means \pm
604 SEM (n=6). Columns with different superscript letters are significantly different at $p \leq 0.05$.

605 **Figure 6:** Photomicrographs of the hepatic sections stained with H&E. (A) Control group
606 showed normal structure, (B) ABM group showed lymphocytic infiltration (*) and hydropic
607 degeneration (curved arrow). The incite showed ballooning of hepatocytes (arrow). The treated
608 groups with SO (C) and AA (D) showed congested bold vessels (thick arrow). (E) The
609 combination group was nearly normal. Scale bar 50 μ m. Incite sale bar 20 μ m.

610 **Figure 7:** Representative photomicrographs of cerebral cortices of experimental groups stained
611 with H&E. (A) Control group showing normal histological structure. (B) ABM group showing
612 dark stained nuclei (arrow), neuropil vacuolation (arrowhead) and hemorrhage (*). Rats treated
613 with SO (C) and AA (D) showing marked reduction of hemorrhage and (E) Combination
614 showing lightly stained vesicular nuclei (arrow). Scale bar 50 um.

615 **Figure 8:** Representative photomicrographs of hippocampus (CA3) of experimental groups
616 stained with H&E. (A) Control group showing outer pleomorphic layer (o), middle pyramidal
617 layer (p) and inner molecular layer (M). (B) Abamectin group showed pyknosis, dark stained
618 pyramidal neurons (arrow) and degenerative changes in the middle pyramidal layer. (C) treated
619 with SO only few pyknotic pyramidal cells (arrow). (D) Rats treated with AA and (E) SO + AA
620 combination showing nearly normal pyramidal cells with euchromatic nuclei (arrow). Scale bar
621 50 um.


622

623


624

625

626




TNF- α concentration (pg/ml)




Relative expression to β -actin


CYP2E1 expression




Caspase-3 expression




p38 MAPK expression



p38 MAPK



β -actin





Control






ABM






ABM+Vit.C

ABM+SO

ABM+AA+SO














Table (1): The primer sequence of the studied rat genes

	Primer sequence
GABA	Forward primer: 5'- TCGGGACCAACCCAACGTGC -3 Reverse primer: 5'- CGTGC TGGCCTGATTGACGCT -3
Abcb1a	Forward primer: 5-ACCAGCGGTCAGTGTGCT-3 Reverse primer: 5-CGGTTGTTTCCTACATTTGC-3
Cytochrome P450 2E1	Forward: 5- TTTGGATCCAATGGGTGATGTTGAG -3 Reverse: 5- TTTGAATTCCTCATTAGTAGCTTTTTTGGAG-3
Caspase-3	Forward primer: 5'- TTC ATT ATT CAG GCC TGC CGA GG -3 Reverse primer: 5'- TTC TGA CAG GCC ATG TCA TCC TCA -3
B-actin	Forward primer: 5'--GGTCGGTGTGAACGGATTTGG -3 Reverse primer: 5'- ATGTAGGCCATGAGGTCCACC-3

Table 2: The chemical composition of sesame oil sample by gas chromatography-mass spectrometry

No.	RT	Name of the compound	Molecular Formula	Molecular weight	Peak Area %
1	3.36	Nonanol	C ₉ H ₂₀ O	144	0.40
2	11.50	1-Tetradecanol	C ₁₄ H ₃₀ O	214	3.85
3	13.30	Undecanal	C ₁₁ H ₂₂ O	170	0.49
4	14.49	Dodecane	C ₁₂ H ₂₆	170	0.59
5	15.93	1-Hexadecanol, 2-methyl-	C ₁₇ H ₃₆ O	256	9.14
6	16.25	Nonadecane	C ₁₉ H ₄₀	268	2.64
7	16.49	Tetradecane, 2,6,10-trimethyl-	C ₁₇ H ₃₆	240	38.41
8	18.19	12-Methyl-E,E-2,13-octadecadien-1	C ₁₉ H ₃₆ O	280	1.81
9	18.68	Octadecane, 6-methyl-	C ₁₉ H ₄₀	268	5.72
10	20.57	Z-8-Methyl-9-tetradecenoic acid	C ₁₅ H ₂₈ O ₂	240	0.96
11	21.36	1-Dodecanol, 3,7,11-trimethyl-	C ₁₅ H ₃₂ O	228	3.14
12	22.24	9-Octadecenoic Acid (Z)-	C ₁₈ H ₃₄ O ₂	282	3.70
13	22.44	Hexadecadienoic Acid, methyl ester	C ₁₇ H ₃₀ O ₂	266	1.29
14	22.84	Docosane	C ₂₂ H ₄₆	310	3.01
15	23.11	E,E,Z-1,3,12-Nonadecatriene-5,14-diol	C ₁₉ H ₃₄ O ₂	294	0.72
16	23.55	2-Dodecen-1-yl(-)succinic anhydride	C ₁₆ H ₂₆ O ₃	266	1.77
17	25.91	9-Hexadecenoic acid	C ₁₆ H ₃₀ O ₂	254	0.51
18	28.16	7-Methyl-Z-tetradecen-1-ol acetate	C ₁₇ H ₃₂ O ₂	268	0.67
19	28.64	Octadecane	C ₁₈ H ₃₈	254	20.52

Table (3): Changes in serum SGPT, SGOT, ALP and direct bilirubin concentrations in different groups

	SGPT (U/L)	SGOT (U/L)	ALP (U/L)	Direct bilirubin (mg/dl)
Control	62.27±2.82 ^a	58.03±4.49 ^a	68.58±3.92 ^a	0.99±0.05 ^a
ABM	79.46±1.88 ^b	86.87±4.55 ^b	125.6±6.03 ^b	2.09±0.10 ^b
ABM+AA	67.16±0.76 ^a	76.30±2.13 ^c	91.08±3.74 ^c	1.64±0.04 ^c
ABM+SO	68.23±1.77 ^a	70.72±3.41 ^{ac}	84.67±3.13 ^{ac}	1.43±0.03 ^c
ABM+AA+SO	68.83±0.49 ^a	67.10±2.62 ^{ac}	80.21±1.94 ^{ac}	1.40±0.06 ^c

Values are expressed as mean ± SEM (n = 6 per group). Values with different letters in a column are significantly different at level $p < 0.0001$. AA: Ascorbic acid, ABM: Abamectin, ALP: Alkaline phosphatase, SGPT: Serum glutamic pyruvic transaminase, SGOT: Serum glutamic oxaloacetic transaminase, SO: Sesame oil

Table (4): Changes in MDA, GSH concentrations and catalase activity in the brain and liver of rats in different groups

	MDA (n.mol/g tissue)		GSH (mg/g tissue)		Catalase (U/g tissue)	
	Brain	Liver	Brain	Liver	Brain	Liver
Control	10.84±0.24 ^a	15.25±0.91 ^a	16±0.12 ^{ac}	16.24±0.12 ^a	0.634±0.01 ^a	0.236±0.01 ^a
ABM	13.20±0.37 ^b	18.63±0.49 ^b	14.12±0.23 ^b	12.22±0.79 ^b	0.461±0.04 ^b	0.323±0.01 ^b
ABM+AA	11.86±0.57 ^{ab}	15.19±0.32 ^a	15.61±0.23 ^a	15.79±0.06 ^a	0.624±0.01 ^a	0.303±0.01 ^{bc}
ABM+SO	11.51±0.52 ^{ab}	15.08±0.51 ^a	15.79±0.14 ^a	15.88±0.23 ^a	0.629±0.01 ^a	0.249±0.02 ^a
ABM+AA+SO	10.99±0.43 ^a	15.24±0.49 ^a	16.68±0.07 ^c	16.64±0.29 ^a	0.630±0.01 ^a	0.222±0.01 ^a

Values are expressed as mean ± SEM (n = 6 per group). Values with different letters in a column are significantly different at level $p < 0.05$. AA: Ascorbic acid, ABM: Abamectin, GSH: Reduced Glutathione, MDA: Malondialdehyde, SO: Sesame oil

