



# Genetic determinism of abdominal fat tissue and relationships with body weight and egg quality in Rhode Island Red

Nicolas Bedere, David Picard Druet, Frédéric Herault, Florian Herry, Sophie Allais, Amandine Varenne, Thierry Burlot, Pascale Le Roy

## ► To cite this version:

Nicolas Bedere, David Picard Druet, Frédéric Herault, Florian Herry, Sophie Allais, et al.. Genetic determinism of abdominal fat tissue and relationships with body weight and egg quality in Rhode Island Red. XIth European symposium on poultry genetics (ESPG), Oct 2019, Prague, Czech Republic. Guarant International spol. s r.o., 2019, Proceedings of the XIth European symposium on Poultry Genetics. hal-02352490

HAL Id: hal-02352490

<https://hal.science/hal-02352490>

Submitted on 6 Nov 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

# GENETIC ARCHITECTURE OF ABDOMINAL FAT TISSUE AND RELATIONSHIPS WITH BODY WEIGHT AND EGG QUALITY IN RHODE ISLAND RED

BEDERE N.<sup>(1)</sup>, PICARD DRUET D.<sup>(1)</sup>, HERAULT F.<sup>(1)</sup>, HERRY F.<sup>(1,2)</sup>, ALLAIS S.<sup>(1)</sup>, VARENNE A.<sup>(2)</sup>, BURLOTT.<sup>(2)</sup>, LE ROY P.<sup>(1)</sup>



**Extending the laying period → trade-off between egg production and safeguarding body reserves in late laying period?**

**In aged roosters (80wk), a dichotomic phenotype has been observed: lean OR fat**

What is the genetic architecture of the presence of abdominal fat in aged rooster



How is it related to other selection criteria

research Project UTOPIGE (FR)

~9,500 Rhode Island Red layers: ...of which 987♂ were also:

- egg quality
- body weight at 70 wk
- phenotyped for abdominal fat tissue at 90 wk
- genotyped on the 600K Affymetrix® Axiom® HD genotyping array


Variance components: REMLf90

GWAS: MULLER

## Variance components

	Abdominal fat ♂	Body weight ♂	Body weight ♀	Egg weight	Egg Shell Colour	Egg Shell Strength	Albumen Height
Abdominal fat ♂	0.20	0.31	NA	NA	NA	NA	NA
Body weight ♂	0.21	0.43	NA	NA	NA	NA	NA
Body weight ♀	0.10	0.67	0.66	0.20	0.20	-0.05	0.01
Egg weight	0.00	0.38	0.40	0.45	0.04	0.04	0.13
Egg Shell Colour	0.02	0.11	0.39	0.16	0.40	0.09	0.11
Egg Shell Strength	-0.03	-0.08	-0.17	0.00	-0.02	0.20	0.07
Albumen Height	-0.05	-0.05	0.00	0.12	0.08	0.11	0.22

## GWAS



The presence/absence of abdominal fat tissue in aged rooster:

- showed a **moderate heritability** estimate
- **Unsignificant genetic correlations** with egg production and little to moderate ones with BW
- BW is easier to measure than AFT but **would not be a good predictor of AFT**
- Further studies undergoing for high-throughput phenotyping in females

