



The Thioredoxin-Regulated α -Amylase 3 of *Arabidopsis thaliana* Is a Target of S-Glutathionylation

Libero Gurrieri, Luca Distefano, Claudia Pirone, Daniel Horrer, David Seung, Mirko Zaffagnini, Nicolas Rouhier, Paolo Trost, Diana Santelia, Francesca Sparla

► To cite this version:

Libero Gurrieri, Luca Distefano, Claudia Pirone, Daniel Horrer, David Seung, et al.. The Thioredoxin-Regulated α -Amylase 3 of *Arabidopsis thaliana* Is a Target of S-Glutathionylation. *Frontiers in Plant Science*, 2019, 10, 10.3389/fpls.2019.00993 . hal-02352493

HAL Id: hal-02352493

<https://hal.science/hal-02352493>

Submitted on 6 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License



The Thioredoxin-Regulated α -Amylase 3 of *Arabidopsis thaliana* Is a Target of S-Glutathionylation

Libero Gurrieri¹, Luca Distefano^{2†}, Claudia Pirone¹, Daniel Horrer², David Seung³, Mirko Zaffagnini¹, Nicolas Rouhier⁴, Paolo Trost¹, Diana Santelia^{2*†} and Francesca Sparla¹

¹ Department of Pharmacy and Biotechnology FaBIT, University of Bologna, Bologna, Italy, ² Department of Plant and Microbial Biology, University of Zürich, Zurich, Switzerland, ³ John Innes Centre, Norwich, United Kingdom, ⁴ Université de Lorraine, Inra, IAM, Nancy, France

OPEN ACCESS

Edited by:

Paula Casati,
National Scientific and Technical
Research Council (CONICET),
Argentina

Reviewed by:

Marcos Ariel Tronconi,
CONICET Center for Photosynthetic
and Biochemical Studies (CEFOBI),
Argentina
Arne Holmgren,
Karolinska Institute (KI), Sweden

*Correspondence:

Diana Santelia
dsantelia@ethz.ch

† Present address:

Luca Distefano and Diana Santelia,
ETH Zürich, Institute of Integrative
Biology, Zurich, Switzerland

Specialty section:

This article was submitted to
Plant Abiotic Stress,
a section of the journal
Frontiers in Plant Science

Received: 27 March 2019

Accepted: 15 July 2019

Published: 31 July 2019

Citation:

Gurrieri L, Distefano L, Pirone C,
Horner D, Seung D, Zaffagnini M,
Rouhier N, Trost P, Santelia D and
Sparla F (2019) The
Thioredoxin-Regulated α -Amylase 3
of *Arabidopsis thaliana* Is a Target
of S-Glutathionylation.
Front. Plant Sci. 10:993.
doi: 10.3389/fpls.2019.00993

Reactive oxygen species (ROS) are produced in cells as normal cellular metabolic by-products. ROS concentration is normally low, but it increases under stress conditions. To stand ROS exposure, organisms evolved series of responsive mechanisms. One such mechanism is protein S-glutathionylation. S-glutathionylation is a post-translational modification typically occurring in response to oxidative stress, in which a glutathione reacts with cysteinyl residues, protecting them from overoxidation. α -Amylases are glucan hydrolases that cleave α -1,4-glucosidic bonds in starch. The *Arabidopsis* genome contains three genes encoding α -amylases. The sole chloroplastic member, AtAMY3, is involved in osmotic stress response and stomatal opening and is redox-regulated by thioredoxins. Here we show that AtAMY3 activity was sensitive to ROS, such as H₂O₂. Treatments with H₂O₂ inhibited enzyme activity and part of the inhibition was irreversible. However, in the presence of glutathione this irreversible inhibition was prevented through S-glutathionylation. The activity of oxidized AtAMY3 was completely restored by simultaneous reduction by both glutaredoxin (specific for the removal of glutathione-mixed disulfide) and thioredoxin (specific for the reduction of protein disulfide), supporting a possible liaison between both redox modifications. By comparing free cysteine residues between reduced and GSSG-treated AtAMY3 and performing oxidation experiments of Cys-to-Ser variants of AtAMY3 using biotin-conjugated GSSG, we could demonstrate that at least three distinct cysteinyl residues can be oxidized/glutathionylated, among those the two previously identified catalytic cysteines, Cys499 and Cys587. Measuring the pK_a values of the catalytic cysteines by alkylation at different pHs and enzyme activity measurement (pK_{a1} = 5.70 ± 0.28; pK_{a2} = 7.83 ± 0.12) showed the tendency of one of the two catalytic cysteines to deprotonation, even at physiological pHs, supporting its propensity to undergo redox post-translational modifications. Taking into account previous and present findings, a functional model for redox regulation of AtAMY3 is proposed.

Keywords: α -amylase 3, post-translational redox modifications, S-glutathionylation, disulfide, cysteine pK_a, glutaredoxin, thioredoxin

INTRODUCTION

Starch is the most abundant non-structural carbohydrate in plants. Mostly depending on its lifetime, starch can be distinguished in storage and transitory starch. Transitory starch is typically produced in photosynthetic organs where it is accumulated during the day and degraded in the following dark period. Transitory starch metabolism has been deeply studied over the last decades and its biosynthetic and degradation pathways have been detailed (Zeeman et al., 2010; Stitt and Zeeman, 2012; Skryhan et al., 2018). Several enzymes emerged as fundamental for leaf starch turnover and when their corresponding genes are knocked out, plants show starch-less or starch-excess phenotypes (Yu et al., 2001; Kötting et al., 2005, 2009; Fulton et al., 2008; Streb et al., 2009; Crumpton-Taylor et al., 2013). Another set of enzymes, the lack of which does not lead to starch-related phenotypes in leaves under standard growth conditions, was found localized in chloroplasts of guard cells (Outlaw, 2003; Horrer et al., 2016) as well as involved in leaf starch degradation pathway occurring in response to abiotic stresses (Zeeman et al., 2004; Valerio et al., 2011; Monroe et al., 2014; Zanella et al., 2016; Thalmann et al., 2016; Horrer et al., 2016).

Among the three α -amylases encoded by the Arabidopsis genome, AtAMY3 is not required for normal starch breakdown in mesophyll cells and its knock-out mutant does not present a starch-excess phenotype (Yu et al., 2005). In contrast to the other α -amylases, AtAMY3 localizes in chloroplast's stroma of both mesophyll and guard cells (Yu et al., 2005; Delatte et al., 2006) and is regulated by thioredoxins (TRXs) through the formation of a disulfide bridge between Cys499 and Cys587 with a midpoint redox potential at pH 7.9 of -329 mV (Seung et al., 2013).

TRXs are small oxidoreductases found in various subcellular compartments. In chloroplasts, TRXs are reduced by electrons provided by photosystem I (PSI) and relayed by Ferredoxin (FDX), FDX:TRX reductase (FTR) and TRX. Through this redox chain, TRX-target enzymes achieve their modulation by light (Michelet et al., 2013; Balsera et al., 2014). The activation of AtAMY3 by TRX-f1 is therefore consistent with its role in starch degradation at day in response to stress (Seung et al., 2013) as well as its presence in guard cells, where starch metabolism follows an opposite rhythm in comparison to mesophyll cells, mobilizing starch in the light to produce malate and/or sucrose, which contribute to increase guard cell osmolarity, turgor and stomatal opening (Vavasseur and Raghavendra, 2005; Lawson et al., 2014; Horrer et al., 2016; Santelia and Lawson, 2016; Santelia and Lunn, 2017).

In addition to a TRX-mediated redox regulation, thiol groups of cysteine residues can be modified by other redox modifications such as S-glutathionylation and S-nitrosylation (for a comprehensive review see Zaffagnini et al., 2019). These redox post-translational modifications (PTMs) are favored by the propensity of a cysteine residue to exist in a deprotonated form (thiolate anion; $-S^-$) even at physiological pH values. Consequently, reactive cysteines are often characterized by pK_a values lower than the pK_a value

of non-reactive cysteine thiol (i.e., below ≈ 7.0 and close to ≈ 8.5 , respectively).

In chloroplasts, environmental changes such as drought, exposure to intense light or high temperature, are rapidly perceived and the production of reactive oxygen species (ROS) increases rapidly (Cruz de Carvalho, 2008; Chaves et al., 2009; Farooq et al., 2009; Sharma et al., 2012; Suzuki et al., 2012; Noctor et al., 2018). Compared to singlet oxygen and superoxide radical, hydrogen peroxide (H_2O_2) has received particular attention, being a long-lived molecule (Mittler and Zilinskas, 1991) that can possibly diffuse over quite long distances acting as signal molecule (Bienert et al., 2007; Sharma et al., 2012). However, at high concentrations, H_2O_2 can act as damaging molecule, rapidly oxidizing thiolate anions and leading to the formation of sulfenylated cysteines ($-SOH$). Under this first oxidative state, sulfenylated cysteines can be further oxidized to sulfinic ($-SO_2H$) and sulfonic ($-SO_3H$) acids (Roos and Messens, 2011; Roos et al., 2013; Trost et al., 2017), both considered irreversible oxidation states leading to protein degradation (Poole, 2015; Zaffagnini et al., 2019). Sulfenic acid can react with a thiol group of a protein or of low molecular weight molecules as reduced glutathione (GSH) leading to protein disulfide or protein S-glutathionylation, respectively (Ito et al., 2003; Dixon et al., 2005; Michelet et al., 2005; Zaffagnini et al., 2012a).

AtAMY3 was found sulfenylated upon H_2O_2 treatment *in vivo* experiments (De Smet et al., 2018), and recombinantly expressed AMY1 of barley was found glutathionylated at Cys95 (Søgaard et al., 1993; Juge et al., 1996). Cys95 of barley AMY1 corresponds to Cys587 in AtAMY3, a residue required for the optimal catalytic rate of the Arabidopsis enzyme, and involved in the formation of the thioredoxin regulated disulfide bridge together with Cys499 (Seung et al., 2013).

Reduced glutathione is one of the redox molecules that, together with ascorbate and a set of ROS-scavenging enzymes, contributes to cellular redox homeostasis (Shigeoka et al., 2002; Mittler et al., 2004; Foyer and Shigeoka, 2011; Sharma et al., 2012; Noctor and Foyer, 2016; Foyer and Noctor, 2016). GSH plays an important role against ROS-induced oxidative damages, as it is able to scavenge H_2O_2 leading to the formation of oxidized glutathione (GSSG). The balance between GSH and GSSG is central for cellular redox state, mirroring the redox state of a cell that can change in response to internal and external stimuli (Foyer and Noctor, 2016). In response to stress, the raise of H_2O_2 concentration causes accumulation of GSSG due to GSH-scavenging effect. The GSSG concentrations may become high enough to allow protein glutathionylation, helping to prevent cysteine overoxidation by other oxidants (Michelet et al., 2005; Noguera-Mazon et al., 2006; Tarrago et al., 2009).

In analogy with TRXs, which reduce regulatory disulfide bridges, glutaredoxins (GRXs) remove glutathione from modified enzymes (Rouhier et al., 2008; Zaffagnini et al., 2012b). GRXs are oxidized by the protein substrates, the activity of which is usually restored upon reduction. The glutathionylated GRX is reduced using another GSH molecule forming GSSG.

In this study, we have expressed the recombinant AtAMY3 and conducted a biochemical characterization of its sensitivity to oxidizing and reducing treatments in the presence of glutathione and H₂O₂. We propose a fine-tuned regulatory mechanism for AtAMY3 activity involving both S-glutathionylation and disulfide bridge formation.

MATERIALS AND METHODS

In silico Analysis of α -Amylases

The amino acid sequences of *Arabidopsis thaliana* α -amylase 3 (UniProtKB: Q94A41; AtAMY3); *A. thaliana* α -amylase 1 (UniProtKB: Q8VZ56; AtAMY1); *A. thaliana* α -amylase 2 (UniProtKB: Q8LFG1; AtAMY2) and *Hordeum vulgare* α -amylase type A isozyme (UniProtKB: P00693; HvAMYA) were aligned with Esript¹ (Robert and Gouet, 2014). Sequence identities among considered α -amylases calculated by Clustal Omega (Sievers et al., 2011) were the following: 46% for AtAMY3 vs. AtAMY1; 49% for AtAMY3 vs. AtAMY2; 47% for AtAMY3 vs. HvAMYA.

Expression and Purification of Recombinant Proteins

Wild-type and mutated forms of AtAMY3 were expressed and purified as described in Seung et al. (2013). Protein purity was assessed by 12.5% SDS-PAGE. Pure recombinant proteins were quantified by their absorbance at 280 nm (Nanodrop; Thermo Fisher Scientific) using molar extinction coefficients and molecular masses calculated from the amino acid sequences of the recombinant enzymes with the online ProtParam tool (Gasteiger et al., 2005). The recombinant *A. thaliana* GRX C5 and poplar GRX S12 were expressed in *Escherichia coli* and purified as described in Couturier et al. (2009, 2011).

Enzyme Activity Assays

The enzymatic activity of AtAMY3 was measured using the α -Amylase Assay Kit (Ceralpha Method) from Megazyme (Megazyme, Ireland) according to the manufacturer's instructions. Briefly, a mixture composed by an equal volume of AtAMY3 in 100 mM Tricine-NaOH, pH 7.9 and the artificial substrate blocked *p*-nitrophenyl maltoheptaoside (B-PNPG7) plus α -glucosidase was incubated at 40°C. After incubation, the reaction was blocked by adding 20-volume of Stopping Reagent (1% Tris, pH 11.0). The absorbance of the samples was evaluated at 400 nm using a spectrophotometer and subtracting the absorbance of blank sample treated under the same condition but without AtAMY3. An extinction coefficient at 400 nm for *p*-nitrophenol of 18.1 mM⁻¹ cm⁻¹ was used.

Oxidative Treatments

All oxidative treatments were performed on freshly reduced AtAMY3. Following 90 min of incubation at 37°C in the presence of 40 mM dithiothreitol (DTT), protein sample was desalted

using NAP-5 column (GE Healthcare) in 100 mM Tricine-NaOH, pH 7.9 and quantified by absorbance at 280 nm using an extinction coefficient of 185420 M⁻¹ cm⁻¹ and molecular weight of 96976.53 Da (Nanodrop; Thermo Fisher Scientific). Sample was brought to the desired concentration ($\approx$ 20 μ M) either by dilution or by concentration through Amicon-Ultra device (Millipore; cut-off 10 kDa).

Inactivation kinetics by H₂O₂ of AtAMY3 were performed by incubating the reduced recombinant enzyme at 25°C in 100 mM Tricine-NaOH, pH 7.9, in the absence (control) or presence (treated sample) of 0.1 mM, 0.25 mM, 0.5 mM and 1 mM H₂O₂. At different time points, the enzyme activity was measured on 1:5 diluted samples using the α -Amylase Assay Kit (Ceralpha Method; Megazyme, Ireland). The data are expressed as percentage of residual activity relatively to the initial activities ($t = 0$) measured on the control samples.

Treatments with H₂O₂, H₂O₂ plus GSH and GSSG were performed by incubating reduced AtAMY3 at 25°C in the presence of 0.1 mM, 0.5 mM or 1 mM H₂O₂ for 1 h; 1 mM H₂O₂ and 5 mM GSH for 1 h; 50 μ M, 0.25 mM, 1 mM or 5 mM GSSG for 30 min. After the treatments, the enzyme activity was measured on 1:5 diluted samples using the α -Amylase Assay Kit (Ceralpha Method; Megazyme, Ireland). The reversibility of the oxidative treatments was tested by incubating the samples with 80 mM DTT for an additional 30 min at 25°C. After the treatments, the enzyme activity was measured on 1:5 diluted samples using the α -Amylase Assay Kit (Ceralpha Method; Megazyme, Ireland).

The data are reported as percentage of activity relative to control samples incubated without oxidants under the same condition.

Reactivation of Glutathionylated AtAMY3 by GSH, GRXs, and TRX

Freshly reduced AtAMY3, obtained as previously described, was incubated with 5 mM GSSG for 90 min at 25°C. After GSSG incubation, sample was desalted through NAP-5 column (GE Healthcare) in 100 mM Tricine-NaOH, pH 7.9 and brought to the desired concentration (20 μ M) through Amicon-Ultra device (Millipore; cut-off 10 kDa). Sample concentration was determined by absorbance at 280 nm (Nanodrop; Thermo Fisher Scientific).

Reactivation treatments of glutathionylated AtAMY3 were performed by incubating samples for 30 min at 25°C in the presence of 50 μ M, 0.2 mM, 1 mM, 2 mM, 5 mM, and 7 mM GSH. Samples were then diluted 5-fold and the activity assayed as described above.

Reactivation assays of GSSG-inhibited AtAMY3 performed in the presence of poplar GRX S12 and *A. thaliana* GRX C5, and commercially available *E. coli* TRX (Sigma-Aldrich; protein id AAA24693), were conducted through 5, 15, and 30 min of incubation of 20 μ M glutathionylated AtAMY3 in the presence of 2 mM GSH with or without 5 μ M GRXs, or in the presence of 0.2 mM DTT with or without 10 μ M TRX. Upon incubation, enzyme activity was measured on 5-fold diluted samples as described above.

¹<http://esript.ibcp.fr>

All data are expressed as percentages of activity relative to the fully reduced sample obtained by incubating glutathionylated AtAMY3 in presence of 80 mM DTT for 30 min at 25°C.

Quantification of Thiol Groups

Protein thiol quantification was performed on both reduced and GSSG-treated AtAMY3. Freshly reduced AtAMY3, obtained by 90 min incubation at 37°C in the presence of 40 mM DTT, was desalted in 100 mM Tricine-NaOH, pH 7.9 with NAP-5 column (GE Healthcare) and then incubated with 5 mM GSSG for additional 90 min at 25°C. After GSSG incubation sample was desalted in 100 mM Tricine-NaOH, pH 7.9 with NAP-5 column (GE Healthcare).

Reduced and GSSG-treated enzyme was incubated in the presence of 0.5 mM 5,5'-dithiobis (2-nitrobenzoic acid) (DTNB) for 1 h at room temperature. The number of free and solvent accessible thiol groups under the two tested conditions, was calculated from the molar ratio between the absorbance at 412 nm [molar extinction coefficient of 14150 M⁻¹ cm⁻¹ for 2-nitro-5-thiobenzoate(thiolate) dianion] and the absorbance at 280 nm (molar extinction coefficient of 185420 M⁻¹ cm⁻¹ for AtAMY3) (Conway et al., 2004).

Biotinylated GSSG Assay

Biotin-conjugated GSSG (BioGSSG) was freshly prepared by incubating for 1 h at room temperature 50 µl of 32 mM GSSG with 50 µl of 48 mM EZ-Link Sulfo-NHS-Biotin (Thermo Fisher Scientific) in 50 mM potassium phosphate buffer pH 7.2. The reaction was stopped by the addition of 35 µl of 0.6 M NH₄HCO₃.

Wild-type AtAMY3 and all single cysteine to serine mutants were desalted in 100 mM Tricine-NaOH, pH 7.9 by NAP-5 columns (GE Healthcare). Protein concentrations were determined by absorbance at 280 nm (Nanodrop; Thermo Fisher Scientific) and 2 µM enzymes were incubated at room temperature in the presence of 2 mM BioGSSG at 25°C. After 1 h incubation, each sample was divided into two aliquots. One aliquot was treated with 80 mM DTT for 30 min to assess the reversibility of the reaction, whereas the second aliquot was transferred into a tube containing SDS-loading buffer 1X without reducing agent and in presence of 100 mM iodoacetamide (IAM) and 20 mM *N*-ethylmaleimide (NEM) to block cysteine reactivity. Negative control samples were incubated with the alkylating agents IAM (100 mM) and NEM (20 mM) for 30 min in the dark before incubation in the presence of 2 mM BioGSSG.

In the BioGSSG assays performed on pre-reduced or pre-oxidized enzymes, the recombinant proteins were incubated with 40 mM DTT or 40 mM *trans*-4,5-Dihydroxy-1,2-dithiane (DTTox) for 4 h at room temperature. Reduced and oxidized enzymes were desalted on a NAP-5 column (GE Healthcare) pre-equilibrated in 100 mM Tricine-NaOH, pH 7.9 before proceeding with BioGSSG analysis as described above.

Following incubations, protein samples were further divided into two aliquots and loaded on two denaturing non-reducing 12.5% SDS-PAGE. One gel was analyzed by Coomassie staining while the second gel was transferred to a nitrocellulose

membrane and analyzed by Western blot using monoclonal anti-biotin antibodies (Sigma-Aldrich) diluted 1:3800. Peroxidase-conjugated secondary antibodies (Sigma-Aldrich) were diluted 1:2000 and were used for the detection by ECL Western Blotting Detection Reagent (GE Healthcare), following the manufacturer's instruction.

Determination of the pK_a of the Catalytic Cysteines

The pK_a values of the active-site cysteines were determined by measuring the pH dependence of the rate of reaction of AtAMY3 as reported in Kallis and Holmgren (1980). Briefly, for each pH value (100 mM Sodium citrate for pH 4.0–5.5; 100 mM MES for pH 6.0–6.5; 100 mM Tris-HCl for pH 7.0–9.0; 100 mM Glycine for pH 10.0–12.0), 8 µM recombinant enzyme was incubated with or without a 10-fold excess of the alkylating reagent IAM with respect to the total thiol content of the protein sample. After 20 min incubation, samples were diluted 5-fold and α-amylase activity was measured with the artificial substrate BPNPG7 (Megazyme). AtAMY3 did not undergo irreversible changes in the analyzed pH range, except for the pH values of 4.0, 11.0, and 12.0, which inhibited AtAMY3 control samples by about 56% on average. The residual activity, expressed as a percentage of inhibition between IAM-treated and untreated samples, was plotted against pH.

Data sets were fitted by non-linear regression using the following equations with one-pK_a or two-pK_a dependence (Karala et al., 2010):

(a) one-pK_a dependence:

$$\% \text{ remaining activity} = 100 - \left(a \times \frac{10^{(\text{pH} - \text{pK}_a)}}{1 + 10^{(\text{pH} - \text{pK}_a)}} \right)$$

(b) two-pK_a dependence:


$$\% \text{ remaining activity} = 100 -$$

$$\left(a \times \frac{10^{(\text{pH} - \text{pK}_{a1})}}{1 + 10^{(\text{pH} - \text{pK}_{a1})}} \right) + b \times \frac{10^{(\text{pH} - \text{pK}_{a2})}}{1 + 10^{(\text{pH} - \text{pK}_{a2})}} \right)$$

RESULTS

GSH Protects AtAMY3 From Irreversible Inactivation Mediated by H₂O₂

AtAMY3 has a well-documented function in the chloroplasts of mesophyll cells in response to osmotic stress, and in guard cells during stomatal opening (Yu et al., 2005; Horrer et al., 2016; Thalmann et al., 2016). Under osmotic stress conditions, ROS production can exceed the scavenging ability of plants (Chaves et al., 2009), becoming harmful molecules that lead to oxidative damage. For this reason, the susceptibility of AtAMY3 to redox modulation was analyzed by measuring the enzymatic activity in the presence of increasing concentrations of H₂O₂ (Figure 1A).



AtAMY3 was inhibited in a dose and time-dependent manner by H₂O₂ treatments, although only with high concentration, i.e., 1 mM H₂O₂, as well as long incubation time, i.e., 1 h, the enzyme activity was completely blocked (Figure 1A). 1 h incubation in the presence of 1 mM H₂O₂ caused oxidation of AtAMY3, which could not be reversed by DTT (Figure 1B).

H₂O₂ is particularly reactive on cysteine thiols, leading to a sequential oxidation of sulfhydryl groups going through sulfenic (–SOH), sulfinic (–SO₂H) and sulfonic (–SO₃H) acid states, the last two of which being irreversible (Supplementary Figure S1A) (Santelia et al., 2015). When cysteines are involved in the catalysis, oxidation affects enzyme activity and irreversible oxidation (i.e., inactivation) can be prevented by thiol modification such as S-glutathionylation. *In vivo* S-glutathionylation mainly occurs by reaction between reduced

GSH and the sulfenic state of cysteine (Supplementary Figure S1A). For this reason AtAMY3 was simultaneously treated with 1 mM H₂O₂ plus 5 mM GSH and the enzyme activity was measured before and after DTT reduction. The simultaneous presence of H₂O₂ and GSH resulted in a completely reversible inhibition of AtAMY3 activity (Figure 1B), suggesting that AtAMY3 could be reversibly modified by glutathione and this may be a mechanism for regulating its activity.

AtAMY3 Activity Is Modulated by GSH:GSSG Ratio and Glutaredoxins Speed Up the Reactivation Process

In vitro protein glutathionylation can be achieved by the addition of GSSG. GSSG spontaneously reacts faster with the deprotonated rather than protonated sulfhydryl group of cysteine residues (Supplementary Figure S1B). To test the effect of GSSG on the enzyme activity, pre-reduced AtAMY3 was incubated for 30 min at 25°C at increasing GSSG concentrations. As expected, GSSG treatment inhibited the enzyme activity in a dose-dependent manner (Figure 2A, black bars) and such inhibitions were completely abolished by a second incubation in the presence DTT (Figure 2A, white bars).

In plants, GSH is the main form of glutathione and only small amounts of GSSG are found inside the cell (Foyer and Noctor, 2011). Being the major endogenous antioxidant, GSH is continuously regenerated from GSSG by glutathione reductase (GR) consuming reducing power (Foyer and Noctor, 2011). However, under stress conditions, concentration of both forms of glutathione varies, often leading to a decrease in the GSH:GSSG ratio (Noctor et al., 2012; Hasanuzzaman et al., 2017). Hence, the effect of GSH on GSSG pre-treated AtAMY3 was tested. As shown in Figure 2B, increasing concentrations of GSH allowed an almost complete restoration of AtAMY3 activity after 30 min incubation, suggesting that the enzyme can be effectively regulated by the simple change of the GSH concentration (Supplementary Figure S1B).

GRXs are redoxins able to control the thiol-based post-translational modifications of target enzymes, specifically reducing glutathione-mixed disulfides (Meyer et al., 2008). The ability of GRXs to reverse GSSG-mediated inhibition of AtAMY3 was assayed by measuring α-amylase activity upon 5 min incubation with GRX S12 and GRX C5 plus 2 mM GSH (Figure 2C). Adding these GRXs led to a recovery of about 80% of the maximal activity, a value close to that obtained after 30 min incubation in the presence of 5–7 mM GSH alone (Figure 2B). Taken together, the data indicate that GRXs play a role in the regulation of AtAMY3 activity speeding up the reactivation of the enzyme (Supplementary Figure S1B).

Both TRX and GRX Are Required to Completely Restore AtAMY3 Activity

Considering that GSSG-inhibited AtAMY3 did not fully restore its activity even in the presence of GRXs (Figure 2C), we speculated that the mixed disulfide bridge occurring between GSSG and the thiol group of one cysteine could be rapidly




FIGURE 2 | Reactivation of GSSG-inhibited AtAMY3 by GSH and effect of GRXs on the reactivation rate. **(A)** Inactivation of AtAMY3 after 1 h incubation with 50 μ M, 0.25 mM, 1 mM, and 5 mM GSSG (black bars). The reversibility of the inhibition was tested by incubating samples for additional 30 min in presence of 80 mM DTT (white bars). Activities are expressed as percentages of reduced sample obtained by incubating glutathionylated AtAMY3 in presence of 80 mM DTT for 30 min at 25°C. Data are reported as mean $\pm$ SD ($n = 3$). Significant reduction in enzyme activity compared to the pre-reduced enzyme was determined based on P -values obtained from Student's t test. $*P < 0.01$. **(B)** Recovery of catalytic activity of AtAMY3 pre-treated with GSSG (black bar) by increasing concentrations of GSH. GSSG-inhibited samples (black bar) were incubated for 30 min at 25°C at different concentrations of GSH (white bars). Activities are expressed as percentages of fully active AtAMY3, obtained by incubating GSSG-inhibited enzyme for 30 min in presence of 80 mM DTT. Data are reported as mean $\pm$ SD ($n = 3$). Significant increase in enzyme activity compared to the GSSG-treated enzyme was determined based on P -values obtained from Student's t test. $*P < 0.01$. **(C)** GSSG-inhibited AtAMY3 (black bar) was incubated with 2 mM GSH in absence (gray bar) or presence (white bars) of GRX S12 or C5. Activities are (Continued)

FIGURE 2 | Continued expressed as percentages of fully active AtAMY3, obtained by incubating GSSG-inhibited enzyme with 80 mM DTT for 30 min. Data are reported as mean $\pm$ SD ($n = 3$). Significant increase in enzyme activity compared to the GSSG-treated enzyme was determined based on P -values obtained from Student's t test. $*P < 0.01$.




FIGURE 3 | TRX-dependent reactivation of GSSG-inhibited AtAMY3 and effect of both GRX S12 and TRX. **(A)** GSSG-treated AtAMY3 (black bar) was incubated in presence of 0.2 mM DTT without (gray bar) or with (white bar) 10 μ M TRX. Activities are expressed as percentage of fully reduced samples obtained after 30 min incubation of GSSG-inhibited AtAMY3 with 80 mM DTT. Data are reported as mean $\pm$ SD ($n = 3$). Significant increase in enzyme activity compared to the GSSG-inhibited enzyme was determined based on P -values obtained from Student's t test. $*P < 0.01$. **(B)** GSSG-inhibited AtAMY3 (black bar) was incubated for 30 min with 2 mM GSH and 5 μ M GRX S12 (gray bar). After 30 min incubation, 0.2 mM DTT was added to the mixture either in the absence (pale gray bar) or in the presence (white bar) of commercially available TRX. Incubation was extended for another 5 min at 25°C before measuring AtAMY3 activity. Activities are expressed as percentage of fully reduced samples obtained after 30 min incubation of GSSG-inhibited AtAMY3 with 80 mM DTT. Data are reported as mean $\pm$ SD ($n = 2$). Significant increase in enzyme activity compared to the GSSG-inhibited enzyme was determined based on P -values obtained from Student's t test. $*P < 0.01$.

attacked by the thiol group ($-SH$) of a second cysteine residue to form a stable disulfide ($-SS-$) and releasing GSH (**Supplementary Figure S1C**). To test this hypothesis GSSG-inhibited AtAMY3 was incubated with 0.2 mM DTT in the presence or absence of *E. coli* TRX. Together with GRXs, TRXs belong to the redoxins family and are well known to efficiently reduce protein

disulfides. As shown in **Figure 3A**, within 5 min incubation TRX partially restores AtAMY3 activity, corroborating the initial hypothesis of the formation of a disulfide bridge in response to GSSG treatment.

To further test whether GSSG treatment led to the formation of both thiol modifications, sequential measurements of the recovery of AtAMY3 activity upon a first incubation with GRX S12 followed by a second incubation with TRX were performed. To ascertain that GRX S12-dependent reduction of AtAMY3 was completed before the addition of TRX, an experiment was performed showing clearly that already after 5 min incubation the reactivation of AtAMY3 by GRX S12 was completed (**Supplementary Figure S2**). Similar to previous results (**Figure 2C**), only 80% of the maximum activity was recovered even after longer incubation (**Supplementary Figure S2**).

Once confirmed that the reactivation of AtAMY3 by GRX S12 was completed after 5 min incubation (**Supplementary Figure S2**), 0.2 mM DTT with or without 10 μ M TRX was added and the incubation extended for 5 min. As shown in **Figure 3B** the activity of AtAMY3 was completely restored only after incubation with both redoxins (i.e., GRX and TRX). Altogether, these results support the conclusion that GSSG could react with AtAMY3 inducing the formation of both glutathionylation and disulfide bond.

At Least Three Different AtAMY3 Cysteine Residues Are Modified by GSSG

As reported in Seung et al. (2013), AtAMY3 is regulated by dithiol/disulfide exchange occurring between the cysteine residues Cys499 and Cys587. To analyze the putative involvement of Cys499 and Cys587 in both PTMs (i.e., dithiol/disulfide exchange and glutathionylation), pre-reduced

and pre-oxidized AtAMY3 were incubated in the presence of BioGSSG (**Figure 4**). This BioGSSG assay detects the formation of a mixed disulfide between the accessible cysteines in the target enzyme and BioGSSG. After the formation of a mixed disulfide, antibiotin antibodies can easily detect biotin. If no reaction occurs between BioGSSG and recombinant enzyme, no signal is obtained.

For comparison, all single cysteine to serine AtAMY3 mutants were subjected to BioGSSG assay (**Supplementary Figure S3**). Both pre-reduced and pre-oxidized wild-type AtAMY3 showed a clear signal that disappeared following incubation with DTT and was prevented by alkylation with IAM and NEM (**Figure 4**). The same was observed for all single Cys to Ser AtAMY3 mutants in their oxidized form (**Supplementary Figure S3**). The positive immunodetection in all samples allowed us to conclude that more than one cysteine residue is subject to S-glutathionylation in AtAMY3 and that Cys499 and Cys587 are not the only cysteine residues targeted by S-glutathionylation. If that was the case, no signal would have been observed in the pre-oxidized AtAMY3 sample.

Albeit not all cysteine residues were found exposed and accessible to DTNB (i.e., out of the nine cysteines present in the primary sequence of AtAMY3 only 7.7 were detected by DTNB) (**Table 1**), just over three cysteine residues were found modified by GSSG, as calculated by the difference of free cysteine in fully

TABLE 1 | 5,5'-dithiobis(2-nitrobenzoic acid) (DTNB) analysis of free thiols in AtAMY3 under reducing or GSSG-treated conditions.

	No. of free thiols (-SH)
Reduced AtAMY3	7.7 $\pm$ 1.1
GSSG-treated AtAMY3	4.2 $\pm$ 1.2

Results are mean $\pm$ SD ($n = 2$).




FIGURE 4 | Effect of BioGSSG incubation on reduced and oxidized AtAMY3. Protein samples (2 μ M) were mixed with 2 mM BioGSSG for 1 h prior to separation on a non-reducing 12.5% SDS-PAGE and transfer to a nitrocellulose membrane. The reversibility of the reaction was assessed by a 30 min incubation with 80 mM DTT. As negative control, 2 μ M protein sample was alkylated with 100 mM IAM and 20 mM NEM before BioGSSG treatment.




FIGURE 5 | pK_a determination of the catalytic cysteines of AtAMY3. AtAMY3 was incubated in different buffers ranging from pH 4 to 12 in presence or absence of IAM before measuring the activity. The residual activity at each pH value was calculated as percentage of inhibition between IAM-treated and untreated samples, and expressed as a function of pH. The obtained curve was fitted by non-linear regression with two- pK_a dependence. Results are mean $\pm$ SD ($n = 3$; SD < 10% are omitted for clarity).

reduced and GSSG-treated AtAMY3 (Table 1), in agreement with BioGSSG assay.

Determination of pK_a Value

Unreactive cysteines are characterized by pK_a values around 8.5, whereas reactive cysteines are typically surrounded by unusual microenvironment that destabilizes the unreactive protonated form of cysteine residues through electrostatic interactions, significantly decreasing their pK_a value (Roos et al., 2013). IAM is a strong alkylating agent able to react specifically with thiolate anion ($-S^-$). When cysteine residues are involved in the catalytic process, the pK_a of catalytic relevant cysteines can thus be determined by measuring the pH dependency of IAM inactivation.

AtAMY3 was treated with IAM at different pH values and the activities were compared with those measured at the same pH but in absence of the alkylating reagent. Experimental data were fit both to a one- pK_a -dependent (Supplementary Figure S4) and to a two- pK_a -dependent event (Figure 5). However, the best fitting returned two pK_a values, the first pK_a was 5.70 ± 0.28 , while the second pK_a was 7.83 ± 0.12 .

The finding of a two inflections curve was also in agreement with the low or even absence of activity measured in the single C499S and C587S AtAMY3 variants (Seung et al., 2013). Moreover, since Cys499 and Cys587 were also responsible for

the TRX-dependent regulation, the present finding supports the involvement of at least one and the same cysteine in both redox modifications.

DISCUSSION

Most organisms constantly produce ROS because of metabolic processes as photosynthesis and respiration. However, unfavorable environmental conditions such as varying temperatures, light intensities and water availability, rapidly modify ROS production. In chloroplasts, several kind of stresses affect photosynthetic efficiency augmenting singlet oxygen (1O_2), superoxide (O_2^-), hydroxyl radical (HO) and hydrogen peroxide (H_2O_2) production. Among them, the long-lived H_2O_2 molecule has received particular attention acting in both signal transduction and regulation of enzyme activity (Bienert et al., 2007; Sharma et al., 2012). Depending on H_2O_2 concentration and on the reactivity of cysteine residues, protein thiols can undergo several reversible or irreversible modifications (Supplementary Figure S1). The formation of disulfide and S-glutathionylation are known mechanisms altering enzyme activity (Zaffagnini et al., 2019).

Differently from other α -amylases, AtAMY3 localizes in the stroma of chloroplast of both guard and mesophyll cells and its




FIGURE 6 | Schematic illustration of the proposed redox-regulatory model of AtAMY3 activity. As suggested by DTNB assay, 3.5 cys residues were modified by GSSG; three of them are highlighted in yellow while the fourth is shown in light yellow. In the model, AtAMY3 (white) is fully active when reduced ($-SH$), but it can be irreversibly inactivated when oxidized to sulfenic (pale gray) or sulfonic (gray) acid forms by H_2O_2 . However, if oxidation occurs in the presence of H_2O_2 and GSH or is mediated by GSSG, AtAMY3 is efficiently protected from overoxidation via S-glutathionylation. S-glutathionylation can lead to a partially (orange) or fully (red) modified enzyme. In case of partial S-glutathionylation (orange), an intramolecular disulfide (blue) can be formed releasing GSH. Thioredoxin (TRX) can activate AtAMY3 by reducing this disulfide (white). Similarly, fully glutathionylated AtAMY3 (red) can be efficiently activated by GSH through a partially (orange) or a fully (red) modified enzyme. In case of partial S-glutathionylation (orange), an intramolecular disulfide (blue) can be formed releasing GSH. Thioredoxin (TRX) can activate AtAMY3 by reducing this disulfide (white). Similarly, fully glutathionylated AtAMY3 (red) can be efficiently activated by GSH through a partially (orange) or a fully (red) modified enzyme.

activity is strictly regulated by TRXf1. Light-driven reduction of the regulatory disulfide bridge between Cys499 and Cys587 activates AtAMY3 (Seung et al., 2013), presumably leading to a diurnal degradation of transitory starch. Hitherto AtAMY3 is a unique α -amylase subject to TRX-regulation, a type of regulation in line with the physiological functions ascribed to AtAMY3, covering stress-induced starch degradation in mesophyll cells and the stimulation of stomatal opening in guard cells (Horner et al., 2016; Thalmann et al., 2016).

However, the stress response implies an increase in the concentration of H₂O₂. The aim of the present study was to elucidate using *in vitro* biochemical approaches whether and how the activity of AtAMY3 is redox-regulated and the catalytic cysteines protected against overoxidation and reactivated.

Already more than 20 years ago, the inhibitory effect caused by S-glutathionylation of the catalytic Cys95 in barley AMY1 was reported (Sogaard et al., 1993). As for the cytosolic isoforms of Arabidopsis, AMY1 from barley lacks an N-terminal ~50 kDa domain (Supplementary Figure S5). However, the C-terminal domain of AtAMY3 and barley AMY1 is highly conserved (46.3% identity and 74.6% similar, based on the comparison between the full-length AMY1 and the last 418 amino acids at the C-terminal of AtAMY3) suggesting that Cys587 (corresponding to Cys95 of barley AMY1) would be target of glutathionylation. Furthermore, a more recent evidence comes from an *in vivo* study that identified the sulfenylated (–SOH) form of AtAMY3 in Arabidopsis cells treated with 1 mM H₂O₂ (De Smet et al., 2018). This first oxidative state of protein thiols can be followed by further oxidation to sulfinic (–SO₂H) and sulfonic (–SO₃H) acids, considered irreversible oxidation (Supplementary Figure S1A). Alternatively, the sulfenylated form can form disulfides reacting with protein or low molecular weight thiols (Supplementary Figure S1A).

For this reason, the effect of H₂O₂, GSH plus H₂O₂ and GSSG were tested on AtAMY3 activity, demonstrating that AtAMY3 can be irreversibly oxidized by H₂O₂ and reversibly glutathionylated by GSH plus H₂O₂ or GSSG (Figures 1B, 2A, 6). Thus, the sulfenylated form of AtAMY3 is target of S-glutathionylation *in vitro*, and this PTM likely protects the enzyme from irreversible oxidation.

At least three cysteine residues were modified by a GSSG treatment (Table 1), two of which, judging from the results obtained by the pK_a analysis (Figure 5), are involved in the catalytic activity. Indeed, this type of analysis returns the pK_a value only of cysteines involved in the enzymatic activity, whereas it cannot detect cysteine residues that do not affect enzyme activity even if they are modified. In Seung et al. (2013), single Cys to Ser mutants of residues 499 and 587 were ~90% less active than wild-type AtAMY3, suggesting that two out of the three GSSG-modified cysteine residues could be Cys499 and Cys587.

The same pair of cysteine residues is also responsible for the TRX-dependent regulation of AtAMY3 (Seung et al., 2013). Taking this into consideration and the fact that both redoxins (i.e., GRX and TRX) were required for a fast

(Figures 2C, 3A) and complete (Figure 3B) reactivation of AtAMY3 activity, the proposed model suggests that during oxidative treatments leading to glutathionylation, partially modified enzyme formed a homodisulfide releasing GSH (Figure 6).

The slow and non-protein-assisted reactivation of AtAMY3 by GSH (Figure 2B) adds complexity to the system. Although varying across tissues and subcellular compartments, in the absence of stress the GSH:GSSG ratio is typically considered very high (Schwarzländer et al., 2016). This could suggest that on the way to recover from stress-induced ROS production, the activity of AtAMY3 would be fine-modulated by the high concentration of GSH. Considering the role of AtAMY3 in stress response (Thalmann et al., 2016), it seems reasonable to assume that the biochemical features of the enzyme equipped AtAMY3 to overcome the burst of oxidative stress. Equally true, however, is that a rapid reactivation of the enzymatic activity is required to the onset of plant response to stress.

In the tangled scenario of the redox regulation, AtAMY3 offers a good example of the cross-talk between redox-dependent protein modifications (i.e., dithiol/disulfide exchange and S-glutathionylation) in the fine-tuning of enzymatic activity. Future research will be required to validate the role of these PTMs *in vivo*.

DATA AVAILABILITY

The raw data supporting the conclusions of this manuscript will be made available by the authors, without undue reservation, to any qualified researcher.

AUTHOR CONTRIBUTIONS

CP, LG, LD, and NR purified the recombinant proteins. CP and LG performed the BioGSSG assays and biochemical characterization. FS and DS coordinated the experiments, supervised the whole research, and wrote the manuscript. All authors discussed the data and reviewed the manuscript.

FUNDING

This research was supported by the Swiss National Science Foundation (grant No. 31003A_166539 to DS). CP was supported by the Department of Pharmacy and Biotechnology – University of Bologna, Programma Marco Polo short term fellowship (Prot. No. 1883, September 15, 2014).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpls.2019.00993/full#supplementary-material>

REFERENCES

- Balsera, M., Uberegui, E., Schürmann, P., and Buchanan, B. B. (2014). Evolutionary development of redox regulation in chloroplasts. *Antioxid. Redox. Signal.* 21, 1327–1355. doi: 10.1089/ars.2013.5817
- Bienert, G. P., Möller, A. L. B., Kristiansen, K. A., Schulz, A., Möller, I. M., Schjoerring, J. K., et al. (2007). Specific aquaporins facilitate the diffusion of hydrogen peroxide across membranes. *J. Biol. Chem.* 282, 1183–1192. doi: 10.1074/jbc.m603761200
- Chaves, M. M., Flexas, J., and Pinheiro, C. (2009). Photosynthesis under drought and salt stress: regulation mechanisms from whole plant to cell. *Ann. Bot. London* 103, 551–560. doi: 10.1093/aob/mcn125
- Conway, M. E., Poole, L. B., and Hutson, S. M. (2004). Roles for cysteine residues in the regulatory CXXC motif of human mitochondrial branched chain aminotransferase enzyme. *Biochemistry* 43, 7356–7364. doi: 10.1021/bi0498050
- Couturier, J., Koh, C. S., Zaffagnini, M., Winger, A. M., Gualberto, J. M., Corbier, C., et al. (2009). Structure-function relationship of the chloroplastic glutaredoxin S12 with an atypical WCSYS active site. *J. Biol. Chem.* 284, 9299–9310. doi: 10.1074/jbc.M807998200
- Couturier, J., Ströher, E., Albetel, A. N., Roret, T., Muthuramalingam, M., Tarrago, L., et al. (2011). Arabidopsis chloroplastic glutaredoxin C5 as a model to explore molecular determinants for iron-sulfur cluster binding into glutaredoxins. *J. Biol. Chem.* 286, 27515–27527. doi: 10.1074/jbc.M111.228726
- Crumpton-Taylor, M., Pike, M., Lu, K. J., Hylton, C. M., Feil, R., Eicke, S., et al. (2013). Starch synthase 4 is essential for coordination of starch granule formation with chloroplast division during Arabidopsis leaf expansion. *New Phytol.* 200, 1064–1075. doi: 10.1111/nph.12455
- Cruz de Carvalho, M. H. (2008). Drought stress and reactive oxygen species: production, scavenging and signaling. *Plant Signal. Behav.* 3, 156–165. doi: 10.4161/psb.3.3.5536
- De Smet, B., Willems, P., Fernandez-Fernandez, A. D., Alseekh, S., Fernie, A. R., Messens, J., et al. (2018). In vivo detection of protein cysteine sulfonylation in plastids. *Plant J.* 97, 765–778. doi: 10.1111/tpj.14146
- Delatte, T., Umhang, M., Trevisan, M., Eicke, S., Thorncroft, D., Smith, S. M., et al. (2006). Evidence for distinct mechanisms of starch granule breakdown in plants. *J. Biol. Chem.* 281, 12050–12059. doi: 10.1074/jbc.m513661200
- Dixon, D. P., Skipsey, M., Grundy, N. M., and Edwards, R. (2005). Stress-induced protein S-glutathionylation in Arabidopsis. *Plant Physiol.* 138, 2233–2244. doi: 10.1104/pp.104.058917
- Farooq, M., Wahid, A., and Lee, D. J. (2009). Exogenously applied polyamines increase drought tolerance of rice by improving leaf water status, photosynthesis and membrane properties. *Acta Physiol. Plant.* 31, 937–945. doi: 10.1007/s11738-009-0307-2
- Foyer, C. H., and Noctor, G. (2011). Ascorbate and glutathione: the heart of the redox hub. *Plant Physiol.* 155, 2–18. doi: 10.1104/pp.110.167569
- Foyer, C. H., and Noctor, G. (2016). Stress-triggered redox signalling: what's in pROSpect? *Plant Cell Environ.* 39, 951–964. doi: 10.1111/pce.12621
- Foyer, C. H., and Shigeoka, S. (2011). Understanding oxidative stress and antioxidant functions to enhance photosynthesis. *Plant Physiol.* 155, 93–100. doi: 10.1104/pp.110.166181
- Fulton, D. C., Stettler, M., Mettler, T., Vaughan, C. K., Li, J., Francisco, P., et al. (2008). b-AMYLASE4, a noncatalytic protein required for starch breakdown, acts upstream of three active b-amylases in Arabidopsis chloroplasts. *Plant Cell* 20, 1040–1058. doi: 10.1105/tpc.107.056507
- Gasteiger, E., Hoogland, C., Gattiker, A., Duvaud, S., Wilkins, M. R., Appel, R. D., et al. (2005). "Protein identification and analysis tools on the ExPASy server," in *The Proteomics Protocols Handbook*, ed. J. M. Walker (New York, NY: Humana Press).
- Hasanuzzaman, M., Nahar, K., Anee, T. I., and Fujita, M. (2017). Glutathione in plants: biosynthesis and physiological role in environmental stress tolerance. *Physiol. Mol. Biol. Plants* 23, 249–268. doi: 10.1007/s12298-017-0422-2
- Horrer, D., Flüttsch, S., Pazmino, D., Matthews, J. S., Thalmann, M., Nigro, A., et al. (2016). Blue light induces a distinct starch degradation pathway in guard cells for stomatal opening. *Curr. Biol.* 26, 362–370. doi: 10.1016/j.cub.2015.12.036
- Ito, H., Iwabuchi, M., and Ogawa, K. (2003). The sugar-metabolic enzymes aldolase and triose-phosphate isomerase are targets of glutathionylation in Arabidopsis thaliana: detection using biotinylated glutathione. *Plant Cell Physiol.* 44, 655–660. doi: 10.1093/pcp/pcg098
- Juge, N., Andersen, J. S., Tull, D., Roepstorff, P., and Svensson, B. (1996). Overexpression, purification, and characterization of recombinant barley alpha-amylases 1 and 2 secreted by the methylotrophic yeast Pichia pastoris. *Protein Expr. Purif.* 8, 204–214. doi: 10.1006/prep.1996.0093
- Kallis, G. B., and Holmgren, A. (1980). Differential reactivity of the functional sulfhydryl groups of cysteine-32 and cysteine-35 present in the reduced form of thioredoxin from Escherichia coli. *J. Biol. Chem.* 255, 10261–10265.
- Karala, A. R., Lappi, A. K., and Ruddock, L. W. (2010). Modulation of an active-site cysteine pKa allows PDI to act as a catalyst of both disulfide bond formation and isomerization. *J. Mol. Biol.* 396, 883–892. doi: 10.1016/j.jmb.2009.12.014
- Kötting, O., Pusch, K., Tiessen, A., Geigenberger, P., Steup, M., and Ritte, G. (2005). Identification of a novel enzyme required for starch metabolism in Arabidopsis leaves: the phosphoglucan, water dikinase. *Plant Physiol.* 137, 242–252. doi: 10.1104/pp.104.055954
- Kötting, O., Santelia, D., Edner, C., Eicke, S., Marthaler, T., Gentry, M. S., et al. (2009). STARCHEXCESS4 is a laforin-like phosphoglucan phosphatase required for starch degradation in Arabidopsis thaliana. *Plant Cell* 21, 334–346. doi: 10.1105/tpc.108.064360
- Lawson, T., Simkin, A. J., Kelly, G., and Granot, D. (2014). Mesophyll photosynthesis and guard cell metabolism impacts on stomatal behaviour. *New Phytol.* 203, 1064–1081. doi: 10.1111/nph.12945
- Meyer, Y., Siala, W., Bashandy, T., Riondet, C., Vignols, F., and Reichheld, J. P. (2008). Glutaredoxins and thioredoxins in plants. *Biochim. Biophys. Acta* 1783, 589–600.
- Michelet, L., Zaffagnini, M., Marchand, C., Collin, V., Decottignies, P., Tsan, P., et al. (2005). Glutathionylation of chloroplast thioredoxin f is a redox signaling mechanism in plants. *Proc. Natl. Acad. Sci. U.S.A.* 102, 16478–16483. doi: 10.1073/pnas.0507498102
- Michelet, L., Zaffagnini, M., Morisse, S., Sparla, F., Pérez-Pérez, M. E., Francia, F., et al. (2013). Redox regulation of the calvin-benson cycle: something old, something new. *Front. Plant Sci.* 4:470. doi: 10.3389/fpls.2013.00470
- Mittler, R., Vanderauwera, S., Gollery, M., and Van Breusegem, F. (2004). Reactive oxygen gene network of plants. *Trends Plant Sci.* 9, 490–498. doi: 10.1016/j.tplants.2004.08.009
- Mittler, R., and Zilinskas, B. A. (1991). Purification and characterization of pea cytosolic ascorbate peroxidase. *Plant Physiol.* 97, 962–968. doi: 10.1104/pp.97.3.962
- Monroe, J. D., Storm, A. R., Badley, E. M., Lehman, M. D., Platt, S. M., Saunders, L. K., et al. (2014). β-AMYLASE1 and β-AMYLASE3 are plastidic starch hydrolases in arabidopsis that appear to be adapted for different thermal, pH, and stress conditions. *Plant Physiol.* 166, 1748–1763. doi: 10.1104/pp.114.246421
- Noctor, G., and Foyer, C. H. (2016). Intracellular redox compartmentation and ros-related communication in regulation and signaling. *Plant Physiol.* 171, 1581–1592. doi: 10.1104/pp.16.00346
- Noctor, G., Mhamdi, A., Chaouch, S., Han, Y., Neukermans, J., Marquez-Garcia, B., et al. (2012). Glutathione in plants: an integrated overview. *Plant Cell Environ.* 35, 454–484. doi: 10.1111/j.1365-3040.2011.02400.x
- Noctor, G., Reichheld, J. P., and Foyer, C. H. (2018). ROS-related redox regulation and signaling in plants. *Semin. Cell. Dev. Biol.* 80, 3–12. doi: 10.1016/j.semcdb.2017.07.013
- Noguera-Mazon, V., Lemoine, J., Walker, O., Rouhier, N., Salvador, A., Jacquot, J. P., et al. (2006). Glutathionylation induces the dissociation of 1-Cys D-peroxiredoxin non-covalent homodimer. *J. Biol. Chem.* 281, 31736–31743.
- Outlaw, W. H. (2003). Integration of cellular and physiological functions of guard cells. *Cr. Rev. Plant Sci.* 22, 503–529. doi: 10.1080/713608316
- Poole, L. B. (2015). The basics of thiols and cysteines in redox biology and chemistry. *Free Rad. Bio. Med.* 80, 148–157. doi: 10.1016/j.freeradbiomed.2014.11.013
- Robert, X., and Gouet, P. (2014). Deciphering key features in protein structures with the new ENDscript server. *Nucl. Acids Res.* 42, W320–W324. doi: 10.1093/nar/gku316
- Roos, G., Foloppe, N., and Messens, J. (2013). Understanding the pK(a) of redox cysteines: the key role of hydrogen bonding. *Antioxid. Redox. Signal.* 18, 94–127. doi: 10.1089/ars.2012.4521
- Roos, G., and Messens, J. (2011). Protein sulfenic acid formation: from cellular damage to redox regulation. *Free Radic. Biol. Med.* 51, 314–326. doi: 10.1016/j.freeradbiomed.2011.04.031

- Rouhier, N., Lemaire, S. D., and Jacquot, J. P. (2008). The role of glutathione in photosynthetic organisms: emerging functions for glutaredoxins and glutathionylation. *Annu. Rev. Plant Biol.* 59, 143–166. doi: 10.1146/annurev.arplant.59.032607.092811
- Santelia, D., and Lawson, T. (2016). Rethinking guard cell metabolism. *Plant Physiol.* 172, 1371–1392. doi: 10.1104/pp.16.00767
- Santelia, D., and Lunn, J. E. (2017). Transitory starch metabolism in guard cells: unique features for a unique function. *Plant Physiol.* 174, 539–549. doi: 10.1104/pp.17.00211
- Santelia, D., Trost, P., and Sparla, F. (2015). New insights into redox control of starch degradation. *Curr. Opin. Plant Biol.* 25, 1–9. doi: 10.1016/j.pbi.2015.04.003
- Schwarzländer, M., Dick, T. P., Meyer, A. J., and Morgan, B. (2016). Dissecting redox biology using fluorescent protein sensors. *Antioxid. Redox. Signal.* 24, 680–712. doi: 10.1089/ars.2015.6266
- Seung, D., Thalmann, M., Sparla, F., Hachem, M. A., Lee, S. K., Issakidis-Bourguet, E., et al. (2013). *Arabidopsis thaliana* AMY3 is a unique redox-regulated chloroplastic α -amylase. *J. Biol. Chem.* 288, 33620–33633. doi: 10.1074/jbc.M113.514794
- Sharma, P., Jha, A. B., Dubey, R. S., and Pessarakli, M. (2012). Reactive oxygen species, oxidative damage, and antioxidative defense mechanism in plants under stressful conditions. *J. Bot.* 2012:26. doi: 10.1155/2012/217037
- Shigeoka, S., Ishikawa, T., Tamoi, M., Miyagawa, Y., Takeda, T., Yabuta, Y., et al. (2002). Regulation and function of ascorbate peroxidase isoenzymes. *J. Exp. Bot.* 53, 1305–1319. doi: 10.1093/jexbot/53.372.1305
- Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., et al. (2011). Fast, scalable generation of high-quality protein multiple sequence alignments using clustal omega. *Mol. Syst. Biol.* 7, 539–545. doi: 10.1038/msb.2011.75
- Skryhan, K., Gurrieri, L., Sparla, F., Trost, P., and Blennow, A. (2018). Redox regulation of starch metabolism. *Front. Plant Sci.* 9:1344. doi: 10.3389/fpls.2018.01344
- Søgaard, M., Andersen, J. S., Roepstorff, P., and Svensson, B. (1993). Electrospray mass spectrometry characterization of post-translational modifications of barley α -amylase 1 produced in yeast. *Biotechnology* 11, 1162–1165. doi: 10.1038/nbt1093-1162
- Stitt, M., and Zeeman, S. C. (2012). Starch turnover: pathways, regulation and role in growth. *Curr. Opin. Plant Biol.* 15, 282–292. doi: 10.1016/j.pbi.2012.03.016
- Streb, S., Egli, B., Eicke, S., and Zeeman, S. C. (2009). The debate on the pathway of starch synthesis: a closer look at low-starch mutants lacking plastidial phosphoglucomutase supports the chloroplast-localized pathway. *Plant Physiol.* 151, 1769–1772. doi: 10.1104/pp.109.144931
- Suzuki, N., Koussevitzky, S., Mittler, R., and Miller, G. (2012). ROS and redox signalling in the response of plants to abiotic stress. *Plant Cell Environ.* 35, 259–270. doi: 10.1111/j.1365-3040.2011.02336.x
- Tarrago, L., Laugier, E., Zaffagnini, M., Marchand, C., Le, Maréchal P, Rouhier, N., et al. (2009). Regeneration mechanisms of *Arabidopsis thaliana* methionine sulfoxide reductases B by glutaredoxins and thioredoxins. *J. Biol. Chem.* 284, 18963–18971. doi: 10.1074/jbc.M109.015487
- Thalmann, M., Pazmino, D., Seung, D., Horrer, D., Nigro, A., Meier, T., et al. (2016). Regulation of leaf starch degradation by abscisic acid is important for osmotic stress tolerance in plants. *Plant Cell* 28, 1860–1878. doi: 10.1105/tpc.16.00143
- Trost, P., Fermani, S., Calvaresi, M., and Zaffagnini, M. (2017). Biochemical basis of sulphenomics: how protein sulphenic acids may be stabilized by the protein microenvironment. *Plant Cell Environ.* 40, 483–490. doi: 10.1111/pce.12791
- Valerio, C., Costa, A., Marri, L., Issakidis-Bourguet, E., Pupillo, P., Trost, P., et al. (2011). Thioredoxin-regulated α -amylase (BAM1) triggers diurnal starch degradation in guard cells, and in mesophyll cells under osmotic stress. *J. Exp. Bot.* 62, 545–555. doi: 10.1093/jxb/erq288
- Vavasseur, A., and Raghavendra, A. S. (2005). Guard cell metabolism and CO₂ sensing. *New Phytol.* 165, 665–682. doi: 10.1111/j.1469-8137.2004.01276.x
- Yu, T. S., Kofler, H., Häusler, R. E., Hille, D., Flügge, U. I., Zeeman, S. C., et al. (2001). The *Arabidopsis* *sex1* mutant is defective in the R1 protein, a general regulator of starch degradation in plants, and not in the chloroplast hexose transporter. *Plant Cell* 13, 1907–1918. doi: 10.1105/tpc.13.8.1907
- Yu, T. S., Zeeman, S. C., Thorncroft, D., Fulton, D. C., Dunstan, H., Lue, W. L., et al. (2005). α -Amylase is not required for breakdown of transitory starch in *Arabidopsis* leaves. *J. Biol. Chem.* 280, 9773–9779. doi: 10.1074/jbc.M413638200
- Zaffagnini, M., Bedhomme, M., Lemaire, S. D., and Trost, P. (2012a). The emerging roles of protein glutathionylation in chloroplasts. *Plant Sci.* 18, 86–96. doi: 10.1016/j.plantsci.2012.01.005
- Zaffagnini, M., Bedhomme, M., Marchand, C. H., Couturier, J. R., Gao, X. H., Rouhier, N., et al. (2012b). Glutaredoxin s12: unique properties for redox signaling. *Antioxid. Redox. Signal.* 16, 17–32. doi: 10.1089/ars.2011.3933
- Zaffagnini, M., Fermani, S., Marchand, C. H., Costa, A., Sparla, F., Rouhier, N., et al. (2019). Redox homeostasis in photosynthetic organisms: novel and established thiol-based molecular mechanisms. *Antioxid. Redox. Signal.* 31, 155–210. doi: 10.1089/ars.2018.7617
- Zanella, M., Borghi, G. L., Pirone, C., Thalmann, M., Pazmino, D., Costa, A., et al. (2016). α -Amylase 1 (BAM1) degrades transitory starch to sustain proline biosynthesis during drought stress. *J. Exp. Bot.* 67, 1819–1826. doi: 10.1093/jxb/erv572
- Zeeman, S. C., Kossmann, J., and Smith, A. M. (2010). Starch: its metabolism, evolution, and biotechnological modification in plants. *Annu. Rev. Plant Biol.* 61, 209–234. doi: 10.1146/annurev-arplant-042809-112301
- Zeeman, S. C., Thorncroft, D., Schupp, N., Chapple, A., Weck, M., Dunstan, H., et al. (2004). Plastidial α -glucan phosphorylase is not required for starch degradation in *Arabidopsis* leaves but has a role in the tolerance of abiotic stress. *Plant Physiol.* 135, 849–858. doi: 10.1104/pp.103.032631

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Gurrieri, Distefano, Pirone, Horrer, Seung, Zaffagnini, Rouhier, Trost, Santelia and Sparla. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.