



**HAL**  
open science

## Identification and expression of microRNAs in european eels *Anguilla anguilla* from two natural sites with different pollution levels

Anthony Bertucci, Fabien Pierron, Tao Ye, Patrice Gonzalez, Patrice Couture, Magalie Baudrimont

### ► To cite this version:

Anthony Bertucci, Fabien Pierron, Tao Ye, Patrice Gonzalez, Patrice Couture, et al.. Identification and expression of microRNAs in european eels *Anguilla anguilla* from two natural sites with different pollution levels. *Environmental Pollution*, 2019, 250, pp.274-283. 10.1016/j.envpol.2019.04.029 . hal-02354043

**HAL Id: hal-02354043**

**<https://hal.science/hal-02354043>**

Submitted on 22 Oct 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Title**

2 Identification and expression of microRNAs in European eels *Anguilla anguilla* from two natural sites  
3 with different pollution levels.

4

5 **Authors:**

6 Anthony Bertucci<sup>1</sup>, Fabien Pierron<sup>1</sup>, Tao Ye<sup>2</sup>, Patrice Gonzalez<sup>1</sup>, Patrice Couture<sup>3</sup> and Magalie  
7 Baudrimont<sup>1</sup>

8

9 **Affiliations and Addresses:**

10 <sup>1</sup> Univ. Bordeaux, UMR EPOC CNRS 5805, 33615 Pessac, France.

11 <sup>2</sup> IGBMC - CNRS UMR 7104 - Inserm U 964, 1 BP 10142, 67404 Illkirch Cedex, France.

12 <sup>3</sup>Institut National de La Recherche Scientifique (INRS), Centre Eau Terre Environnement, 490 rue De  
13 La Couronne, Québec, QC G1K 9A9, Canada.

14

15 **Corresponding author:**

16 Anthony Bertucci

17 Phone number: +33 (0)5 56 22 39 30

18 Fax number: +33 (0)5 40 70 85 04

19 Email: [anthony.bertucci@u-bordeaux.fr](mailto:anthony.bertucci@u-bordeaux.fr)

20 **Abstract**

21 MicroRNAs (miRNAs) are a class of small non-coding RNA that control multiple biological processes  
22 through negative post-transcriptional regulation of gene expression. Recently a role of miRNAs in the  
23 response of aquatic organisms to environmental toxicants emerged. Toxicant-induced changes in  
24 miRNA expression might then represent novel biomarkers to evaluate the health status of these  
25 organisms. In this study, we aimed to identify the miRNA repertoire in the liver of the European eel  
26 *Anguilla anguilla* and to compare their differential expression between a polluted site located in the  
27 Gironde Estuary and a pristine site in Arcachon Bay (France).

28 A total of 299 mature miRNAs were identified. In polluted water, 19 miRNAs were up-regulated and  
29 22 were down-regulated. We predicted that these differentially expressed miRNAs could target 490  
30 genes that were involved in ribosome biogenesis, response to hormones, response to chemical and  
31 chromatin modification. Moreover, we observed only few examples (29) of negative correlation  
32 between the expression levels of miRNAs and their targets suggesting that, in the system studied,  
33 miRNAs might not only regulate gene expression directly by degrading mRNA but also by inhibiting  
34 protein translation or by regulating other epigenetic processes.

35 This study is the first example of *in situ* investigation of the role of miRNAs in the response of a fish  
36 species to water quality. Our findings provide new insights into the involvement of epigenetic  
37 mechanisms in the response of animals chronically exposed to pollution and pave the way for the  
38 utilization of miRNAs in aquatic ecotoxicology.

39

40 **Capsule**

41 This work explores the repertoire of small non-coding RNAs (microRNAs) and its response to water  
42 pollution in the endangered European eel.

43

44 **Keywords**

45 MicroRNA; *Anguilla anguilla*; Aquatic ecotoxicology; Epigenetics; Biomarker

46

## 47 **Introduction**

48 MicroRNAs (miRNAs) are a class of small non-coding RNA <sup>1,2</sup>. These 20-24 nucleotides-long  
49 sequences associate with the 3'-untranslated region (3'-UTR) of target messenger RNAs (mRNAs),  
50 and post-transcriptionally regulate the expression of numerous genes by mediating translational  
51 repression or mRNA degradation <sup>3,4</sup>. Owing to this ability to regulate gene expression without  
52 involving DNA sequence changes, miRNAs are hence considered as epigenetic factors like histone  
53 modification and DNA methylation. In mammals, more than 50% of mRNAs are predicted to be the  
54 subject of miRNA-mediated control <sup>5</sup>. One miRNA may regulate hundreds of target mRNAs, and one  
55 mRNA may contain multiple binding sites for multiples miRNAs <sup>6</sup>, thus resulting in a complex  
56 regulatory network. Although miRNAs are involved in regulation of almost all cellular processes, such  
57 as development, growth, apoptosis, immunity and maintenance of tissue-specific function <sup>7</sup>,  
58 mechanistic aspects of this regulation are not fully understood. In mammals, the aberrant expression of  
59 miRNAs has been linked to various diseases and cancer <sup>8,9</sup> and to the inheritance of stress-induced  
60 phenotypes across generations <sup>10</sup>. Toxic environmental factors such as nanoparticles, organic  
61 pollutants and metals can alter miRNA expression <sup>11,12</sup>. These changes in miRNAs expression may be  
62 part of the adaptive responses of organisms to pollutants and of the mechanisms of toxicity and disease  
63 aetiology.

64 Advances and accessibility of sequencing technologies, coupled with the development of microRNAs  
65 dedicated tools are improving the sensitivity of analyses and the ability to detect lowly abundant small  
66 RNAs <sup>13-16</sup> even in non-model species. The latest release of the miRNA database (miRBase 21)  
67 contains 28,645 hairpin precursors from 223 species <sup>17</sup>. In aquatic organisms, the role of miRNAs has  
68 been investigated in response to various environmental stresses, such as immune challenge, metals,  
69 salinity, hypoxia or acidic water <sup>18-24</sup>.

70 As a catadromous fish species, the European eel *Anguilla anguilla* (Anguillidae; Teleostei) exhibits a  
71 complex life cycle including marine (spawning, larval phase and sexual maturation) and continental  
72 (somatic growth) environments. Moreover, *A. anguilla* is a panmictic species and its unique life cycle  
73 makes it particularly vulnerable to pollution. After spawning in the Sargasso Sea, eel larvae drift back

74 toward European and North African continental waters following oceanic currents such as the Gulf  
75 Stream and the North Atlantic Drift. There, they first metamorphose into glass eel (unpigmented)  
76 before reaching the juvenile growth phase (yellow eel) in continental water bodies where they remain  
77 up to 15-20 years, often moving across different habitat types. This stage ends with a second  
78 metamorphosis and maturing eels (silver eels) will migrate back to the Sargasso Sea to reproduce and  
79 die <sup>25,26</sup>. For their unusual life cycle and vulnerability to pollution, eels are considered as sentinel  
80 species in ecotoxicology <sup>27,28</sup>. Moreover, the species has been considered as critically endangered since  
81 2008 (IUCN red list) <sup>29</sup>. Its catadromous migratory behaviour, long life, serious habitat reduction,  
82 migration barriers, pollution, human-introduced diseases, overfishing, as well as climatic events may  
83 be amongst the causes of the catastrophic collapse of the European eel population observed over the  
84 past decades <sup>30-32</sup>. A better understanding of the biology of this species may come from the use of  
85 high-throughput (next generation) sequencing technologies that have recently provided transcriptomic  
86 <sup>33-36</sup> and genomic <sup>37-39</sup> information. However, like in many non-model species, the non-coding portion  
87 of these data has been overlooked in spite of the emerging importance of their role in gene expression  
88 regulation. So far, in *A. anguilla*, microRNAs were only predicted from *de novo* assembled  
89 transcriptome <sup>33</sup> while in the related species *A. marmorata* (marbled eel), microRNAs were  
90 specifically sequenced and their expression pattern assessed in response to salinity <sup>20</sup>.

91 In previous studies, the responses of wild immature yellow eels to pollution was investigated in 8  
92 locations presenting a broad contamination gradient in France and Canada <sup>34,40-42</sup>. These studies  
93 identified Arcachon Bay and the Gironde Estuary as the cleanest and the most contaminated French  
94 sites, respectively. We used samples from these two highly contrasted sites in order i) to identify  
95 evolutionarily conserved as well as novel miRNAs in the European eel *Anguilla anguilla* by using  
96 next generation sequencing, ii) to compare their expression between a pristine and a polluted site and  
97 iii) to link miRNA and mRNA levels. Comprehension of gene regulation by miRNAs may give new  
98 insights into a neglected mechanism of response to environmental pollution in aquatic organisms.

99

## 100 **Material and Methods**

101 **Sample collection** – European eels *Anguilla anguilla* (Linnaeus, 1758) were collected from two  
102 sampling sites with a contrasted contamination profile. The Gironde estuary (GE, 45° 12.110'N, 0°  
103 43.579'W) is considered as a highly impacted site while Arcachon Bay (AB, 44° 41.300'N, 1°  
104 1.650'W) represents a pristine environment. Six individuals were collected on each site. The liver was  
105 dissected, transferred in RNA-later solution and placed at 4°C overnight before storage at -20°C until  
106 RNA extraction. These samples were part of a previous study carried out by our group <sup>34</sup>.

107 **Condition indices, tissue composition and trace metal analyses** – Information concerning the 12  
108 individuals used in the present study (biometry, tissue composition and trace metals) were extracted  
109 from previously published data <sup>34,41,43,44</sup>. Fulton index and HSI were calculated as (total weight /  
110 length<sup>3</sup>) x 100 and as (liver weight / total weight) x 100, respectively (weight is in grams and length is  
111 in centimeters). The silvering status of the eels was assessed by the calculation of the ocular index (OI)  
112 according to the Pankhurst method <sup>45</sup>. Statistical differences between the two sites were assessed either  
113 with Student t tests or Wilcoxon tests after checking assumptions of normality and homoscedasticity.

114 **Small RNA isolation and deep sequencing** – Total RNA were isolated from individual liver tissues  
115 following the Trizol™ Reagent (Invitrogen™) protocol. MicroRNA were then purified by using the  
116 PureLink™ MiRNA Isolation kit (Invitrogen™) according to the manufacturer's instructions. Total  
117 RNA and MicroRNA concentration and quality were assessed by measuring the absorbance at 260nm  
118 and 280nm, and migration on a TBE-urea 12% polyacrylamide gel. The NEBNext® Small RNA  
119 Library Prep Set for Illumina® (New Englan BioLabs®) was used to prepare the sequencing libraries.  
120 Fragment size (around 150 bp) and quality of the individual libraries were checked on a 2100  
121 Bioanalyzer® (Agilent ®) with a High Sensitivity chip. The 12 individually tagged libraries were then  
122 sequenced using a HiSeq 4000 sequencer at the GenomeEast microarray and sequencing platform  
123 (Illkirch, France).

124 **MicroRNA discovery analysis** – The 50bp raw reads from each of the 12 samples were quality  
125 filtered (Q > 30) and adaptors (5' AGATCGGAAGAGCACACGTCTGAACTCCAGTCAC 3') were  
126 removed. The *Anguilla anguilla* genome <sup>37</sup> (GenBank accession number: AZBK00000000.1) was used  
127 as a reference genome. High quality, adaptor processed reads were aligned against the eel reference  
128 genome using the miRDeep2 package <sup>14,46</sup> with default parameters. Our strategy was similar to the one

129 used for the discovery and characterization of miRNA in the salmon *Salmo salar*<sup>47</sup>. Shortly,  
130 miRDeep2 provides a list of putative miRNA precursors with their corresponding mature and star  
131 sequences. Each precursor is associated with a score that is a measurement of the posterior probability  
132 that a predicted sequence is a true miRNA gene. Each score corresponds to a signal-to-noise ratio that  
133 estimates total miRNA reported/mean estimated false positive miRNA over 100 rounds of permuted  
134 controls. We used a miRDeep2 score of 1 that yielded a signal-to-noise ratio of 10:1 as a cut-off  
135 threshold. MiRNAs over this score were further compared by BLAST searches to all known stem-loop  
136 sequences from all species deposited in miRBase, release 21<sup>17</sup>. Only miRNA with reads matching  
137 both arms (5p and 3p) of their precursors were considered. MicroRNA with an E-value < 1.e-6 were  
138 annotated as a true *Anguilla anguilla* miRNA and named accordingly to its ortholog in miRBase.  
139 Sequences with no significant match in miRBase were used as queries in a BLAST analysis against  
140 the nr/nt and refseqRNA databases in GenBank<sup>48</sup>, the functional small RNA database<sup>49</sup> and the small  
141 RNA families in Rfam<sup>50</sup> to exclude other kind of small RNA. Finally, the remaining miRNAs were  
142 aligned with the reference genome. Sequences with an E-value < 1e-6 against >5 loci were considered  
143 as repeats and discarded. Final sequences were then considered as novel miRNAs. The nomenclature  
144 rules used for the miRNA naming were in agreement with guidelines from miRBase<sup>51,52</sup>.

145 **Transcriptome annotation** – The transcriptome assembly was previously annotated based on  
146 similarity (E value  $\leq 1e-10$ ) with known proteins from the swissprot and nr protein databases using the  
147 BLASTx algorithm<sup>34</sup>. For the present work, we enriched this annotation with Gene Ontology (GO)  
148 assignment and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis that were  
149 performed using the BLAST2GO software<sup>53,54</sup> and the GhostKOALA server<sup>55</sup>, respectively.

150 Enrichment tests were performed with the Fisher exact test procedure and a correction for multiple  
151 testing using the Benjamini and Hochberg method. A Pvalue threshold of 0.01 was used. Redundancy  
152 in GO terms was reduced by the GO Trimming method<sup>56</sup> using a uniqueness threshold of 50%. For a  
153 matter of clarity, only GO and KEGG categories containing  $\geq 5$  sequences were considered.

154 **Differential expression of transcripts** – Differences in transcription levels between the two study  
155 sites were assessed with the RNAseq data extracted from Baillon *et al*<sup>34</sup> for the exact same individual

156 eels. Normalization and expression analysis were performed using the DESeq2 package <sup>57</sup>. *P*-values  
157 for differential gene expression were corrected for multiple testing using the Benjamini and Hochberg  
158 method, and a false discovery rate (FDR) threshold of 0.05 was used.

159 **Expression analysis of Eel miRNAs** – The copy number of known and novel mature miRNAs were  
160 counted with the quantifier module in miRDeep2 (additional files 1 and 3). Normalization and  
161 expression analysis were performed using the DESeq2 package <sup>57</sup>. The false discovery rate (FDR)  
162 threshold was 0.05. In order to validate these results, the expression profiles of 10 mature miRNAs  
163 were assessed using qRT-PCR (additional file 4). Reverse transcriptions were performed by using the  
164 miScript II RT system (Qiagen®). Real Time PCR Analyses were performed on a Roche LightCycler®  
165 480 Instrument II by using the miScript PCR Starter Kit (Qiagen®). Cycling conditions were 95°C for  
166 15 seconds followed by 40 cycles of 15 seconds at 94°C, 30 seconds at 55°C and 30 seconds at 70°C.

167 **Target prediction** – The sequences and annotation of protein-coding transcripts were retrieved from  
168 Baillon *et al* <sup>34</sup>. The target genes of mature eel miRNAs were predicted using the miRanda software <sup>58</sup>  
169 that is based on the local alignment between the query miRNA sequences and the reference  
170 transcriptome and on the thermodynamic stability of miRNA/mRNA duplexes (parameters used were  
171 score  $\geq 160$ , free energy  $\leq -25$ kcal/mol).

172

## 173 **Results**

174 **Trace metals analysis** – Differences between individuals collected in both sites are presented in Table  
175 1. First, AB and GE sites differed in salinity and dissolved oxygen concentrations. We observed no  
176 difference in Fulton condition factor between the two sites. However, the Hepato-Somatic Index (HSI)  
177 of AB eels was higher than GE eels. No difference was detected in OI and with a value below the  
178 threshold value of 6.5 <sup>45</sup>, all the individuals were indeed at the yellow eels stage. As expected from  
179 previous results <sup>41</sup>, trace metal concentrations were significantly higher in the liver of eels from GE,  
180 with the exceptions of Cr and Zn that showed no difference and Ni that was higher in AB. Cadmium  
181 showed the largest difference with a concentration more than 200-times higher in the liver of GE  
182 individuals in comparison to fish from AB.



183 **Small RNA sequencing** – The sequencing of the 12 sRNA libraries led to a total of 362,831,354 raw  
184 reads out of which 318,968,039 (87.91%) were kept for the following analyses after quality filtering  
185 ( $Q > 30$ ) and adaptor removal (detailed statistics are given in Table 2). With the exception of the  
186 sample AB03 (59.35%), between 72.12% (AB06) and 89.40% (GE02) of these reads were 21-23  
187 nucleotides (nt) in length (Figure 1). Another population of 28-30 nt was identified, representing on  
188 average 12.99% and 2.36% of the reads in AB and GE samples, respectively. These two short reads  
189 populations represented between 81.70% and 90.76% of the total high-quality reads. Up to 87.73% of  
190 these reads could be mapped against the reference genome for miRNA prediction (Table 2). The raw  
191 sequencing data from this study were submitted to the NCBI Gene Expression Omnibus<sup>59</sup> under GEO  
192 Accession GSE109689.

193 **Identification of known and novel miRNAs in *A. anguilla*** – A total of 820 putative miRNA  
194 precursors were predicted by the miRDeep2 analysis (additional file 1) amongst which 474 had a score  
195 above 1 and reads on both harms, *i.e.* were associated with both a 3p and a 5p sequence (Figure 2).  
196 Amongst those, 230 pre-miRNA provided a match against a stem-loop sequence in miRBase (E-value  
197  $< 1e-6$ ) and were considered *Anguilla anguilla* orthologs of an evolutionary conserved miRNA gene  
198 and named in accordance with the nomenclature rules. Most of these conserved miRNAs (181 out of  
199 230) matched with miRNAs from other fish species in miRBase. They corresponded to 158 unique  
200 mature miRNAs from 96 evolutionary conserved families. The remaining 244 putative precursors  
201 were further analyzed by BLAST searches against the reference genome and other RNA databases.  
202 That allowed us to identify 145 potential novel pre-miRNAs that were named based on their genome  
203 location. Redundancy was lower since 141 unique novel mature miRNAs were identified. Taken  
204 together we thus identified a total of 375 miRNA precursors with their respective mature and star  
205 sequences. The total number of unique mature miRNAs was 299 (detailed list provided in additional  
206 file 2). In all libraries, aan-mir-122 was on average 14 times more expressed than the second most  
207 abundant mature miRNA (ranging from ~4 in AB06 to ~25 times in GE05). Other abundant miRNAs  
208 included aan-mir-21, aan-mir-22 and aan-let-7a (additional file 3).

209 **Prediction of transcriptomic targets** – In order to determine the biological processes potentially  
210 regulated by the 299 mature microRNAs we identified, we predicted their target genes in the

211 *anguilla*'s transcriptome assembly<sup>34</sup>. The *A. anguilla* transcriptome contains 18,113 contigs with high  
212 homology with known sequences (BLASTx, Evalue  $\leq 1e-10$ ). Although 3'UTR regions are the most  
213 important regulatory sites for post-transcriptional regulation of gene expression by miRNAs, we  
214 considered whole length transcripts. 242 miRNAs showed a potential to hybridize within the  
215 transcriptome wherein 3,637 contigs were identified as potential targets (additional file 4). As  
216 expected, one miRNA could target several transcripts and one transcript could be targeted by several  
217 miRNAs. For instance, the novel miRNA aan-scaffold-2762 could hybridize with up to 271 targets  
218 and the transcript contig\_15454, encoding a homologue of E3 ubiquitin-protein ligase UBR2 (Uniprot  
219 accession number Q8I WV8), was potentially the target of 8 different miRNAs. The entire set of  
220 predicted mRNA targets was further functionally categorized through GO annotation and KEGG  
221 pathway analysis. No GO term, nor KEGG pathway, was significantly enriched in the list of target  
222 genes compared to the transcriptome background, suggesting that miRNAs might affect any biological  
223 process.

224 **Differentially expressed mature miRNAs between the two sites** – One objective of this study was  
225 to evaluate the response of miRNAs to differing levels of contamination. A multidimensional scaling  
226 of the count data showed that despite inter-individual variability, samples are well separated according  
227 to their site of origin (additional file 5). We identified 41 mature miRNAs with differential expression  
228 (DE) in GE compared to the reference site (AB). Nineteen (19) miRNAs were up-regulated, while 22  
229 were down-regulated (Table 3). The most up-regulated and down-regulated miRNAs were  
230 respectively mir-29a (2.77 times more expressed in GE) and mir-462 (3.97 times less expressed in  
231 GE). These results were validated by qRT-PCR ( $r^2 = 0.88$ , additional file 5). Amongst all the DE  
232 miRNAs, 35 were likely to target 490 sequences in *A. Anguilla*'s transcriptome (Table 3). The targets  
233 of up-regulated miRNAs were functionally enriched in transcripts encoding proteins involved in  
234 ribosome biogenesis. This process is hence expected to decrease in GE. Biological processes affected  
235 by down-regulated miRNAs were related to the response to hormones (*e.g.* GO:0016568), response to  
236 chemicals (*e.g.* GO:0071417) and chromatin organisation (*e.g.* GO:0006325) (Figure 3). These  
237 processes are then expected to increase in GE.

238 **Links between DE miRNAs and DE transcripts** – Based on previous data <sup>34</sup>, we identified 1,554  
239 down-regulated and 1,343 up-regulated genes in the Gironde Estuary compared to the Arcachon Bay  
240 (additional file 7). Down-regulated genes contained a significant proportion of sequences related to  
241 lipid metabolism (GO:0006629 and GO:0044255), response to chemical (GO:0001101 and  
242 GO:0009410), primary metabolism, development (GO:0051216 and GO:0032836) and hormone  
243 metabolism (GO:0042445). Up-regulated transcripts were involved in processes such as transport  
244 (GO:0006810, GO:0051649 and GO:0032940), development (GO:0007423, GO:0007389,  
245 GO:0048840 and GO:0035889) and response to stimulus (GO:0010035 and GO:0071216) (Figure 3).  
246 Due to the action mode of miRNAs, their expression patterns generally showed an inverse relationship  
247 with those of their mRNA targets. We compared the list of DE genes with the lists of DE miRNAs and  
248 their targets. Out of the 35 DE miRNAs that had putative targets in the transcriptome, 20 of them were  
249 able to hybridize to 49 (out of 490) DE mRNAs targets, forming 52 miRNA-mRNA pairs with both  
250 positive and negative relationships in their expression values ( $\log_2FC$ ). In detail, there were 29 (out of  
251 52) negative miRNA-mRNA interactions (Spearman's  $\rho = -0.69$ , Pvalue =  $3.46e-05$ ) involving 18 DE  
252 miRNAs and 29 DE mRNAs. Whilst a single DE miRNA could target up to 4 different DE transcripts,  
253 each DE transcript was targeted by only one DE miRNA. Many of these DE transcripts were involved  
254 in biological processes cited above, such as immune response and lipid metabolism (Table 4).

255

## 256 **Discussion**

257 The increasing evidence that the expression of miRNAs is affected by several known toxicants as well  
258 as oxidative and other forms of cellular stress certainly suggests an important role of miRNAs in  
259 toxicology, which could provide a link between environmental influences and gene expression <sup>60</sup>. The  
260 use of miRNAs in environmental toxicology is expanding, particularly in human health <sup>8,11,12</sup>. This  
261 interest mainly results from the strong potential of miRNAs to become robust, minimally invasive and  
262 non-expensive biomarkers for various diseases <sup>61</sup>. In the same manner, examples of miRNA-related  
263 work are more frequent in aquatic organisms exposed to environmental stressors. One can cite some  
264 recent publication in bivalves <sup>22,62,63</sup>, crustaceans <sup>64,65</sup> or teleost fish <sup>20,21,66,67</sup> for instance. To our

265 knowledge, our work is the first *in situ* study using miRNA in an aquatic organism. MicroRNAs are  
266 being characterized in an increasing number of teleost fish species where they repress target mRNA  
267 abundance or translation in a highly conserved manner across most eukaryotes <sup>68</sup>.

268 In this study, we conducted a comprehensive identification of the microRNA repertoire in the liver of  
269 the European eel *A. anguilla* and its response to environmental pollution. This organ was chosen  
270 owing to its role in pollutant metabolism, accumulation and toxicity. Only trace metal concentrations  
271 were reported here since they were measured in the liver. However, one should note that eels from the  
272 Gironde Estuary also showed higher levels of persistent organic pollutants (Poly-Chlorinated  
273 Biphenyls, Organo-Chlorine Pesticides, and Poly-Brominated Diphenyl Ethers) in their muscle  
274 compared to eels from the Arcachon Bay, as evidenced by previous analyses <sup>34,43</sup>. Owing to panmixia  
275 and proximity between the two sites, this contrast is most likely due to water quality rather than  
276 population differences.

277 The size distribution of the small RNA libraries revealed that 22-23 nt microRNAs were dominant,  
278 followed by small RNAs of 28-29 nt in length (Figure 1). This distribution is similar to what was  
279 found in the sister species *A. marmorata* <sup>20</sup>. It is interesting to note that, in both eel species, the relative  
280 abundance of 22-23 nt and 28-29 nt varies depending on environmental conditions, which could  
281 represent a new path of investigation in the future. These 28-29 nt sRNA might correspond to piRNA  
282 (piwi-interacting RNA that have a length of 24-32 nt <sup>1,69</sup>), another smallRNA silencing system that  
283 plays a role in germline specification, gametogenesis, transposon silencing, genome integrity, and  
284 stem cell maintenance across the animal phylogeny <sup>70</sup>.

285 We identified a total of 299 mature miRNA in eel liver, of which 158 were known and 141 were  
286 novel. This number is comparable to what is currently available in the miRBase database for the  
287 model species *Danio rerio* (375 mature sequences). This identification benefited from the availability  
288 of a sequenced genome of *A. anguilla* <sup>37</sup>. Only mature miRNAs were considered in our analysis since  
289 they eventually regulate gene expression at the post-transcriptional level. In all libraries, mir-122 was  
290 the most abundant mature sequence. This was not surprising since it is the most abundant liver  
291 miRNA in vertebrates (>70% of the total hepatic miRNA expression) with exquisite tissue specificity.

292 Mir-122 is mainly involved in lipid homeostasis <sup>71</sup>. Its expression level increased by 2.34-fold in  
293 polluted water (Table 3). Such accumulation is associated to injured hepatocytes and is a biomarker of  
294 liver toxicity in zebrafish <sup>72</sup> and mammals <sup>73</sup>. The alteration of lipid metabolism was also evidenced by  
295 prevalence of related Gene Ontology terms amongst down-regulated transcripts (Figure 3). In farmed  
296 tilapia (GIFT, *Oreochromis niloticus*), the level of mir-122 was negatively correlated to Cd stress and  
297 promotes hepatic antioxidant defence <sup>74</sup>. This discrepancy with the present work, where Cd was the  
298 metal showing the largest difference between the two sites, could either be due to the action of other  
299 stressors or to taxa specific mechanisms of miRNA regulation. The two other abundant miRNAs, mir-  
300 21 and mir-22, were also shown to play a role in the development of hepatic diseases <sup>75,76</sup>. Liver  
301 damage was confirmed by a series of transcriptomic changes described previously <sup>34</sup>. This first set of  
302 results shows the high tissue specificity of our approach and confirms the potential of miRNAs in  
303 aquatic ecotoxicology.

304 We chose to study two contrasted sites both in terms of organic and metallic contaminants <sup>34,40</sup>. These  
305 sites also differed in salinity (19.9‰ and 7.78‰ for Arcachon Bay and Gironde Estuary, respectively)  
306 and oxygen percentage (60% vs. 86.5%) whereas the temperature was relatively similar (21.9°C vs.  
307 21.3°C) (Table 1). With a dissolved oxygen concentration of 4.68 mg/l, site AB faces moderate  
308 hypoxia according to the European Water Framework Directive (WFD, 2000/60/EC). In such a  
309 complex context, it is obviously challenging to identify causing agents. However, since contamination  
310 of aquatic ecosystems typically includes many pollutants that present different mechanisms of action  
311 likely interacting together and under the influence of natural factors, this situation might provide more  
312 realistic information as well as a first insight into the role of miRNAs in a polluted environment. In  
313 fish, salinity and hypoxia were previously shown to affect the miRNA regulatory machinery <sup>20,21,77</sup>. In  
314 our study, only 5 miRNAs (miR-19b, miR-139, miR99a, miR-454 and miR-221) and 3 miRNAs (miR-  
315 125a, miR-181c and miR-30b) were found to respond to salinity and hypoxia, respectively, in the  
316 same way than in the literature cited above. Moreover, eels were shown to be able to cope with much  
317 more severe hypoxia, down to about 30% O<sub>2</sub> saturation <sup>78</sup>. Then, the differential expression of the  
318 majority of the miRNA we identified could most likely be linked to the presence of pollutants. The

319 expression level of 19 miRNAs increased in the polluted site while the expression of 22 miRNAs was  
320 higher in AB. This relatively small number of DE miRNA is similar to what was found in other fish  
321 species exposed to environmental stressors<sup>19-21,24</sup> and could be due to the relatively low number of  
322 samples per site and inter-individual variability (additional file 5) that should be taken into account in  
323 future studies. The most expressed known miRNAs in AB (>3 fold) were miR-462, miR-15e and miR-  
324 23a, and the most up-regulated known miRNAs in GE (>2 fold) were miR-29a, miR-122 (discussed  
325 above) and miR-203. As previously, the fold change values measured here are in agreement with what  
326 can be found for Teleosts in the literature<sup>19-21,24</sup>. The role of these miRNAs could be inferred from  
327 previous studies.

328 Despite the highly conserved mode of action of miRNAs, caution must be taken when comparing fish  
329 and mammals as only a small fraction of miRNA-mRNA interactions (~10%) is conserved<sup>79</sup> and  
330 identification of conserved targets between species remains challenging<sup>80</sup>. For instance, in fish, miR-  
331 462 was shown to be upregulated after hypoxia treatment in liver<sup>81,82</sup> or virus infection<sup>18,83</sup>. MiR-462  
332 is only described in teleost fish and is actually an orthologue of the human miR-191 that plays a role in  
333 cell cycle regulation, but not in viral infection<sup>83</sup>. The role of miR-23a instead seems to be conserved.  
334 It was identified as an essential regulator of adipocyte differentiation of which increased expression  
335 might reduce lipid accumulation and triglyceride content in adipocytes<sup>84</sup>. This role was confirmed in  
336 our study by the identification of a transcript encoding a very long-chain acyl-CoA synthetase  
337 (Uniprot accession O14975, additional file 5) as a putative target of miR-23a. In human, miR-29a is  
338 known to regulate cell proliferation, differentiation and apoptosis as several of its targets are  
339 oncogenes or anti-apoptotic genes<sup>85</sup>. Amongst the predicted mRNA targets of miR-29a, we identified  
340 a thyroid receptor-interacting protein homologue (Uniprot accession Q15643, additional file 5) of  
341 which degradation could reduce gene transcription in response to thyroid hormone while miR-122  
342 predicted targets included transcripts encoding proteins involved in lipid synthesis, energy generation,  
343 platelet activation, apoptosis and transcription regulation. The role of miR-203 remains unclear as its  
344 overexpression might promote both apoptosis and tumor growth, as well as host-virus interactions<sup>86</sup>.

345 This difference in lipid metabolism is not confirmed by our observation as fish from the two sites

346 showed no significant difference regarding their muscular lipid content (table 1), contrary to previous  
347 studies that showed that eels exposed to Cd <sup>87</sup> and organic pollutants <sup>88</sup> increased their fat consumption  
348 and/or reduced their energy reserves. We can here hypothesize that miRNA regulation in the liver  
349 contributes to maintain a significant production of triglycerides and their export to the main site of  
350 lipid storage in eels, *i.e.* the muscle <sup>87</sup>. At the opposite, the lower HSI values in fish from the estuary  
351 could indicate an impairment in the accumulation of energy reserves under the form of carbohydrate  
352 such as glycogen in hepatocytes.

353 More comprehensive information was gained from the functional analysis of miRNA targets in *A.*  
354 *Anguilla*'s transcriptome (Figure 3). Based on gene ontology of these targets, changes in miRNA  
355 expression might result in reduced ribosome biogenesis and increased response to hormone and  
356 chemical, and chromatin modification. The effect on ribosome biogenesis could be evidenced by a  
357 significantly reduced level of total proteins in the liver of GE eels (Table 1). One could expect to find  
358 some similar functions in the predicted targets of up-regulated miRNAs and down-regulated  
359 transcripts, and conversely. In spite, we found no relation between enriched gene ontology in miRNA  
360 targets and mRNA, as well as a small number of miRNA-mRNA pairs with negative correlation  
361 (Table 4). This situation was also observed in a comparable study carried out in *Pelteobagrus vachelli*  
362 exposed to hypoxia <sup>21</sup> in which the authors identified only 97 negative correlations out of 308 miRNA-  
363 mRNA pairs. In the same study, and in the same manner as our results, most of the DE mRNA (43 out  
364 of 60) were target by a single DE miRNA. This highlights that the identification of functional miRNA-  
365 targets interaction is still challenging and/or false positives might still be frequent with *in silico*  
366 analyses <sup>89</sup>. MicroRNAs are obviously not the only mode of regulation of gene expression, however  
367 based on the lack of correspondence between miRNA targets and DE transcripts, we could  
368 hypothesize that the degradation of messenger RNA was not the major mode of action of miRNAs  
369 here and that they act more likely by inhibiting mRNA translation into proteins, instead. This  
370 inhibition might be reinforced by the targeting of mRNA involved in ribosome assembly and function.  
371 This result is in accordance with those obtained in juvenile zebrafish exposed to the antibacterial agent  
372 triclosan where "structural constituent of ribosome" and "ribosome biogenesis" are amongst the  
373 significantly enriched annotation of differentially expressed miRNAs target genes <sup>90</sup>. Which might

374 suggest a conserved mode of action of miRNA in response to pollutants in fish. Another way by which  
375 miRNA could regulate gene expression is through chromatin modification, as suggested by the  
376 presence of 28 epigenetic regulators in miRNA targets (additional file 8). In human, miRNA and  
377 epigenetic regulation are not entirely separable and a strong interplay exists, but the mechanisms of  
378 cooperation are still unclear<sup>91,92</sup>. Compared to mammals, the study of epigenetics in teleost fish has  
379 received less attention so far<sup>67</sup>. Pollutant-induced changes in epigenetic marks were already  
380 documented<sup>42,93,94</sup>. For instance, transcriptomic changes in the liver of wild yellow perch (*Perca*  
381 *flavescens*) exposed to cadmium and copper may be driven through histone modifications<sup>95</sup>. The  
382 present work suggests that similar effects in eels could be mediated by miRNA regulation.  
383 The European eel spends most of its lifetime (about 8-15 years) in European estuarine and freshwater  
384 basins in a stage known as yellow eels. During this period, eels accumulate a considerable amount of  
385 lipid reserves, which are fundamental for the success of the long oceanic migration since during this  
386 period they do not feed. Before initiating their oceanic migration to reproduce in the Sargasso Sea,  
387 they metamorphose into silver eel<sup>25,26</sup>. Laboratory and field studies have shown that eels can  
388 bioaccumulate high concentrations of environmental contaminants, which have been shown to cause  
389 mortality, growth delay, reproductive alterations, tumours, malformations, nervous and endocrine  
390 disturbances and immunological changes<sup>28</sup>. In the present work, we evidenced a role of miRNAs in  
391 the regulation of genes involved in all these processes, reinforcing the potential of miRNAs as  
392 biomarkers of yellow eel's health status in a polluted environment. MiRNAs involved in lipid  
393 metabolism (like miR-122) and endocrine function might be of particular interest owing to the critical  
394 importance of these biological processes in the reproductive success of *A. anguilla*. Although lipid  
395 storage might not be absolutely necessary to initiate metamorphosis (silvering<sup>96</sup>), a sufficient energy  
396 reserve is needed to cover the migratory needs, ensure gonadal maturation and reproduction.  
397 Moreover, morphological and physiological changes that take place during the silvering phase are  
398 initiated and regulated by hormonal changes<sup>28</sup>.  
399 The present study was carried out on wild organisms, in a multi-stress context. In such conditions, it is  
400 difficult to establish significant relationships between the expression level of one particular miRNA



401 and one particular contaminant. Further experiments carried out with more individuals, in controlled  
402 conditions, different organs and/or at different developmental stages are thus needed to fully reveal the  
403 potential of miRNAs as biomarkers.

404

## 405 **Conclusion**

406 To the best of our knowledge, this study is the first example of high-throughput sequencing of  
407 microRNAs in wild aquatic organisms. We were able to identify both conserved and putative novel  
408 miRNAs and their targets at the transcriptomic level in the liver of European eels from two sites with  
409 contrasted pollution profiles. We showed that some miRNA were differentially expressed in  
410 organisms experiencing chronic exposure to pollution and that these changes have the potential to  
411 affect important biological processes such as lipid and hormone metabolism or protein synthesis.  
412 Moreover, our work also revealed promising aspects of epigenetics in teleost fish, as miRNAs might  
413 regulate gene expression by interacting with other epigenetic mechanisms such as chromatin  
414 modification. Part of the biological processes affected by miRNAs for which expression levels were  
415 altered were similar to previous transcriptome studies, confirming the potential role of miRNAs as  
416 early molecular markers for monitoring stress responses in exposed animals.

417

## 418 **Funding**

419 MicroRNA libraries' quality check was performed at the Genome Transcriptome Facility of Bordeaux  
420 (grants from the Conseil Régional d'Aquitaine 20030304002FA and 20040305003FA, from the  
421 European Union FEDER 2003227 and from Investissements d'Avenir ANR-10-EQPX-16-01).  
422 Sequencing was performed by the IGBMC Microarray and Sequencing platform, a member of the  
423 'France Génomique' consortium (ANR-10-INBS-0009). This work was supported by the Regional  
424 Council of Aquitaine, the Agence Nationale de la Recherche of France (ANR Immortel ANR-10-  
425 INTB-1704) and a strategic project grant of the Natural Sciences and Engineering Research Council of  
426 Canada to Patrice Couture, jointly funded by the Innovative Project grant of the UMR EPOC 5805.

427

## 428 **References**

- 429 1. Großhans, H. & Filipowicz, W. Molecular biology: The expanding world of small RNAs. *Nature*  
430 **451**, 414–416 (2008).
- 431 2. Yasuda, J. & Hayashizaki, Y. The RNA continent. *Adv. Cancer Res.* **99**, 77–112 (2008).
- 432 3. Bartel, D. P. MicroRNAs: Genomics, Biogenesis, Mechanism, and Function. *Cell* **116**, 281–297  
433 (2004).
- 434 4. Chekulaeva, M. & Filipowicz, W. Mechanisms of miRNA-mediated post-transcriptional regulation  
435 in animal cells. *Curr. Opin. Cell Biol.* **21**, 452–460 (2009).
- 436 5. Friedman, R. C., Farh, K. K.-H., Burge, C. B. & Bartel, D. P. Most mammalian mRNAs are conserved  
437 targets of microRNAs. *Genome Res.* **19**, 92–105 (2009).
- 438 6. Lim, L. P. *et al.* Microarray analysis shows that some microRNAs downregulate large numbers of  
439 target mRNAs. *Nature* **433**, 769–773 (2005).
- 440 7. Bushati, N. & Cohen, S. M. microRNA functions. *Annu. Rev. Cell Dev. Biol.* **23**, 175–205 (2007).
- 441 8. Huang, J.-T., Wang, J., Srivastava, V., Sen, S. & Liu, S.-M. MicroRNA Machinery Genes as Novel  
442 Biomarkers for Cancer. *Front. Oncol.* **4**, (2014).
- 443 9. Cho, W. C. S. MicroRNAs in cancer - from research to therapy. *Biochim. Biophys. Acta* **1805**, 209–  
444 217 (2010).
- 445 10. Gapp, K. *et al.* Implication of sperm RNAs in transgenerational inheritance of the effects of early  
446 trauma in mice. *Nat. Neurosci.* **17**, 667–669 (2014).
- 447 11. Yu, H. W. & Cho, W. C. The role of microRNAs in toxicology. *Arch. Toxicol.* **89**, 319–325 (2015).
- 448 12. Hou, L., Wang, D. & Baccarelli, A. Environmental chemicals and microRNAs. *Mutat. Res.* **714**,  
449 105–112 (2011).
- 450 13. Creighton, C. J., Reid, J. G. & Gunaratne, P. H. Expression profiling of microRNAs by deep  
451 sequencing. *Brief. Bioinform.* **10**, 490–497 (2009).
- 452 14. Friedländer, M. R. *et al.* Discovering microRNAs from deep sequencing data using miRDeep. *Nat.*  
453 *Biotechnol.* **26**, 407–415 (2008).
- 454 15. Li, Y. *et al.* Performance comparison and evaluation of software tools for microRNA deep-  
455 sequencing data analysis. *Nucleic Acids Res.* **40**, 4298–4305 (2012).
- 456 16. Eminaga, S., Christodoulou, D. C., Vigneault, F., Church, G. M. & Seidman, J. G. Quantification of  
457 microRNA Expression with Next-Generation Sequencing. *Curr. Protoc. Mol. Biol. Ed. Frederick M*  
458 *Ausubel Al* **0 4**, Unit-4.17 (2013).
- 459 17. Kozomara, A. & Griffiths-Jones, S. miRBase: annotating high confidence microRNAs using deep  
460 sequencing data. *Nucleic Acids Res.* **42**, D68–D73 (2014).
- 461 18. Andreassen, R. *et al.* Identification of differentially expressed Atlantic salmon miRNAs responding  
462 to salmonid alphavirus (SAV) infection. *BMC Genomics* **18**, 349 (2017).
- 463 19. Wang, L., Bammler, T. K., Beyer, R. P. & Gallagher, E. P. Copper-induced deregulation of  
464 microRNA expression in the zebrafish olfactory system. *Environ. Sci. Technol.* **47**, 7466–7474  
465 (2013).
- 466 20. Wang, X. *et al.* MicroRNA-Sequence Profiling Reveals Novel Osmoregulatory MicroRNA  
467 Expression Patterns in Catadromous Eel *Anguilla marmorata*. *PLOS ONE* **10**, e0136383 (2015).
- 468 21. Zhang, G. *et al.* Integrated analysis of mRNA-seq and miRNA-seq in the liver of *Pelteobagrus*  
469 *vachelli* in response to hypoxia. *Sci. Rep.* **6**, srep22907 (2016).
- 470 22. Zhou, Z. *et al.* The identification and characteristics of immune-related microRNAs in haemocytes  
471 of oyster *Crassostrea gigas*. *PLoS One* **9**, e88397 (2014).
- 472 23. Lau, K. *et al.* Identification and expression profiling of microRNAs in the brain, liver and gonads of  
473 marine medaka (*Oryzias melastigma*) and in response to hypoxia. *PLoS One* **9**, e110698 (2014).
- 474 24. Kure, E. H. *et al.* Molecular responses to toxicological stressors: profiling microRNAs in wild  
475 Atlantic salmon (*Salmo salar*) exposed to acidic aluminum-rich water. *Aquat. Toxicol. Amst. Neth.*  
476 **138–139**, 98–104 (2013).
- 477 25. Feunteun, E. Management and restoration of European eel population (*Anguilla anguilla*): An  
478 impossible bargain. *Ecol. Eng.* **18**, 575–591 (2002).

- 479 26. Guimarães, L., Gravato, C., Santos, J., Monteiro, L. S. & Guilhermino, L. Yellow eel (*Anguilla*  
480 *anguilla*) development in NW Portuguese estuaries with different contamination levels.  
481 *Ecotoxicology* **18**, 385–402 (2009).
- 482 27. Belpaire, C., Goemans, G., Geeraerts, C., Quataert, P. & Parmentier, K. Pollution fingerprints in  
483 eels as models for the chemical status of rivers. *ICES J. Mar. Sci.* **65**, 1483–1491 (2008).
- 484 28. Geeraerts, C. & Belpaire, C. The effects of contaminants in European eel: a review. *Ecotoxicology*  
485 **19**, 239–266 (2010).
- 486 29. IUCN. *Anguilla anguilla*: Jacoby, D. & Gollock, M.: The IUCN Red List of Threatened Species 2014:  
487 e.T60344A45833138. (2013). doi:10.2305/IUCN.UK.2014-1.RLTS.T60344A45833138.en
- 488 30. Knights, B. A review of the possible impacts of long-term oceanic and climate changes and fishing  
489 mortality on recruitment of anguillid eels of the Northern Hemisphere. *Sci. Total Environ.* **310**,  
490 237–244 (2003).
- 491 31. *Spawning Migration of the European Eel*. (Springer Netherlands, 2009). doi:10.1007/978-1-4020-  
492 9095-0
- 493 32. Bevacqua, D., Melià, P., Gatto, M. & De Leo, G. A. A global viability assessment of the European  
494 eel. *Glob. Change Biol.* **21**, 3323–3335 (2015).
- 495 33. Coppe, A. *et al.* Sequencing, de novo annotation and analysis of the first *Anguilla anguilla*  
496 transcriptome: EelBase opens new perspectives for the study of the critically endangered  
497 European eel. *BMC Genomics* **11**, 635 (2010).
- 498 34. Baillon, L. *et al.* Transcriptome profile analysis reveals specific signatures of pollutants in Atlantic  
499 eels. *Ecotoxicol. Lond. Engl.* **24**, 71–84 (2015).
- 500 35. Pujolar, J. M. *et al.* Surviving in a toxic world: transcriptomics and gene expression profiling in  
501 response to environmental pollution in the critically endangered European eel. *BMC Genomics*  
502 **13**, 507 (2012).
- 503 36. Podgorniak, T. *et al.* Differences in brain gene transcription profiles advocate for an important  
504 role of cognitive function in upstream migration and water obstacles crossing in European eel.  
505 *BMC Genomics* **16**, 378 (2015).
- 506 37. Henkel, C. V. *et al.* Primitive Duplicate Hox Clusters in the European Eel’s Genome. *PLoS ONE* **7**,  
507 (2012).
- 508 38. Minegishi, Y., Henkel, C. V., Dirks, R. P. & Thillart, G. E. E. J. M. van den. Genomics in Eels —  
509 Towards Aquaculture and Biology. *Mar. Biotechnol.* **14**, 583–590 (2012).
- 510 39. Callol, A. *et al.* An Enriched European Eel Transcriptome Sheds Light upon Host-Pathogen  
511 Interactions with *Vibrio vulnificus*. *PLOS ONE* **10**, e0133328 (2015).
- 512 40. Laporte, M. *et al.* RAD sequencing reveals within-generation polygenic selection in response to  
513 anthropogenic organic and metal contamination in North Atlantic Eels. *Mol. Ecol.* **25**, 219–237  
514 (2016).
- 515 41. Pannetier, P. *et al.* A comparison of metal concentrations in the tissues of yellow American eel  
516 (*Anguilla rostrata*) and European eel (*Anguilla anguilla*). *Sci. Total Environ.* **569–570**, 1435–1445  
517 (2016).
- 518 42. Pierron, F., Daffe, G., Lambert, P., Couture, P. & Baudrimont, M. Retrotransposon methylation  
519 and activity in wild fish (*A. anguilla*): A matter of size. *Environ. Pollut.* **245**, 494–503 (2019).
- 520 43. Patey, G., Couillard, C. M., Pierron, F., Baudrimont, M. & Couture, P. Biotransformation,  
521 antioxidant and histopathological biomarker responses to contaminants in European and  
522 American yellow eels from the Gironde and St. Lawrence estuaries. *Chemosphere* **188**, 292–303  
523 (2017).
- 524 44. Caron, A. *et al.* Organic and inorganic contamination impacts on metabolic capacities in  
525 American and European yellow eels. *Can. J. Fish. Aquat. Sci.* **73**, 1557–1566 (2016).
- 526 45. Pankhurst, N. W. Relation of visual changes to the onset of sexual maturation in the European  
527 eel *Anguilla anguilla* (L.). *J. Fish Biol.* **21**, 127–140 (1982).
- 528 46. Friedländer, M. R., Mackowiak, S. D., Li, N., Chen, W. & Rajewsky, N. miRDeep2 accurately  
529 identifies known and hundreds of novel microRNA genes in seven animal clades. *Nucleic Acids*  
530 *Res.* **40**, 37–52 (2012).

- 531 47. Andreassen, R., Worren, M. M. & Høyheim, B. Discovery and characterization of miRNA genes in  
532 Atlantic salmon (*Salmo salar*) by use of a deep sequencing approach. *BMC Genomics* **14**, 482  
533 (2013).
- 534 48. Benson, D. A. *et al.* GenBank. *Nucleic Acids Res.* **41**, D36–D42 (2013).
- 535 49. Mituyama, T. *et al.* The Functional RNA Database 3.0: databases to support mining and  
536 annotation of functional RNAs. *Nucleic Acids Res.* **37**, D89–D92 (2009).
- 537 50. Nawrocki, E. P. *et al.* Rfam 12.0: updates to the RNA families database. *Nucleic Acids Res.* **43**,  
538 D130–D137 (2015).
- 539 51. Ambros, V. *et al.* A uniform system for microRNA annotation. *RNA* **9**, 277–279 (2003).
- 540 52. Griffiths-Jones, S., Grocock, R. J., van Dongen, S., Bateman, A. & Enright, A. J. miRBase: microRNA  
541 sequences, targets and gene nomenclature. *Nucleic Acids Res.* **34**, D140–144 (2006).
- 542 53. Conesa, A. *et al.* Blast2GO: a universal tool for annotation, visualization and analysis in functional  
543 genomics research. *Bioinformatics* **21**, 3674–3676 (2005).
- 544 54. Götz, S. *et al.* High-throughput functional annotation and data mining with the Blast2GO suite.  
545 *Nucleic Acids Res.* **36**, 3420–3435 (2008).
- 546 55. Kanehisa, M., Sato, Y. & Morishima, K. BlastKOALA and GhostKOALA: KEGG Tools for Functional  
547 Characterization of Genome and Metagenome Sequences. *J. Mol. Biol.* **428**, 726–731 (2016).
- 548 56. Jantzen, S. G., Sutherland, B. J., Minkley, D. R. & Koop, B. F. GO Trimming: Systematically  
549 reducing redundancy in large Gene Ontology datasets. *BMC Res. Notes* **4**, 267 (2011).
- 550 57. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-  
551 seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).
- 552 58. Enright, A. J. *et al.* MicroRNA targets in *Drosophila*. *Genome Biol.* **5**, R1 (2003).
- 553 59. Edgar, R., Domrachev, M. & Lash, A. E. Gene Expression Omnibus: NCBI gene expression and  
554 hybridization array data repository. *Nucleic Acids Res.* **30**, 207–210 (2002).
- 555 60. Lema, C. & Cunningham, M. J. MicroRNAs and their implications in toxicological research.  
556 *Toxicol. Lett.* **198**, 100–105 (2010).
- 557 61. Schraml, E., Hackl, M. & Grillari, J. MicroRNAs and toxicology: A love marriage. *Toxicol. Rep.* **4**,  
558 634–636 (2017).
- 559 62. Bao, Y., Zhang, L., Dong, Y. & Lin, Z. Identification and comparative analysis of the *Tegillarca*  
560 *granosa* haemocytes microRNA transcriptome in response to Cd using a deep sequencing  
561 approach. *PLoS One* **9**, e93619 (2014).
- 562 63. Xu, F. *et al.* Identification of conserved and novel microRNAs in the Pacific oyster *Crassostrea*  
563 *gigas* by deep sequencing. *PLoS One* **9**, e104371 (2014).
- 564 64. Chen, S., McKinney, G. J., Nichols, K. M., Colbourne, J. K. & Sepúlveda, M. S. Novel Cadmium  
565 Responsive MicroRNAs in *Daphnia pulex*. *Environ. Sci. Technol.* **49**, 14605–14613 (2015).
- 566 65. Chen, S., Nichols, K. M., Poynton, H. C. & Sepúlveda, M. S. MicroRNAs are involved in cadmium  
567 tolerance in *Daphnia pulex*. *Aquat. Toxicol. Amst. Neth.* **175**, 241–248 (2016).
- 568 66. Lee, J., Kho, Y., Kim, P.-G. & Ji, K. Exposure to bisphenol S alters the expression of microRNA in  
569 male zebrafish. *Toxicol. Appl. Pharmacol.* **338**, 191–196 (2018).
- 570 67. Best, C. *et al.* Epigenetics in teleost fish: From molecular mechanisms to physiological  
571 phenotypes. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* (2018).  
572 doi:10.1016/j.cbpb.2018.01.006
- 573 68. Bizuayehu, T. T. & Babiak, I. MicroRNA in teleost fish. *Genome Biol. Evol.* **6**, 1911–1937 (2014).
- 574 69. Wang, J. *et al.* Current Research on Non-Coding Ribonucleic Acid (RNA). *Genes* **8**, (2017).
- 575 70. Iwasaki, Y. W., Siomi, M. C. & Siomi, H. PIWI-Interacting RNA: Its Biogenesis and Functions. *Annu.*  
576 *Rev. Biochem.* **84**, 405–433 (2015).
- 577 71. Willeit, P., Skroblin, P., Kiechl, S., Fernández-Hernando, C. & Mayr, M. Liver microRNAs: potential  
578 mediators and biomarkers for metabolic and cardiovascular disease? *Eur. Heart J.* **37**, 3260–3266  
579 (2016).
- 580 72. Vliegthart, A. D. B. *et al.* Retro-orbital blood acquisition facilitates circulating microRNA  
581 measurement in zebrafish with paracetamol hepatotoxicity. *Zebrafish* **11**, 219–226 (2014).

- 582 73. Vliegthart, A. D. B. *et al.* Comprehensive microRNA profiling in acetaminophen toxicity  
583 identifies novel circulating biomarkers for human liver and kidney injury. *Sci. Rep.* **5**, (2015).
- 584 74. Qiang, J. *et al.* miR-122 promotes hepatic antioxidant defense of genetically improved farmed  
585 tilapia (GIFT, *Oreochromis niloticus*) exposed to cadmium by directly targeting a metallothionein  
586 gene. *Aquat. Toxicol.* **182**, 39–48 (2017).
- 587 75. Yang, F., Hu, Y., Liu, H.-X. & Wan, Y.-J. Y. MiR-22-silenced Cyclin A Expression in Colon and Liver  
588 Cancer Cells Is Regulated by Bile Acid Receptor. *J. Biol. Chem.* **290**, 6507–6515 (2015).
- 589 76. Caviglia, J. M. *et al.* MicroRNA-21 and Dicer are dispensable for hepatic stellate cell activation  
590 and the development of liver fibrosis. *Hepatology. Baltim. Md* (2017). doi:10.1002/hep.29627
- 591 77. Sun, S., Xuan, F., Ge, X., Zhu, J. & Zhang, W. Dynamic mRNA and miRNA expression analysis in  
592 response to hypoxia and reoxygenation in the blunt snout bream ( *Megalobrama amblycephala* ).  
593 *Sci. Rep.* **7**, 12846 (2017).
- 594 78. Cruz-Neto, A. P. & Steffensen, J. F. The effects of acute hypoxia and hypercapnia on oxygen  
595 consumption of the freshwater European eel. *J. Fish Biol.* **50**, 759–769 (1997).
- 596 79. Xu, J. *et al.* The evolution of evolvability in microRNA target sites in vertebrates. *Genome Res.* **23**,  
597 1810–1816 (2013).
- 598 80. Berezikov, E. Evolution of microRNA diversity and regulation in animals. *Nat. Rev. Genet.* **12**,  
599 846–860 (2011).
- 600 81. Huang, C.-X. *et al.* The zebrafish miR-462/miR-731 cluster is induced under hypoxic stress via  
601 hypoxia-inducible factor 1 $\alpha$  and functions in cellular adaptations. *FASEB J. Off. Publ. Fed. Am.*  
602 *Soc. Exp. Biol.* **29**, 4901–4913 (2015).
- 603 82. Huang, C.-H. *et al.* Involvement of the miR-462/731 cluster in hypoxia response in *Megalobrama*  
604 *amblycephala*. *Fish Physiol. Biochem.* **43**, 863–873 (2017).
- 605 83. Schyth, B. D. *et al.* Two Virus-Induced MicroRNAs Known Only from Teleost Fishes Are  
606 Orthologues of MicroRNAs Involved in Cell Cycle Control in Humans. *PLOS ONE* **10**, e0132434  
607 (2015).
- 608 84. Shen, L. *et al.* MicroRNA-23a regulates 3T3-L1 adipocyte differentiation. *Gene* **575**, 761–764  
609 (2016).
- 610 85. Kriegel, A. J., Liu, Y., Fang, Y., Ding, X. & Liang, M. The miR-29 family: genomics, cell biology, and  
611 relevance to renal and cardiovascular injury. *Physiol. Genomics* **44**, 237–244 (2012).
- 612 86. Deng, B. *et al.* MiRNA-203 suppresses cell proliferation, migration and invasion in colorectal  
613 cancer via targeting of EIF5A2. *Sci. Rep.* **6**, 28301 (2016).
- 614 87. Pierron, F. *et al.* Impairment of lipid storage by cadmium in the European eel (*Anguilla anguilla*).  
615 *Aquat. Toxicol. Amst. Neth.* **81**, 304–311 (2007).
- 616 88. Sancho, E., Ferrando, M. & Andreu, E. Effects of sublethal exposure to a pesticide on levels of  
617 energetic compounds in *Anguilla anguilla*. *J. Environ. Sci. Health Part B* **33**, 411–424 (1998).
- 618 89. Tabas-Madrid, D. *et al.* Improving miRNA-mRNA interaction predictions. *BMC Genomics* **15**, S2  
619 (2014).
- 620 90. Wang, F., Liu, F. & Chen, W. Exposure to triclosan changes the expression of microRNA in male  
621 juvenile zebrafish (*Danio rerio*). *Chemosphere* **214**, 651–658 (2019).
- 622 91. Osella, M., Riba, A., Testori, A., Corà, D. & Caselle, M. Interplay of microRNA and epigenetic  
623 regulation in the human regulatory network. *Front. Genet.* **5**, (2014).
- 624 92. Wang, X., Zheng, G. & Dong, D. Coordinated action of histone modification and microRNA  
625 regulations in human genome. *Gene* **570**, 277–281 (2015).
- 626 93. Head, J. A., Dolinoy, D. C. & Basu, N. Epigenetics for ecotoxicologists. *Environ. Toxicol. Chem.* **31**,  
627 221–227 (2012).
- 628 94. Vandegehuchte, M. B. & Janssen, C. R. Epigenetics and its implications for ecotoxicology.  
629 *Ecotoxicol. Lond. Engl.* **20**, 607–624 (2011).
- 630 95. Pierron, F. *et al.* Effects of chronic metal exposure on wild fish populations revealed by high-  
631 throughput cDNA sequencing. *Ecotoxicology* **20**, 1388–1399 (2011).
- 632 96. Svedäng, H. & Wickström, H. Low fat contents in female silver eels: indications of insufficient  
633 energetic stores for migration and gonadal development. *J. Fish Biol.* **50**, 475–486 (2005).

634  
635  
636

### **Supporting information**

637 Appendix S1: miRNAs prediction by miRDeep2.  
638 Appendix S2: List of the 375 conserved and novel miRNA in *A. Anguilla*.  
639 Appendix S3: Table of raw microRNA-seq reads count.  
640 Appendix S4: Potential microRNA target prediction by MiRanda.  
641 Appendix S5: Multidimensional scaling plot of the gene expression profiles obtained with the DESeq2  
642 package for RNA-seq and miRNA-seq data.  
643 Appendix S6: RT-qPCR validation of DEseq2 results.  
644 Appendix S7: Anguilla anguilla's transcriptome annotation and expression analysis.  
645 Appendix S8: Example of down-regulated miRNA targeting transcripts involved in chromatin  
646 modification.

		Arcachon Bay	Gironde Estuary	
<b>Environmental variables</b>	<b>Temperature (°C)</b>	21.9	21.3	
	<b>Salinity (‰)</b>	19.90	7.78	
	<b>O<sub>2</sub> (%)</b>	60.0	86.5	
	<b>O<sub>2</sub> (mg/l)</b>	4.68	7.33	
<b>Condition indices</b>	<b>Fulton</b>	0.156 ± 0.013	0.163 ± 0.014	
	<b>HSI</b>	1.20 ± 0.11	0.85 ± 0.10	*
	<b>OI</b>	5.39 ± 1.42	5.28 ± 0.48	
<b>Tissue composition</b>	<b>Liver proteins (mg/g ww)</b>	138.43 ± 6.41	117.66 ± 4.93	*
	<b>Muscle lipids (% dw)</b>	35.3 ± 5.29	49.76 ± 8.72	
<b>Metals (µg/g dw)</b>	<b>Ag</b>	0.14 ± 0.07	1.16 ± 0.45	**
	<b>As</b>	6.31 ± 1.82	8.40 ± 1.24	*
	<b>Cd</b>	0.05 ± 0.01	11.55 ± 5.29	**
	<b>Cr</b>	1.24 ± 0.88	0.87 ± 0.89	
	<b>Cu</b>	65.07 ± 20.89	95.71 ± 19.90	*
	<b>Ni</b>	0.40 ± 0.25	0.15 ± 0.08	*
	<b>Pb</b>	0.26 ± 0.13	1.68 ± 0.76	**
	<b>Se</b>	13.42 ± 6.54	58.01 ± 9.23	***
	<b>Zn</b>	212.05 ± 51.79	230.11 ± 31.69	
	<b>Hg</b>	0.22 ± 0.08	1.45 ± 0.78	**

647

648 Table 1. Environmental variables, condition indices, tissue composition, and metal concentrations in  
649 the liver (mean ± SE, n=6) of European eels sampled in Arcachon Bay and Gironde Estuary. HSI:  
650 hepato-somatic index, OI: ocular index, ww: wet weight, dw: dry weight. (\*, \*\* and \*\*\* symbols are  
651 Pvalue ≤0.05, ≤0.01 and ≤0.001, respectively)

Sample	Average read size (bp)	N. raw reads	Adaptor dimers (%)	Bases >Q30 (%)	Adaptor-clipped reads (%)	N. clean reads	Mapped reads (%)
AB01	152	26 358 156	2.78	97.18	89.95	22 858 432	78.67
AB02	153	33 128 781	1.90	97.35	91.85	29 568 983	72.95
AB03	154	31 170 977	2.09	97.19	92.32	27 946 975	57.50
AB04	153	23 589 174	2.00	97.17	86.17	19 630 656	72.24
AB05	154	32 383 114	1.45	97.21	86.25	27 176 296	74.61
AB06	154	36 886 852	2.00	97.13	82.04	29 121 827	72.89
GE01	154	27 871 224	2.61	97.42	92.59	24 925 687	87.73
GE02	155	30 141 909	1.13	97.44	94.47	27 999 536	87.45
GE03	155	26 984 661	0.79	97.36	92.49	24 603 024	85.79
GE04	156	32 705 673	0.83	97.41	91.15	29 347 359	87.53
GE05	157	28 617 828	0.70	97.43	91.55	25 872 244	87.14
GE06	156	32 993 005	0.77	97.37	91.95	29 917 020	86.65
<b>Total</b>		<b>362 831 354</b>				<b>318 968 039</b>	

652

653 Table 2: Summary of samples sequenced for discovery of *Anguilla anguilla* miRNA genes. AB, Arcachon Bay; GE, Gironde Estuary.

654

655



miRNA name	FC in GE	n targets	miRNA name	FC in GE	n targets
aan-mir-462	-3.97	2	aan-mir-29a	2.77	2
aan-mir-15e	-3.44	25	aan-scaffold_1	2.69	0
aan-mir-23a	-3.41	1	aan-mir-122	2.34	15
aan-scaffold_951	-3.38	70	aan-scaffold_3557	2.31	8
aan-mir-125a	-2.75	38	aan-mir-203	2.04	1
aan-mir-221	-2.7	17	aan-mir-101a	1.87	0
aan-mir-23b	-2.65	0	aan-mir-101b	1.81	1
aan-scaffold_3084	-2.57	0	aan-mir-194	1.78	2
aan-mir-181c	-2.32	16	aan-mir-139	1.78	19
aan-mir-16b	-2.3	25	aan-mir-19c	1.69	0
aan-mir-16a	-2.29	21	aan-let-7c	1.66	13
aan-mir-454	-2.26	4	aan-mir-15b	1.65	83
aan-mir-150	-2.17	36	aan-mir-17	1.53	10
aan-mir-30b	-2.02	1	aan-mir-21	1.51	2
aan-mir-15d	-1.87	27	aan-mir-152a	1.5	2
aan-mir-92a	-1.85	12	aan-mir-19b	1.49	1
aan-scaffold_11090	-1.77	17	aan-mir-19a	1.42	1
aan-mir-451a	-1.72	0	aan-mir-99	1.34	8
aan-scaffold_8003	-1.7	1	aan-mir-722	1.32	1
aan-mir-30a	-1.68	6			
aan-mir-15c	-1.58	3			
aan-mir-22a	-1.29	27			

656


657 Table 3: Relative expression and number of target transcripts of differentially expressed miRNA  
658 (FDR $\leq$ 0.05) in GE compared to AB groups.

miRNA		Target					
Name	log <sub>2</sub> FC	Name	log <sub>2</sub> FC	Hit	Description	Evalue	Metabolic pathway
aan-let-7c	0.73	contig_15984	-1.78	P31643	Sodium- and chloride-dependent taurine transporter	2e-98	Neurotransmitter transporter
aan-mir-122	1.23	contig_21530	-0.89	Q7YQL6	Oligophrenin-1	0	Membrane trafficking
aan-mir-139	0.84	contig_17347	-1.23	P17453	Bactericidal permeability-increasing protein	3e-59	Immune response
		contig_22852	-1.00	Q7ZXB8	Polyadenylate-binding protein 2-B	1e-71	mRNA surveillance pathway
		contig_9114	-1.53	Q5RDH6	Transferrin receptor protein 1	2e-20	Positive regulation of T and B cell proliferation
aan-mir-15b	0.73	contig_16649	-2.39	P12276	Fatty acid synthase	0	Fatty acid metabolism
		contig_20278	-0.79	P26038	Moesin	0	Tight junction
		contig_35159	-0.93	A1L1F4	Sister chromatid cohesion protein PDS5 homolog A	0	Chromatin organisation
aan-mir-17	0.62	contig_27803	-1.20	Q0P4F7	Acyl-CoA synthetase family member 2, mitochondrial	1e-145	Lipid biosynthesis
aan-scaffold_3557	1.19	contig_35648	-0.69	Q6PEI3	Phosphatase and actin regulator 4	6e-84	Cytoskeleton regulation
aan-mir-125a	-1.46	contig_18202	0.86	Q5ZIK9	Coatomer subunit epsilon	3e-127	Membrane trafficking
aan-mir-150	-1.12	contig_23625	0.93	Q9D9K3	Cell death regulator Aven	6e-23	Cell growth and death
		contig_6617	0.86	Q96A54	Adiponectin receptor protein 1	2e-72	Glucose and lipid metabolism
		contig_15851	1.98	Q8K4T3	STE20-related kinase adapter protein beta	6e-84	Signal transduction
aan-mir-15e	-1.77	contig_39346	0.93	Q9CZV8	F-box/LRR-repeat protein 20	0	Ubiquitin system
aan-mir-16a	-1.20	contig_15454	0.56	Q8I WV8	E3 ubiquitin-protein ligase UBR2	0	Ubiquitin system
aan-mir-16b	-1.20	contig_18374	0.70	P46940	Ras GTPase-activating-like protein IQGAP1	1e-152	Cytoskeleton regulation
		contig_21720	1.52	A5PJN2	ERO1-like protein alpha	0	Protein processing in endoplasmic reticulum
aan-mir-181c	-1.21	contig_22779	0.94	Q7ZTU9	T-box transcription factor TBX2b	0	Signal transduction
aan-mir-221	-1.44	contig_18895	1.65	Q9H2F3	3 beta-hydroxysteroid dehydrogenase type 7	4e-122	Lipid metabolism
aan-mir-22a	-0.37	contig_7998	0.89	Q4R372	F-box only protein 25	1e-170	Signal transduction
aan-mir-30a	-0.75	contig_14191	0.85	P27544	Ceramide synthase 1	9e-112	Lipid metabolism
aan-mir-92a	-0.89	contig_5968	0.99	Q6EDY6	Leucine-rich repeat-containing protein 16A	9e-108	membrane trafficking
aan-scaffold_11090	-0.83	contig_21680	0.75	Q2L969	Metaxin-2	9e-95	Mitochondrial biogenesis
		contig_35030	1.12	Q9QZR5	Homeodomain-interacting protein kinase 2	0	Cell growth and death

aan-scaffold_951	-1.71	contig_14686	0.77	Q9NYF5	Protein FAM13B	2e-135	Signal transduction
		contig_8899	1.42	P60467	Protein transport protein Sec61 subunit beta	5e-21	Secretion system
		contig_16299	1.13	A5PN09	Ubiquitin carboxyl-terminal hydrolase 20	9e-128	Ubiquitin system
		contig_38623	0.76	A2AQ19	RNA polymerase-associated protein RTF1 homolog	6e-167	Transcription machinery


659

660 Table 4: Negative relationship between the relative expression of DE miRNA and their DE targets. Up-regulated and Down-regulated genes (FDR≤0.05) in  
661 GE vs. AB are indicated in red and green, respectively.



662


663 Figure 1: Length distribution of high-quality reads in Illumina libraries. Left panel: Arcachon Bay (AB), Right panel: Gironde Estuary (GE).



664

665 Figure 2: Pie chart showing the distribution of the 820 miRNA sequences predicted by miRDeep2.

666 N=number of sequences in each category.



**Figure 3.** GO enrichment analysis ( $p\text{-value} \leq 0.01$  ;  $\geq 5$  sequences) of the transcript targets of differentially expressed miRNAs (Left panel) and differentially expressed mRNAs (Right panel), **A:** targets of up-regulated miRNAs. **B:** down-regulated mRNAs, **C:** targets of down-regulated miRNAs, **D:** up-regulated mRNAs. X-axis, enrichment score ( $-\log_{10}$  Pvalue) ; Y-axis, Biological processes. The number of sequences in each category is presented on the right of the corresponding bar.

667  
 668 Figure 3: GO enrichment analysis ( $p\text{-value} \leq 0.01$ ;  $\geq 5$  sequences) of the transcript targets of differentially expressed miRNAs (Left panel) and differentially  
 669 expressed mRNAs (Right panel), A: targets of up-regulated miRNAs. B: down-regulated mRNAs, C: targets of down-regulated miRNAs, D: up-regulated  
 670 mRNAs. X-axis, enrichment score ( $-\log_{10}$  Pvalue); Y-axis, Biological processes. The number of sequences in each category is presented on the right of the  
 671 corresponding bar.

