



Toxicity mechanisms of cobalt oxide particles (Co_3O_4 P) on human lung cells: impact of solubilization.

Veronique Malard, Carole Bresson, Chiara Ubaldi, Pascale Delangle, Anne van Der Meeren, David Lemaire, Carine Darolles, Thierry Orsière, Richard Ortega, C. Berthomieu

► To cite this version:

Veronique Malard, Carole Bresson, Chiara Ubaldi, Pascale Delangle, Anne van Der Meeren, et al.. Toxicity mechanisms of cobalt oxide particles (Co_3O_4 P) on human lung cells: impact of solubilization.. BIO Web of Conferences, 2019, 14, pp.05010. 10.1051/bioconf/20191405010 . hal-02355003

HAL Id: hal-02355003

<https://hal.science/hal-02355003>

Submitted on 8 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Toxicity mechanisms of cobalt oxide particles ($\text{Co}_3\text{O}_4\text{P}$) on human lung cells: impact of solubilization

Véronique Malard^{1,*}, Carole Bresson², Chiara Ubaldi³, Pascale Delangle⁴, Anne Van der Meeren⁵, David Lemaire¹, Carine Darolles⁶, Thierry Orsière³, Richard Ortega⁷, and Catherine Berthomieu¹

¹Laboratory of Protein-Metal Interactions (LIPM), Institute of Biosciences and Biotechnologies of Aix Marseille (BIAM), UMR7265 CEA – CNRS - Aix Marseille Univ, CEA Cadarache, F-13108 Saint-Paul-lez-Durance, France

²Den – Service d'Etudes Analytiques et de Réactivité des Surfaces (SEARS), CEA, Université Paris-Saclay, F-91191, Gif sur Yvette, France

³ Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale (IMBE), Aix Marseille Univ, CNRS, IRD, Univ Avignon, Equipe Biomarqueurs, Environnement Santé, Faculté de Médecine Timone, Marseille, France

⁴INAC-SyMMES, Univ. Grenoble Alpes, CEA, CNRS, 38000 Grenoble, France

⁵ Laboratoire de Radiotoxicologie, CEA, Paris-Saclay University, 91297 Arpajon – France

⁶ CEA, DRF, Biosciences and biotechnologies institute (BIAM), Bagnols-sur-Cèze, France

⁷Centre d'Etudes Nucléaires de Bordeaux Gradignan (CENBG), CNRS, IN2P3, Université de Bordeaux, UMR 5797, F-33170 Gradignan, France

* Corresponding author: veronique.malard@cea.fr

The wide use of cobalt in industrial applications, including the nuclear industry, could lead to accidental or chronic occupational exposure mainly by inhalation. Cobalt may be encountered in industry in a variety of chemical forms, including metal dusts, oxides (Co_3O_4), and soluble salts and could also be present in fragments of irradiated fuel. ^{60}Co is an important activation product produced in nuclear power plants. The potential toxicity of poorly soluble cobalt (II, III) oxide particles ($\text{Co}_3\text{O}_4\text{P}$) is a consequence of the long retention time (years) in lungs of the gamma-ray emitting radionuclide. Lung damage may be expected from both chronic radiation exposure, and chemical toxicity of Co.

As a first approach to evaluate the chemical toxicity of cobalt particles, stable $^{59}\text{Co}_3\text{O}_4$ particles were used. We have demonstrated that cobalt oxide particles were readily endocytosed by the BEAS-2B human lung cells via the clathrin-dependent pathway. Despite their very low solubility in the culture medium, cobalt particles were partially solubilized at low pH within lysosomes, leading to the release of cobalt ions. Using ion beam microanalysis, solubilized cobalt was detected within the cytoplasm and the nucleus. As expected with these low-solubility particles, the intracellular solubilized cobalt content was low compared with the intracellular particulate cobalt content in the parts-per-thousand range or below. However, this minute fraction of intracellular solubilized cobalt was found to be responsible for the overall toxicity [1].

The genotoxic potential of poorly soluble $\text{Co}_3\text{O}_4\text{P}$ was then investigated in the BEAS-2B cell line, which exhibits the highest homology in gene expression pattern with primary non

tumor lung cells. Cytokinesis-block micronucleus assay and comet assay revealed that $\text{Co}_3\text{O}_4\text{P}$ had a genotoxic potential following exposure at non cytotoxic concentrations. Indeed, enhanced micronuclei formation and both primary and oxidative DNA damage were noted following exposure to increasing concentrations of $\text{Co}_3\text{O}_4\text{P}$. The scoring of $\gamma\text{-H2Ax}$ foci demonstrated that $\text{Co}_3\text{O}_4\text{P}$ were able to generate double DNA strand breaks. The involvement of oxidative stress was confirmed by the inhibition of foci formation in the presence of the ROS scavenger N-acetylcysteine. Comparison of the effects induced by $\text{Co}_3\text{O}_4\text{P}$ with those triggered by Co^{2+} , indicated that the genotoxicity of $\text{Co}_3\text{O}_4\text{P}$ was not solely due to the released soluble fraction [2].

To go further, a project aiming at broadening the therapeutic offer in case of accidental contamination with insoluble cobalt particles was recently undertaken. This project, dealing with $^{60}\text{Co}_3\text{O}_4$, uses a transversal approach ranging from chemistry (to select ligands able to enhance particle solubilization), to decorporation studies in rodents. Preliminary results obtained in the framework of this collaborative project will be presented.

References

- [1] Ortega R, Bresson C, Darolles C, Gautier C, Roudeau S, Perrin L, Janin M, Floriani M, Aloin V, Carmona A, Malard V. *Low-solubility particles and a Trojan-horse type mechanism of toxicity: the case of cobalt oxide on human lung cells*. Part Fibre Toxicol. 2014; 11 (1) 14.
- [2] Ubaldi, C., Orsiere, T., Darolles, C., Aloin, V., Tassistro, V., George, I., & Malard, V. *Poorly soluble cobalt oxide particles trigger genotoxicity via multiple pathways*. Part Fibre Toxicol. 2016, 13 (1), 5.

Results presented here were obtained in the frame of the EDF-CEA collaboration.