



Treatment-Induced Cardiotoxicity in Breast Cancer: A Review of the Interest of Practicing a Physical Activity

Angeline Ginzac, Judith Passildas, Emilie Gadéa, Catherine Abrial, Ioana Molnar, Romain Trésorier, Martine M. Duclos, Emilie E. Thivat, Xavier Durando

► To cite this version:

Angeline Ginzac, Judith Passildas, Emilie Gadéa, Catherine Abrial, Ioana Molnar, et al.. Treatment-Induced Cardiotoxicity in Breast Cancer: A Review of the Interest of Practicing a Physical Activity. *Oncology*, 2019, 96 (5), pp.223-234. 10.1159/000499383 . hal-02355131

HAL Id: hal-02355131

<https://hal.science/hal-02355131>

Submitted on 8 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**TREATMENT-INDUCED CARDIOTOXICITY IN BREAST CANCER: A REVIEW OF THE INTEREST
OF PRACTISING A PHYSICAL ACTIVITY**

Angeline GINZAC^{1*}, Judith PASSILDAS¹, Emilie GADEA², Catherine ABRIAL¹, Ioana MOLNAR¹,
Romain TRESORIER³, Martine DUCLOS^{4,5}, Emilie THIVAT¹, Xavier DURANDO¹

¹ Université Clermont Auvergne, Centre Jean Perrin, INSERM, U1240 Imagerie Moléculaire et
Stratégies Théranostiques, Clermont-Ferrand, France

² Centre Hospitalier Emile Roux, Le Puy en Velay, France

³ Centre Jean Perrin, Clermont-Ferrand, FR-63000, France

⁴ INRA, UMR 1019, UNH, CRNH Auvergne, Clermont-Ferrand, France

⁵ Centre Hospitalier Universitaire Clermont-Ferrand, Service de Médecine du Sport et des
Explorations Fonctionnelles, CRNH, INRA, Université Clermont Auvergne, Clermont-Ferrand,
France

*CRNH : Centre de Recherche en Nutrition Humaine ; INRA : Institut National de la Recherche
Agronomique ; INSERM : Institut National de la Santé et de la Recherche Médicale ; UNH : Unité de
Nutrition Humaine ; UMR : Unité Mixte de Recherche*

Short title: Physical activity practice during breast cancer treatment

*Corresponding author:

Angeline GINZAC
Centre Jean Perrin
Division de Recherche Clinique
58, rue Montalembert
CLERMONT-FERRAND Cedex 1, Puy-de-Dôme, 63011, France
Tel: 0463663337
Fax: 0473278029
Email: angeline.ginzac@clermont.unicancer.fr

Keywords: HER2-positive breast cancer, cardiotoxicity, anthracycline, trastuzumab, physical activity

ABSTRACT

Physical activity is known to prevent cancer occurrence and to decrease the risk of breast cancer. At diagnosis of breast cancer, fewer than half of the patients reach the international recommendation for physical activity. However, breast cancer patients, and particularly HER2+ breast cancer patients, are exposed to treatment-induced cardiotoxicity because of a side effect of two molecules used in standard therapy to treat these tumors: anthracycline and trastuzumab. Cardiotoxicity can sometimes lead to the treatment being discontinued and even to the development of cardiovascular diseases. Exercise is known to protect the cardiovascular system in the healthy population. Consequently, being physically active during treatment appears as a way to prevent the negative impact of cancer treatment on the heart in this population. In particular, aerobic exercising could provide a protective effect against treatment-induced cardiotoxicity.

A supervised physical activity program seems to be the best way for breast cancer patients to be active during their period of treatment. However, there is very little information available to patients, and in particular a lack of guidelines, on exercising. Interventional trials that have been conducted on this topic are very heterogeneous and no standard recommendations have been made available for cancer patients thus far. An effective physical activity program needs to take each patient's barriers and motivations into account in order to encourage the practice of physical activity throughout treatment. To ensure the success of the program, it is essential to facilitate adherence and especially to maintain motivation. Further studies are needed to determine what practice guidelines oncologists should give their patients.

INTRODUCTION

Physical inactivity is the fourth mortality risk worldwide [1]. Regardless of the level of physical activity, reducing sedentary time decreases mortality, and also the risk of developing cardiovascular diseases, type 2 diabetes, obesity and certain cancers [2]. In 2010, the World Health Organization (WHO) issued international physical activity recommendations for the healthy population, by age group [3]. Over the following years, these recommendations were extended to breast cancer patients through different frameworks and by different authorities, for example, the *Plan Cancer* in France [4] and the American Cancer Society [5,6].

The link between high body mass index (BMI) at breast cancer diagnosis and poor prognosis has been well established, and highlights the importance of being physically active [7,8]. Physical activity from diagnosis has become a routine recommendation in breast cancer treatment, especially because of its many benefits on quality-of-life [9,10]. Physical activity also improves survival rates. Indeed, a meta-analysis has shown that being active after diagnosis is linked to a 34% reduction in breast cancer deaths, a 41 % reduction in deaths from all causes and a 24 % reduction in disease recurrence [11]. Another meta-analysis evaluating 5- and 10-year survival rates showed that the higher the level of physical activity, the better were the survival rates [12]. Indeed, the 5-year survival rate was 97 % for patients who engaged in 3 to 9 or more metabolic equivalent task (MET) hours per week, and 93 % for those who engaged in fewer than 3 MET-h/weeks. The 10-year survival rate of patients who engaged in 9 or more MET-h/week was 92 %, whereas for patients engaging in 3 to 8.9 MET-h/week it was 89 %, and 86 % for those who practiced for fewer than 3 MET-h/week. However, at diagnosis, the majority of patients are insufficiently active [13–15] and maintaining an adequate level of physical activity during treatment can be difficult on account of the numerous barriers encountered by patients. The most frequent barriers identified over the last decades are treatment side effects and the management of everyday routine activities. In addition to improving the outcomes of cancer, physical activity could limit some of the toxicity of anticancer treatments. There is strong evidence that meeting the international physical activity recommendation (i.e. 150 to 300 minutes of moderate-intensity or 75 to 150 minutes of vigorous-intensity aerobic activity on a weekly basis) reduces the incidence and recurrence of cardiometabolic disease [16,17]. It can thus be hypothesized that patients who develop

treatment-induced cardiotoxicity could draw benefit from a physical activity intervention.

In order to promote physical activity during treatment, interventional programs are offered to patients.

Since 2010, the literature on this topic has been developing, and the number of trials involving

physical activity interventions has multiplied. Nevertheless, the question arises as to what is the

optimal program, particularly in terms of duration, type of exercise or intensity.

The aim of this review was to explore treatment-induced cardiotoxicity and identify the interest,

barriers and facilitators of physical activity practice during treatment in order to mitigate cardiac

dysfunction.

I. BREAST CANCER AND CARDIOTOXICITY

1. Anthracycline-induced cardiotoxicity

Anthracyclines belong to the standard chemotherapy treatment for early breast cancer. Nevertheless,

one limitation of this treatment is that it can cause cumulative, dose-dependent cardiac damage, mainly

cardiomyopathy and heart failure [18,19].

Patients treated with anthracycline-based chemotherapy are five times more at risk of developing

cardiotoxicity than those treated without anthracycline [20]. Some meta-analyses have also shown that

the risk of severe cardiotoxicity is higher in patients under anthracycline-based chemotherapy than

among those treated with taxanes without anthracycline [21,22]. Cardiotoxicity is defined by a left

ventricular ejection fraction (LVEF) < 50 % or a decrease of 10 % in the LVEF compared to baseline

[23–26]. It is the first cause of heart failure among women and is responsible for 15-20 % of cases of

treatment discontinuation [27].

Two main hypotheses have been put forward to explain the potential mechanisms implicated in

anthracycline-induced cardiotoxicity. Some authors have suggested that the reactive oxygen species

(ROS) generated by anthracycline treatment cause damage to the DNA and changes in protein and

lipid levels, ultimately leading to cardiomyocyte death [28–30]. It has also been shown that

doxorubicin-induced cardiotoxicity is linked to the activation of p53 and the apoptotic pathway,

leading to a loss of myocardia mass, which contributes to cardiotoxicity [31].

2. *Trastuzumab-induced cardiotoxicity*

Approximately 20 % of breast cancer patients overexpress the human epidermal growth factor receptor 2 (HER2) [32]. This characteristic of the tumour has been associated with poorer disease-free and overall survival [32,33]. One of the standard treatments for this specific cancer is a targeted therapy, trastuzumab (Herceptin®), which can be administered alone [23,34] or in combination with chemotherapy [35,36]. Trastuzumab is a recombinant humanized monoclonal anti-body that binds selectively to subdomain IV of the extracellular ErbB2 domain, which downregulates HER2 expression [37]. Adjuvant treatment with trastuzumab reduces by half the rate of recurrence and by one third the mortality rate [38].

It is estimated that 1 % to 4 % of patients treated with trastuzumab develop heart failure [39,40] and nearly 10 % have asymptomatic LVEF reduction [35,41]. The cumulate incidence rate for major cardiac events has been estimated at 6.6 % for sequential therapy (anthracycline and trastuzumab) and 5.1 % for trastuzumab alone [42]. A meta-analysis has shown that patients treated with trastuzumab were about five times more at risk of developing congestive heart failure in early breast cancer compared to patients not treated with this targeted therapy [43]. Nevertheless, if trastuzumab is discontinued at the time of the first cardiac symptoms, the damage can be reversible [44].

It can be hypothesized that trastuzumab-induced cardiotoxicity is linked to antibody-dependent cellular cytotoxicity (ADCC), leading to an immune-mediated destruction of cardiomyocytes or to a disruption of HER2 signalling, which is essential for myocardial contractility, see for example [44,45]. It is also important to note that HER2, the target of trastuzumab, is expressed in adult cardiomyocytes and is implicated in cardiomyocyte development [45–49]. When trastuzumab blocks HER2 signalling, it prevents the repair mechanisms in cardiomyocytes [50].

3. *Cardiotoxicity associated with anthracyclines and trastuzumab treatment*

When trastuzumab and anthracycline-based chemotherapy are associated, the incidence of cardiotoxicity is 27 % against 13 % when trastuzumab is associated with paclitaxel [51]. More generally, the risk of cardiac failure is higher when trastuzumab is administered after anthracycline-based chemotherapy than after non-anthracycline chemotherapy [22]. Patients treated with

trastuzumab and chemotherapy are twice as likely to develop heart failure or LVEF decrease [52]. Cardiotoxicity can occur acutely but it can also lead to cardiovascular disease development in the long term. A retrospective study compared the incidence of cardiotoxicity in the five years following treatment associating anthracycline and trastuzumab or anthracycline alone. Among the 29.6 % of the patients treated with anthracyclines alone, cardiotoxicity at five years involved 4.3 %, while for the 3.5 % of population who received an association of the two drugs, the proportion was 20.1 %, more than 4.5 times greater [53].

Today, new targeted anti-HER2 therapies are available, such as pertuzumab, trastuzumab emtansine (T-DM1), lapatinib, neratinib and afatinib, but they do not have an indication for early breast cancer. A study was conducted to determine the cardiac tolerance of pertuzumab. The TRYPHAENA study assessed cardiac tolerance of the concomitant or sequential addition of pertuzumab and trastuzumab to docetaxel with or without anthracycline in neoadjuvant treatment [54]. The results did not show an increased incidence of cardiac dysfunction. However, more data is needed to confirm the results of this study.

A recent literature review concluded that the cardiac impact of these novel drugs was lower than that of trastuzumab, and they mainly led to an asymptomatic decrease of the LVEF [55]. The mechanisms responsible for the lower cardiotoxicity of these novel drugs are still not completely understood. However, lapatinib is known to activate the adenosine monophosphate kinase (AMPK) pathway, which could protect heart cells against tumor necrosis factor alpha (TNF- α)-induced cell death by inhibiting its stimulation [45].

II. PHYSICAL ACTIVITY TO PREVENT TREATMENT-RELATED CARDIOTOXICITY IN BREAST CANCER

1. Benefits of being physically active for breast cancer patients

The amount of moderate-to-vigorous physical activity is inversely associated with cardiovascular mortality, and the incidence of cardiovascular disease, stroke and heart failure [16]. Physical activity is also a cornerstone in cardiometabolic chronic diseases such as diabetes or high blood pressure. Among

healthy subjects, regular physical activity reduces the incidence of cardiac events such as hypertension or coronary heart disease [1]. In addition, among breast cancer patients, physical activity can to some extent improve cardiopulmonary function (increased LVEF, improved heart rate and systolic blood pressure, etc.) [56–66], suggesting that physical activity could prevent some of the potential negative effects of treatment-related cardiotoxicity in HER2+ breast cancer subtypes.

A recent literature review highlighted the interest in implementing an exercise intervention program in the population at risk for cardiotoxicity [67], as is the case with HER2+ breast cancer patients, especially because breast cancer patients can also have pre-existing comorbidities, such as diabetes and hypertension, that predispose them to cardiotoxicity [67,68]. The mechanisms explaining the benefit of physical activity on cardiovascular function are still not clearly known. A preclinical study conducted on a murine model showed that physical exercising for 5 consecutive days protected the cardiomyocyte mitochondria from the oxidative damage and dysfunction induced by doxorubicin [69]. Indeed, the authors found that in physical exercising conditions, there was a reduction in doxorubicin-induced ROS release from the cardiac mitochondria. In addition, aerobic training induced endogenous antioxidant mechanisms that protect the heart against ROS [70]. A study conducted on 2973 non-metastatic breast cancer patients assessed the relationship between exposure to physical exercise and cardiovascular events [71]. The results, with a median follow-up of 8.6 years, showed a reduction of 23 % in the risk of cardiovascular events for patients that followed physical activity recommendations (i.e. ≥ 9 MET-h/week.) after the end of their treatment (≥ 9 MET-h/week) compared to those who did not follow the recommendations.

2. Physical activity interventions improving cardiopulmonary capacities

Haykowsky *et al.* were the first to specifically study the impact of physical activity on trastuzumab-induced cardiotoxicity. They conducted a trial on 17 HER2+ breast cancer patients. A supervised physical activity intervention took place in the course of the first 4 months of trastuzumab administration. The intervention was composed of 3 aerobic sessions of 30 to 60 minutes per week on

205 an electrically braked cycle ergometer at 60 to 90% of the oxygen peak consumption (VO_{2peak}). The
206 results of the trial concluded that although the aerobic activity took place during the chemotherapy,
207 trastuzumab induced left ventricular cavity dilation and a significant decrease in the ventricular
208 ejection fraction [72]. Nevertheless, according to the authors, the relatively low program adherence
209 (59 %) could explain the absence of a cardioprotective effect of physical activity. Indeed, they
210 highlighted the fact that exercise adherence was predictive of change in peak oxygen consumption and
211 that patients need to attend ≥ 55 % of the prescribed sessions to derive any benefit on cardiopulmonary
212 function. Other studies assessed the effect of physical activity interventions during treatment on the
213 cardiovascular system and cardiorespiratory function among non-HER2+ breast cancer patients (Table
214 1). The studies were conducted at different moments in treatment: during adjuvant chemotherapy
215 [73,74], during other adjuvant treatments [65–70], or after the end of breast cancer treatment [59].
216 Rates of adherence to the program were higher than in Haykowsky's study, ranging from 66 % to 98.4
217 %, and an improvement in cardiopulmonary function was rapidly observed after the physical activity
218 program was started. For example, after 8 or 12 weeks of an aerobic exercise intervention, an increase
219 in the VO_{2peak} and a decrease in systolic blood pressure were observed [63,66]. Furthermore, patients
220 following a 12-week aerobic intervention enhanced their VO_{2max} by 2.6 ± 3.5 mL/min/kg [60] and
221 there was an increase of 2.37 ± 0.68 mL/min/kg for those who attended 27 weeks of an aerobic and
222 reinforcement program [58]. Overall, randomized controlled clinical trials have shown that patients
223 taking part in a physical activity program have better cardiopulmonary outcomes than those who do
224 not take part [57–60]. Phase II of Hornsby's randomized study was the only one conducted on HER2+
225 patients treated with neoadjuvant chemotherapy (+ aerobic training) [60]. The intervention consisted
226 in a one-to-one supervised cycle ergometry session (3 sessions/week) covering the 12 weeks of
227 neoadjuvant chemotherapy, with progressive intensification of both duration and intensity of the
228 exercise. There was no resistance training offered in this trial. However, cardiopulmonary function
229 improved in the intervention group, unlike the control group, in which it decreased. According to De
230 Luca, a supervised program (24 weeks, 2 sessions of 90 min per week, aerobic and strength training)
231 at least six months after completion of all cancer therapies, improved VO_{2max} by about 38.8 % from
232 baseline to the end of the program [59]. One limitation of these studies was the small numbers of

patients included (N = 20). Studies on larger populations are required. Recently, it has been shown that a 16-week physical activity intervention for physically inactive overweight or obese breast cancer patients after treatment improves physical fitness (N = 91) [61]. The authors highlighted the high adherence rate (96 %) and noted that this could be linked to a reduction in the patients' barriers to physical activity. Indeed, patients were free to go to one-to-one supervised exercise sessions every day of the week from 5 am to 8 pm. Furthermore, parking and bus passes were made available in order to facilitate access to the facility.

3. Ongoing interventional clinical trials for HER2+ breast cancer patients

At present, only two French trials, including our own, are studying the effect of an exercise program on cardiotoxicity among HER2+ patients. CARDAPAC (NCT02433067) is a phase II, prospective, randomized, multicenter study conducted by a research team in Besançon [76]. It aims to evaluate the impact on cardiac function of aerobic exercise among HER2+ patients exclusively treated with trastuzumab. The exercising regime is individualized and consists in 45-minutes of endurance exercises (of moderate and high intensity), 3 times a week for 3 months. The program starts after the end of the chemotherapy and radiotherapy if indicated, and patients are monitored for 6 months. The primary objective of CARDAPAC is to evaluate the incidence rate of cardiotoxicity, defined as a decrease in the LVEF to less than 50 % or an absolute drop in the LVEF of 10 % from baseline.

Our own study, the APACAN2 (NCT02963363) trial, aims to evaluate the feasibility of home-based adapted physical activity for HER2+ breast cancer patients treated with neoadjuvant chemotherapy and anti-HER2 targeted therapy. It is a prospective, non-randomized, interventional study. The physical activity program takes place precisely during the period of the anthracycline-based chemotherapy. It comprises both aerobic and resistance exercises. The main objective of the study is to show the feasibility of a physical activity intervention conducted at home, for patients treated with neoadjuvant chemotherapy and anti-HER2 targeted therapy, in order to comply with the international recommendations defined by the WHO (i.e. at least 150 minutes of moderate-intensity aerobic physical activity or at least 75 minutes of vigorous-intensity activity or an equivalent combination of moderate-to-vigorous intensity activity throughout the week [3]). There is a lack of data concerning

physical activity practice from the beginning of breast cancer treatment in case of neoadjuvant chemotherapy, and this study will allow the feasibility of home-based adapted physical activity to be assessed (aerobic and resistance training) for the first time in this early period of treatment.

According to the data, the need for HER2+ breast cancer patients is to increase or maintain their level of physical activity until they reach the international recommendations. It also enables their sedentary time to be reduced in order to offset treatment-induced cardiotoxicity. The intervention for this patient subgroup needs to be initiated as soon as possible in order to anticipate and prevent any cardiac events. A cardiopulmonary exercise test enables the intensity of physical activity to be adjusted according to each patient's capacities and to ensure that there is no contraindication to exercise.

It is essential to integrate aerobic exercising into the program because of the benefits on the cardiovascular system. It enables cardiopulmonary function, notably VO_{2max} , to be relatively rapidly improved among breast cancer patients (**Table 1**). Nevertheless, muscular strength is also required because increase in muscle mass is linked to a decrease in fat mass, and thus contributes to better body composition [73]. Furthermore, De Luca *et al.* among breast cancer survivors showed that resistance training limited body mass gain, which is often associated with an increase in fat mass and a loss of muscle mass [59]. Indeed, adjuvant breast cancer treatments, such as endocrine therapy, are linked to weight gain mainly corresponding to fat mass gain in the abdominal region [77,78] and physical activity practice could improve these negative long term side effects.

A particularity of the physical activity program is that it can offer support in many forms to encourage practice [79–81]. For example, a pedometer can be given to patients to measure the distance that needs to be reached every week. It can even be a pedometer connected to a smartphone application. Patients could thus see their daily activity and sedentary time, or even monitor their personalized objectives and their progress. The advantage of this tool is that their sports instructor can also see their patient's physical activity and adapt the program according to the patient's physical fitness.

III. BARRIERS AND FACILITATORS TO PRACTICE: HOW CAN PHYSICAL ACTIVITY PROGRAMS BE OPTIMISED?

The success of physical activity interventions lies in the participation of patients. There is therefore a real challenge to identify the barriers and facilitators in order to optimize adherence to a physical activity program.

1. Barriers to exercising

Possible barriers to practice are numerous and can be physical, psychological or even organisational. It is essential to identify these barriers so that they can be overcome for the purpose of improving physical activity practice. Some factors hindering physical activity practice have already been identified.

Barriers linked to disease or treatment side effects are among the most important. Indeed, Courneya *et al.* studied barriers to physical activity in a three-arm randomized trial (aerobic exercise training *versus* resistance exercise training *versus* usual care) and reported that side effects were the reason for more than 50 % of missed supervised exercise sessions on offer in their trial during chemotherapy [82]. Patients mentioned mainly fatigue, nausea, and not feeling well on the day of the chemotherapy administration as barriers to physical activity, since it interfered with their ability to exercise. Nevertheless, it should be considered that before treatment initiation, patients can have comorbidities that already affect their ability to engage in physical activity [83]. Further to this, the notion of self-image should also be considered. Changes in appearance and body image are also barriers to physical activity because patients can be wearing a wig or a prosthesis [84]. This makes the practice of certain activities more complicated, such as swimming for example, because of the close-fitting swimsuits [85]. It adds a further barrier to exercise, especially because of the embarrassment patients might feel in public. Organisational barriers are the second theme among limitations to physical activity practice. Firstly, the distance between home or the workplace and the sports facility frequently appears as a considerable limitation [83,86–88]. What is more, infrastructures are often located in cities, so that access is more difficult for those who live in the countryside. Nevertheless, even for patients living in the city, the risk of encountering traffic problems should also be taken into account as a barrier [86]. When the disease occurs, patients struggle to maintain a balance between work and family life and give more value to the time spent with family and/or friends than doing exercise [84]. They want to

take responsibility for their families as they did before diagnosis, and this can also be a barrier to exercise [88,89]. Patients who did not have time to practice any physical activity before, find it all the more difficult after diagnosis [90,91], especially as a lack of willpower is also a considerable barrier [82,89–91]. Furthermore, being able to continue to go to work is a real need for certain patients, to fight emotional distress and to feel that they are in control. However, this requires a lot of energy and patients are often too tired to engage in a physical activity on top of everything [84]. Furthermore, it has been shown that socioeconomic status also influences physical activity. Indeed, women with a higher level of education are more likely to work full time, and alongside to have a lower moderate exercise level. Thus, employment status could interfere with leisure time and reduce the opportunity to engage in exercising [92].

The psychological aspect is another theme highlighted in certain studies [85–87,93,94]. Overprotection by the patients' entourage has been recognised as a limitation to physical exercise. Furthermore, some patients do not know that they can exercise during treatment or are afraid that it might worsen their symptoms. This raises the problem of the lack of information on physical activity that should be provided by the medical staff [93,94]. And although they may feel uncomfortable exercising in front of healthy people, as mentioned above, patients also report the importance of support to increase their motivation [89,92–94].

It has been shown that the more barriers there are for patients, the lower their level of physical activity [82,95], and also that reducing barriers to exercise is correlated with an increase in physical activity levels. Indeed, Ottenbacher *et al.* showed that patients who reduced their barriers from baseline to 1 year increased their physical activity levels by 91 minutes per week [91]. Facilitating access to training sessions (flexible times, car or bus access, etc.) evidences better adherence rates [61].

2. How can patients be encouraged to be physically active during treatment?

Identifying barriers to physical activity provides an understanding of how to encourage patients to be physically active. First of all, receiving information about the benefits of physical activity during treatment is highly important [83]. Some studies have shown that patients who are aware of the benefits of exercise during treatment are more active than those who do not have this information [96].

Other studies have shown that the influence of the physician proves to be the most decisive incentive among patients to practice a physical activity: by providing reassurance on potential health issues (restoring confidence in their health, reassurance about fear of injury and fatigue), along with the physician's involvement and interest (specific request and monitoring of the patient's physical activity) [97,98]. All these elements point to the importance of information to patients and the training of physicians/oncologists on the effects of physical activity. Finally, a personal background of physical activity is also important, because it is easier for patients to continue a physical activity practice during treatment if they were already active before. For other patients, diagnosis appears as a trigger and a motivational factor that motivates them to take care of themselves [83].

Another key point in promoting physical activity is to personalize the program for each patient. A randomized controlled study, after a one-year follow-up, showed that patients in the tailored-intervention arm adhered better to the diet and exercise recommendations than the control group who were only given standardized diet and exercise brochures [99]. This trial underlines the importance of personalizing the intervention for each patient in order to facilitate enrolment on a training program. It is also essential to define realistic goals for each patient so as to enhance their feeling of self-efficacy [100,101], especially since it has been shown that patients who are less inclined to take part in a physical activity program are also those who feel unable to carry it through [102].

Social support plays a favourable role in physical activity levels. Patients feel comfortable when exercising with other breast cancer women and developing relationships is a source of motivation [85,86].

Lastly, the emergence of mobile applications and connected devices promoting diet and physical activities has upgraded the classic approach to practice. Using new technologies through applications for smartphones or other electronic devices has proved to be a motivation for patients [103]. It has been shown that a physical activity program associated with receiving weekly review emails and access to an e-counsellor for advice on exercise and physical activity is more efficient than a standard physical activity program for improving levels of physical activity among breast cancer patient survivors [104]. Furthermore, new information and communication technologies appear as a good way to establish regular patient follow-up, which is paramount during a physical activity program.

CONCLUSION

As a result of the side effects of the treatment they receive, HER2+ breast cancer patients are particularly exposed to cardiac events during or after therapy. Over the years, the idea has emerged that being active during treatment could present several advantages for patients, and studies have confirmed an improvement in numerous aspects of patient condition during and after treatment (quality of life, asthenia, treatment compliance, cardiac function, etc.). Trials that have been conducted on this issue are very heterogeneous and no standard recommendations are available for HER2+ breast cancer patients. Nevertheless, the importance of practising aerobic exercise is obvious on account of its benefits for the heart. It is essential to facilitate adherence, in particular, so as to maintain patient motivation. The practice of physical activity should be viewed as leisure and integrated into the patient's daily life. Further studies are certainly required to determine how to integrate physical activity practice into HER2+ breast cancer patients' treatment plans. Studies on cost-effectiveness are also needed to assess the medical- and economic consequence of physical activity practice during treatment, which could promote a return to professional activity, thereby reducing the long-term costs.

STATEMENTS

Acknowledgement: None

Statement of ethics: The authors have no ethical conflicts to disclose.

Disclosure of statement: The authors have no conflicts of interest to declare.

Funding sources: Not applicable

Author contributions: AG, ET, EG, XD wrote the original draft of the paper. MD, JP, IM, CA and RT reviewed the paper and suggested draft changes. All authors have read and approved the final version of the manuscript.

REFERENCES

- OMS | Activité physique. WHO [cited 2016 Apr 4];Available from: <http://www.who.int/dietphysicalactivity/pa/fr/>
- Biswas A, Oh PI, Faulkner GE, Bajaj RR, Silver MA, Mitchell MS, et al.: Sedentary time and its association with risk for disease incidence, mortality, and hospitalization in adults: a systematic review and meta-analysis. *Ann Intern Med* 2015;162:123–132.

403 3 WHO: Global Recommendations on Physical Activity for Health. Geneva, World Health
404 Organization, 2010, [cited 2018 May 7].Available from:
405 <http://www.ncbi.nlm.nih.gov/books/NBK305057/>

406 4 INCa: Objectif 8 : Réduire les risques de séquelles et de second cancer - Les 17 objectifs du Plan
407 | Institut National Du Cancer 2014 [cited 2016 May 26];Available from: [http://www.e-](http://www.e-cancer.fr/Plan-cancer/Plan-cancer-2014-2019-priorites-et-objectifs/Les-17-objectifs-du-Plan2/Objectif-8-Reduire-les-risques-de-sequelles-et-de-second-cancer)
408 [cancer.fr/Plan-cancer/Plan-cancer-2014-2019-priorites-et-objectifs/Les-17-objectifs-du-](http://www.e-cancer.fr/Plan-cancer/Plan-cancer-2014-2019-priorites-et-objectifs/Les-17-objectifs-du-Plan2/Objectif-8-Reduire-les-risques-de-sequelles-et-de-second-cancer)
409 [Plan2/Objectif-8-Reduire-les-risques-de-sequelles-et-de-second-cancer](http://www.e-cancer.fr/Plan-cancer/Plan-cancer-2014-2019-priorites-et-objectifs/Les-17-objectifs-du-Plan2/Objectif-8-Reduire-les-risques-de-sequelles-et-de-second-cancer)

410 5 Rock CL, Doyle C, Demark-Wahnefried W, Meyerhardt J, Courneya KS, Schwartz AL, et al.:
411 Nutrition and physical activity guidelines for cancer survivors. *CA Cancer J Clin* 2012;62:243–
412 274.

413 6 Doyle C, Kushi LH, Byers T, Courneya KS, Demark-Wahnefried W, Grant B, et al.: Nutrition and
414 physical activity during and after cancer treatment: an American Cancer Society guide for
415 informed choices. *CA Cancer J Clin* 2006;56:323–353.

416 7 Thivat E, Théronel S, Lapirot O, Abrial C, Gimbergues P, Gadéa E, et al.: Weight change during
417 chemotherapy changes the prognosis in non metastatic breast cancer for the worse. *BMC*
418 *Cancer* 2010;10:648.

419 8 Gadéa É, Thivat É, Wang-Lopez Q, Viala M, Paulon R, Planchat É, et al.: [Poor prognostic value of
420 weight change during chemotherapy in non-metastatic breast cancer patients: causes,
421 mechanisms involved and preventive strategies]. *Bull Cancer (Paris)* 2013;100:865–870.

422 9 Mishra SI, Scherer RW, Geigle PM, Berlanstein DR, Topaloglu O, Gotay CC, et al.: Exercise
423 interventions on health-related quality of life for cancer survivors. *Cochrane Database Syst Rev*
424 2012;CD007566.

425 10 Zhang X, Li Y, Liu D: Effects of exercise on the quality of life in breast cancer patients: a
426 systematic review of randomized controlled trials. *Support Care Cancer Off J Multinatl Assoc*
427 *Support Care Cancer* 2018; DOI: 10.1007/s00520-018-4363-2

428 11 Ibrahim EM, Al-Homaidh A: Physical activity and survival after breast cancer diagnosis: meta-
429 analysis of published studies. *Med Oncol* 2010;28:753–765.

430 12 Holmes MD, Chen WY, Feskanich D, Kroenke CH, Colditz GA: Physical activity and survival after
431 breast cancer diagnosis. *JAMA* 2005;293:2479–2486.

432 13 Blanchard CM, Courneya KS, Stein K, American Cancer Society's SCS-II: Cancer survivors'
433 adherence to lifestyle behavior recommendations and associations with health-related quality
434 of life: results from the American Cancer Society's SCS-II. *J Clin Oncol Off J Am Soc Clin Oncol*
435 2008;26:2198–2204.

436 14 Coups EJ, Ostroff JS: A population-based estimate of the prevalence of behavioral risk factors
437 among adult cancer survivors and noncancer controls. *Prev Med* 2005;40:702–711.

438 15 Eakin EG, Youlden DR, Baade PD, Lawler SP, Reeves MM, Heyworth JS, et al.: Health behaviors
439 of cancer survivors: data from an Australian population-based survey. *Cancer Causes Control*
440 *CCC* 2007;18:881–894.

- 441 16 Arem H, Moore SC, Patel A, Hartge P, Berrington de Gonzalez A, Visvanathan K, et al.: Leisure
442 time physical activity and mortality: a detailed pooled analysis of the dose-response
443 relationship. *JAMA Intern Med* 2015;175:959–967.
- 444 17 Kyu HH, Bachman VF, Alexander LT, Mumford JE, Afshin A, Estep K, et al.: Physical activity and
445 risk of breast cancer, colon cancer, diabetes, ischemic heart disease, and ischemic stroke
446 events: systematic review and dose-response meta-analysis for the Global Burden of Disease
447 Study 2013. *BMJ* 2016;354:i3857.
- 448 18 Yu AF, Jones LW: Breast cancer treatment-associated cardiovascular toxicity and effects of
449 exercise countermeasures. *Cardio-Oncol Lond Engl* 2016;2. DOI: 10.1186/s40959-016-0011-5
- 450 19 Swain SM, Whaley FS, Ewer MS: Congestive heart failure in patients treated with doxorubicin: a
451 retrospective analysis of three trials. *Cancer* 2003;97:2869–2879.
- 452 20 Smith LA, Cornelius VR, Plummer CJ, Levitt G, Verrill M, Canney P, et al.: Cardiotoxicity of
453 anthracycline agents for the treatment of cancer: systematic review and meta-analysis of
454 randomised controlled trials. *BMC Cancer* 2010;10:337.
- 455 21 Mantarro S, Rossi M, Bonifazi M, D'Amico R, Blandizzi C, La Vecchia C, et al.: Risk of severe
456 cardiotoxicity following treatment with trastuzumab: a meta-analysis of randomized and cohort
457 studies of 29,000 women with breast cancer. *Intern Emerg Med* 2016;11:123–140.
- 458 22 Chen T, Xu T, Li Y, Liang C, Chen J, Lu Y, et al.: Risk of cardiac dysfunction with trastuzumab in
459 breast cancer patients: a meta-analysis. *Cancer Treat Rev* 2011;37:312–320.
- 460 23 Smith I, Procter M, Gelber RD, Guillaume S, Feyereislova A, Dowsett M, et al.: 2-year follow-up
461 of trastuzumab after adjuvant chemotherapy in HER2-positive breast cancer: a randomised
462 controlled trial. *Lancet Lond Engl* 2007;369:29–36.
- 463 24 Zamorano Jose Luis, Lancellotti Patrizio, Rodriguez Muñoz Daniel, Aboyans Victor, Asteggiano
464 Riccardo, Galderisi Maurizio, et al.: 2016 ESC Position Paper on cancer treatments and
465 cardiovascular toxicity developed under the auspices of the ESC Committee for Practice
466 Guidelines. *Eur J Heart Fail* 2017;19:9–42.
- 467 25 Procter M, Suter TM, de Azambuja E, Dafni U, van Dooren V, Muehlbauer S, et al.: Longer-term
468 assessment of trastuzumab-related cardiac adverse events in the Herceptin Adjuvant (HERA)
469 trial. *J Clin Oncol Off J Am Soc Clin Oncol* 2010;28:3422–3428.
- 470 26 Piccart-Gebhart MJ, Procter M, Leyland-Jones B, Goldhirsch A, Untch M, Smith I, et al.:
471 Trastuzumab after adjuvant chemotherapy in HER2-positive breast cancer. *N Engl J Med*
472 2005;353:1659–1672.
- 473 27 McArthur HL, Chia S: Cardiotoxicity of trastuzumab in clinical practice. *N Engl J Med*
474 2007;357:94–95.
- 475 28 Chen B, Peng X, Pentassuglia L, Lim CC, Sawyer DB: Molecular and cellular mechanisms of
476 anthracycline cardiotoxicity. *Cardiovasc Toxicol* 2007;7:114–121.
- 477 29 Simůnek T, Stérba M, Popelová O, Adamcová M, Hrdina R, Gersl V: Anthracycline-induced
478 cardiotoxicity: overview of studies examining the roles of oxidative stress and free cellular iron.
479 *Pharmacol Rep PR* 2009;61:154–171.

480 30 Wallace KB: Adriamycin-induced interference with cardiac mitochondrial calcium homeostasis.
481 Cardiovasc Toxicol 2007;7:101–107.

482 31 Zhu W, Soonpaa MH, Chen H, Shen W, Payne RM, Liechty EA, et al.: Acute doxorubicin
483 cardiotoxicity is associated with p53-induced inhibition of the mammalian target of rapamycin
484 pathway. Circulation 2009;119:99–106.

485 32 Slamon DJ, Clark GM, Wong SG, Levin WJ, Ullrich A, McGuire WL: Human breast cancer:
486 correlation of relapse and survival with amplification of the HER-2/neu oncogene. Science
487 1987;235:177–182.

488 33 Cobleigh MA, Vogel CL, Tripathy D, Robert NJ, Scholl S, Fehrenbacher L, et al.: Multinational
489 study of the efficacy and safety of humanized anti-HER2 monoclonal antibody in women who
490 have HER2-overexpressing metastatic breast cancer that has progressed after chemotherapy
491 for metastatic disease. J Clin Oncol Off J Am Soc Clin Oncol 1999;17:2639–2648.

492 34 Baselga J, Carbonell X, Castañeda-Soto N-J, Clemens M, Green M, Harvey V, et al.: Phase II study
493 of efficacy, safety, and pharmacokinetics of trastuzumab monotherapy administered on a 3-
494 weekly schedule. J Clin Oncol Off J Am Soc Clin Oncol 2005;23:2162–2171.

495 35 Slamon DJ, Leyland-Jones B, Shak S, Fuchs H, Paton V, Bajamonde A, et al.: Use of
496 chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that
497 overexpresses HER2. N Engl J Med 2001;344:783–792.

498 36 Marty M, Cognetti F, Maraninchi D, Snyder R, Mauriac L, Tubiana-Hulin M, et al.: Randomized
499 phase II trial of the efficacy and safety of trastuzumab combined with docetaxel in patients with
500 human epidermal growth factor receptor 2-positive metastatic breast cancer administered as
501 first-line treatment: the M77001 study group. J Clin Oncol Off J Am Soc Clin Oncol
502 2005;23:4265–4274.

503 37 Cuello M, Ettenberg SA, Clark AS, Keane MM, Posner RH, Nau MM, et al.: Down-regulation of
504 the erbB-2 receptor by trastuzumab (herceptin) enhances tumor necrosis factor-related
505 apoptosis-inducing ligand-mediated apoptosis in breast and ovarian cancer cell lines that
506 overexpress erbB-2. Cancer Res 2001;61:4892–4900.

507 38 Romond EH, Perez EA, Bryant J, Suman VJ, Geyer CE, Davidson NE, et al.: Trastuzumab plus
508 adjuvant chemotherapy for operable HER2-positive breast cancer. N Engl J Med
509 2005;353:1673–1684.

510 39 Sengupta PP, Northfelt DW, Gentile F, Zamorano JL, Khandheria BK: Trastuzumab-induced
511 cardiotoxicity: heart failure at the crossroads. Mayo Clin Proc 2008;83:197–203.

512 40 Farolfi A, Melegari E, Aquilina M, Scarpi E, Ibrahim T, Maltoni R, et al.: Trastuzumab-induced
513 cardiotoxicity in early breast cancer patients: a retrospective study of possible risk and
514 protective factors. Heart Br Card Soc 2013;99:634–639.

515 41 Fried G, Regev T, Moskovitz M: Trastuzumab-related cardiac events in the treatment of early
516 breast cancer. Breast Cancer Res Treat 2013;142:1–7.

517 42 Thavendiranathan P, Abdel-Qadir H, Fischer HD, Camacho X, Amir E, Austin PC, et al.: Breast
518 Cancer Therapy-Related Cardiac Dysfunction in Adult Women Treated in Routine Clinical
519 Practice: A Population-Based Cohort Study. J Clin Oncol Off J Am Soc Clin Oncol 2016;34:2239–
520 2246.

521 43 Moja L, Tagliabue L, Balduzzi S, Parmelli E, Pistotti V, Guarneri V, et al.: Trastuzumab containing
522 regimens for early breast cancer. *Cochrane Database Syst Rev* 2012;CD006243.

523 44 Albin A, Cesana E, Donatelli F, Cammarota R, Bucci EO, Baravelli M, et al.: Cardio-oncology in
524 targeting the HER receptor family: the puzzle of different cardiotoxicities of HER2 inhibitors.
525 *Future Cardiol* 2011;7:693–704.

526 45 Spector NL, Yarden Y, Smith B, Lyass L, Trusk P, Pry K, et al.: Activation of AMP-activated protein
527 kinase by human EGF receptor 2/EGF receptor tyrosine kinase inhibitor protects cardiac cells.
528 *Proc Natl Acad Sci U S A* 2007;104:10607–10612.

529 46 Fuller SJ, Sivarajah K, Sugden PH: ErbB receptors, their ligands, and the consequences of their
530 activation and inhibition in the myocardium. *J Mol Cell Cardiol* 2008;44:831–854.

531 47 Zhao YY, Sawyer DR, Baliga RR, Opel DJ, Han X, Marchionni MA, et al.: Neuregulins promote
532 survival and growth of cardiac myocytes. Persistence of ErbB2 and ErbB4 expression in neonatal
533 and adult ventricular myocytes. *J Biol Chem* 1998;273:10261–10269.

534 48 Iwamoto R, Yamazaki S, Asakura M, Takashima S, Hasuwa H, Miyado K, et al.: Heparin-binding
535 EGF-like growth factor and ErbB signaling is essential for heart function. *Proc Natl Acad Sci U S*
536 *A* 2003;100:3221–3226.

537 49 De Keulenaer GW, Doggen K, Lemmens K: The vulnerability of the heart as a pluricellular
538 paracrine organ: lessons from unexpected triggers of heart failure in targeted ErbB2 anticancer
539 therapy. *Circ Res* 2010;106:35–46.

540 50 Cote GM, Sawyer DB, Chabner BA: ERBB2 inhibition and heart failure. *N Engl J Med*
541 2013;368:876.

542 51 Seidman A, Hudis C, Pierri MK, Shak S, Paton V, Ashby M, et al.: Cardiac dysfunction in the
543 trastuzumab clinical trials experience. *J Clin Oncol Off J Am Soc Clin Oncol* 2002;20:1215–1221.

544 52 Dahabreh IJ, Linardou H, Siannis F, Fountzilas G, Murray S: Trastuzumab in the adjuvant
545 treatment of early-stage breast cancer: a systematic review and meta-analysis of randomized
546 controlled trials. *The Oncologist* 2008;13:620–630.

547 53 Bowles EJA, Wellman R, Feigelson HS, Onitilo AA, Freedman AN, Delate T, et al.: Risk of heart
548 failure in breast cancer patients after anthracycline and trastuzumab treatment: a retrospective
549 cohort study. *J Natl Cancer Inst* 2012;104:1293–1305.

550 54 Schneeweiss A, Chia S, Hickish T, Harvey V, Eniu A, Hegg R, et al.: Pertuzumab plus trastuzumab
551 in combination with standard neoadjuvant anthracycline-containing and anthracycline-free
552 chemotherapy regimens in patients with HER2-positive early breast cancer: a randomized
553 phase II cardiac safety study (TRYPHAENA). *Ann Oncol Off J Eur Soc Med Oncol* 2013;24:2278–
554 2284.

555 55 Sendur MAN, Aksoy S, Altundag K: Cardiotoxicity of novel HER2-targeted therapies. *Curr Med*
556 *Res Opin* 2013;29:1015–1024.

557 56 Schneider CM, Hsieh CC, Sprod LK, Carter SD, Hayward R: Effects of supervised exercise training
558 on cardiopulmonary function and fatigue in breast cancer survivors during and after treatment.
559 *Cancer* 2007;110:918–925.

560 57 Fairey AS, Courneya KS, Field CJ, Bell GJ, Jones LW, Mackey JR: Effects of exercise training on
561 fasting insulin, insulin resistance, insulin-like growth factors, and insulin-like growth factor
562 binding proteins in postmenopausal breast cancer survivors: a randomized controlled trial.
563 Cancer Epidemiol Biomark Prev Publ Am Assoc Cancer Res Cosponsored Am Soc Prev Oncol
564 2003;12:721–727.

565 58 Cornette T, Vincent F, Mandigout S, Antonini MT, Leobon S, Labrunie A, et al.: Effects of home-
566 based exercise training on VO2 in breast cancer patients under adjuvant or neoadjuvant
567 chemotherapy (SAPA): a randomized controlled trial. Eur J Phys Rehabil Med 2016;52:223–232.

568 59 De Luca V, Minganti C, Borriore P, Grazioli E, Cerulli C, Guerra E, et al.: Effects of concurrent
569 aerobic and strength training on breast cancer survivors: a pilot study. Public Health
570 2016;136:126–132.

571 60 Hornsby WE, Douglas PS, West MJ, Kenjale AA, Lane AR, Schwitzer ER, et al.: Safety and efficacy
572 of aerobic training in operable breast cancer patients receiving neoadjuvant chemotherapy: a
573 phase II randomized trial. Acta Oncol Stockh Swed 2014;53:65–74.

574 61 Dieli-Conwright CM, Courneya KS, Demark-Wahnefried W, Sami N, Lee K, Sweeney FC, et al.:
575 Aerobic and resistance exercise improves physical fitness, bone health, and quality of life in
576 overweight and obese breast cancer survivors: a randomized controlled trial. Breast Cancer Res
577 BCR 2018;20:124.

578 62 Foucaut A-M, Berthouze-Aranda SE, Touillaud M, Kempf-Lépine A-S, Baudinet C, Meyrand R, et
579 al.: Reduction of health risk factors through an adapted physical activity program in patients
580 with breast cancer. Support Care Cancer Off J Multinatl Assoc Support Care Cancer
581 2014;22:1097–1104.

582 63 Kim C-J, Kang D-H, Smith BA, Landers KA: Cardiopulmonary responses and adherence to
583 exercise in women newly diagnosed with breast cancer undergoing adjuvant therapy. Cancer
584 Nurs 2006;29:156–165.

585 64 Kolden GG, Strauman TJ, Ward A, Kuta J, Woods TE, Schneider KL, et al.: A pilot study of group
586 exercise training (GET) for women with primary breast cancer: feasibility and health benefits.
587 Psychooncology 2002;11:447–456.

588 65 Leach HJ, Danyluk JM, Nishimura KC, Culos-Reed SN: Benefits of 24 versus 12 weeks of exercise
589 and wellness programming for women undergoing treatment for breast cancer. Support Care
590 Cancer Off J Multinatl Assoc Support Care Cancer 2016;24:4597–4606.

591 66 Noble M, Russell C, Kraemer L, Sharratt M: UW WELL-FIT: the impact of supervised exercise
592 programs on physical capacity and quality of life in individuals receiving treatment for cancer.
593 Support Care Cancer Off J Multinatl Assoc Support Care Cancer 2012;20:865–873.

594 67 Sturgeon KM, Ky B, Libonati JR, Schmitz KH: The effects of exercise on cardiovascular outcomes
595 before, during, and after treatment for breast cancer. Breast Cancer Res Treat 2014;143:219–
596 226.

597 68 Jones LW, Haykowsky MJ, Swartz JJ, Douglas PS, Mackey JR: Early Breast Cancer Therapy and
598 Cardiovascular Injury. J Am Coll Cardiol 2007;50:1435–1441.

599 69 Kavazis AN, McClung JM, Hood DA, Powers SK: Exercise induces a cardiac mitochondrial
600 phenotype that resists apoptotic stimuli. Am J Physiol Heart Circ Physiol 2008;294:H928–935.

601 70 Ascensão A, Ferreira R, Magalhães J: Exercise-induced cardioprotection--biochemical,
602 morphological and functional evidence in whole tissue and isolated mitochondria. *Int J Cardiol*
603 2007;117:16–30.

604 71 Jones LW, Habel LA, Weltzien E, Castillo A, Gupta D, Kroenke CH, et al.: Exercise and Risk of
605 Cardiovascular Events in Women With Nonmetastatic Breast Cancer. *J Clin Oncol Off J Am Soc*
606 *Clin Oncol* 2016;34:2743–2749.

607 72 Haykowsky MJ, Mackey JR, Thompson RB, Jones LW, Paterson DI: Adjuvant Trastuzumab
608 Induces Ventricular Remodeling Despite Aerobic Exercise Training. *Clin Cancer Res*
609 2009;15:4963–4967.

610 73 Courneya KS, Segal RJ, Mackey JR, Gelmon K, Reid RD, Friedenreich CM, et al.: Effects of aerobic
611 and resistance exercise in breast cancer patients receiving adjuvant chemotherapy: a
612 multicenter randomized controlled trial. *J Clin Oncol Off J Am Soc Clin Oncol* 2007;25:4396–
613 4404.

614 74 Travier N, Velthuis MJ, Steins Bisschop CN, van den Buijs B, Monninkhof EM, Backx F, et al.:
615 Effects of an 18-week exercise programme started early during breast cancer treatment: a
616 randomised controlled trial. *BMC Med* 2015;13:121.

617 75 Courneya KS, McKenzie DC, Mackey JR, Gelmon K, Friedenreich CM, Yasui Y, et al.: Subgroup
618 effects in a randomised trial of different types and doses of exercise during breast cancer
619 chemotherapy. *Br J Cancer* 2014;111:1718–1725.

620 76 Jacquinet Q, Meneveau N, Chatot M, Bonnetain F, Degano B, Bouhaddi M, et al.: A phase 2
621 randomized trial to evaluate the impact of a supervised exercise program on cardiotoxicity at
622 3 months in patients with HER2 overexpressing breast cancer undergoing adjuvant treatment
623 by trastuzumab: design of the CARDAPAC study. *BMC Cancer* 2017;17:425.

624 77 Ginzac A, Émilie T, Marie-Ange M-R, Pascale D-L, Isabelle VP, Judith P, et al.: Weight evolution
625 during endocrine therapy for breast cancer in postmenopausal patients: impact of initial fat
626 mass percentage and previous adjuvant treatments. *Clin Breast Cancer DOI:*
627 10.1016/j.clbc.2018.06.010

628 78 Battisti S, Guida FM, Coppa F, Vaccaro DM, Santini D, Tonini G, et al.: Modification of abdominal
629 fat distribution after aromatase inhibitor therapy in breast cancer patients visualized using 3-D
630 computed tomography volumetry. *Clin Breast Cancer* 2014;14:365–370.

631 79 Zhu J, Ebert L, Liu X, Chan SW-C: A mobile application of breast cancer e-support program
632 versus routine Care in the treatment of Chinese women with breast cancer undergoing
633 chemotherapy: study protocol for a randomized controlled trial. *BMC Cancer* 2017;17:291.

634 80 Coughlin SS, Besenyi GM, Bowen D, De Leo G: Development of the Physical activity and Your
635 Nutrition for Cancer (PYNC) smartphone app for preventing breast cancer in women. *mHealth*
636 2017;3:5.

637 81 Cheong IY, An SY, Cha WC, Rha MY, Kim ST, Chang DK, et al.: Efficacy of Mobile Health Care
638 Application and Wearable Device in Improvement of Physical Performance in Colorectal Cancer
639 Patients Undergoing Chemotherapy. *Clin Colorectal Cancer* 2018;17:e353–e362.

- 640 82 Courneya KS, McKenzie DC, Reid RD, Mackey JR, Gelmon K, Friedenreich CM, et al.: Barriers to
641 supervised exercise training in a randomized controlled trial of breast cancer patients receiving
642 chemotherapy. *Ann Behav Med Publ Soc Behav Med* 2008;35:116–122.
- 643 83 Henriksson A, Arving C, Johansson B, Igelström H, Nordin K: Perceived barriers to and
644 facilitators of being physically active during adjuvant cancer treatment. *Patient Educ Couns*
645 2016;99:1220–1226.
- 646 84 Husebø AML, Karlsen B, Allan H, Søreide JA, Bru E: Factors perceived to influence exercise
647 adherence in women with breast cancer participating in an exercise programme during
648 adjuvant chemotherapy: a focus group study. *J Clin Nurs* 2015;24:500–510.
- 649 85 Emslie C, Whyte F, Campbell A, Mutrie N, Lee L, Ritchie D, et al.: “I wouldn’t have been
650 interested in just sitting round a table talking about cancer”; exploring the experiences of
651 women with breast cancer in a group exercise trial. *Health Educ Res* 2007;22:827–838.
- 652 86 Wurz A, St-Aubin A, Brunet J: Breast cancer survivors’ barriers and motives for participating in a
653 group-based physical activity program offered in the community. *Support Care Cancer Off J*
654 *Multinatl Assoc Support Care Cancer* 2015;23:2407–2416.
- 655 87 Heffernan K, Murphy H, McLeod J, Mutrie N, Campbell A: Understanding barriers to exercise
656 implementation 5-year post-breast cancer diagnosis: a large-scale qualitative study. *Health*
657 *Educ Res* 2013;28:843–856.
- 658 88 Loh SY, Lee SY, Quek KF, Murray L: Barriers to participation in a randomized controlled trial of
659 Qigong exercises amongst cancer survivors: lessons learnt. *Asian Pac J Cancer Prev APJCP*
660 2012;13:6337–6342.
- 661 89 Charlier C, Van Hoof E, Pauwels E, Lechner L, Spittaels H, De Bourdeaudhuij I: The contribution
662 of general and cancer-related variables in explaining physical activity in a breast cancer
663 population 3 weeks to 6 months post-treatment. *Psychooncology* 2013;22:203–211.
- 664 90 Loh SY, Chew S-L, Lee S-Y: Barriers to exercise: perspectives from multiethnic cancer survivors
665 in Malaysia. *Asian Pac J Cancer Prev APJCP* 2011;12:1483–1488.
- 666 91 Ottenbacher AJ, Day RS, Taylor WC, Sharma SV, Sloane R, Snyder DC, et al.: Exercise among
667 breast and prostate cancer survivors--what are their barriers? *J Cancer Surviv Res Pract*
668 2011;5:413–419.
- 669 92 Steinhilber L, Geyer S, Sperlich S: Health behavior change among breast cancer patients. *Int J*
670 *Public Health* 2013;58:603–613.
- 671 93 Rogers LQ, Matevey C, Hopkins-Price P, Shah P, Dunnington G, Courneya KS: Exploring social
672 cognitive theory constructs for promoting exercise among breast cancer patients. *Cancer Nurs*
673 2004;27:462–473.
- 674 94 Browall M, Mijwel S, Rundqvist H, Wengström Y: Physical Activity During and After Adjuvant
675 Treatment for Breast Cancer: An Integrative Review of Women’s Experiences. *Integr Cancer*
676 *Ther* 2018;17:16–30.
- 677 95 Perna FM, Craft L, Carver CS, Antoni MH: Negative affect and barriers to exercise among early
678 stage breast cancer patients. *Health Psychol Off J Div Health Psychol Am Psychol Assoc*
679 2008;27:275–279.

- 680 96 Jones LW, Courneya KS, Fairey AS, Mackey JR: Effects of an oncologist's recommendation to
681 exercise on self-reported exercise behavior in newly diagnosed breast cancer survivors: a
682 single-blind, randomized controlled trial. *Ann Behav Med Publ Soc Behav Med* 2004;28:105–
683 113.
- 684 97 Lanhers C, Duclos M, Guttman A, Coudeyre E, Pereira B, Ouchchane L: General Practitioners'
685 Barriers to Prescribe Physical Activity: The Dark Side of the Cluster Effects on the Physical
686 Activity of Their Type 2 Diabetes Patients. *PloS One* 2015;10:e0140429.
- 687 98 Duclos M, Dejager S, Postel-Vinay N, di Nicola S, Quéré S, Fiquet B: Physical activity in patients
688 with type 2 diabetes and hypertension--insights into motivations and barriers from the MOBILE
689 study. *Vasc Health Risk Manag* 2015;11:361–371.
- 690 99 Demark-Wahnefried W, Clipp EC, Lipkus IM, Lobach D, Snyder DC, Sloane R, et al.: Main
691 Outcomes of the FRESH START Trial: A Sequentially Tailored, Diet and Exercise Mailed Print
692 Intervention Among Breast and Prostate Cancer Survivors. *J Clin Oncol* 2007;25:2709–2718.
- 693 100 Pinto BM, Frierson GM, Rabin C, Trunzo JJ, Marcus BH: Home-based physical activity
694 intervention for breast cancer patients. *J Clin Oncol Off J Am Soc Clin Oncol* 2005;23:3577–
695 3587.
- 696 101 Demark-Wahnefried W, Morey MC, Clipp EC, Pieper CF, Snyder DC, Sloane R, et al.: Leading the
697 Way in Exercise and Diet (Project LEAD): intervening to improve function among older breast
698 and prostate cancer survivors. *Control Clin Trials* 2003;24:206–223.
- 699 102 Rabin C, Simpson N, Morrow K, Pinto B: Behavioral and psychosocial program needs of young
700 adult cancer survivors. *Qual Health Res* 2011;21:796–806.
- 701 103 Goode AD, Lawler SP, Brakenridge CL, Reeves MM, Eakin EG: Telephone, print, and Web-based
702 interventions for physical activity, diet, and weight control among cancer survivors: a
703 systematic review. *J Cancer Surviv Res Pract* 2015;9:660–682.
- 704 104 Hatchett A, Hallam JS, Ford MA: Evaluation of a social cognitive theory-based email
705 intervention designed to influence the physical activity of survivors of breast cancer.
706 *Psychooncology* 2013;22:829–836.

707

708

709 **FIGURE LEGENDS**

710 Table 1: Physical activity interventions reporting cardiopulmonary outcomes among breast cancer
711 patients during and after treatment

712 *Adj: Adjuvant; AET: aerobic exercise training; CT: chemotherapy; Ctrl: Control; HR_{max}: maximum*
713 *heart rate; HT: Hormone therapy; NA: Not applicable; NACT: Neoadjuvant chemotherapy; ND: Not*
714 *done; RCT: Radiochemotherapy; RET: Resistance exercise training; RM: repetition maximum; RPE:*
715 *Rate of perceived exertion; RT: Radiotherapy; UC: usual care; VO_{2max}: maximum peak oxygen*
716 *consumption*

STUDY	N	TREATMENT	MODALITY	PROGRAM DURATION	FREQUENCY	ACTIVITY	INTENSITY	ADHERENCE	IMPROVEMENT
Courneya KS, 2007	82	Adj CT	NA	NA	NA	UC	NA	NA	NA
	82		Group (supervised)	17 weeks (± 4 months)	3 sessions/week	RET	60% to 70% of their estimated one repetition maximum	68%	↑ Body strength ↑ Lean body mass
	78					AET	From 60 to 80% of VO_{2max}	72%	↑ Aerobic fitness
Fairey AS, 2003	25 Intervention group)	Completed surgery, RT and/or CT with or without current HT use	Group	15 weeks	3 sessions/week	AET	70-75% of VO_{2max}	98.4%	Between group difference for VO_{2max} : + 0.29 L/min (p<0.001)
	28 (Ctrl group)		No training	NA	NA	NA	NA	NA	
Cornette T, 2016	22 (Intervention group)	NACT or Adj CT and RT	At home	27 weeks	2 sessions/week 1 session/week	AET RET	> 3 METs	88%	VO_{2max} : + 1.83 \pm 0.68 ml/min/kg
	22 (Ctrl group)		NA	NA	NA	UC	NA	NA	VO_{2max} : - 1.31 \pm 0.65 mL/min/kg

De Luca V, 2016	10 (Intervention group)	Surgery + all cancer therapies completed at least 6 months before	Group (supervised)	24 weeks	2 sessions/week	AET RET	70% to 80% of estimated HR _{max} A load of 40% to 60% of estimated 1RM	ND	↑ Vo _{2max} + 38.8 % between baseline and end of intervention
	10 (Ctrl group)		NA	NA	NA	NA	NA	NA	No significant change
Hornsby, 2014	10 (Intervention group)	NACT	One-to-one supervised sessions	12 weeks	2 to 3 sessions/week	AET	Moderate to high intensity	66%	VO _{2max} : + 2.6 ± 3.5 mL/min/kg
	10 (Ctrl group)	NA	NA	NA	NA	UC	NA	NZ	VO _{2max} : - 1.5 ± 2.2 mL/min/kg
Travier, 2015	102 (Intervention group)	Adj CT	Group	18 weeks	2 sessions/week	AET RET	Based on the heart rate at the ventilatory threshold as determined during baseline cardiopulmonary exercise test	83%	VO _{2max} : - 0.7 mL/min/kg
	102 (Ctrl group)		NA	NA	NA	UC	NA	NA	NA
Dieli- Conwright CM, 2018	46 (Intervention group)	< 6 months post-treatment (chemo or radiotherapy)	One-to-one supervised sessions	16 weeks	3 sessions/week (2 AET+RET and 1 AET)	AET RET	Moderate-vigorous intensity	96 %	↑ Vo _{2max} ↓ Resting heart rate

	45 (Control group)		NA	NA	NA	NA	NA	NA	No change in physical fitness
Foucaut AM, 2014	61	Adj RCT	Group	3 months	1 session/week 1 session/week	AET RET	Moderate to vigorous	80%	VO _{2max} : + 0;6 ml/min/kg
Kim CJ, 2006	41	Adj therapy	Group	8 weeks	2 sessions/week	AET RET	Moderate	78.3%	↑ Vo _{2peak}
Kolden, 2012	40	Adj therapy	Group	16 weeks	3 sessions/week	AET RET Flexibility	40-60% of estimated maximal aerobic capacity	88%	↑ aerobic capacity
Leach, 2016	102	currently treated by CT or CT/RT discontinued within 3 months	Group or home-based exercise according to patient's preference	24 weeks	2 sessions/week 1 session/week 5-7 sessions/week	AET RET Flexibility	Mild to vigorous	80.6% (12 week assessment) 42% (24 week assessment)	VO _{2max} : + 1.9 mL/min/kg
Noble, 2012	575	Surgery or CT or RT or HT	Group	12 weeks	2 sessions/week	AET RET Stretching	RPE 11-13	69%	↓ heart rate response ↓ systolic blood pressure ↓ RPE