



Cold Water Immersion Enhanced Athletes' Wellness and 10-m Short Sprint Performance 24-h After a Simulated Mixed Martial Arts Combat

Montassar Tabben, Mohammed Ihsan, Nihel Ghoul, Jeremy Coquart, Anis Chaouachi, Helmi Chaabene, Claire Tourny, Karim Chamari

► To cite this version:

Montassar Tabben, Mohammed Ihsan, Nihel Ghoul, Jeremy Coquart, Anis Chaouachi, et al.. Cold Water Immersion Enhanced Athletes' Wellness and 10-m Short Sprint Performance 24-h After a Simulated Mixed Martial Arts Combat. *Frontiers in Physiology*, 2018, 9, 10.3389/fphys.2018.01542 . hal-02355413

HAL Id: hal-02355413

<https://hal.science/hal-02355413>

Submitted on 14 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Cold Water Immersion Enhanced Athletes' Wellness and 10-m Short Sprint Performance 24-h After a Simulated Mixed Martial Arts Combat

Montassar Tabben^{1*}, Mohammed Ihsan¹, Nihel Ghoul², Jeremy Coquart², Anis Chaouachi^{3,4,5}, Helmi Chaabene^{6,7}, Claire Tourny² and Karim Chamari¹

¹ ASPETAR, Orthopaedic and Sports Medicine Hospital, Doha, Qatar, ² UFR STAPS, CETAPS, Normandie University – University of Rouen, Rouen, France, ³ Research Laboratory "Sport Performance Optimization," National Centre of Medicine and Science in Sport, Tunis, Tunisia, ⁴ Sports Performance Research Institute, AUT University, Auckland, New Zealand, ⁵ PVF Football Academy, Hang Yen, Vietnam, ⁶ Division of Training and Movement Sciences, Research Focus Cognition Sciences, University of Potsdam, Potsdam, Germany, ⁷ High Institute of Sports and Physical Education Kef, University of Jendouba, Tunis, Tunisia

OPEN ACCESS

Edited by:

Robert James Aughey,
Victoria University, Australia

Reviewed by:

David Andrew Low,
Liverpool John Moores University,
United Kingdom
Giovanni Messina,
University of Foggia, Italy

*Correspondence:

Montassar Tabben
Montassar.Tabben@aspetar.com

Specialty section:

This article was submitted to
Exercise Physiology,
a section of the journal
Frontiers in Physiology

Received: 27 April 2018

Accepted: 15 October 2018

Published: 01 November 2018

Citation:

Tabben M, Ihsan M, Ghoul N, Coquart J, Chaouachi A, Chaabene H, Tourny C and Chamari K (2018) Cold Water Immersion Enhanced Athletes' Wellness and 10-m Short Sprint Performance 24-h After a Simulated Mixed Martial Arts Combat. *Front. Physiol.* 9:1542. doi: 10.3389/fphys.2018.01542

Objective: The aim of the present study was to examine the effect of Cold Water Immersion (CWI) on the recovery of physical performance, hematological stress markers and perceived wellness (i.e., Hooper scores) following a simulated Mixed Martial Arts (MMA) competition.

Methods: Participants completed two experimental sessions in a counter-balanced order (CWI or passive recovery for control condition: CON), after a simulated MMA competition (3 × 5-min MMA rounds separated by 1-min of passive rest). During CWI, athletes were required to submerge their bodies, except the trunk, neck and head, in the seated position in a temperature-controlled bath (~10°C) for 15-min. During CON, athletes were required to be in a seated position for 15-min in same room ambient temperature. Venous blood samples (creatinine kinase, cortisol, and testosterone concentrations) were collected at rest (PRE-EX, i.e., before MMA), immediately following MMA (POST-EX), immediately following recovery (POST-R) and 24 h post MMA (POST-24), whilst physical fitness (squat jump, countermovement-jump and 5- and 10-m sprints) and perceptual measures (well-being Hooper index: fatigue, stress, delayed onset muscle soreness (DOMS), and sleep) were collected at PRE-EX, POST-R and POST-24, and at PRE-EX and POST-24, respectively.

Results: The main results indicate that POST-R sprint (5- and 10-m) performances were 'likely to very likely' ($d = 0.64$ and 0.65) impaired by prior CWI. However, moderate improvements were in 10-m sprint performance were 'likely' evident at POST-24 after CWI compared with CON ($d = 0.53$). Additionally, the use of CWI 'almost certainly' resulted in a large overall improvement in Hooper scores ($d = 1.93$). Specifically, CWI 'almost certainly' resulted in improved sleep quality ($d = 1.36$), stress ($d = 1.56$) and perceived fatigue ($d = 1.51$), and 'likely' resulted in a moderate decrease in DOMS ($d = 0.60$).

Conclusion: The use of CWI resulted in an enhanced recovery of 10-m sprint performance, as well as improved perceived wellness 24-h following simulated MMA competition.

Keywords: recovery, combat sports, MMA, delayed onset muscle soreness, stress, fatigue

INTRODUCTION

Mixed Martial Arts (MMA) is characterized by a combination of fighting styles inspired from other combat sports, resulting in an attractive conglomerate of fighting techniques, i.e., strikes, clinch and grappling, and submission (Amtmann et al., 2008; del Vecchio et al., 2011; La Bounty et al., 2011). Following massive media exposure, particularly with the Ultimate Fighting Championship® (UFC) events, its worldwide popularity has grown exponentially, resulting in MMA being one of the most increasingly participated sport at amateur, right through to professional levels (Ghoul et al., 2017). MMA combat is highly intensive in nature, and has previously shown to induce significant fatigue and muscle damage, which was largely persistent throughout a 24-h period. (Amtmann et al., 2008; Ghoul et al., 2017). Results showed that simulated MMA competition induced high HR, blood lactate and RPE (181.3 bpm, 13.6 mmol/L, and 6.9 – scale of 10, respectively) (Ghoul et al., 2017). Moreover, a single MMA combat or training session has been shown to increase salivary and blood cortisol levels, as well as urinary neopterin content, indicating profound physiological stress (Ghoul et al., 2017; Lindsay et al., 2017). As such, it is essential that appropriate recovery strategies are undertaken, to ensure that training and/or completion demands are adequately met.

Cold water immersion (CWI) is a recovery modality typically undertaken following endurance (Vaile et al., 2008) and/or team-sport activities (Rowell et al., 2011). More recently, the use of CWI as a post-exercise recovery strategy has gained considerable popularity within combat sports (Lindsay et al., 2015, 2017). Indeed, CWI seems effective to promote recovery by ameliorating effects of exercise-induced muscle damage (EIMD) and reducing muscle oedema via several mechanisms associated with localized cooling, hydrostatic pressure and the redistribution of blood flow (Ihsan et al., 2016). Moreover, CWI has been purported to confer analgesic effects, which has shown to reduce delayed onset muscle soreness (DOMS), and consequently improve perceptual recovery (Corbett et al., 2011). Recently, there has been increased interest in the efficacy of this recovery modality following MMA-specific activity. For instance, Lindsay et al. (2017) demonstrated reduced inflammation and oxidative stress following an intense MMA training session when CWI was incorporated as a recovery intervention. However, further research investigating the effects of CWI on the recovery of physical performance following simulated combat are warranted. Indeed, given that MMA tournaments are highly demanding in nature, with requirements to compete every 1–2 day(s), the recovery of physical performance should be a crucial outcome measure. Moreover, given that perceptual recovery has been regarded as an important component of recovery, and has

shown to be associated with changes in physical performances (Ihsan et al., 2017), it is also imperative to examine whether CWI would potentially be an effective modality to improve perceptual recovery within MMA-specific context. As such, the current study aimed to understand how physiological and perceptual outcomes are simultaneously affected by recovery CWI amongst MMA fighters. The primary purpose of this investigation was to examine the immediate and post 24-h effect of CWI and passive recovery on physical performance following simulated MMA competition. Secondly, we examined changes in the hematological stress markers (creatinine kinase, cortisol, and testosterone) associated with recovery. Finally, we studied the perceptual effects of CWI by including wellness measures (i.e., Hooper scores). It was hypothesized that compared to passive recovery, CWI would allow for a better recovery in performance, perceptual, and physiological measures.

MATERIALS AND METHODS

Participants

Twelve male well-trained MMA athletes (mean \pm standard deviation for age: 26.5 ± 5.0 years, height: 182.0 ± 7.4 cm, and body mass: 86.2 ± 10.9 kg) participated in this study. All participants competed in local or regional competitions once per month and were regularly training (technical, tactical, and/or physical sessions) 3–4 times per week. Participants were informed about the experimental procedures, potential risks and benefits, and signed an informed consent document before the commencement of the study. All participants were instructed to avoid drinking coffee, tea, cola, or alcohol 24 h preceding the experimental session. All testing procedures were in line with the latest Declaration of Helsinki and the protocol was approved by the institutional IRB-board Ethics committee.

Experimental Design

One week before the study commencement, the participants undertook a familiarization session where they accustomed to all experimental procedures involved. In a randomized, cross-over manner (repeated measures), participants completed two experimental sessions, conducted 72 h apart. The experimental sessions were conducted at the same time of day (8–10 am), where temperature and humidity were consistent across the sessions (temperature ~ 29 – 30°C ; relative humidity ~ 38 – 40%).

The experimental protocol involved four testing time-points (**Figure 1**), during which physical performance measures and blood sampling was undertaken (described below); (i) prior to simulated MMA combat (MMAs) (PRE-EX), (ii) immediately following MMAs (POST-EX), (iii) immediately

following recovery (POST-R), and (iv) 24 h post MMAs (POST-24). Shortly after arriving on day 1, height was assessed to the nearest 0.1 cm using a stadiometer (Holtain Ltd., Crymch, United Kingdom) and body mass was measured to the nearest 0.1 kg using a body composition monitor Tanita BC 545-N (Tanita Health Equipment, Hong Kong, China). Following the completion of a 15-min standardized warm-up and PRE-EX measurements, athletes performed the MMAs protocol. The MMAs consists of 3 × 5-min MMA rounds separated by 1-min of passive rest. They were divided into pairs with a difference of body mass of no more than 10%. All combats were judged by qualified MMA referees according to the official rules. Additionally, as during an official competition, athletes' coaches were present and they consistently provided guidance, technical advice, and verbal encouragement. No interruption of the combats was permitted even when submissions occurred to ensure the continuity of the fights. Following the completion of the MMAs, POST-EX was undertaken. Three minutes later, athletes performed, one of the following recovery modalities for 15-min in a randomized, counter-balanced manner: CWI or passive recovery (i.e., control modality: CON). Testing measures were immediately undertaken following recovery intervention (i.e., POST-R) and POST-24 was undertaken 24 h following MMAs.

Recovery Modalities

- (1) CWI: Athletes were required to submerge their bodies, except the trunk, neck and head, in the seated position in a temperature-controlled bath (~10°C) for 15-min. (Lindsay et al., 2017).
- (2) Passive recovery (i.e., control modality: CON): Athletes were required to be in a seated position for 15-min with minimal movement at the same room ambient temperature. (Lindsay et al., 2017).

Experimental Measures

Heart Rate Monitoring

Heart rate was continuously recorded using a chest monitor during the sparing bouts and the subsequent rest phase. Recorded data were downloaded on a computer using Polar software (Polar Precision Performance SW5.20, Polar Electro, Kempele, Finland). All irregular heartbeats were automatically identified and replaced with interpolated adjacent R-R interval values with the Polar Software. (Nunan, 2006)

Blood Lactate Concentration

Blood lactate concentration was measured at PRE-EX, POST-EX and POST-24. A fingertip capillary blood sample (5 µL) was collected and analyzed for lactate concentration using a calibrated Lactate Pro analyzer (Arkray®, Tokyo, Japan).

Perceptual Responses

The well-being Hooper index based on ratings relative to (i) fatigue, (ii) stress, (iii) whole-body DOMS, and (iv) sleep has been used as a cost-effective strategy for prevention and early detection of non-functional overreaching/overtraining syndromes (Hooper et al., 1995). The participants were asked to

subjectively rate the quality of their prior night-sleep, their fatigue level, stress, and DOMS on a scale of 1–7 before each testing day. 1 was anchored as the positive and 7 the negative end of the continuum for all perceptual variables.

Hooper scores were analyzed as the sum of the four scores as well as for each of the four variables separately. (Hooper et al., 1995).

Physical Performances

Squat jump (SJ) and countermovement jump (CMJ)

SJ and CMJ performances were undertaken to determine athletes' explosive power (de Freitas et al., 2017). Jumps were assessed using an infrared jump system (Optojump, Microgate®, Bolzano, Italy) interfaced with a microcomputer. Participants were not allowed to use their arms, which were positioned on their hips throughout the jumps. The SJ was completed first followed by the CMJ, with two attempts allowed for each jump type. A 30-s passive recovery was imposed between each jump, which allowed the tester to record vertical jump performance, and to reset the systems for recording of the next trial. A third trial was completed if jump heights varied > 5%. The highest score relative to each type of jump was retained for further analysis.

10-m sprint time

The 10-m sprint assessment was undertaken 2 min following the completion of the jumps tests, and was performed on an outdoor synthetic 20-m sprint court. Participants started in standing position with the toe of the preferred foot positioned forward behind a starting line (i.e., 0.5 m). Time was taken automatically when the athlete voluntarily initiated the test. Sprint times were recorded using telemetric photoelectric cells (Racetime 2, Microgate®, Bolzano, Italy; accuracy of 0.01 s) placed 0.7 m above the ground. Three attempts were given with 3-min walk-back recovery in-between attempts and the run. The fastest 10-m time (and corresponding 5-m split time) was selected for analysis.

Blood sampling and analysis

Venous blood samples (8 mL) were obtained from an antecubital vein in a rested and seated position at PRE-EX, POST-EX, POST-R, and POST-24 to determine serum creatine kinase (CK), cortisol, and testosterone concentrations. Blood was collected into EDTA containing tubes and serum and plasma samples were obtained by centrifuging (3,000 rpm for 15 min at 4°C) and were stored at –80°C until analysis. All analyses were completed 1 week following the experiment, and performed by a qualified medical personal. All analyzers were calibrated according to the manufacturers' instructions prior to use. Plasma creatine kinase (CK) activity was used as a marker of muscle damage and was measured spectrophotometrically in an Advia 1800 analyzer (Siemens Healthcare, Erlangen, Germany). Testosterone and cortisol levels were determined by using a solid-phase radioimmunoassay procedure (Immunotech RIA, Beckman Coulter®, Brea, United States). The intra-assay coefficients of variation (CVs) for each measurement were established at 3.0% for the CK, 1.6% for Hb and Ht, 2.1% for Cortisol, and 4.3 % for testosterone.

Statistical analysis

Data was analyzed using effect sizes and magnitude-based inferences, where within-group and between-group comparisons were undertaken using specifically designed Excel spreadsheets (Hopkins, 2006; Hopkins et al., 2009). These spreadsheets calculated the standardized differences or effect sizes (d) with 90% confidence limits (90% CL), and d values of 0.2, <0.5 , and ≥ 0.8 were interpreted as small, moderate and large effects, respectively (Cohen, 1988). Magnitude of differences were considered substantial if there was at least a 75% likelihood that the effect was greater than the smallest worthwhile change (i.e., $d = 0.2$), and qualitatively evaluated as 75–<95%: likely; ≥ 95 –99%: very likely; and ≥ 99 –100%: almost certain. If the chance of substantially higher or lower differences were both $>5\%$, the true difference was assessed as ‘unclear’ (Hopkins et al., 2009).

RESULTS

The intensity of the simulated combat was consistent across conditions, as HR (CON: 186 ± 15 vs. CWI: 183 ± 15 bpm, $d = 0.2$) and $[La^-]$ following the combat bouts (CON: 13.3 ± 1.7 vs. CWI: 13.1 ± 1.3 mmol·L⁻¹, $d = 0.17$) were similar between CON and CWI.

Changes in performance for the 5-m and 10-m sprint, SJ and CMJ are presented in **Figure 1**. CWI resulted in slower 5-m ($d = 0.64$) and 10-m ($d = 0.91$) sprint time compared with CON at POST-R. In contrast, CWI resulted in improved 10-m sprint performance ($d = 0.53$) at POST-24 compared with CON. Compared with PRE-EX, small to moderate decreases in SJ and CMJ were observed at POST-R both CON and CWI, with no differences (i.e., unclear) between conditions at all time-points (**Figure 1C**).

Changes in CK, cortisol, testosterone and cortisol/testosterone ratio are presented in **Figure 2**. The simulated combat resulted in small to large increases in cortisol, testosterone and cortisol/testosterone ratio during the POST-EX and POST-R periods, which receded to levels similar to PRE-EX by POST-24 in both conditions. In contrast, peak concentrations for CK were observed at POST-24, whilst, differences between CON and CWI for all blood variables were unclear throughout the time points.

Total and individual Hooper scores were similar between conditions at PRE-EX. However, at POST-24, moderate to large differences between CON and CWI were noted for changes in total Hooper scores ($d = 1.93$), DOMS ($d = 0.60$), sensations of fatigue ($d = 1.51$), sleep quality ($d = 1.36$) and perceived stress ($d = 1.56$), with CWI having a positive effect on the exercise-mediated changes within these variables.

DISCUSSION

The main findings of the present study indicate that POST-R sprint (5- and 10-m) performances were ‘likely to very likely’ (**Figures 1A,B**, $d = 0.64$ and 0.65) impaired by prior CWI. However, moderate improvements in 10-m sprint performance were ‘likely’ evident at POST-24 after CWI compared with

CON (**Figure 1B**, $d = 0.53$; -1.24 to 0.18). Additionally, the use of CWI ‘almost certainly’ resulted in a large overall improvement in Wellness/Hooper scores (**Figure 3**, $d = 1.93$; -2.60 to -1.25). Specifically, CWI ‘almost certainly’ resulted in improved sleep quality ($d = 1.36$; -2.04 to -0.68), stress ($d = 1.56$; -2.24 to -0.88) and perceived fatigue ($d = 1.51$; -2.19 to -0.84), and ‘likely’ resulted in a moderate decrease in DOMS ($d = 0.60$; -1.27 to 0.08). These findings as such render CWI as a suitable recovery intervention following MMA combat, particularly where the restoration of physical performance is required within a 24-h period. Within CON, small to moderate decreases in 10-m sprint, as well as CMJ and SJ performances were observed following (POST-24) simulated MMA combat (**Figure 1**). Such observations are in agreement with previous studies where sustained decrements in explosive, neuro-muscular performances were observed following a MMA training session (Lindsay et al., 2017) and simulated MMA competition (Ghoul et al., 2017). This decrement in muscular performance may be in-part explained by activity-induced muscle damage, likely sustained due to the collision, as well as time-sustained eccentric and/or isometric contractions during grappling moves, locks and holds performed during MMA activity. For instance, within rugby codes, both the severity of muscle damage, and the decrease in lower body neuro-muscular performances measured just after games/training sessions, have been shown to be associated to the intensity and frequency of collisions. (Tavares et al., 2017).

While a recent study (Lindsay et al., 2017) focused on the influence of CWI on the inflammatory/oxidative stress response following MMA combat, the recovery in physical performance has yet to be characterized. Accordingly, this is the first study to investigate the influence of CWI on the recovery of physical performance up to 24-h following simulated MMA combat. The present findings demonstrate that CWI enhanced the recovery of 10-m sprint performance at 24-h post simulated combat compared with CON (**Figure 1**). However, it is pertinent to note that 5-m sprint, as well as both jump performances were not influenced by CWI when compared to CON (**Figure 1**). It is likely that these performance measures were too brief and/or their changes influenced by CWI were too small to be detected. Alternatively, muscle recovery mediated by CWI might have little influence on the initial rate of force development (which are likely greater determinants of 5-m and jump performances) compared with extended force development, which would be better related to 10-m sprint performance.

Within the current study, it would be reasonable to consider that CWI might have improved the recovery of some performance measures (i.e., 10-m sprint) through reducing the extent of muscle damage and/or inflammation. While such reasoning would agree well with previous work documenting the beneficial effects of CWI on exercise-induced muscle damage (Pournot et al., 2010; Rowsell et al., 2011; Bleakley et al., 2012; Ihsan et al., 2016; Machado et al., 2016), the accompanying hematological data demonstrating no influence of CWI (**Figure 2**) clearly indicates an alternative mechanism by which sprint performance was improved. In the present study, the improved performance could potentially be explained by the






FIGURE 1 | Changes in performances during 5-m sprint (A), 10-m sprint (B), squat jump (C) and countermovement jump (D) prior to exercise (PRE-EX), immediately after recovery (POST-R) and 24-h following exercise (POST-24) according to recovery modalities: passive recovery (CON) or cold water immersion (CWI). Differences are indicated by the size of the effect (S, small; M, moderate; L, large), accompanied by symbols denoting within (*CON, +CWI) and between group (#CON vs. CWI) differences. All reported differences denote $\geq 75\%$ likelihood that the stated effect is substantially different from the smallest worthwhile change ($d = 0.2$).

reduced sensation of DOMS (Figure 3), independent of muscle damage. Indeed, perception of DOMS seems to be important to the recovery of exercise performance, as muscle pain independent of muscle damage has been shown to impair force generating capacity (Graven-Nielsen et al., 2002). Accordingly, CWI has been suggested to modulate the sensation of DOMS, and by extension muscle function, through limiting oedema (Ihsan et al., 2013; Choo et al., 2016) and/or through activating cutaneous receptors that mediate analgesia (Knowlton et al., 2013; Ihsan et al., 2016). Counterintuitive to the notion of recovery, CWI resulted in moderate and large decrements in POST-R for 5-m and 10-m sprint performances, respectively (Figure 1). Indeed, low intramuscular temperatures have been purported to result in delayed action potential generation, cross-bridge cycling and/or impaired fiber contractile properties. Moreover, the dynamic contractile force of the leg extensors has been shown to decrease by 4–6 % for every 1°C reduction in intramuscular temperatures (Bergh and Ekblom, 1979). However, the authors feel that these detrimental effects are likely transient in nature, and would likely recede following the rewarming of the musculature. Nevertheless, practitioners should proceed with caution when utilizing CWI between sprint/power based performance bouts scheduled close to each other.

The data also demonstrates an overall improvement in wellness measures 24 h following CWI (Figure 3), which

could also in-part account for the improved recovery in sprint performance. This has been supported by recent findings that have demonstrated strong temporal associations between changes in wellness with the recovery of neuromuscular function and match running performances within team-sport athletes (Johnston et al., 2013; Ihsan et al., 2017). However, we acknowledge that the mechanisms underpinning improvements in post-exercise wellness following CWI are not entirely understood. It is suggested that improvements in perceptual recovery following CWI is modulated through changes in cardiac parasympathetic activity. For instance, enhanced parasympathetic activity following post-exercise CWI has been shown to be associated with reduced muscle soreness and perception of fatigue, as well as improved sense of relaxation (Stanley et al., 2012; Stanley et al., 2013; Choo et al., 2018). Additionally, perceptual recovery following CWI may also be related to alterations within the cerebral dopaminergic and noradrenergic systems, which have been implicated in the development of fatigue and may potentially be ameliorated following exposure to a cooling stimulus (Mundel et al., 2007; Roelands et al., 2013; Ihsan et al., 2016). However, the Hooper wellness battery also included ratings of sleep quality, which was seemingly improved following the use of CWI (Figure 3). This finding is somewhat in contrast with Robey et al. (2013), who demonstrated no influence of CWI



on salivary melatonin, polysomnography or actigraphy-derived sleep measures. The reasons for such discrepancies are unclear, but perhaps there may be some dissociation between objective and subjective sleep measures, at least following the use of CWI use.

Although the simulated MMA combat protocol resulted in increased CK within both conditions at POST-24, there was no effect of CWI on CK concentration in the bloodstream (Figure 2). This finding is in agreement with some reports (Corbett et al., 2011; Pointon et al., 2011) but not all, as several investigations have reported decreased blood CK following CWI over a 24- to 72-h period (Eston and Peters, 1999; Ascensao et al., 2011; Minett et al., 2013). Such discrepancies within the literature may be due to differences in sampling time, nature of exercise and level of athletes. It was also suggested that CWI might facilitate physiological and perceptual recovery through altering the hormonal milieu. The current findings do not support this suggestion, as no changes in cortisol, testosterone or cortisol-to-testosterone ratio were observed following CWI (Figure 2). The present study findings are similar to Halson et al. (2008), who found no changes in cortisol or testosterone concentrations in response to post-exercise CWI. However, in the current study, we have characterized these hormonal changes for over a 24-h recovery period to elucidate potential mechanisms if any, as opposed to a 40-min post-exercise window undertaken in the study of Halson et al. (2008).

The present findings render CWI modality as a suitable recovery intervention during tournament scenarios or during intensive training camps/periods where athletes are required to compete regularly and the recovery/restoration of physical ability is paramount. The current data also advocates caution with

regards to utilizing this recovery modality when performance bouts are closely scheduled together. In such scenarios, a warm-up exercises to facilitate the re-warming of the musculature are certainly warranted, and further studies should investigate this particular point.

We acknowledge several limitations associated with this study. Firstly, we were unable to monitor recovery exceeding 24 h, which could have been valuable in understanding the effects of CWI on longer term recovery. However, given that our participants were well-trained athletes, a 24 h recovery period is reasonable, considering their general ability to recover quicker, as well as their availability to commit to controlled experimental studies. Moreover, the lack of sport-specific performance variables could be considered a limitation as well. However, the choice of sprints and jumps, although general, are better controlled for, easily replicable and enables better objectivity when comparing with other studies.

CONCLUSION

In summary, the present study examined the influence of post-exercise CWI on the recovery of physical performance, hematological stress markers and perceived wellness following a simulated MMA combat match. The use of CWI resulted in an enhanced recovery of 10-m sprint performance, as well as improved perceived wellness 24 h following simulated

MMA combat. However, sprint/power ability was impaired when undertaken shortly after CWI. As such, active warm-ups are certainly warranted if CWI is administered in between performance bouts that are closely scheduled together. Hematological stress markers were not influenced by CWI, indicating that the physical and perceptual recovery was dissociated from the hormonal markers examined.

AUTHOR CONTRIBUTIONS

MT, NG, JC, and CT conceived and planned the experiments. MT, NG, JC, HC, and CT carried out the experiments. MT, MI, NG, JC, AC, CT, and KC contributed to the interpretation of the results. MT took the lead in writing the manuscript. All authors provided critical feedback and helped shape the research, analysis and manuscript.

ACKNOWLEDGMENTS

The authors thank Machida Mixed Martial Arts Nabeul Team and Sensei Fehri Ounaies for the help concerning the implementation of the experimentation. They thank Dr. Rim Sioud and Mr. Mohamed Hedi Mahfoudhi for their help during the experimentation. They also thank “Dragon Bleu France” for their support by providing the equipment.

REFERENCES

- Amtmann, J. A., Amtmann, K. A., and Spath, W. K. (2008). Lactate and rate of perceived exertion responses of athletes training for and competing in a mixed martial arts event. *J. Strength Cond. Res.* 22, 645–647. doi: 10.1519/JSC.0b013e318166018e
- Ascensao, A., Leite, M., Rebelo, A. N., Magalhaes, S., and Magalhaes, J. (2011). Effects of cold water immersion on the recovery of physical performance and muscle damage following a one-off soccer match. *J. Sports Sci.* 29, 217–225. doi: 10.1080/02640414.2010.526132
- Bergh, U., and Ekblom, B. (1979). Influence of muscle temperature on maximal muscle strength and power output in human skeletal muscles. *Acta Physiol. Scand.* 107, 33–37. doi: 10.1111/j.1748-1716.1979.tb06439.x
- Bleakley, C., McDonough, S., Gardner, E., Gdavid, B., Jty, H., and Davison, G. (2012). Cold-water immersion for preventing and treating muscle soreness after exercise. *Cochrane Database Syst. Rev.* 15:CD008262. doi: 10.1002/14651858.CD008262.pub2
- Choo, H. C., Nosaka, K., Peiffer, J. J., Ihsan, M., Yeo, C. C., and Abbiss, C. R. (2016). Peripheral blood flow changes in response to postexercise cold water immersion. *Clin. Physiol. Funct. Imaging* 38, 46–55. doi: 10.1111/cpf.12380
- Choo, H. C., Nosaka, K., Peiffer, J. J., Ihsan, M., Yeo, C. C., and Abbiss, C. R. (2018). Effect of water immersion temperature on heart rate variability following exercise in the heat. *Kinesiology* 50, 67–74.
- Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences*, 2nd Edn. Hillsdale, MI: Erlbaum Associates.
- Corbett, J., Barwood, M. J., Lunt, H. C., Milner, A., and Tipton, M. J. (2011). Water immersion as a recovery aid from intermittent shuttle running exercise. *Eur. J. Sport Sci.* 12, 509–514. doi: 10.1080/17461391.2011.570380
- de Freitas, V. H., Ramos, S. P., Bara-Filho, M. G., Freitas, D. G., Coimbra, D. R., Cecchini, R., et al. (2017). Effect of cold water immersion performed on successive days on physical performance, muscle damage, and inflammatory, hormonal, and oxidative stress markers in volleyball players. *J. Strength Cond. Res.* doi: 10.1519/JSC.0000000000001884 [Epub ahead of print].
- del Vecchio, F. B., Hirata, S. M., and Franchini, E. (2011). A review of time-motion analysis and combat development in mixed martial arts matches at regional level tournaments. *Percept. Mot. Skills* 112, 639–648. doi: 10.2466/05.25.PMS.112.2.639-648
- Eston, R., and Peters, D. (1999). Effects of cold water immersion on the symptoms of exercise-induced muscle damage. *J. Sports Sci.* 17, 231–238. doi: 10.1080/026404199366136
- Ghoul, N., Tabben, M., Miarka, B., Tourny, C., Chamari, K., and Coquart, J. (2017). Mixed martial arts induces significant fatigue and muscle damage up to 24 hours post-combat. *J. Strength Cond. Res.* doi: 10.1519/JSC.0000000000002078 [Epub ahead of print].
- Graven-Nielsen, T., Lund, H., Arendt-Nielsen, L., Danneskiold-Samsøe, B., and Bliddal, H. (2002). Inhibition of maximal voluntary contraction force by experimental muscle pain: a centrally mediated mechanism. *Muscle Nerve* 26, 708–712. doi: 10.1002/mus.10225
- Halson, S. L., Quod, M. J., Martin, D. T., Gardner, A. S., Ebert, T. R., and Laursen, P. B. (2008). Physiological responses to cold water immersion following cycling in the heat. *Int. J. Sports Physiol. Perform.* 3, 331–346. doi: 10.1123/ijspp.3.3.331
- Hooper, S. L., Mackinnon, L. T., Howard, A., Gordon, R. D., and Bachmann, A. W. (1995). Markers for monitoring overtraining and recovery. *Med. Sci. Sports Exerc.* 27, 106–112. doi: 10.1249/00005768-199501000-00019
- Hopkins, W. G. (2006). Spreadsheets for analysis of controlled trials with adjustment for a subject characteristic. *Sportscience* 10, 46–50.
- Hopkins, W. G., Marshall, S. W., Batterham, A. M., and Hanin, J. (2009). Progressive statistics for studies in sports medicine and exercise science. *Med. Sci. Sports Exerc.* 41, 3–13. doi: 10.1249/MSS.0b013e3181818cb278
- Ihsan, M., Tan, F., Sahrom, S., Choo, H. C., Chia, M., and Aziz, A. R. (2017). Pre-game perceived wellness highly associates with match running performances during an international field hockey tournament. *Eur. J. Sport Sci.* 17, 593–602. doi: 10.1080/17461391.2017.1301559
- Ihsan, M., Watson, G., and Abbiss, C. R. (2016). What are the physiological mechanisms for post-exercise cold water immersion in the recovery from prolonged endurance and intermittent exercise? *Sports Med.* 46, 1095–1109. doi: 10.1007/s40279-016-0483-3

- Ihsan, M., Watson, G., Lipski, M., and Abbiss, C. R. (2013). Influence of postexercise cooling on muscle oxygenation and blood volume changes. *Med. Sci. Sports Exerc.* 45, 876–882. doi: 10.1249/MSS.0b013e31827e13a2
- Johnston, R. D., Gabbett, T. J., and Jenkins, D. G. (2013). Influence of an intensified competition on fatigue and match performance in junior rugby league players. *J. Sci. Med. Sport* 16, 460–465. doi: 10.1016/j.jsams.2012.10.009
- Knowlton, W. M., Palkar, R., Lippoldt, E. K., McCoy, D. D., Baluch, F., Chen, J., et al. (2013). A sensory-labeled line for cold: TRPM8-expressing sensory neurons define the cellular basis for cold, cold pain, and cooling-mediated analgesia. *J. Neurosci.* 33, 2837–2848. doi: 10.1523/JNEUROSCI.1943-12.2013
- La Bounty, P., Campbell, B. I., Galvan, E., Cooke, M., and Antonio, J. (2011). Strength and conditioning considerations for mixed martial arts. *Strength Cond. J.* 33, 56–67. doi: 10.1519/SSC.0b013e3182044304
- Lindsay, A., Carr, S., Cross, S., Petersen, C., Lewis, J. G., and Gieseg, S. P. (2017). The physiological response to cold-water immersion following a mixed martial arts training session. *Appl. Physiol. Nutr. Metab.* 42, 529–536. doi: 10.1139/apnm-2016-0582
- Lindsay, A., Lewis, J., Gill, N., Gieseg, S. P., and Draper, N. (2015). Effect of varied recovery interventions on markers of psychophysiological stress in professional rugby union. *Eur. J. Sport Sci.* 15, 543–549. doi: 10.1080/17461391.2015.1029982
- Machado, A. F., Ferreira, P. H., Micheletti, J. K., De Almeida, A. C., Lemes, Í. R., Vanderlei, F. M., et al. (2016). Can water temperature and immersion time influence the effect of cold water immersion on muscle soreness? A systematic review and meta-analysis. *Sports Med.* 46, 503–514. doi: 10.1007/s40279-015-0431-7
- Minett, G. M., Duffield, R., Billaut, F., Cannon, J., Portus, M. R., and Marino, F. E. (2013). Cold-water immersion decreases cerebral oxygenation but improves recovery after intermittent-sprint exercise in the heat. *Scand. J. Med. Sci. Sports* 24, 656–666. doi: 10.1111/sms.12060
- Mundel, T., Bunn, S. J., Hooper, P. L., and Jones, D. A. (2007). The effects of face cooling during hyperthermic exercise in man: evidence for an integrated thermal, neuroendocrine and behavioural response. *Exp. Physiol.* 92, 187–195. doi: 10.1113/expphysiol.2006.034934
- Nunan, D. (2006). Development of a sports specific aerobic capacity test for karate - a pilot study. *J. Sports Sci. Med.* 5, 47–53.
- Pointon, M., Duffield, R., Cannon, J., and Marino, F. (2011). Cold water immersion recovery following intermittent-sprint exercise in the heat. *Eur. J. Appl. Physiol.* 112, 2483–2494. doi: 10.1007/s00421-011-2218-3
- Pournot, H., Bieuzen, F., Duffield, R., Lepretre, P.-M., Cozzolino, C., and Hausswirth, C. (2010). Short term effects of various water immersions on recovery from exhaustive intermittent exercise. *Eur. J. Appl. Physiol.* 111, 1287–1295. doi: 10.1007/s00421-010-1754-6
- Robey, E., Dawson, B., Halson, S., Gregson, W., King, S., Goodman, C., et al. (2013). Effect of evening postexercise cold water immersion on subsequent sleep. *Med. Sci. Sports Exerc.* 45, 1394–1402. doi: 10.1249/MSS.0b013e318287f321
- Roelands, B., De Koning, J., Foster, C., Hettinga, F., and Meeusen, R. (2013). Neurophysiological determinants of theoretical concepts and mechanisms involved in pacing. *Sports Med.* 43, 301–311. doi: 10.1007/s40279-013-0030-4
- Rowell, G. J., Coutts, A. J., Reaburn, P., and Hill-Haas, S. (2011). Effect of post-match cold-water immersion on subsequent match running performance in junior soccer players during tournament play. *J. Sports Sci.* 29, 1–6. doi: 10.1080/02640414.2010.512640
- Stanley, J., Buchheit, M., and Peake, J. M. (2012). The effect of post-exercise hydrotherapy on subsequent exercise performance and heart rate variability. *Eur. J. Appl. Physiol.* 112, 951–961. doi: 10.1007/s00421-011-2052-7
- Stanley, J., Peake, J. M., and Buchheit, M. (2013). Consecutive days of cold water immersion: effects on cycling performance and heart rate variability. *Eur. J. Appl. Physiol.* 113, 371–384. doi: 10.1007/s00421-012-2445-2
- Tavares, F., Smith, T. B., and Driller, M. (2017). Fatigue and recovery in rugby: a review. *Sports Med.* 47, 1515–1530. doi: 10.1007/s40279-017-0679-1
- Vaile, J., Halson, S., Gill, N., and Dawson, B. (2008). Effect of hydrotherapy on recovery from fatigue. *Int. J. Sports Med.* 29, 539–544. doi: 10.1055/s-2007-989267

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Tabben, Ihsan, Ghoul, Coquart, Chaouachi, Chaabene, Tourny and Chamari. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.