



HAL
open science

Poplar rotation coppice at a trace element-contaminated phytomanagement site: A 10-year study revealing biomass production, element export and impact on extractable elements

Michel Chalot, Olivier Girardclos, Lisa Ciadamidaro, Cyril Zappellini, Loïc Yung, Alexis Durand, Stéphane Pfendler, Isabelle Lamy, Vincent Driget, Damien Blaudez

► To cite this version:

Michel Chalot, Olivier Girardclos, Lisa Ciadamidaro, Cyril Zappellini, Loïc Yung, et al.. Poplar rotation coppice at a trace element-contaminated phytomanagement site: A 10-year study revealing biomass production, element export and impact on extractable elements. *Science of the Total Environment*, 2020, 699, pp.134260. <10.1016/j.scitotenv.2019.134260>. <hal-02355647>

HAL Id: hal-02355647

<https://hal.science/hal-02355647v1>

Submitted on 13 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

1 Poplar rotation coppice at a trace element-contaminated
2 phytomanagement site: a 10-year study revealing biomass production,
3 element export and impact on extractable elements

4

5 **Michel Chalot^{a,b}, Olivier Girardclos^a, Lisa Ciadamidaro^a, Cyril Zappellini^a, Loic Yung^a,**
6 **Alexis Durand^a, Stéphane Pfendler^a, Isabelle Lamy^c, Vincent Driget^a, Damien Blaudez^d**

7

8 ^aUniversité de Bourgogne Franche-Comté, CNRS, Laboratoire Chrono-environnement, F-
9 25250, Montbéliard, France

10 ^bUniversité de Lorraine, F-54000, Nancy, France.

11 ^c INRA, AgroParisTech, UMR1402 ECOSYS, Ecotoxicology division, F-78026 Versailles
12 cedex, France

13 ^dUniversité de Lorraine, CNRS, LIEC, F-54000, Nancy, France

14

15 *Corresponding author: Laboratoire Chrono-Environnement, UMR 6249, Université de
16 Bourgogne Franche-Comté, 4 place Tharradin, BP71427, F-25211 Montbéliard, France

17 Email: michel.chalot@univ-fcomte.fr

18

19 **Abstract**

20 Growing lignocellulosic crops on marginal lands could compose a substantial proportion of
21 future energy resources. The potential of poplar was explored, by devising a field trial of two
22 hectares in 2007 in a metal-contaminated site to quantify the genotypic variation in the growth
23 traits of 14 poplar genotypes grown in short-rotation coppice and to assess element transfer
24 and export by individual genotypes. Our data led us to conclusions about the genotypic
25 variations in poplar growth on a moderately contaminated site, with the Vesten genotype
26 being the most productive. This genotype also accumulated the least amounts of trace
27 elements, whereas the Trichobel genotype accumulated up to $170 \text{ mg Zn kg}^{-1} \text{ DW}$ in the
28 branches, with large variation being exhibited among the genotypes for trace element (TE)
29 accumulation. Soil element depletion occurred for a range of TEs, whereas the soil content of
30 major nutrients and the pH remained unchanged or slightly increased after 10 years of poplar
31 growth. The higher TE content of bark tissues compared with the wood and the higher
32 proportion of bark in branches compared with the wood led us to recommend that only stem
33 wood be harvested, instead of the whole tree, which will enable a reduction in the risks
34 encountered with TE-enriched biomass in the valorization process.

35

36

37

38 **Keywords:** export of trace elements, biomass yield, phytomanagement, poplar genotypes

39

40 **1. Introduction**

41 Renewable resources are the focal point of European Union (EU) initiatives to establish
42 energy self-sufficiency in the EU. Woody biomass is a renewable resource and is almost
43 carbon-neutral (Zanchi et al., 2012). Using wood as an energy source is certainly not new;
44 however, second-generation biofuels utilize woody biomass as raw materials to develop liquid
45 fuels, *i.e.*, ethanol and diesel within biorefinery projects. Lignocellulosic perennial crops are
46 indeed recognized as feedstock for platform chemicals and lignin-based building blocks
47 (Rohde et al., 2018). The EU has been, and remains, the major global source of demand as a
48 result of its predicted targets for renewable energy. This demand is largely met by the forest
49 resources of the UE and supplemented by imports from the United States, Canada and Russia.
50 Growing lignocellulosic crops on contaminated or marginal lands could make up a substantial
51 proportion of future energy resources (Gelfand et al., 2013). Efforts have been made in the
52 past ten years to link phytotechnologies to the production of valuable biomass products and to
53 the fate of contaminants during industrial processes (Ancona et al., 2019; Asad et al., 2017;
54 Bert et al., 2017; Chalot et al., 2012; Dastyar et al., 2019; Delplanque et al., 2013; Šyc et al.,
55 2012). Despite these efforts, phytotechnologies have not been commercially applied to date
56 (Conesa et al., 2012; Gerhardt et al., 2017) and the use of gentle remediation options (GRO)
57 as practical onsite remedial opportunities is still limited in Europe, particularly at trace-
58 element-contaminated sites (Cundy et al., 2016). This lack of application is mostly due to the
59 lack of long-term field assessments under realistic conditions. A recent article discussed how
60 phytotechnologies could be upgraded from a proven technology to a widely accepted practice
61 (Gerhardt et al., 2017), including the implementation of large-scale field trials in long-term
62 experiments.

63

64 Among useful crops, poplar species and hybrids are considered low-input woody crops,
65 with recognized adaptation capacities to a wide range of site and climatic conditions (Navarro
66 et al., 2014; Verlinden et al., 2013; Yáñez et al., 2019). A useful list of publications on the use
67 of poplar as a renewable resource is provided by the Food and Agriculture Organization of the
68 United Nations (FAO, 2016). This document, to which recent reviews on the use of poplar
69 biomass for platform chemicals (Rohde et al., 2018) and thermal energy (Pandey et al., 2016)
70 can be added, listed the numerous industrial processes that may consider poplar biomass.

71 Poplars also readily grow on marginal lands where biomass yields can fall within the
72 range 8-12 tons per hectare per year (Ciadamidaro et al., 2017; Verlinden et al., 2013). The
73 use of *Populus* species has been proposed for phytomanagement purposes because these
74 woody species are characterized by fast-growth, deep-rooting systems and can be managed in
75 short-rotation coppicing plantations (Ciadamidaro et al., 2017; Laureysens et al., 2005, 2004;
76 Marmioli et al., 2011; Migeon et al., 2009; Rohde et al., 2018; Vangronsveld et al., 2009;
77 Yáñez et al., 2019). The genus *Populus* has traditionally been divided into six sections on the
78 basis of the flower and leaf characters, with approximately 30–40 species occurring
79 worldwide, mainly in the Northern Hemisphere (Polle and Douglas, 2010). Different
80 genotypes of the same poplar species may exhibit variable tolerance to trace elements (TE)
81 because of intraspecific genetic dissimilarity (Castiglione et al., 2009). For instance, a large
82 variation in poplar genotypic responses to leachate irrigation has been found (Zalesny et al.,
83 2007), and high variability occurs in foliar trace element concentrations within a selection of
84 14 poplar genotypes (Pottier et al., 2015).

85 A number of works have been dedicated to the study of poplar responses to soil
86 contamination in controlled pot or hydroponic (Cicatelli et al., 2010; De Oliveira and Tibbett,
87 2018; Langer et al., 2012; Lingua et al., 2008; Migeon et al., 2012), tank (Zalesny and Bauer,
88 2007), or lysimeter (Dimitriou and Aronsson, 2011, 2010) set ups. Such experiments are

89 needed to study fundamental mechanisms (contaminant transfer between soil-aboveground
90 and below ground plant tissues, interactions between plant and microbes) but may not reflect
91 how trees will behave in natural situations in the long term. Field trials have been
92 implemented, and short-term data are available for a few field experiments (Castiglione et al.,
93 2009; French et al., 2006; Gamalero et al., 2012; Lettens et al., 2011; Sabatti et al., 2014).
94 However, long-term studies are scarce (Evangelou et al., 2012; Foulon et al., 2016a, 2016b;
95 Mertens et al., 2007) but are crucial to the study of long-term impacts of tree plantation on
96 pedological and biological soil characteristics.

97 The PHYTOPOP project aimed to explore the potential of poplar for
98 phytomanagement purposes, by setting up a field trial of two hectares in 2007 to produce
99 enough biomass for further valorization and estimate productivity at a realistic scale. The
100 study was based on the utilization of a large panel of poplar cultivars, planted in a short
101 rotation coppice (SRC) scheme. During the project, the harvesting process was also
102 investigated from an economic point of view, and analyzed poplar behavior in a combustion
103 unit (Chalot et al., 2012). The PHYTOPOP biomasses were further used to assess the fate of
104 TE during pyrolysis (Bert et al., 2017) and bioethanol production (Asad et al., 2017). The
105 mechanisms controlling metal content in the leaves and the expression of genes that may be
106 involved in metal transport were also studied (Migeon et al., 2010; Pottier et al., 2015).
107 However, in none of these papers we evaluated biomass production at this specific site. We
108 indeed wanted to obtain long-term data, which are lacking in the literature, to provide some
109 recommendations to end-users.

110 The present paper aimed to determine the impacts of a PHYTOPOP plantation on soil
111 characteristics, TE accumulation and poplar yields after one rotation. The specific objectives
112 were 1) to quantify the genotypic variation in the growth traits of 14 poplar genotypes grown

113 in SRC systems at a TE contaminated site and 2) to assess element transfer and export by the
114 different genotypes.

115 **2. Materials and methods**

116 **2.1. Site description and experimental design**

117 The experimental site is located in the Northwest region of Paris at Pierrelaye
118 (49°01'47.6"N, 2°10'34.2"E). The whole area has been irrigated with raw wastewaters from
119 the end of the 19th century until the beginning of the 21st century to increase the soil
120 fertilization potential and has been dedicated to vegetable gardening for more than one
121 century (Baize et al., 2002, Dère et al., 2007, Lamy et al., 2006). However, this resulted in the
122 accumulation of TE such as Pb, Cu, Zn and Cd in the first 30-50 cm of soil layer. This leads
123 to poly-metallic pollution characterized by TE concentrations that could reach levels more
124 than 10-fold greater than those of a non irrigated local reference soil.

125 Poplars were planted in April 2007 as a short-rotation coppice (SRC) at densities of
126 1000 stems per ha for bioenergy production as part of the PHYTOPOP project, as fully
127 described in previous papers (Foulon et al., 2016a, 2016b; Pottier et al., 2015). Based on the
128 soil characteristics at the experimental site, 14 poplar genotypes were chosen that belong to
129 the two Aigeros and Tacamahaca sections and to 4 species *P. deltoides*, *P. deltoides* x *P.*
130 *nigra*, *P. trichocarpa* and *P. trichocarpa* x *P. maximowiczii*. Each of the 14 genotypes was
131 grown on two independent plots of 389 m², with 49 poplars per plot (7 rows of 7 trees), with
132 spaces of 3.50 m per line and 2.85 m per row. Details on the genotypes are provided in Table
133 S1.

134

135 **2.2. Soil sampling and chemical analyses**

136 In April 2007 and January 2011, 25 topsoil samples were taken from all poplar
137 genotypes with an auger (4 cm in diameter), to a depth of 20-25 cm, each being taken at 50

138 cm from the planted poplar. The 25 samples were mixed in a large container; then, 2 to 3 kg
139 of the mixture (approximately 50%) was taken to the laboratory as a composite sample and
140 analyzed. Soil samples were further collected using the same procedure in October 2013 and
141 October 2016 for the Skado genotype only. This genotype, together with the I214 genotype,
142 has been selected for a subsequent field trial in 2011 within the BIOFILTREE project
143 (Ciadamidaro et al., 2017), given preliminary growth assessments in 2010.

144 Initial (2007) and later 2011 soil analyses were performed by the INRA Soil Analysis
145 Laboratory (LAS-INRA, Arras, France), which is accredited by COFRAC (French
146 accreditation board). The soil pH (soil: water ratio = 1:5) was measured according to NF ISO
147 10390. For initial soils, the total Zn, Pb, Cu, and Cd were determined according to NF X31-
148 147 (tri-acid HF–HCl–HNO₃ digestion). In addition, TE availability in soils was determined
149 using two extracting reagents: an organic chelating agent, diethylenetriamine-pentaacetic acid
150 (DTPA), and a neutral salt, ammonium nitrate (NH₄NO₃). The NH₄NO₃ extraction was
151 selected to estimate the easily exchangeable pool of elements (Pruess, 1998) and has been
152 widely used (McLaughlin et al., 2000; Pérez-de-Mora et al., 2006; Madejón et al., 2006). Soil
153 was homogenized in a solution of NH₄NO₃ 1 M (1/2.5 m/v) (DIN 19730, 1997). The aim of
154 the DTPA extraction was to quantify exchangeable metals and also metals associated with
155 organic matter and with Mn, Fe, and Al oxy-hydroxides (Baize, 2003). Extraction was done in
156 triethanolamine-CaCl₂-DTPA (0.005 M) solution at pH = 7.3 (NF X 31-121). After
157 extraction, the total, DTPA-, and NH₄NO₃-extractable element concentrations in solution
158 were obtained with inductively coupled plasma mass spectrometry analyses.

159 The third and fourth samplings were carried out in October 2013 and October 2016 (as
160 described above) but only under the Skado genotype, thus allowing 7 and 10 years,
161 respectively, of poplar growth. To determine the TE extractable fractions of these soil
162 samples, 5 g of 2-mm sieved soil was dried at 60°C for 48 h and incubated with 50 ml of

163 10 mM calcium chloride under agitation (40 rpm) for 2 h at room temperature. The mixture
164 was first filtered with ash-free filters and subsequently passed through a 0.45- μm mesh and
165 acidified at 2% (v/v) with HNO_3 prior to inductively coupled plasma atomic emission
166 spectrometry (ICP-AES, Thermo Fischer Scientific, Inc., Pittsburg, USA) analysis. The 2007
167 and 2011 samples were also reanalyzed by the same analytical procedure. The methods for
168 measuring other physicochemical characteristics are those described by Zappellini et al.
169 (2015).

170 **2.3. Biomass harvest and measurements**

171 In the experimental field, diameter at breast height (dbh) and tree total height (H_{tot}) were
172 measured in October 2013, allowing 7 years of growth after establishment of the plantation
173 (one rotation), on all individual trees from the 14 plots (*i.e.*, 25 trees per plot, excluding the
174 border trees). Allometric equations were calculated per genotype to identify genotypic
175 variability, using trunk and branches dry weight (DW) of a tree as a function of stem dbh. The
176 plot estimates were further extrapolated to determine DW per ha and per year. The proportion
177 of bark dry mass of total biomass (wood and bark) was calculated for samples of various
178 diameters. Allometric power equations between diameter and bark proportion were used to
179 estimate whole tree bark dry biomass. The plot estimates were extrapolated to produce DW
180 per ha and per year.

181 To determine the best fitting allometric equation, a subset of trees was selected using an
182 optimal design approach based on the best subset of trees, allowing the statistical model
183 parameters to be estimated, as described in the “statistical analysis” section below. Damaged
184 trees were removed at this selection step. Per genotype, between 5 and 7 trees were required
185 to meet the statistical criteria on the 2013 observed dbh and H_{tot} , considering the allometric
186 assumption that these two variables are related to DW. In February 2014, selected trees were
187 harvested, and fresh biomasses were determined for the two compartments, trunks (primary

188 axis up to the top of tree) and branches (all secondary axes). Three trunk sections were
189 collected on each tree at their basis (DM_b), at half of total height (DM_m) and at the tree
190 summit, at a diameter of approximately 4 cm (DM_s); branches were also collected (Fig. S1).
191 Fresh samples were weighed immediately after harvest and dried in a forced air oven at 50°C
192 until reaching constant weights. The dry biomass was then measured. The total trunk
193 (trunkDW in kilograms) and branch (branchDW in kilograms) dry weights were calculated
194 from fresh weight (FW in kilograms) data and desiccation rates (DR), considering the samples
195 mentioned above, using the following equations (1-2):

196 (1) $\text{trunkFW (kg)} * (\text{DR}_b + \text{DR}_m + \text{DR}_s) / 3 = \text{trunkDW (kg)}$

197 (2) $\text{branchFW (kg)} * \text{DR}_{br} = \text{branchDW (kg)}$

198 For allometric data, the following best fitting equation was used (3):

199 (3) $DW = a * dbh^b$

200 where DW is the aboveground biomass (excluding leaves) production (kg DW tree^{-1}),
201 dbh is the diameter at breast height (in cm), and a and b are the regression coefficients that
202 were shown to fit the material properly for poplar (Johansson, 1999; Picard et al., 2012;
203 Morhart et al., 2013).

204 **2.4. Element concentrations in biomasses**

205 Poplar samples (branches, trunks, wood, and bark) were used for element analysis
206 using previously published procedures (Ciadamidaro et al., 2017). Poplar samples were
207 washed with distilled water, dried at 70°C for at least 48 h, and ground using a stainless-steel
208 mill. The studied elements (Ca, Cd, Cr, Cu, Fe, K, Mg, Mn, P, Pb and Zn) were determined
209 by ICP-AES following the acid digestion with concentrated HNO_3 (DigiPREP system). All
210 samples were run together with certified reference materials, including oriental basma tobacco
211 leaves (INCT-OBTL-5, LGC Promochem, Molsheim, France) and a loamy clay soil
212 (CRM052, LGC Promochem, Molsheim, France). The percentage of recovery of the studied

213 elements ranged between 77 and 110%. The element export from the soil was further
214 evaluated by each poplar genotype by calculating the quantity of elements transferred to the
215 woody biomass per hectare and per year.

216

217 **2.5. Statistical analyses**

218 For allometric data, all statistical analyses were performed using R 3.4.1 software (R
219 Development Core Team, 2015). Optimal subsets of trees were selected per genotype using R
220 package AlgDesign (Wheeler 2014), which allowed estimation of statistical model parameters
221 without bias and with minimum variance of the relationship between dbh and H_{tot} , thereby
222 maximizing the criteria established earlier (Fedorov, 1972). After providing the DW data,
223 graphical exploration allowed for checking the sampling and drying procedure bias and
224 showed the presence of heteroscedasticity in the nonlinear relationship between dbh and
225 branchDW or trunkDW, as is the case for many taxa (Picard et al., 2012). For the model
226 calibration, goodness of fit was evaluated by the coefficient of determination (R^2), and
227 heteroscedasticity was assessed by plotting residuals against observed values and testing their
228 normality (Shapiro-Wilk test). Logarithm transformation of data and power form
229 ($DW=a*dbh^b$) with a and b as the regression coefficients were the most relevant models.
230 When regressions were performed, slopes and intercepts were tested for differences among
231 genotypes by analysis of covariance. Interactions between the dbh and genotypes have been
232 considered but were never statistically significant in modeling. The two biomass components,
233 branchDW and trunkDW, were modeled separately as two dependent variables without
234 consideration of possible additivity (Vega-Nieva et al., 2015). However, when applying the
235 models, the sum of the two components resulted only in an average of 2.3% higher than the
236 observed total tree dried biomass.

237 The effects of genotype and position of the sample in the branch or trunk wood on
238 element concentrations and quantity of TE exported were tested with a two-way analysis of
239 variance. All tests were considered significant when $p < 0.05$. Prior to analysis, outliers were
240 discarded, and the normality of the variable or of the residual values was tested with the
241 Shapiro-Wilk test, and homoscedasticity, with the Bartlett test. The design was a randomized
242 block design with genotype, branches and trunk as the fixed factors and plot as a random
243 factor. For the selected genotype Skado, the effect of sample position on element
244 concentration was tested for the branch, trunk wood and trunk bark.

245 **3. Results and discussion**

246 **3.1. Soil characteristics and short-term impact of the poplar plantation**

247 The pedogeochemical characteristics of the entire Pierrelaye area has been thoroughly
248 studied (Lamy et al., 2006; van Oort et al., 2008), and the authors concluded that irrigation
249 with wastewaters for more than 100 years has deeply modified the macromorphology and the
250 chemistry of the soil. The surface horizons of the Pierrelaye Luvisol have indeed accumulated
251 large amounts of organic matter and have been markedly enriched with TE. The total Zn, Pb,
252 Cu and Cd at the Pierrelaye site for the initial PHYTOPOP plots sampled in 2007 values were
253 20- to 317-fold higher than the natural pedogeochemical background values, as shown in
254 Table 1, and are in the range of the TE content and surface content spatial heterogeneity
255 recorded in that contaminated area (Lamy et al., 2006).

256
257 The impact of the poplar plantation on the TE pools and pH was first evaluated after 4
258 growing seasons for all genotype plots, by comparing the ammonium nitrate -TE extractable
259 fractions in 2007 and 2011, as well as the pH values (Table 2). Only the 3 major TE—Cd, Cu
260 and Zn—are provided (Pb values fell under the detection limits). The NH_4NO_3 -TE extractable
261 fraction decreased in all plots between the two sampling periods (Table 2), whereas the

262 DTPA-TE extractable fraction slightly increased for most poplar genotypes (Table S2). As the
263 data provided are from composite samples, statistical analysis could not be performed on this
264 dataset. The pH, which is an essential parameter in phytomanagement practices either evolved
265 slightly upward or remained unchanged.

266 **3.2. Longer-term impact of the poplar plantation**

267 Our work was further pursued on the poplar Skado plot only. This genotype was chosen
268 in 2011, together with the I214 genotype, for a new phytomanagement plantation at the same
269 site within the BIOFILTREE project (Ciadamidaro et al., 2017). Soil samples were collected
270 in 2013 and 2016, from which the pH and the CaCl₂-extractable element fractions were
271 measured. The 2007 and 2011 soil samples from under the Skado genotype were also
272 reanalyzed; these samples had been stored while dry, with the same extraction and analytical
273 methods used for the 2013 and 2016 soil samples as on earlier samples. The pH analysis after
274 7 and 10 years of growth under the Skado genotype confirmed the lack of impact of the
275 poplar trees on this key soil parameter (Fig. 1). A depletion of the CaCl₂-extractable fraction
276 for Cd, Cu, Mn, Ni and Zn was also evident between 2007 and 2016 (Fig. 1), which confirms
277 the first series of analyses (2007-2011) performed with the NH₄NO₃-extractant (Table 2) for
278 Cd, Cu and Zn. The Mn and Ni were not chosen as target elements in the analyses of the 2007
279 and 2011 soil samples. Other elements, such as K Mg, S and Si, increased significantly with
280 time, which confirmed some of our previous data (Foulon et al., 2016b). Other elements (Na,
281 Al, and Fe) followed the same trends; however, the changes were not statistically significant.

282 The depletion of the NH₄NO₃ (2007-2011, first series of analysis)- or CaCl₂ (2007-
283 2011-2013-2016, second series of analysis)-extractable TE fractions was not observed for the
284 major elements and was low in absolute terms but high in relative terms, especially for Mn.
285 Two main hypotheses could explain this decrease in the long term to confirm a depletion of
286 the soil exchangeable compartment, with both hypotheses involving processes attributed to

287 changes in TE speciation under the land use: i) an increase in leaching in the soil profile under
288 trees without replenishment of the exchangeable pool of TE or ii) an increase in retention in
289 the soil due to the so-called process of “natural attenuation.” In the first case, a leaching of TE
290 had been already observed in a soil profile of this area. But due to the constitution of the
291 Luvisols, mobile TE were found trapped in the Bt horizon mainly constituted of clay before
292 reaching the water table. Indeed, Dère et al (2007) observed that exogenous Zn and Cu but not
293 Pb were found down to the base of the studied solum (i.e. around 1 m), mostly readsorbed on
294 deep horizon’s constituents. In the second case, the TEs that are easily phytoavailable would
295 have been taken up by poplar roots. As no obvious differences were observed between
296 planted and unplanted poplar (naturally revegetated) plots (data not shown), it can be
297 hypothesized that the herbaceous layer under the poplar trees played a major role in changing
298 levels in the CaCl₂-extractable fraction. However, these depletions for Cd, Cu and Mn under
299 the adjacent maize plots was noticed in one of our previous studies (Foulon et al., 2016a),
300 which may indicate an overall phenomenon of land use with plant species efficient in the use
301 of atmospheric carbon, which then resulted in high amounts of the soil organics able to
302 change TE speciation rather than from the poplar having a specific impact on its underlying
303 vegetation. Indeed, the indirect effect of the newly developed vegetation (poplar trees or the
304 adjacent herbaceous layer), by regularly enriching the Pierrelaye soil with fresh organic
305 material other than from the wastewaters, could result in the alteration of the distribution of
306 TE in the various soil extractable fractions. Previous works have demonstrated the importance
307 of leaf litter on the nutrient cycles and bioavailability of TE (Ciadamidaro et al., 2013, 2014),
308 and hence on the element export capacity of trees. Litter or litter components represent the
309 main route of entry of nutrients in the soil and is one of the key points of the recycling of
310 organic matter (OM) and nutrients, and, in forest ecosystems, the main source of OM and
311 nutrients for the topsoil (Kuzyakov and Domanski, 2000). In particular, it can cover between

312 60 and 80% of the nutritional requirements of the tree per year. We did not attempt to
313 determine the TOC content of the soils under the various poplar clones, but future projects at
314 this experimental site would need to address this key feature in the management of TE
315 contaminated soils.

316 Some changes between 2007 and 2011 indeed occurred in the DTPA fraction in poplar
317 or unplanted (naturally revegetated) soils (Table S2). The small increase in the DTPA
318 fraction, which tends to mimic an increase in the available TE pool, is unlikely to be solely
319 responsible for the displacement of the extractable fraction, as these fractions were of much
320 greater magnitude, but it may have partially contributed to it.

321 The lack of change in the pH is an important indicator and could be associated with soil
322 stabilization. It has to consider, however, that such soils rich in organic carbon have a high
323 buffering effect against changes in pH, and data from a longer term could be useful for
324 drawing conclusions. Due to these buffering effects, changes in soil pH are less important
325 than those of TE availability seen either through DTPA, CaCl₂ or ammonium nitrate
326 extractions. But the decrease in salt-exchangeable PTE compared to the increase in DPTA-
327 extractable PTE is consistent with a change in PTE speciation in favor with PTE
328 complexation with freshly organic matter brought by the aboveground vegetation.

329 Nevertheless, such an indication is important to take into consideration when looking
330 for future management of the site, as a reforestation of the Pierrelaye area is now being
331 planned for the next few years. In a previous study, the soil pH under poplar stands of various
332 genotypes, ages and tree density did not differ significantly from adjacent arable fields (Weih
333 et al., 2003).

334 **3.3. Genetic variability of poplar productivity at Pierrelaye**

335 H_{tot} and dbh are frequently used as predictor variables, relating an easily measurable
336 parameter to that of a less obtainable dependent factor such as tree dry weight. Both

337 parameters are known to be strongly indicative of aboveground woody biomass (Morhart et
338 al., 2013; Zianis, 2008) and are part of the equations detailed in the material and methods
339 section. Individual genotypic allometric equations allowed establishing relationships between
340 the diameter measured at breast height (dbh) in 2013 and total dry biomasses of harvested
341 trunk and branches respectively, for the 14 poplar genotypes grown at the Pierrelaye site
342 (Fig. 2). Although the equations gave similar curves, the differences between the poplar
343 genotypes was evident and showed that among the poplar genotypes retained in the present
344 study, some variability occurred in the dry weight distribution between trunk and branches
345 and/or wood density.

346 The significance of our datasets was further tested (Table 3). This analysis allowed us to
347 conclude that the genotype effect was significant for the trunkDW parameter, whereas for the
348 branchDW parameter, the regression coefficients were not statistically significant. As a
349 consequence, specific equations for each genotype were used to calculate the trunkDW,
350 whereas a single equation for all genotypes was used to calculate the branchDW. The
351 proportion of branchDW in the total biomass was significantly correlated ($r = 0.9$) with the
352 “b” regression coefficients of the trunkDW equations, indicating that these different equations
353 reflected the architectural variability within a genotype. These data further allowed us to
354 estimate the productivity (expressed as oven dry tons or odt $\text{ha}^{-1} \text{year}^{-1}$) of each genotype,
355 which was estimated in 2013 (7-year-old plantation).

356 Over the entire 7-year rotation, huge genotypic variability was observed, with the
357 Vesten genotype producing the largest biomass ($8.5 \text{ odt ha}^{-1} \text{year}^{-1}$) in the SRC production
358 scheme, whereas the Dorskamp genotype produced 2.8 times less biomass (Fig. 3). In Europe,
359 the mean modeled yields for poplars ranged between 1.5 and $11.7 \text{ odt ha}^{-1} \text{year}^{-1}$ for Aspen
360 (Aylott et al., 2008; Johansson, 1999; Morhart et al., 2013; Verlinden et al., 2013). There was
361 no clear relationship between the genetic origin and production potential. The Bakan and

362 Skado (TxM) genotypes showed a similar yield of approximately 7 odt ha⁻¹ year⁻¹ and were
363 among the most productive genotypes, whereas the DxN genotypes exhibited large variations
364 ranging from 3.1 (Dorskamp) to 8.5 (Vesten) odt ha⁻¹ year⁻¹. Our data contrast with those
365 from a previous study (Verlinden et al., 2013), where clustering of the poplar genotypes was
366 clearly determined by genetic origin, with the poplar plantation being established on a normal,
367 uncontaminated soil. Interestingly, in that study, the Bakan and Skado genotypes showed very
368 similar biomass production to that of our study, approximately 5.75 odt ha⁻¹ year⁻¹. The
369 Vesten genotype in our study was more productive than that in the Verlinden study
370 (Verlinden et al., 2013). The trees were, however, harvested after only 2 growing seasons in
371 this previous study. The Dorskamp genotype showed notably lower biomass productivity in
372 our study than the other 13 genotypes. However, this genotype experienced more rabbit
373 feeding than the other genotypes, and the mortality was much higher for that genotype (Table
374 S1). A high variability in growth and in cutting quality (especially cutting diameter) during
375 the first years is likely due to weed pressure and management practices (Broeckx et al., 2012).
376 However, at our site, all plots received the same appropriate treatment for weed management,
377 and cutting diameters were homogenous between all genotypes that were planted.

378 The dry branch proportions ranged from 26 to 40% of the total tree biomass (Fig. 3).
379 The highest proportions were found for the D (Dvina and Lena) and DxN (Triplo and Vesten)
380 poplar genotypes, whereas the four T (Trichobel and Fritzi Pauley) or TxM (Bakan and
381 Skado) genotypes exhibited the lowest branch proportions. The branch proportion in the TxM
382 genotypes is close to that reported (22.2%) for another genotype (hybrid 275) of the same
383 hybrid family (Morhart et al., 2013) grown in standard conditions. The quantification of
384 branch dry biomass is an important factor, as it may greatly influence the overall quality of
385 the final product. This is even more important in the context of phytomanagement, as the
386 proportion of bark increased with decreasing diameter (see results below and Fig. 4).

387 **3.4. Macronutrient concentrations in wood and branches of poplar genotypes**

388 Macronutrient concentrations (expressed as mg kg⁻¹ DW) were determined in poplar
389 branches and bark-removed trunks (wood) after seven growing seasons (Table 4).
390 Macronutrients were measured at three heights (summit, middle, and base), as described in
391 the material and methods section, and the data provided are the averages of samples from the
392 three sections.

393 Overall, branches accumulated more Ca, K, Mg, and P than woody tissues (Table 4).
394 The highest Ca concentrations in branches were measured for Bakan, which was 1.8 times
395 significantly higher than for Vesten. This ratio was 1.6 for the wood of Dorskamp compared
396 to that of Triplo (Table 4). The highest ratio for K in the wood was 1.4 between the Trichobel
397 and Koster genotypes, when there was no effect of the genotype on the K concentration
398 (ANOVA) in the branches (Table 4). The Mg concentration was about 1.7 times higher in the
399 branches of Dvina or Triplo compared to Trichobel or Vesten. The Dvina, Dorskamp, and
400 Soligo genotypes also had the highest Mg levels in their wood, and significantly higher than
401 in Skado or Bakan, for example (Table 4). For most poplar genotypes, the K and P
402 concentrations were the highest in the upper sections of the trunk (not shown). For instance, K
403 and P doubled in the summit sections of the Bakan and Fritzi Pauley trunks compared to the
404 basal sections.

405 According to a study of Obernberger et al. (2006), typical woody biomasses from
406 deciduous species (from uncontaminated soils) contained, on average, 1200 mg kg⁻¹ Ca, 800
407 mg kg⁻¹ K, 200 mg kg⁻¹ Mg, and 100 mg kg⁻¹ P for the major elements. The K, Mg, and P
408 concentrations were, on average, 1.5 to 5-fold higher than those given by Obernberger et al.
409 (2006), whereas Ca concentrations were in the same range.

410 **3.5. Trace element concentrations in trunk and branches of poplar genotypes**

411 The Bakan and Dorskamp genotypes accumulated significantly more Cd and/or Zn than
412 the Fritzi Pauley, Triplo and Lena genotypes, both in the wood (Table 5) and in the branches
413 (Table 6). The largest variations measured for these two TE were of the order of two times.
414 For Cu, the highest values were measured for Dorskamp, they were significantly different
415 from the Cu values measured for Fritzi Pauley. There were no significant variations in Cr and
416 Pb concentrations in wood between genotypes (Table 5). Cr and Pb values were the lowest of
417 all elements mentioned. For Fe, the highest levels were measured for the Lena genotype, 2.6
418 and 2.3 times higher in the wood (Table 5) and branches (Table 6), respectively, when
419 compared to the I214 genotype. Mn accumulated the most in the branches of Dvina, Flevo
420 and Dorskamp and in the wood of Dvina, Koster, Bakan, while the Fritzi Pauley, I214 or
421 Trichobel genotypes accumulated significantly 1.8 (wood) to 2.7 (branches) times less.

422 Even if it is difficult to obtain clear trends for all elements, it should be noted that the
423 Dorskamp, Bakan and Dvina genotypes often appeared as the most accumulating genotypes
424 (Mn, Cu, Fe, Zn, and Cd) for wood or branches. On the contrary genotypes such as Fritzi
425 Pauley, Trichobel or Triplo were more often quoted as the less accumulating genotypes.
426 Although the Fritzi Pauley and Trichobel genotypes belong to the same species, the
427 Dorskamp, Bakan and Dvina genotypes belong to three different species (Table S1). It is
428 therefore difficult to establish a link between the accumulation of elements and the taxonomic
429 relationships of the genotypes.

430 Branches accumulated more Cd, Cu, Fe, Mn and Zn than woody tissues (Tables 5 and
431 6). Pb and Cr concentrations were within narrow ranges, and no clear significant tendency
432 was observed between the wood and branches. The largest variations between branches and
433 woods were observed for Mn, with an order of magnitude difference observed for most
434 genotypes. The major contaminants Cu, Zn and Cd were 3- to 4-fold higher in branches

435 compared with wood. For most poplar genotypes, the TE concentrations were the highest in
436 the upper sections of the trunk (not shown). For instance, Fe doubled in the summit sections
437 of the Bakan and Fritzi Pauley trunks compared to the basal sections. The Pb concentration
438 was always the highest in the basal section of the trunk, probably reflecting a simple
439 contamination by aerosols, rather than an active transport, as already detailed in Migeon et al.
440 (2009). No such clear tendency was observed for the other elements.

441 Metal content in woody biomass can greatly affect the industrial process, as it may
442 interfere with the thermal degradation by acting as catalysts. A previous paper reported for
443 instance that Cl ions may change metal behavior by increasing metal volatilization (Bert et al.,
444 2017). For TE, 0.1 mg kg⁻¹ Cd, 1 mg kg⁻¹ Cr, 2 mg kg⁻¹ Cu, 25 mg kg⁻¹ Fe, 83 mg kg⁻¹ Mn, 2
445 mg kg⁻¹ Pb, and 10 mg kg⁻¹ Zn are the concentrations usually reported (Oberberger et al.,
446 2006). In our study, Cu, Cr, Fe, Mn and Pb concentrations in poplar wood, which range
447 between 0.1 and 3 mg kg⁻¹, are so low that they will not be discussed further herein. Cd
448 concentrations were also very low in wood (< 0.5 mg kg⁻¹); however, the volatile nature of Cd
449 and its potential dissemination into the atmosphere when the biomass is burned should be
450 considered (Chalot et al., 2012; Evangelou et al., 2012). When compared to the present
451 dataset, Zn was accumulated at a much higher rate in a previous study, although large
452 variations were observed among the poplar genotypes (Castiglione et al., 2009). The total Zn
453 concentration (879-950 mg kg⁻¹) was also higher compared to our soil data (Table 1). The
454 authors also did not mention whether bark was included or not in the stem analysis.

455 In a previous paper (Pottier et al., 2015), data on the leaf concentrations were provided
456 from the same trees and on changes during seasons. These previous analyses demonstrated
457 large variations in the TE concentrations of leaf tissues among the various genotypes.
458 Phytomanagement of TE-contaminated sites strongly relies on the capacity of the plant cover
459 to tolerate high concentrations of TE. Salicaceous species respond to that constraint by

460 storing TE in their aboveground biomass (Migeon et al., 2009), with potential sequestration of
461 TEs within cells in the organelles. In poplar, this is illustrated by the sequestration of Mn in a
462 Golgi-associated compartment through the transport protein PtMPT11 (Peiter et al., 2007) and
463 by the vacuolar sequestration of Zn through the transport protein PtMTP1 (Blaudez et al.,
464 2003). The potential function of other target transport genes were further underscore in TE
465 management (Pottier et al., 2015) and their potential use in phytomanagement using *Populus*
466 species. *PtNRAMP1* may, for instance, represent an interesting target for genetic engineering
467 of poplar and modulation of leaf TE content.

468 **3.6. Element accumulation in bark tissues**

469 Given the overall contribution of the bark tissues in element distribution within the
470 whole tree, our study was further focused on this aspect on the Skado genotype. Bark was
471 separated from the wood at different heights of the trunk and compared to branches. The bark
472 content averaged 10-12% of dry trunk biomass, whereas branches < 2 cm in diameter are
473 more than 30% bark (Fig. 4).

474 Element concentrations were further determined in bark collected on the trunk from
475 Skado (Table 7). Bark accumulated 2 to 14 times more Ca, Fe, Mn, and Zn than debarked-
476 wood. Differences were also observed for Cd, K and Mg, but to a lesser extent. Other
477 elements such as Cr, Cu and Pb did not accumulate differently in bark and wood, with ratios
478 close to one. Very similar ratios between wood and bark were also measured in a field
479 experiment where poplar were grown on TE-contaminated soils (Evangelou et al., 2012).

480 In a previous study, the authors concluded that the quality of the final product is
481 strongly impacted by the amount of bark in the biomass feedstock (Morhart et al., 2013). As
482 wood chips from SRC are being used as recycling materials to enhance soil fertility, concern
483 is growing over the fate of TE from bark tissues (Vandecasteele et al., 2013). This is even
484 more meaningful for biomass harvested at phytomanagement sites, where the presence of TE

485 in the bark tissues may have an impact on the industrial processes. Evangelou and
486 collaborators (Evangelou et al., 2012) also concluded that harvesting biomass with smaller
487 bark proportion in relation to trunk is desirable.

488

489 **3.7. Element export by poplar genotypes**

490 Given the biomass production data and the element contents, the amount of exported
491 element (expressed as $\text{kg}^{-1} \text{ha}^{-1} \text{year}^{-1}$ or $\text{g}^{-1} \text{ha}^{-1} \text{year}^{-1}$) was calculated after 7 growing
492 seasons. The most productive genotypes facilitated the highest export of elements (Table 8).
493 As a consequence of the variation in the element accumulations and biomass produced, the
494 export capacities were highly variable among the genotypes. The Bakan, Triplo and Vesten
495 genotypes showed higher export capacities, with Ca and Cd export greater than $20 \text{ kg}^{-1} \text{ha}^{-1}$
496 year^{-1} and $3 \text{ g}^{-1} \text{ha}^{-1} \text{year}^{-1}$, respectively. Conversely, the Flevo and Dorskamp genotypes
497 exported much less of these elements. These findings were not related to the *Populus* species,
498 as the Bakan and Triplo, for instance, do not belong to the same species (Table S1).

499 The ability to export these elements is an important consideration in phytomanagement,
500 as the concentration of contaminants in the harvested part will greatly determine their
501 potential valorization. Additionally, the export of nutrients is also a key consideration. The
502 export potential was clearly greater for branches for Ca, Mn and Fe and to a lesser extent for
503 K, P, Mg, Zn, Cd compared with the trunk. This difference was directly linked to the higher
504 bark content of branches and higher content in these elements (Table 7). Conversely, wood
505 tissues showed greater Cu, Cr and Pb export capacities, related to their equal amounts in
506 wood and bark samples (Table 8). Previous studies have suggested that harvesting poplar as
507 the whole tree from short-term rotations should cause soil nutrient depletion and induce
508 degradation of soil and decreased biomass production (Ge et al., 2015). However, such a
509 decrease in soil nutrient contents was not observed at our site when comparing total (not
510 shown) or ammonium nitrate-extractable (Fig. 1) fractions of major elements in the 2007 and

511 2011, 2013 and 2016 samplings, perhaps because their levels were already high enough. The
512 enrichment of the Pierrelaye soils with many cations, such as K and Mg, in poplar plots may
513 indicate a thorough mobilization that occurs through microbial activities that are stimulated
514 by the presence of poplar trees (Foulon et al., 2016b, 2016a). Conversely, whether these
515 microbial communities are indeed expressing the capacity to mobilize these elements from the
516 bound fraction is unknown.

517

518 **4. Conclusions**

519 Our set of data clearly emphasizes genotypic variations in the growth of poplar at a
520 moderately contaminated site, as well as in element accumulation, and hence in nutrient
521 export capacities. The present dataset highlighted that soil element depletion occurred for a
522 range of TE, whereas the soil content of macronutrients remained unchanged or increased
523 after 10 years of poplar growth. Given i) the greater TE content of bark tissues compared with
524 wood and ii) the higher proportion of bark in branches compared with trunk, it can be
525 recommended choosing stem wood-only harvesting instead of whole-tree harvesting, which
526 will allow the risks encountered with TE-enriched biomass in the valorization process to be
527 reduced. In the long term, this approach may also allow only limited nutrient losses. However,
528 the impact of element-enriched remnants on physicochemical and biological soil properties
529 requires further investigation. Future research programs should also expand this type of
530 investigation based on the use of various genotypes on a broader range of site conditions to
531 further enhance the uptake of phytotechnologies by stakeholders.

532 **5. Acknowledgements**

533 This work was supported by the Agence Nationale de la Recherche (ANR) under the
534 PRECODD (ANR-06-ECOT-O15-01-PHYTOPOP) and the Blanc International (ANR-10-
535 INTB-1703-01-BIOFILTREE) projects.

536 **6. References**

- 537 Ancona, V., Barra Caracciolo, A., Campanale, C., De Caprariis, B., Grenni, P., Uricchio,
538 V.F., Borello, D., 2019. Gasification treatment of poplar biomass produced in a
539 contaminated area restored using plant assisted bioremediation. *J. Environ. Manage.*
540 239, 137–141. <https://doi.org/10.1016/j.jenvman.2019.03.038>
- 541 Asad, M., Menana, Z., Ziegler-Devin, I., Bert, V., Chalot, M., Herzig, R., Mench, M., Brosse,
542 N., 2017. Pretreatment of trace element-enriched biomasses grown on phytomanaged
543 soils for bioethanol production. *Ind. Crops Prod.* 107, 63–72.
544 <https://doi.org/10.1016/j.indcrop.2017.05.028>
- 545 Aylott, M.J., Casella, E., Tubby, I., Street, N.R., Smith, P., Taylor, G., 2008. Yield and spatial
546 supply of bioenergy poplar and willow short-rotation coppice in the UK. *New Phytol.*
547 178, 358–370. <https://doi.org/10.1111/j.1469-8137.2008.02396.x>
- 548 Baize, D., Lamy, I., vanOort, F., Dère, C., Chaussod, R., Sappin-Didier, V., Bermond, A.,
549 Bourgeois S., Schmitt, C., and Schwartz, C. 2002. 100 years spreading of urban
550 wastewater on market- garden soils close to Paris (France): subsequent impacts and
551 hazards. *CDRom Transactions 17th World Congress of Soil Science, Bangkok,*
552 *Thailand, CDRom paper 204, pp 204-1-204-10.*
- 553 Baize, D., 2003. Modélisation empirique du transfert du cadmium et du zinc des sols vers les
554 grains de blé tendre. *Étude Gest. Sols* 22.
- 555 Bert, V., Allemon, J., Sajet, P., Dieu, S., Papin, A., Collet, S., Gaucher, R., Chalot, M.,
556 Michiels, B., Raventos, C., 2017. Torrefaction and pyrolysis of metal-enriched poplars
557 from phytotechnologies: Effect of temperature and biomass chlorine content on metal
558 distribution in end-products and valorization options. *Biomass Bioenergy* 96, 1–11.
559 <https://doi.org/10.1016/j.biombioe.2016.11.003>

560 Blaudez, D., Kohler, A., Martin, F., Sanders, D., Chalot, M., 2003. Poplar metal tolerance
561 protein 1 confers zinc tolerance and is an oligomeric vacuolar zinc transporter with an
562 essential leucine zipper motif. *Plant Cell* 15, 2911–2928.
563 <https://doi.org/10.1105/tpc.017541>

564 Broeckx, L.S., Verlinden, M.S., Ceulemans, R., 2012. Establishment and two-year growth of
565 a bio-energy plantation with fast-growing *Populus* trees in Flanders (Belgium): Effects
566 of genotype and former land use. *Biomass Bioenergy* 42, 151–163.
567 <https://doi.org/10.1016/j.biombioe.2012.03.005>

568 Castiglione, S., Todeschini, V., Franchin, C., Torrigiani, P., Gastaldi, D., Cicatelli, A.,
569 Rinaudo, C., Berta, G., Biondi, S., Lingua, G., 2009. Clonal differences in survival
570 capacity, copper and zinc accumulation, and correlation with leaf polyamine levels in
571 poplar: A large-scale field trial on heavily polluted soil. *Environ. Pollut.* 157, 2108–
572 2117. <https://doi.org/10.1016/j.envpol.2009.02.011>

573 Chalot, M., Blaudez, D., Rogaume, Y., Provent, A.-S., Pascual, C., 2012. Fate of trace
574 elements during the combustion of phytoremediation wood. *Environ. Sci. Technol.* 46,
575 13361–13369. <https://doi.org/10.1021/es3017478>

576 Ciadamidaro, L., Madejon, P., Puschenreiter, M., Madejon, E., 2013. Growth of *Populus alba*
577 and its influence on soil trace element availability. *Sci. Total Environ.* 454-455, 337-
578 347. <http://dx.doi.org/10.1016/j.scitotenv.2013.03.032>

579 [Ciadamidaro, L., Madejon, E., Robinson, B., Madejon, P., 2014. Soil plant interactions of](https://doi.org/10.1016/j.jenvman.2013.11.010)
580 [*Populus alba* in contrasting environments. *J. Environ. Manag.* 132, 329-337.](https://doi.org/10.1016/j.jenvman.2013.11.010)
581 <https://doi.org/10.1016/j.jenvman.2013.11.010>

582 Ciadamidaro, L., Girardclos, O., Bert, V., Zappelini, C., Yung, L., Foulon, J., Papin, A., Roy,
583 S., Blaudez, D., Chalot, M., 2017. Poplar biomass production at phytomanagement

584 sites is significantly enhanced by mycorrhizal inoculation. *Environ. Exp. Bot.* 139,
585 48–56. <https://doi.org/10.1016/j.envexpbot.2017.04.004>

586 Cicutelli, A., Lingua, G., Todeschini, V., Biondi, S., Torrigiani, P., Castiglione, S., 2010.
587 Arbuscular mycorrhizal fungi restore normal growth in a white poplar clone grown on
588 heavy metal-contaminated soil, and this is associated with upregulation of foliar
589 metallothionein and polyamine biosynthetic gene expression. *Ann. Bot.* 106, 791–802.
590 <https://doi.org/10.1093/aob/mcq170>

591 Conesa, H.M., Evangelou, M.W.H., Robinson, B.H., Schulin, R., 2012. A critical view of
592 current state of phytotechnologies to remediate soils: still a promising tool? *Sci. World*
593 *J.* 2012, 1–10. <https://doi.org/10.1100/2012/173829>

594 Cundy, A.B., Bardos, R.P., Puschenreiter, M., Mench, M., Bert, V., Friesl-Hanl, W., Müller,
595 I., Li, X.N., Weyens, N., Witters, N., Vangronsveld, J., 2016. Brownfields to green
596 fields: Realising wider benefits from practical contaminant phytomanagement
597 strategies. *J. Environ. Manage.* 184, 67–77.
598 <https://doi.org/10.1016/j.jenvman.2016.03.028>

599 Dastyar, W., Raheem, A., He, J., Zhao, M., 2019. Biofuel production using thermochemical
600 conversion of heavy metal-contaminated biomass (HMCB) harvested from
601 phytoextraction process. *Chem. Eng. J.* 358, 759–785.
602 <https://doi.org/10.1016/j.cej.2018.08.111>

603 De Oliveira, V.H., Tibbett, M., 2018. Tolerance, toxicity and transport of Cd and Zn in
604 *Populus trichocarpa*. *Environ. Exp. Bot.* 155, 281–292.
605 <https://doi.org/10.1016/j.envexpbot.2018.07.011>

606 Delplanque, M., Collet, S., Del Gratta, F., Schnuriger, B., Gaucher, R., Robinson, B., Bert,
607 V., 2013. Combustion of *Salix* used for phytoextraction: The fate of metals and

608 viability of the processes. *Biomass Bioenergy* 49, 160–170.
609 <https://doi.org/10.1016/j.biombioe.2012.12.026>

610 Dère C., Lamy I, Jaulin A., Cornu S. 2007. Long-term fate of exogenous metals in a sandy
611 Luvisol subjected to intensive irrigation with raw wastewater. *Environmental*
612 *Pollution*, 145, 31-40.

613 Dimitriou, I., Aronsson, P., 2011. Wastewater and sewage sludge application to willows and
614 poplars grown in lysimeters—Plant response and treatment efficiency. *Biomass*
615 *Bioenergy* 35, 161–170. <https://doi.org/10.1016/j.biombioe.2010.08.019>

616 Dimitriou, I., Aronsson, P., 2010. Landfill leachate treatment with willows and poplars –
617 Efficiency and plant response. *Waste Manag.* 30, 2137–2145.
618 <https://doi.org/10.1016/j.wasman.2010.06.013>

619 Evangelou, M.W.H., Deram, A., Gogos, A., Studer, B., Schulin, R., 2012. Assessment of
620 suitability of tree species for the production of biomass on trace element contaminated
621 soils. *J. Hazard. Mater.* 209–210, 233–239.
622 <https://doi.org/10.1016/j.jhazmat.2012.01.008>

623 FAO, 2016. Poplars and other fast-growing trees - renewable resources for future green
624 economies. publications listed in country progress reports. 25th Session of the
625 International Poplar Commission, Berlin, Germany, 13-16 September 2016.
626 International Poplar Commission Working Paper IPC/16. FAO, Rome.

627 Fedorov, V., 1972. *Theory of optimal experiments*. Academic Press, San Diego, Calif, USA,
628 3rd edition.

629 Foulon, J., Zappellini, C., Durand, A., Valot, B., Blaudez, D., Chalot, M., 2016a. Impact of
630 poplar-based phytomanagement on soil properties and microbial communities in a
631 metal-contaminated site. *FEMS Microbiol. Ecol.* 92, fiw163.
632 <https://doi.org/10.1093/femsec/fiw163>

633 Foulon, J., Zappellini, C., Durand, A., Valot, B., Girardclos, O., Blaudez, D., Chalot, M.,
634 2016b. Environmental metabarcoding reveals contrasting microbial communities at
635 two poplar phytomanagement sites. *Sci. Total Environ.* 571, 1230–1240.
636 <https://doi.org/10.1016/j.scitotenv.2016.07.151>

637 French, C.J., Dickinson, N.M., Putwain, P.D., 2006. Woody biomass phytoremediation of
638 contaminated brownfield land. *Environ. Pollut.* 141, 387–395.
639 <https://doi.org/10.1016/j.envpol.2005.08.065>

640 Gamalero, E., Cesaro, P., Cicatelli, A., Todeschini, V., Musso, C., Castiglione, S., Fabiani,
641 A., Lingua, G., 2012. Poplar clones of different sizes, grown on a heavy metal
642 polluted site, are associated with microbial populations of varying composition. *Sci.*
643 *Total Environ.* 425, 262–270. <https://doi.org/10.1016/j.scitotenv.2012.03.012>

644 Ge, X., Tian, Y., Tang, L., 2015. Nutrient distribution indicated whole-tree harvesting as a
645 possible factor restricting the sustainable productivity of a poplar plantation system in
646 China. *PLOS ONE* 10, e0125303. <https://doi.org/10.1371/journal.pone.0125303>

647 Gelfand, I., Sahajpal, R., Zhang, X., Izaurralde, R.C., Gross, K.L., Robertson, G.P., 2013.
648 Sustainable bioenergy production from marginal lands in the US Midwest. *Nature*
649 493, 514.

650 Gerhardt, K.E., Gerwing, P.D., Greenberg, B.M., 2017. Opinion: Taking phytoremediation
651 from proven technology to accepted practice. *Plant Sci.* 256, 170–185.
652 <https://doi.org/10.1016/j.plantsci.2016.11.016>

653 Johansson, T., 1999. Biomass equations for determining fractions of European aspen growing
654 on abandoned farmland and some practical implications. *Biomass Bioenergy* 10.

655 Kuzyakov, I., Domanski, G., 2000. Carbon input by plants into the soil. Review. *J. Plant*
656 *Nutrit. Soil Sci.* 163, 421–431. [http://doi.org/10.1002/1522-
657 2624\(200008\)163:4<421::AID-JPLN421>3.0.CO;2-R](http://doi.org/10.1002/1522-2624(200008)163:4<421::AID-JPLN421>3.0.CO;2-R)

658 Lamy, I., van Oort, F., Dère, C., Baize, D., 2006. Use of major- and trace-element correlations
659 to assess metal migration in sandy Luvisols irrigated with wastewater. *Eur. J. Soil Sci.*
660 57, 731–740. <https://doi.org/10.1111/j.1365-2389.2005.00765.x>

661 Langer, I., Santner, J., Krpata, D., Fitz, W.J., Wenzel, W.W., Schweiger, P.F., 2012.
662 Ectomycorrhizal impact on Zn accumulation of *Populus tremula* L. grown in
663 metalliferous soil with increasing levels of Zn concentration. *Plant Soil* 355, 283–297.
664 <https://doi.org/10.1007/s11104-011-1098-y>

665 Laureysens, I., Blust, R., De Temmerman, L., Lemmens, C., Ceulemans, R., 2004. Clonal
666 variation in heavy metal accumulation and biomass production in a poplar coppice
667 culture: I. Seasonal variation in leaf, wood and bark concentrations. *Environ. Pollut.*
668 131, 485–494. <https://doi.org/10.1016/j.envpol.2004.02.009>

669 Laureysens, I., De Temmerman, L., Hastir, T., Van Gysel, M., Ceulemans, R., 2005. Clonal
670 variation in heavy metal accumulation and biomass production in a poplar coppice
671 culture. II. Vertical distribution and phytoextraction potential. *Environ. Pollut.* 133,
672 541–551. <https://doi.org/10.1016/j.envpol.2004.06.013>

673 Lettens, S., Vandecasteele, B., De Vos, B., Vansteenkiste, D., Verschelde, P., 2011. Intra- and
674 inter-annual variation of Cd, Zn, Mn and Cu in foliage of poplars on contaminated
675 soil. *Sci. Total Environ.* 409, 2306–2316.
676 <https://doi.org/10.1016/j.scitotenv.2011.02.029>

677 Lingua, G., Franchin, C., Todeschini, V., Castiglione, S., Biondi, S., Burlando, B.,
678 Parravicini, V., Torrigiani, P., Berta, G., 2008. Arbuscular mycorrhizal fungi
679 differentially affect the response to high zinc concentrations of two registered poplar
680 clones. *Environ. Pollut.* 153, 137–147. <https://doi.org/10.1016/j.envpol.2007.07.012>

681 Madejon, E., Pérez-de-Mora, A., Felipe, E., Burgos, P., Cabrera, F., 2006. Soil amendments
682 reduce trace element solubility in a contaminated soil and allow regrowth of natural
683 vegetation. *Environ. Pollut.* 139, 40-52. [http://doi.org/ 10.1016/j.envpol.2005.04.034](http://doi.org/10.1016/j.envpol.2005.04.034)

684 Marmioli, M., Pietrini, F., Maestri, E., Zacchini, M., Marmioli, N., Massacci, A., 2011.
685 Growth, physiological and molecular traits in *Salicaceae* trees investigated for
686 phytoremediation of heavy metals and organics. *Tree Physiol.* 31, 1319–1334.
687 <https://doi.org/10.1093/treephys/tpr090>

688 McLaughlin, M.J., Hamon, R.E., McLaren, R.G., Speir, T.W., Rogers, S.L., 2000. Review: A
689 bioavailability-based rationale for controlling metal and metalloid contamination of
690 agricultural land in Australia and New Zealand. *Austr. J. Soil Res.* 38, 1037-1086.
691 [http://doi.org/ 10.1071/SR99128](http://doi.org/10.1071/SR99128)

692 Mertens, J., Van Nevel, L., De Schrijver, A., Piesschaert, F., Oosterbaan, A., Tack, F.M.G.,
693 Verheyen, K., 2007. Tree species effect on the redistribution of soil metals. *Environ.*
694 *Pollut.* 149, 173–181. <https://doi.org/10.1016/j.envpol.2007.01.002>

695 Migeon, A., Blaudez, D., Wilkins, O., Montanini, B., Campbell, M.M., Richaud, P., Thomine,
696 S., Chalot, M., 2010. Genome-wide analysis of plant metal transporters, with an
697 emphasis on poplar. *Cell. Mol. Life Sci.* 67, 3763–3784.
698 <https://doi.org/10.1007/s00018-010-0445-0>

699 Migeon, A., Richaud, P., Guinet, F., Blaudez, D., Chalot, M., 2012. Hydroponic screening of
700 poplar for trace element tolerance and accumulation. *Int. J. Phytoremediation* 14, 350–
701 361. <https://doi.org/10.1080/15226514.2011.620651>

702 Migeon, A., Richaud, P., Guinet, F., Chalot, M., Blaudez, D., 2009. Metal accumulation by
703 woody species on contaminated sites in the north of France. *Water. Air. Soil Pollut.*
704 204, 89–101. <https://doi.org/10.1007/s11270-009-0029-5>

705 Morhart, C., Sheppard, J., Spiecker, H., 2013. Above ground leafless woody biomass and
706 nutrient content within different compartments of a *P. maximowiczii* × *P. trichocarpa*
707 Poplar Clone. *Forests* 4, 471–487. <https://doi.org/10.3390/f4020471>

708 Navarro, A., Facciotto, G., Campi, P., Mastrorilli, M., 2014. Physiological adaptations of five
709 poplar genotypes grown under SRC in the semi-arid Mediterranean environment.
710 *Trees* 28, 983–994. <https://doi.org/10.1007/s00468-014-1012-3>

711 Obernberger, I., Brunner, T., Barnthaler, G., 2006. Chemical properties of solid biofuels—
712 significance and impact. *Biomass Bioenergy* 30, 973–982.
713 <https://doi.org/10.1016/j.biombioe.2006.06.011>

714 Pandey, V.C., Bajpai, O., Singh, N., 2016. Energy crops in sustainable phytoremediation.
715 *Renew. Sustain. Energy Rev.* 54, 58–73. <https://doi.org/10.1016/j.rser.2015.09.078>

716 Peiter, E., Montanini, B., Gobert, A., Pedas, P., Husted, S., Maathuis, F.J.M., Blaudez, D.,
717 Chalot, M., Sanders, D., 2007. A secretory pathway-localized cation diffusion
718 facilitator confers plant manganese tolerance. *Proc. Natl. Acad. Sci. U. S. A.* 104,
719 8532–8537. <https://doi.org/10.1073/pnas.0609507104>

720 Pérez-de-Mora, A., Madejon, E., Burgos, P., Cabrera, F., 2006. Trace element availability and
721 plant growth in a mine-spill contaminated soil under assisted natural remediation I.
722 *Soils. Sci. Total Environ.* 363, 28–37. <http://doi.org/10.1016/j.scitotenv.2005.10.015>

723 Picard, N., Saint-André, L., Henry, M., 2012. Manuel de construction d'équations
724 allométriques pour l'estimation du volume et la biomasse des arbres De la mesure de
725 terrain à la prédiction. Organisation des Nations Unies pour l'alimentation et
726 l'agriculture (FAO), Rome, Italy.

727 Polle, A., Douglas, C., 2010. The molecular physiology of poplars: paving the way for
728 knowledge-based biomass production: Editorial. *Plant Biol.* 12, 239–241.
729 <https://doi.org/10.1111/j.1438-8677.2009.00318.x>

730 Pottier, M., García de la Torre, V.S., Victor, C., David, L.C., Chalot, M., Thomine, S., 2015.
731 Genotypic variations in the dynamics of metal concentrations in poplar leaves: A field
732 study with a perspective on phytoremediation. *Environ. Pollut.* 199, 73–82.
733 <https://doi.org/10.1016/j.envpol.2015.01.010>

734 Pruess, A., 1998. Action values for mobile (NH₄NO₃-extractable) trace elements in soils
735 based on the German national standard DIN 19730. Presented at the Proceedings of
736 the Third International Conference on Biogeochemistry of Trace Elements, Paris.

737 R Development Core Team, 2015. R: a Language and Environment for Statistical Computing.
738 R Foundation for Statistical Computing, Vienna Austria.

739 Rohde, V., Hahn, T., Wagner, M., Böringer, S., Tübke, B., Brosse, N., Dahmen, N.,
740 Schmiedl, D., 2018. Potential of a short rotation coppice poplar as a feedstock for
741 platform chemicals and lignin-based building blocks. *Ind. Crops Prod.* 123, 698–706.
742 <https://doi.org/10.1016/j.indcrop.2018.07.034>

743 Sabatti, M., Fabbrini, F., Harfouche, A., Beritognolo, I., Mareschi, L., Carlini, M., Paris, P.,
744 Scarascia-Mugnozza, G., 2014. Evaluation of biomass production potential and
745 heating value of hybrid poplar genotypes in a short-rotation culture in Italy. *Ind. Crops*
746 *Prod.* 61, 62–73. <https://doi.org/10.1016/j.indcrop.2014.06.043>

747 Šyc, M., Pohořelý, M., Kameníková, P., Habart, J., Svoboda, K., Punčochář, M., 2012.
748 Willow trees from heavy metals phytoextraction as energy crops. *Biomass Bioenergy*
749 37, 106–113. <https://doi.org/10.1016/j.biombioe.2011.12.025>

750 van Oort, F., Jongmans, A.G., Lamy, I., Baize, D., Chevallier, P., 2008. Impacts of long-term
751 waste-water irrigation on the development of sandy Luvisols: consequences for metal
752 pollutant distributions. *Eur. J. Soil Sci.* 59, 925–938. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2389.2008.01047.x)
753 [2389.2008.01047.x](https://doi.org/10.1111/j.1365-2389.2008.01047.x)

754 Vandecasteele, B., Willekens, K., Zwertvaegher, A., Degrande, L., Tack, F.M.G., Du Laing,
755 G., 2013. Effect of composting on the Cd, Zn and Mn content and fractionation in
756 feedstock mixtures with wood chips from a short-rotation coppice and bark. *Waste*
757 *Manag.* 33, 2195–2203. <https://doi.org/10.1016/j.wasman.2013.06.014>

758 Vangronsveld, J., Herzig, R., Weyens, N., Boulet, J., Adriaensen, K., Ruttens, A., Thewys, T.,
759 Vassilev, A., Meers, E., Nehnevajova, E., van der Lelie, D., Mench, M., 2009.
760 Phytoremediation of contaminated soils and groundwater: lessons from the field.
761 *Environ. Sci. Pollut. Res.* 16, 765–794. <https://doi.org/10.1007/s11356-009-0213-6>

762 Vega-Nieva, D.J., Valero, E., Picos, J., Jiménez, E., 2015. Modeling the above and
763 belowground biomass of planted and coppiced *Eucalyptus globulus* stands in NW
764 Spain. *Ann. For. Sci.* 72, 967–980. <https://doi.org/10.1007/s13595-015-0493-6>

765 Verlinden, M.S., Broeckx, L.S., Van den Bulcke, J., Van Acker, J., Ceulemans, R., 2013.
766 Comparative study of biomass determinants of 12 poplar (*Populus*) genotypes in a
767 high-density short-rotation culture. *For. Ecol. Manag.* 307, 101–111.
768 <https://doi.org/10.1016/j.foreco.2013.06.062>

769 Wheeler, B., 2014. AlgDesign: Algorithmic Experimental Design. R package version 1.1-7.3.
770 <https://CRAN.R-project.org/package=AlgDesign>.

771 Weih, M., Karacic, A., Munkert, H., Verwijst, T., Diekmann, M., 2003. Influence of young
772 poplar stands on floristic diversity in agricultural landscapes (Sweden). *Basic Appl.*
773 *Ecol.* 4, 149–156. <https://doi.org/10.1078/1439-1791-00157>

774 Yáñez, M.A., Zamudio, F., Espinoza, S., Ivković, M., Guerra, F., Espinosa, C., Baettig, R.M.,
775 2019. Genetic variation and growth stability of hybrid poplars in high-density short-
776 rotation coppice stands in central Chile. *Biomass Bioenergy* 120, 84–90.
777 <https://doi.org/10.1016/j.biombioe.2018.11.011>

778 Zalesny, J.A., Zalesny, R.S., Coyle, D.R., Hall, R.B., 2007. Growth and biomass of *Populus*
779 irrigated with landfill leachate. *For. Ecol. Manag.* 248, 143–152.
780 <https://doi.org/10.1016/j.foreco.2007.04.045>

781 Zalesny, R.S., Bauer, E.O., 2007. Evaluation of *Populus* and *Salix* continuously irrigated with
782 landfill leachate I. Genotype-specific elemental phytoremediation. *Int. J.*
783 *Phytoremediation* 9, 281–306. <https://doi.org/10.1080/15226510701476461>

784 Zanchi, G., Pena, N., Bird, N., 2012. Is woody bioenergy carbon neutral? A comparative
785 assessment of emissions from consumption of woody bioenergy and fossil fuel. *GCB*
786 *Bioenergy* 4, 761–772. <https://doi.org/10.1111/j.1757-1707.2011.01149.x>

787 Zappelini, C., Karimi, B., Foulon, J., Lacercat-Didier, L., Maillard, F., Valot, B., Blaudez, D.,
788 Cazaux, D., Gilbert, D., Yergeau, E., Greer, C., Chalot, M., 2015. Diversity and
789 complexity of microbial communities from a chlor-alkali tailings dump. *Soil Biol.*
790 *Biochem.* 90, 101–110. <https://doi.org/10.1016/j.soilbio.2015.08.008>

791 Zianis, D., 2008. Predicting mean aboveground forest biomass and its associated variance.
792 *For. Ecol. Manag.* 256, 1400–1407. <https://doi.org/10.1016/j.foreco.2008.07.002>

793

794

795

796 Figure captions

797 **Fig. 1.** CaCl₂-TE extractable fraction in Pierrelaye soils collected under the Skado genotypes.
798 Values are means and were measured in samples collected in 2007, 2011, 2013 and 2016. The
799 different letters above the element (n = 3) and pH (n=3) values represent significant
800 differences (p<0.05, Kruskal-Wallis). ns : not significant.

801

802 **Fig. 2.** Comparison of measured data (2013) and modeled relationships for trunkDW (kg) and
803 dbh (cm), considering genotypic variability for the 14 poplar genotypes grown at the
804 Pierrelaye site. Data shown in color are for the Skado (red) and Trichobel (green) genotypes.

805

806 **Fig. 3.** Productivity (odt ha⁻¹ year⁻¹) for the 14 poplar genotypes grown at the Pierrelaye site
807 after one rotation (2013). The proportion of branches to trunk are provided (right axis). D: *P.*
808 *deltoides*; DxN: *P. deltoides* x *P. nigra*; T: *P. trichocarpa*; TxM: *P. trichocarpa* x *P.*
809 *maximowiczii*.

810

811 **Fig. 4.** Relationship between bark content (%) and branch diameter for the Skado genotype.

812

813

814 Supplemental materials

815

816 **Fig. S1.** Variables measured in the field for individual trees and location of sampling for dry
817 matter and element concentration (in blue)

818

819 **Table S1**

820 Poplar genotypes grown at the experimental field of Pierrelaye D: *P. deltoides*; DxN: *P.*

821 *deltoides* x *P. nigra*; T: *P. trichocarpa*; TxM: *P. trichocarpa* x *P. maximowiczii*. B: Belgium;

822 I: Italy, NL: the Netherlands.

823

824 **Table S2**

825 DTPA-TE extractable fraction in Pierrelaye soils collected under the various poplar
826 genotypes. Values presented were measured in 2007 and 2011 and the changes occurring
827 between the two samplings are provided.

828

829

830

831

832