

Birth seasonality in the Baka Pygmies

Laura Piqué-Fandiño, Miquel Hernández, Fernando V Ramirez

▶ To cite this version:

Laura Piqué-Fandiño, Miquel Hernández, Fernando V Ramirez. Birth seasonality in the Baka Pygmies. European Society for the Study of Human Evolution, Sep 2019, Liège, Belgium. hal-02359579

HAL Id: hal-02359579 https://hal.science/hal-02359579v1

Submitted on 12 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

BIRTH SEASONALITY IN THE BAKA PYGMIES

Laura Piqué-Fandiño¹, Miquel Hernández¹, Fernando V Ramirez Rozzi²

1. Department of Evolutionary Biology, Ecology and Environmental Sciences, University of Barcelona, Spain. 2. UMR 7206 CNRS MNHS UP Ecoanthropologie Musée de l'Homme, Paris, France.

INTRODUCTION

Patterns in births have been observed in all kind of human populations [1-3], but less studies have been focused on smaller hunter-gatherer populations. The patterns differ across populations, and are usually explained by environmental variables, in combination with social and economic factors [4]. The two main weather



The Baka Pygmies in Moange le Bosquet are a semi-nomadic population living in equatorial Africa with an economy based on hunts and gathers. During the year they develop different economical activities. Pygmies share a common ancestor and splitted from Bantu-speaking populations at around 60,000 years BP [5] and present a particular growth, slower rate of growth during infancy for Baka, which explains their pygmy phenotype. [6-8]

variables that have been related with fluctuations in births are **temperature** (behavioural and biological) and **rainfall** (farming seasons, scarcity of food). Also cultural

factors such as marriages, labour migration, religion festivities or holidays, among others, have been related to birth patterns in western populations.



OBJECTIVES — MATERIALS & METHODS

To assess the presence of a specific pattern of birth seasonality in the Baka Pygmies from Moange le Bosquet, Cameroon.



1,067 BIRTH BIRTH

The slopes graph represents the increase or decrease of births from one month to the following.Weassigned1ifthenumberofbirths increased, -1 if it decreased and 0 if it did not

change. When the line is in the negative area, the number of births decrease, and it increases in the positive area. If it crosses the 0 going from negative to positive, it's a minimum, and from positive to negative, it's a maximum. The method was designed to minimize the effect of long term change in birth number over the years fluctuations, by looking only whether birth numbers increase or decrease between months.



Image 1. Baka Pygmy individual, taken by Dr. Ramirez Rozzi with informed consent from the individual.

RESULTS



There is a **significant birth pattern** in the Baka Pygmy population of Moange le Bosquet (P<0,05)

Months of **rainfall** peaks (May and October), correspond with peaks in birth (Fig. 1). The mechanisms through which rainfall affects births are unknown, but are not likely to be related to the farming cycle or food scarce because of the Baka's lifestyle.

Year-round oscillation in **temperature is too small** (2,7° C) to affect conceptions or births. At least through the mechanisms described in previous literature.

Cultural factors like marriages or holidays are **ruled out** as factors causing the birth seasonality because of the characteristics of the population.



Figure 1. Monthly slope showing the increase /decrease of number of births whithin months. Monthly precipitation (mm) and temperature (°C) means from 1961-1990 from Batouri weather station. Seasons of different economical activities carried out by the Baka. The **economical activities** developed by the Baka, such as hunting seasons and collecting wild mango, could affect season of birth.

CONCLUSIONS

The pattern of birth in the Baka Pygmy population in Moange le Bosquet, in South East Cameroon is seasonal, and it seems to be related to

rainfall, and probably with economical activities.

1. Cogwill, U. M. (1996). The Season of Birth in Man. Man, 1 (2), 232-240

- 2. Lam, D. A. & Miron, J. J. A. (1991). Seasonality of Births in Human Populations. Biodemography and Social Biology, 38 (1-2), 57-78
- 3. Leslie, P. W. & Fry, P. H. (1989). Extreme Seasonality of Births among Nomadic Turkana Pastoralists. American Journal of Physical Anthropology, 79 (1), 103-115.
- 4. Bailey, R. C., Jenike, M. R., Elliston, P. T., Bantley, G. R., Harrigan, M. A., & Peacock, N. R. (1992). The Ecology of Birth Seasonality among Agriculturalists in Central Africa. Journal of Biosocial Science, 24 (3), 393-412.
- 5. Verdu, P. et al. (2009). Origins and Genetic Diversity of Pygmy Hunter-Gatherers from Western Central Africa. Current Biology, 19, 312-318.
- 6. Ramirez Rozzi, F. V., Koudou, Y., Froment, A., Le Bouc, Y. & Botton, J. (2015). Growth pattern from birth to adulthood in African pygmies of known age. Nature Communications, 6:7772.
- 7. Ramirez Rozzi, F. V. (2016). Diversity in tooth eruption and life history in humans: illustration from a Pygmy population. Scientific reports, 6, 27405.
- 8. Ramirez Rozzi, F. V. (2018). Reproduction in the Baka pygmies and drop in their fertility with the arrival of alcohol. Proceedings of the National Academy of Sciences of the United States of America, 115 (27), 1-9.

CONTACT

Laura Piqué-Fandiño - l_pique@hotmail.com Miquel Hernández - mhernandez@ub.edu Fernando V Ramirez Rozzi - fernando.ramirez-rozzi@mnhn.fr



Acknowledgements

We thank Dr. Samuel Pavard for his great contribution to the study, as well as his advice and recommendations.