



HAL
open science

Exposure to *Bordetella pertussis* adenylate cyclase toxin affects integrin-mediated adhesion and mechanics in alveolar epithelial cells

Christelle Angely, Ngoc-Minh Nguyen, Sofia Andre Dias, Emmanuelle Planus, Gabriel Pelle, Bruno Louis, Marcel Filoche, Alexandre Chenal, Daniel Ladant, Daniel Isabey

► To cite this version:

Christelle Angely, Ngoc-Minh Nguyen, Sofia Andre Dias, Emmanuelle Planus, Gabriel Pelle, et al.. Exposure to *Bordetella pertussis* adenylate cyclase toxin affects integrin-mediated adhesion and mechanics in alveolar epithelial cells. *Biology of the Cell*, 2017, 109 (8), pp.293-311. 10.1111/boc.201600082 . hal-02360591

HAL Id: hal-02360591

<https://hal.science/hal-02360591>

Submitted on 7 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Exposure to *Bordetella pertussis* adenylate cyclase toxin affects integrin-mediated adhesion and mechanics in alveolar epithelial cells

ANGELY Christelle^{a,b,c,*}, NGUYEN Ngoc Minh^{a,b,c,*}, ANDRE DIAS Sofia^{a,b,c}, PLANUS

Emmanuelle^f, PELLE Gabriel^{a,b,c,d}, LOUIS Bruno ^{a,b,c}, FILOCHE Marcel^{a,b,c,e}, CHENAL Alexandre^g,

LADANT Daniel^g, ISABEY Daniel^{a,b,c}.

^a Inserm, U955, Equipe 13, Biomécanique & Appareil Respiratoire: une approche multi-échelle, 8, rue du Général Sarrail, F-94010 Créteil Cedex, France ;

^b Université Paris Est, UMRS 955, UPEC, 8, rue du Général Sarrail, F-94010 Créteil Cedex, France ;

^c CNRS, ERL 7240, 8, rue du Général Sarrail, F-94010 Créteil Cedex, France ;

^d AP-HP, Groupe Hospitalier H. Mondor - A. Chenevier, Service des Explorations Fonctionnelles, 51, Avenue du Maréchal de Lattre de Tassigny, F-94010 Créteil Cedex, France ;

^e Laboratoire de Physique de la Matière Condensée, Ecole Polytechnique, CNRS, Université Paris Saclay, 91128 Palaiseau, France;

^f Institute for Advanced Biosciences (IAB), Centre de Recherche UGA / Inserm U1209 / CNRS UMR 5309, Site Santé Grenoble - Allée des Alpes - 38700 La Tronche, France ;

^g Unité de Biochimie des Interactions Macromoléculaires (CNRS UMR 3528), Département de Biologie Structurale et Chimie, Institut Pasteur, 75014 Paris, France.

Key words: Adenylate cyclase, toxin, cell invasion, integrins, cytoskeleton, adhesion, cellular stiffness, rupture force, cyclic adenosine monophosphate

Corresponding author:

Daniel Isabey

E13, IMRB, UMR 955

Faculté de Médecine

8, rue du Général Sarrail

94010 CRETEIL

FRANCE

Tel.: 33 1 49 81 37 00

Fax: 33 1 48 98 17 77

* The two first authors A. C. and N. N-M have equally contributed

A.C. acquired and analyzed data, and wrote the paper, N. N-M conceived and performed the experiments, analyzed data and wrote the paper; A.D. S. provided assistance in cell experiments and corrected the paper; P.E. provide expertise on cell biology and corrected the paper; P.G., L.B., F.M. provided physical expertise, mathematical computation and paper proofreading, C.A. and L.D. provided the expertise, purified the toxin and proofread the paper, and I.D. designed and coordinated the study and wrote the paper.

Abstract

Background: The adenylate cyclase (CyaA) toxin is a major virulent factor of *Bordetella pertussis*, the causative agent of whooping cough. CyaA toxin is able to invade eukaryotic cells where it produces high levels of cyclic adenosine monophosphate (cAMP) affecting cellular physiology. Whether CyaA toxin can modulate cell matrix adhesion and mechanics of infected cells remains largely unknown.

Results: In this study, we use a recently proposed Multiple bond Force Spectroscopy (MFS) with an Atomic Force Microscope to assess the early phase of cell adhesion (maximal detachment and local rupture forces) and cell rigidity (Young's modulus) in alveolar epithelial cells (A549) for toxin exposure <1hour. At 30min of exposure, CyaA toxin has a minimal effect on cell viability (> 95%) at the physiological CyaA concentration of 0.5nM, but a significant effect ($\approx 81\%$) at 10nM. MFS performed on A549 for 3 different concentrations (0.5, 5, 10 nM) demonstrates that CyaA toxin significantly affects both cell adhesion (detachment forces are decreased) and cell mechanics (Young's modulus is increased). CyaA toxin (at 0.5 nM) assessed at 3 indentation/retraction speeds (2, 5, 10 $\mu\text{m/s}$) significantly affects global detachment forces, local rupture events and Young modulus compared to control conditions while an enzymatically inactive variant CyaAE5 has no effect. These results reveal the loading rate dependence of the multiple bonds newly formed between the cell and integrin-specific coated probe as well as the individual bond kinetics which are only slightly affected by the patho-physiological dose of CyaA toxin. Finally, theory of multiple bond force rupture enables us to deduce the bond number N which is reduced by a factor of 2 upon CyaA exposure ($N \approx 6$ versus $N \approx 12$ in control conditions).

Conclusions: MFS measurements demonstrate that adhesion and mechanical properties of A549 are deeply affected by exposure to the CyaA toxin but not to an enzymatically inactive variant. This indicates that the alteration of cell mechanics triggered by CyaA is a consequence of the increase in intracellular cAMP in these target cells.

Significance: These results suggest that mechanical and adhesion properties of the cells appear as pertinent markers of cytotoxicity of CyaA toxin.

Introduction

The adenylate cyclase toxin (CyaA) is one of the main virulent factors of the bacteria *Bordetella pertussis* (Bp) - the whooping cough agent (Khelef *et al.*, 1992; Ladant and Ullmann, 1999; Carbonetti, 2010). CyaA toxin possesses the remarkable property of translocating its catalytic domain into the cytosol of eukaryotic cells where it is activated by an endogenous protein, calmodulin (CaM) to produce supra-physiological levels of cyclic adenosine monophosphate (cAMP) which in turn alter cell physiology (Ladant and Ullmann, 1999; Vojtova *et al.*, 2006; Carbonetti, 2010). The process by which the catalytic domain is translocated across the plasma membrane is called internalization or intoxication (Ladant and Ullmann, 1999). There are several experimental evidences supporting that internalization occurs through a direct translocation of CyaA catalytic domain across the plasma membrane of target cells (Veneziano *et al.*, 2013). First, increased levels of cAMP can be detected within seconds after adding CyaA toxin (by contrast, entry of toxins through endocytosis and vesicular trafficking would require a much longer exposure time). Accordingly, CyaA toxin plays a key role in the early stages of respiratory tract colonization. Second, its entry is independent of both acidification of endocytic vesicles and cytochalasin D treatment (Mouallem *et al.*, 1990), an inhibitor of F-actin polymerization which affects actin-related functions (Qualmann *et al.*, 2000). Although the exact process of toxin internalization is not fully understood, it is considered to be a two-step process implicating: (1) binding to the cell surface through interaction with integrin receptor CD11b/CD18 (or $\alpha_M\beta_2$) (Guermontprez *et al.*, 2001) or through direct contact with the lipid membrane (Gray *et al.*, 2004), and (2) translocation of the catalytic domain in which CyaA might use the

electrical field across the plasma membrane as a driving force to enter target cells (Otero *et al.*, 1995; Veneziano *et al.*, 2013).

CyaA toxin plays a key role in the early stages of respiratory tract colonization (Vojtova *et al.*, 2006; Carbonetti, 2010). To disable the innate immune responses, CyaA toxin targets immune cells such as macrophages and neutrophils that express the main toxin receptor, the β_2 -integrin receptor CD11b/CD18 (or $\alpha_M\beta_2$) (Guermontprez *et al.*, 2001) and rapidly increases the intracellular cAMP concentration. In macrophages, inactivation of the GTPase RhoA by CyaA toxin leads to actin cytoskeleton rearrangements including membrane ruffling and alterations of phagocytic functions (Kamanova *et al.*, 2008). In addition, CyaA can induce apoptosis associated with caspase activation and thus promote the activation of cell death cascade in macrophages cells (Gueirard *et al.*, 1998; Ahmad *et al.*, 2016). However, CyaA toxin also invades a broad range of cells that do not express CD11b/CD18, although the functional and patho-physiological consequences of CyaA cytotoxicity are not fully understood. In particular, CyaA toxin is able to invade the epithelial cells of the conductive respiratory tract (Eby *et al.*, 2010). In cultured epithelial cells free of the $\alpha_M\beta_2$ receptor, CyaA causes cell rounding (Ohnishi *et al.*, 2008). Besides, CyaA toxin is assumed to induce apoptosis associated with caspase activation in a variety of cells including lung cancer cell lines (Carbonetti, 2010). To our knowledge, the cytotoxic effect of CyaA toxin in alveolar epithelial cells remains not documented.

Alveolar epithelial cells can also be directly exposed to bacterial invasion when the airway defense system is depleted, e.g., secondary to the impairment of mucociliary transport caused by particular patho-physiological conditions (chronic obstructive pulmonary disease and asthma) and/or specific genetic abnormalities. Even when structural

cilia defects are not identified, cilia function can still be impaired by various exposures to toxins. These toxins can affect cilia function directly or can modify the secretions and viscosity of the mucus layer which is transported by the mucociliary escalator (Yaghi and Dolovich, 2016). Whether CyaA invasion is able to modify the distal respiratory alveolar epithelium during *B. pertussis* colonization remains presently unknown.

In the present study, we explore the mechanical and adhesion properties of alveolar epithelial cells (A549 cell lines) to quantify the short term response to CyaA cytotoxicity at the cellular and molecular levels. We used Atomic Force Microscopy (AFM) and an improved analysis of cell-matrix adhesion named Multiple bond Force Spectroscopy (MFS) described in a companion paper (Nguyen *et al.*, 2017). The model-system used in the present study is a culture of A549 cells exposed to various concentrations of CyaA toxin. Three values of CyaA concentration are tested (0.5, 5, 10 nM) which is a range covering, for the lowest dose tested, the patho-physiological conditions (i.e., up to 100 ng/ml - about 0.6 nM - at the bacterial cell interface according to Eby *et al.*, 2013) and for the two higher dose tested the supra-physiological conditions already used to compensate the lack of $\alpha_M\beta_2$ integrin receptors in presently tested A549 cells. Here, we explore whether this range of CyaA concentration affects the adhesion and mechanical properties (i.e. cell rigidity - Young's modulus - and in cell adhesion strength - detachment and rupture forces) of alveolar epithelial cells (A549) using MFS and AFM. The MFS results along with cell imaging provide consistent experimental evidences that CyaA toxin has a significant role on cytoskeleton and adhesion structures from the smallest dose of CyaA toxin tested, i.e., 0.5 nM, not only in terms of number of associated bonds but also in terms of energy of individual integrin-ligand bonds. Present results suggest that mechanical and adhesive properties appear as pertinent markers of the cytotoxic effect of CyaA in non inflammatory cells.

Material and Methods

Preparation of A549 alveolar epithelial cells

A549 are human alveolar epithelial cells (ATCC) issued from a human lung carcinoma but having the phenotype of alveolar type II epithelial cells. When forming a monolayer, A549 cells adopt a phenotype consistent with type II AECs in vivo and, importantly do not functionally differentiate in culture from type I AECs. They have the ability to form adherent and tight junctions when grown to confluence, and they offer many advantages for studying in vitro the patho-physiological response of alveolar epithelial cells. These cells express a wide variety of cytokines, growth factors and receptors and notably several transmembrane receptors of the integrin type: β_1 , α_3 , α_6 , α_5 and α_2 (Massin *et al.*, 2004). In addition, integrin subunits α_v associated with integrin subunits β_3 and β_6 are expressed in A549 cells (Merilahti *et al.*, 2016). These integrin receptors bind the synthetic peptide containing the RGD sequence present in many extracellular matrix components. Indeed, the specificity of RGD-binding integrins include five integrin subunits α_v ($\alpha_v\beta_1$, $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_v\beta_6$, and $\alpha_v\beta_8$), two β_1 integrin subunits ($\alpha_5\beta_1$ and $\alpha_8\beta_1$), and $\alpha_{11b}\beta_3$ (Humphries *et al.*, 2006), most of these subunits being expressed by A549 cells but at a level which depends on the passage number: the lower the passage number, the higher the level of integrin expression (Sean *et al.*, 2010). Here, passage number stays in the low range ($\approx 12-16$). A cyclic RGD peptide is used instead of proteins because peptide generates no immune response, no infection risk, better stability, more simple orientation and single adhesion motif (Hersel *et al.*, 2003).

Cells were cultured in T25 flasks (TPP) with 5ml of complete medium containing DMEM-Glutamax High glucose medium (Gibco Life technologies), 1% Penicillin/Streptomycin (Sigma-Aldrich) and 10% Fetal Bovin Serum ou FBS (Sigma-Aldrich) under controlled atmosphere (37°C, 5% CO₂, 95% air). To subconfluence, cells were washed with phosphate-buffer saline without calcium and magnesium ions or PBS^{-/-} (Gibco Life technologies), then trypsinized with 0.5ml Trypsin-EDTA (Sigma-Aldrich) and incubated at controlled atmosphere for 2 minutes. After that, trypsin was inactivated by 5ml of fresh complete medium.

For subculturing in T25 flasks, a split ratio of 1/10 was used. For experiments, we used a density of $7 \cdot 10^5$ cells and seeded on petri-dish (TPP Ø34mm) coated with fibronectin at 10ng/mm². The culture of A549 reached confluence in 24 hours in complete medium. Once cells reach confluence state, - i.e., practically when cells cover more than 90% of the surface - the culture medium is replaced by a culture medium without serum and with 0.5% of BSA first to stop cell proliferation, second to eliminate all not known compounds present in the serum than might interfere with the CyaA treatment, and third to increase the specificity of bead binding at the cell surface. 30 min before experiments, cells were incubated in serum free medium supplemented with 0.5% Bovin Serum Albumin or BSA to block the non-specific binding (Planus *et al.*, 1999).

Production and purification of CyaA and inactive variant CyaAE5

CyaA and the enzymatically inactive variant CyaAE5 (resulting from a LQ dipeptide insertion between D188 and I189 in the catalytic core of the enzyme (Ladant *et al.*, 1992) were expressed in *E. coli* and purified to near homogeneity by previously established procedures (Guermontprez *et al.*, 2001; Karst *et al.*, 2014). Succinctly, the inclusion bodies were solubilized in 50 ml of 20mM Hepes, 8M urea, pH 7.4 by overnight solubilization

under constant stirring at 4°C. A centrifugation for 20 min at 1200 rpm enables to extract the supernatant which is supplemented with 0.14M NaCl and then incubated for 1 hour at RT with 75 ml of Q-Sepharose resin equilibrated with 20 mM Hepes, 140 NaCl, 8M urea, pH 7.4. The Q-Sepharose resin, retaining the CyaA protein, is loaded onto a column. After an extensive wash to eliminate contaminants, the CyaA protein is eluted in 20mM Hepes, 500 mM NaCl, 8M urea, pH 7.4. After dilution of the eluate in 20mM Hepes and 8M urea, it is loaded onto a second Q-sepharose column. Washing and elution are operated in the same conditions as the first time. This step allows to obtain a concentrated CyaA protein which is deluted 5 times then with 20mM Hepes, 1M NaCl, pH 7.4 and loads onto a 70ml phenyl-sepharose column and washes with 20mM Hepes, 1M NaCl, with Hepes 20mM again and then with 50% isopropanol. After an extensive wash, the toxin is eluated with 20mM Hepes and 8M urea. In a final step, the eluate is applied onto a sephacryl 500 (GE healthcare, HIPREP 26/60) equilibrated in 20mM Hepes and 8M urea. CyaA batches are pooled and concentrated by ultrafiltration and stored at -20°C in 20mM Hepes and 8M urea. CyaA toxin concentrations are determined by spectrophotometry using a molecular extinction coefficient at 280nm, $E_{m_{280}} = 143590 \text{ M}^{-1}\text{cm}^{-1}$, which is computed from the CyaA sequence on the ProtParam server (<http://web.expasy.org/protparam/>). The purity of CyaA batches is higher than 90% as judged by SDS PAGE analysis and contained less than 1 EU of LPS/ μg of protein as determined by a standard LAL assay (Lonza). Finally, CyaA is refolded into a urea-free, monomeric and functional holo-state (Karst *et al.*, 2014) on a BioSEC5 21.2 * 300mm with a 5 mL G25-SF as pre-column. The refolding efficiency is around $40 \pm 5 \%$ (population monomer / total population of proteins). The biological functions, i.e., hemolysis, translocation of ACD and cAMP production, are routinely assayed as described in Karst

2014 (Karst *et al.*, 2014) and Cannella 2017 (Cannella *et al.*, 2017). The aliquots of monomeric species of CyaA are stored at -20°C in 20 mM Hepes, 150 mM NaCl, 2 mM CaCl_2 , pH 7.4.

Handling of CyaA and CyaAE5 toxins

For our experiments, CyaA and CyaAE5 toxins are diluted into a solution containing 97% of BSA-DMEM without red phenol medium, 2% of buffer and 1% of CaCl_2 , of the total volume, which is essential for the activation of these toxins. CyaA/CyaAE5 mixture is added 15 minutes prior to the experiment. Due to the sequential character of AFM experiments, the CyaA/CyaAE5 exposure time varies from cell to cell in the range from 15 min to approximately 60 min. We used a range of concentration of CyaA/CaAE5 toxin from 0.5 to 10 nM.

cAMP assays

A549 cells are plated at $3 \cdot 10^5$ cells per well and exposed to control conditions or to 3 CyaA concentrations (0.5, 5 and 10nM) or to 3 CyaAE5 concentrations (0.5, 5 and 10nM) for 30 min. Intracellular cAMP concentration has been determined with the Cyclic AMP Competitive ELISA kit (Invitrogen) according to the manufacturer's instructions.

Viability test

The trypan blue is a vital stain used to selectively color dead cells which is used only for the viability test and not for AFM experiments. The colorant penetrates and stays only in dead cells which are not able to eject it. Indeed, this mechanism requires energy that live cells provide by mitochondria to eject it. Practically, the trypan blue (Sigma-Aldrich) is added to cell suspension which was previously exposed or not to CyaA toxin (0.5, 5 and 10 nM). Then, specifically for the cell counting procedure, we discriminated live cells (no color) and dead cells (in blue) by counting with Malassez Single cell to obtain viability percentage: $(\text{live cell number}/\text{total cell number}) \times 100$.

Actin cytoskeleton and focal adhesion staining

A549 cells are plated at 100,000 cells/mL on coverslip (\varnothing 18mm). After 24h at 37°C and 5% CO₂, cells reach 60% of confluence and are exposed during 30 min to different conditions: control (complete medium and CyaA activation buffer) and CyaA toxin (toxin and CyaA activation buffer) at 3 different CyaA concentrations: 0.5, 5 and 10 nM. Then, cells are washed 3 times with PBS. Cells are right after fixed 10 min with 4%-paraformaldehyde. This short phase occurs with diluted solution of paraformaldehyde which greatly limits the impact on observed structures. Cells are washed 3 times with PBS and then permeabilized by 0.3% Triton during 10 min. This step allows antibodies to access to intracellular antigens and requires the use of a detergent such as Triton X-100. Cells washed 3 times with PBS are hereafter incubated for 30 min with PBS containing 1% of Bovine Serum Albumin (BSA) in order to block non-specific sites. After removing blocking solution, cells are incubated 1 h at room temperature (RT) in diluted phospho-tyrosine PY99 primary antibody solution (sc-7020, Santa Cruz Biotechnology, dilute at 1/150) to stain the focal adhesion points. Cells are then

washed 3 times with a solution containing PBS and 0.02% Triton and incubated 1 hour at RT in dark in diluted fluorochrome-conjugated secondary antibody (Alexa-488, LifeTechnology, dilute at 1/1000) to reveal in the green colour, the staining of focal adhesion points. After a new cell washing repeated 3 times with PBS-0.02% Triton cells are incubated in diluted Phalloidin Tetramethylrhodamine B-isothiocyanate solution (Sigma, dilute at 1/2000) to stain F-actin in the red colour. Coverslides were mounted on a slide with the cell side down in Prolong (LifeTechnology). Staining structures are observed and images are acquired on an Axio Imager confocal microscope (Zeiss) at 63× magnification. To analyze cellular images, the mean level of fluorescence of F-actin is determined in each cell by automatic quantification by ImageJ software relatively to the fluorescent background level of each image. After subtraction, we obtain the effective mean fluorescence of the cells. To determine the number of focal adhesion points, the phospho-tyrosin PY99 level is analyzed using a thresholding method and the automatic particle analyzer in ImageJ software. On these same coverslides, confocal microscope with the LSM5 Pascal software was used to perform z-stack images of fluorescent cellular planes every 0.5 μm and then 3D-reconstructions to estimate cell height for each tested condition.

AFM measurements by integrin-specific cellular probing

Force spectroscopy (FS) was conducted using commercial NanoWizard 1 AFM (JPK Instruments) on cells showing a characteristic shape of adherent epithelium without any blebs, or cellular damage (cells in ball shape or cells partially or totally detached from the culture, are not chosen, See SM1). The AFM is settled on the AxioObserver Z1 inverted microscope (Zeiss) equipped with a Zeiss EC-Plan-NEOFLUAR 20x lens. The microscope is

placed on a stabilization table TL-150 (Table stable Ltd.) so that most vibration sources are minimized. The CellHesion module (JPK Instruments) comprising an electronic controller and a piezo-driven stage enables a vertical displacement up to 100 μ m. The DirectOverlay™ system (JPK Instruments) uses the accuracy of the AFM closed-loop scanning system to enable a perfect match of the optical and AFM images in such way that the cellular position of the FS maneuvers can be selected from the optical image (Fig. SM1). FS measurements are performed on the cell membrane above the relatively flat nuclear region, i.e., where the cell thickness is always the highest (>3.5 μ m, see SM2) thus minimizing the influence of the substrate (Domke and Radmacher, 1998) and the influence of intracellular variability, while the contact geometry rather well defined. A unique force curve per cell is intentionally performed in order to minimize cell adaptation response that might be induced by a repeated mechanical stimulation (Tamada *et al.*, 2004). For each condition tested, about 40-60 different cells are selected.

FS measurements are carried out with a 6.6 μ m spherical probe SquBe CP-CONT-SiO-C (Nanoandmore) mounted on a cantilever whose stiffness is in the range of 0.02-0.77 N/m (average cantilever stiffness value: 0.5 N/m). To take into account this variability, all cantilevers are systematically calibrated prior to FS experiment by the thermal noise oscillator method (Butt and Jaschke, 1995). The spherical shape of the probe is chosen to avoid any membrane injury.

AFM cantilever, after a preliminary cleaning with Triton 0.5% and ethanol, is systematically treated with air-based plasma (Harrick Plasma Cleaner-PDC-002-HPCE). This step makes the probe surface hydrophilic and thus energetically available for contact with the aqueous coating solution. The probe is then sterilized under UV exposure for 25 min functionalized with 2.5mg/ml RGD solution and finally stored at 4°C for 1 night. Before being

mounted on the cantilever holder, the probe is thoroughly rinsed with PBS in order to eliminate unbound RGD peptides.

A typical FS cycle comprises 3 phases depicted in Fig. 1A: (i) Approach: the piezo moves the stage upward so that the cell is brought into contact and the probe deforms the cytoplasm until a point of maximum effort is reached, i.e., contact force = 5 nN; (ii) Contact: the stage is immobilized during a predetermined contact time which is set at 2s to enable studying adhesion in its early phase; These contact conditions were chosen after preliminary experiments which permitted to define the best compromise between a sufficient number of rupture events and complete bond rupture during the retraction period; (iii) Retraction (detailed in Fig. 1B): at the end of the contact time, the piezo moves the stage downward at the same constant speed as approach, the probe is progressively separated from the cell, the bond(s) created at the extremity of cellular extensions break randomly thereby generating sudden steps throughout the retraction force-curve. It should be indicated here that the binding system generated during such short contact time (2s + a few seconds of retraction maneuver duration, see Fig. 1A) corresponds to the early phase of integrin binding and thus is markedly different from the mature cell-substrate focal adhesion system observed by immunostaining (see previous paragraph).

The cantilever speed is kept constant at 3 predetermined values: $v=2, 5,$ and $10 \mu\text{m/s}$. This speed range is purposely limited (i) toward low values to minimize the duration of FS experiments (maximum half a minute per cell) and thus to avoid cell morphological changes, and (ii) towards high values to guaranty quasi-static conditions and minimal hydrodynamic friction on the probe. Indeed, due to the moderate range of approach and separation speeds used in this study, the hydrodrag is negligible (a few % maximum) for the 2 lowest speeds

tested (2 and 5 $\mu\text{m/s}$) and, is smaller than 10pN for the highest speed tested (10 $\mu\text{m/s}$). This leads to an error on the measured rupture forces lower than 10%, letting the hydrodrag correction unnecessary in present experimental conditions. Closed-loop system is activated in order to precisely control the z-movement. The z-movement of 35 μm from contact point leads to the total separation of the probe from cells in most of presently studied cases. The sampling rate is set at 1 kHz allowing capturing rapid change in signal.

Multiple and single bond analysis

We consider that the cellular adhesion system created between the adherent cell and the coated probe - presently a 6.6 μm -diameter spherical probe coated for specific integrin binding - is a complex bond system, i.e., containing a number of N constitutive integrin-ligand junctions with individual energy barriers. The Kramer-Smoluchowski theory (Evans, 1998, 2001; Tsukasaki *et al.*, 2007) predicts that the global lifetime needed to transit across such a complex energy barrier can be expressed as the sum of the individual times needed to transit across each energy barriers. The inverse of the global lifetime is the global dissociation rate, noted $K_{off}(f)$ which can thereby be expressed in terms of individual dissociation rates. Assuming that individual barrier height are all identical meaning similar integrin-ligand bonding due to the integrin-specific probe coating treatment, the individual dissociation rates are thus equivalent and noted $k_{off}(f)$. Accordingly:

$$K_{off}(f) = \frac{1}{\sum_{n=1}^N [k_{off}(f)]^{-1}} \quad 1$$

Considering two typical configurations of bonds working non cooperatively, (i.e, breaking randomly), namely zipper and parallel as previously done (Evans, 2001; Williams, 2003), we

recently demonstrated (Isabey *et al.*, 2013) that the global lifetime can be written in the form of a generalized Bell's model (Bell, 1978) with a prefactor called global natural dissociation rate K_{off}^0 and an exponent representing the separating force f normalized by a global force scale noted F_β :

$$K_{off}(f) = K_{off}^0 \exp\left(\frac{f}{F_\beta}\right) \quad 2$$

For each individual bond, the bond kinetic parameters are classically given by a similar expression:

$$k_{off}(f) = k_{off}^0 \exp\left(\frac{f}{f_\beta}\right) \quad \text{with} \quad k_{off}^0 = \frac{1}{t_D} \exp\left(-\frac{E_b}{k_B T}\right) \quad \text{and} \quad f_\beta = \frac{k_B T}{x_\beta} \quad 3$$

k_{off}^0 represents the natural dissociation rate of an individual bond which is related to the height of the chemical energy barrier E_b in the absence of force, $k_B T$ is a thermal energy scale ($= 4 \text{ pN} \times \mu\text{m}$), t_D is the Brownian time ($= 10^{-9}$ - 10^{-10} s), x_β is the distance (in the nm range) at which the energy barrier is located (called transition state).

The relationship between the global adhesion parameters: K_{off}^0 , F_β and the individual adhesion parameters: k_{off}^0 , f_β , depends on the bond configuration. Parallel configuration appears to be the appropriate bond model for force spectroscopy experiments (Nguyen *et al.*, 2017; Isabey *et al.*, 2013). Indeed, during AFM experiments, the force applied on the bonds created between the bead and the cell is exerted perpendicularly to the plane of cell culture, and the force, f applied on the collective bond, has more chance to be homogeneously distributed between existing bonds and force distribution likely remains

uniform up to the total rupture of the bond system (Sulchek *et al.*, 2005; Rankl *et al.*, 2008).

For parallel bonds, K_{off}^0 depends on the force level f uniformly distributed on all bonds:

$$K_{off}^0 \approx \left(\frac{k_{off}^0}{\ln N} \right)_{F \rightarrow 0} \quad \text{or} \quad K_{off}^0 \approx \left(k_{off}^0 \sqrt{2} \left(\frac{f}{Nf_\beta} \right)^{0.5} \right)_{N \geq 2; F \neq 0} \quad 4$$

Eq. 4 means that K_{off}^0 decreases slightly with N and $F_\beta = Nf_\beta$, i.e., F_β is proportional to N which means that an increase in bond number induces an exponential increase in the global bond lifetime, and therefore an exponential decrease in the global dissociation rate, $K_{off}(f)$ (Eq. 2).

In addition, during AFM experiments, bonds can normally be stretched under a constant loading rate $r_f = cste = df/dt$, the profile of dissociation kinetics under force may be obtained (Evans and Kinoshita, 2007), leading to the calculation of the most probable rupture force for individual bonds. For complex bonds, we used the analytical expressions of $K_{off}(f)$ proposed for the parallel configuration in (Isabey *et al.*, 2013), leading for to the following expression of the global rupture force F^* (also called in the literature: “maximal detachment force”):

$$F^* = F_\beta \ln \left[\frac{r_f}{K_{off}^0 F_\beta \left(\frac{2}{2 + F_\beta / F^*} \right)} \right] \approx F_\beta \ln \left[\frac{r_f}{K_{off}^0 F_\beta} \right] \quad N \geq 2 \quad 5$$

Eq. 5 shows that the rupture force of a complex parallel bond, F^* , is approximately proportional to the logarithm of the loading rate as for single bonds (Evans and Ritchie, 1997; Evans, 1998). It should be indicated here that the role of the surrounding mechanical

environment on strength of a collective bond is actually taken into account through the loading rate dependency described by Eq. 5. Noteworthy, F^* increases almost proportionally with N through $F_\beta = Nf_\beta$ if an appropriate estimate of the loading rate can be performed. If this can be done, the $F^* - \ln[r_f]$ relationships enables to calculate global reference force, F_β , and beyond an estimate of the bond number N if an appropriate estimate of the individual reference force f_β can also be performed. Since a similar expression, i.e.,

$$f^* \approx f_\beta \ln \left[\frac{r_f}{k_{off}^0 f_\beta} \right] \text{ (Evans and Ritchie, 1997; Evans, 1998) can be applied to the analysis of}$$

individual rupture events (except that the loading rate r_f is the actual loading rate preceding the single bond rupture), the slope of the linear relationship $f^* - \ln[r_f]$ enables to estimate the reference force, f_β . Moreover, the value of the natural dissociation rate K_{off}^0 can be obtained from the intercept between this regression line and, e.g., the vertical axis ($\left[F^* \right]_{\ln r_f=0} = -f_\beta \ln(K_{off}^0 F_\beta)$). A similar approach used for the single bond leads to an estimate of k_{off}^0 .

The adhesion reinforcement permitted by the collective (parallel) bond configuration can thereby be evaluated through a characterization of bond number by comparing the $\ln(r_f)$ -dependence of rupture forces (i) in a multiple bond system (F^*) and (ii) in the individual bonds f^* which properly constitute the tested multiple bond structure. This comparison can be performed by AFM throughout a given force spectroscopy curve to the extent that this curve contains - not one as in standard single bond force spectroscopy (Taubenberger *et al.*, 2007) - but several bond rupture events as found in present

experiments (See records of FS curves obtained for 3 different CyaA concentrations in SM3) . This requires that a multiple bond system has been created between the AFM probe and the cell (Fig. 1B). Such a method called *multiple bond force spectroscopy* (MFS) is described in a companion paper (Nguyen *et al.*, 2017) and is used in the present study to evaluate the effect of CyaA toxin on cell mechanics and cell adhesion in its early phase.

Determination of loading rate and adhesion parameters from force-distance curves

Based on Evans and Ritchie (Evans and Ritchie, 1997, 1999) the loading rate can be estimated from the product of the retraction speed v (constant in a given AFM experiments) by the effective stiffness:

$$r_f = \frac{df}{dt} = \frac{df}{dx} \frac{dx}{dt} = k_{eff} v \quad 6$$

The parameter $k_{eff} = \left[\frac{1}{k_{cant}} + \frac{1}{k_{cell}} \right]^{-1}$ account for the AFM cantilever stiffness k_{cant} on one side of the bond, and on the stiffness k_{cell} of the cellular structure connected on the other side (cell cortex, tensed actin filaments...etc) (Odorico *et al.*, 2007). In present experiments, $k_{cant} \gg k_{cell}$ leading to $k_{eff} \approx k_{cell}$ hence:

$$r_f = k_{cell} v \quad 7$$

In present experiments, the cellular stiffness k_{cell} was determined at global and local scales:

(i) At the global cellular scale, from the product of a characteristic cell deformation scale (calculated cell-by-cell as indicated in SM7) and of the Young's modulus E_{cell} whose value is estimated at small deformation, i.e., in the initial portion of the curve (cell indentation / cell height $\approx 0.5\mu\text{m} / 5\mu\text{m} \approx 10\%$ deformation) using the Hertz model (Kuznetsova *et al.*, 2007) and assuming an infinitely rigid probe material. If F is the known

vertical indentation force, h the indentation depth, E_{cell} is obtained from:

$$F \approx \frac{16\sqrt{R}}{9} E_{cell} h^{3/2}. \text{ Note that in the low deformation range, indentation and retract curves}$$

are supposed to lead to similar values of Young's modulus (Bulychev *et al.*, 1975).

(ii) At the local cellular/molecular level by the slope of a straight line fitted through the final part of the retract force-displacement curve preceding the point of bond rupture (see insert in Fig. 1B) as earlier proposed by several authors (Li *et al.*, 2003; Taubenberger *et al.*, 2007). This slope provides an estimate of the local cellular/molecular stiffness right before local bond rupture. The local loading rate is obtained by Eq. 6, assuming that the retraction speed v is preserved throughout any of the local cellular extensions.

Global and local adhesion properties are determined directly from the retraction curve. The global properties are the detachment force F^* defined by the maximal absolute value of force, and the work of adhesion W_{adh} defined as the area delimited by the retraction curve and the x axis (Fig. 1B). The work of adhesion W_{adh} is a convolution of cell mechanics and cell adhesion energies needed to reach complete separation of the probe from the cell. In that sense, W_{adh} integrates all mechanical and rupture events contributing to the link between the probe and the cell. The local bond rupture events are quantified from the height of the force steps visible on the retraction curve and by the negative slope preceding the rupture event (Fig. 1B, insert). Each force step represents the individual bond rupture events, each bond contributing to the global adhesion properties characterized by F^* . The slope gives the local rigidity of the subcellular system to which the individual bonds are attached.

Statistical tests and plots

Comparison of the mean values of mechanical and adhesion parameters between the different biological conditions tested was performed with statistical software package (Statistica v7.1, StatSoft, France) using Kruskal-Wallis non parametric test. A 1-, 2-, 3-star significance corresponds to P-values smaller than 0.05, 0.01 and 0.001 respectively depicted by black horizontal lines. Statistical plot type used in this article is box-whisker: dot in box marks the mean value, whisker bars the \pm SD values, and horizontal bar in box gives the median value. Black horizontal lines show statistical significance. In bar graph, error bar are \pm SEM values.

Results

Viability of cells exposed to CyaA toxin

We first demonstrate that, in human alveolar epithelial cells (A549) exposed to CyaA toxin for times between 15 min and 60 min, a significant dose-dependent increase in cAMP of order(s) of magnitude is observed for the 3 tested CyaA concentrations, while a CyaA variant lacking enzymatic activity, CyaAE5, has no effect on intracellular cAMP levels (Fig. 2A).

Second, viability of A549 human alveolar epithelial cells (AEC) is measured for different durations (15, 30, 60 min) and for different concentrations of the CyaA toxin (Fig. 2B). These results are compared to the viability of cells exposed to an enzymatically inactive variant of CyaA, CyaAE5 (Fig. 2C). We find a dose-dependent decrease in cell viability for cells exposed to the active CyaA toxin but a wide majority of cells remain viable because of the

limited duration of MFS experiments. The CyaA toxin effect on cell viability becomes significant only at the highest concentration tested (10 nM). By contrast, CyaAE5 does not modify the cell viability at all concentration or exposure duration tested. Most importantly, exposure of the A549 cells to the lowest concentration (0.5nM) of CyaA toxin within limited periods of time (30 min on the average in present MFS experiments), leads to negligible cell death (viability >95%) and in any case to non-significant differences with control conditions. For supra-physiological concentrations of CyaA, cell viability at 30 min of CyaA exposure decreases to 87%-viability at 5 nM and 81%-viability at 10 nM. Based on these results, accurate quantifications of adhesion in exposed cells are performed for CyaA concentrations for which a maximum number of cells remain alive.

CyaA effects on F-actin and focal adhesion structures

Fig. 3 shows fluorescent images of F-actin network (Fig. 3D), focal adhesions (Fig. 3E), and merge (Fig. 3F), in A549 cells exposed to CyaA toxin for 30 min of duration at a low concentration of 0.5 nM (at which a maximum number of cells remain alive) by comparison to control conditions (Fig. 3A, B, C respectively). Quantifications of fluorescent F-actin and number of focal adhesion points are presented in Fig. 3G and H. We find significant differences in F-actin level and average number of adhesion points between cells exposed to CyaA toxin and the control group. By comparison to the cells exposed to the lowest concentration tested (Fig. 3), the two highest CyaA concentrations tested (5 and 10nM) (see SM4) induce marked concentration-dependent alteration on F-actin structure with disappearance of stress fibers (see arrows in SM4A, D and bar graphs in SM5G) and loss in focal adhesion points (see dotted arrows in SM4B, E and bar graphs in SM5H). Confocal images of the fluorescent F-actin structures (Figure 3 and SM4) obtained at 30 min exposure

showed that this concentration-dependent remodeling of the cytoskeleton and adhesion points results in a significant increase in cell height (see SM2).

CyaA effect on cell adhesion and cell mechanics assessed by MFS

MFS is performed on top of living adherent cells (only one force-distance curve performed per cell) whose viability is assessed cell-by-cell by their characteristic shape evaluated on transmission images (and not by the trypan test that was used only for data showed in Fig. 2B and C - see Material and Methods and SM1). Exposure to increasing concentrations of CyaA toxin deeply modified the recorded force-distance curves, especially because at the highest concentrations where some detached cells (see SM1) are susceptible to generate a perturbation in the cantilever motion, as shown in SM3C with 10 nM CyaA. Before studying CyaA toxin effects on cell adhesion and mechanics, we verified in control experiments that: (i) the RGD coating was stable throughout the time of AFM experiments by changes in the maximal detachment force. These changes considered at different times and for different retraction speeds remained limited to the intercellular variability and not significant (see SM5A), (ii) the probe coating was specific by comparing the values of maximal detachment forces for specific and non-specific coating conditions and verifying that the RGD-coating condition leads to the highest adhesion forces (see SM5B), and that (iii) the CyaA dilution buffer used to CyaA condition (see Material and Methods) had no effect on the measured cellular parameters if CyaA toxin is not present (results not shown).

The effects of various concentrations of CyaA toxin (0.5, 5, or 10 nM) are studied (at $v=5\mu\text{m/s}$) on alveolar epithelial cells for exposed to the toxin for less than 60 min maximum, (30 min on the average). Results are presented in Fig. 4 and in SM6. Each point corresponds to a different cell. Cell exposure to CyaA toxin results in significant effects on cell adhesion:

the maximal detachment force F^* and the work of adhesion W_{adh} are significantly decreased compared to control conditions suggesting adhesion weakening at the apical cell surface in presence of CyaA toxin. By comparison, the enzymatically inactive variant of CyaA, CyaAE5, had no effect.

Interestingly, the effect of CyaA on Young modulus appears more significant at the lowest CyaA concentration (0.5 nM) as no significant change in E_{cell} was observed at 5 nM CyaA. This might be due to some compensatory effects occurring for intermediate levels of CyaA concentration, the reinforced alteration in actin structure counterbalancing the mechanical effects observed at the lower dose of CyaA toxin. The large variations in mechanical properties shown in Fig. 4C provide an idea of the mechanical variability which could in turns affect the variability in loading rate through Eq. 6 (SM7) and beyond the variability in F^* through Eq. 5 (Fig. 4A and SM7).

Global effects of CyaA at different indentation / retraction speeds

In order to assess the role of the loading rate on measured cellular parameters F^* , W_{adh} and E_{cell} , we first need to extend the range of loading rates by testing various indentation - retraction speeds. To do so, we only used the CyaA concentration (0.5 nM), and compare the results for the 3 different retraction speeds ($v=2, 5, 10 \mu\text{m/s}$) (see Fig. 5 and also values in SM7). Depending on the speed tested, CyaA exposure leads to significant differences (compared to control conditions) for maximal detachment force F^* , work of adhesion W_{adh} , and Young modulus E_{cell} . The cellular effects observed at the 3 speeds appear consistent: adhesion parameters F^* and W_{adh} are decreased meaning that newly formed adhesion is weakened, and Young modulus parameter E_{cell} is increased meaning

that CyaA toxin tends to stiffen the cells. The large dispersion of data which is associated to the intercellular variability renders difficult the observation of a significant speed-dependent effect on measured parameters, hence the need to consider the cell-by-cell values of cell stiffness and beyond its effect on the actual value of loading rate, proper to each cell.

The effect of CyaA on adhesion and mechanical parameter dependence of loading rate

The use of 3 retraction speeds in the range 2-10 $\mu\text{m/s}$ has permitted to extend over about 2 decades the range of global loading rates (Fig. 6). The loading rate is calculated cell-by-cell using Eq. 7. The cellular stiffness k_{Cell} which enters in the calculation of the global loading rate r_f is deduced from: $k_{\text{Cell}} = E_{\text{cell}} \times \delta$ where the characteristic distance δ is estimated cell-by-cell from the ratio: $\delta = W_{\text{adh}} / F^*$ (see in SM7 the mean $\pm\text{SD}$ values of E_{cell} , δ and r_f obtained on each tested cell populations (n) at CyaA concentration of 0.5 nM). To characterize the relationships between F^* and the global loading rate r_f , two methods of data analysis are used and results are presented in Fig. 6 and upper part of Table 1: (i) a linear regression performed on all data points obtained at the 3 tested speeds, each point representing a different cell (see the continuous line in Fig. 6), (ii) a linear regression performed on the 3 mean $\pm\text{SD}$ values of F^* and obtained at each one of the 3 retraction speeds tested ($v=2, 5$ and $10 \mu\text{m/s}$). The first method (large number method) is based on the law of large numbers which states that because of large number of data points, the found relationship is close to the expected relationship predicted by the probability theory for multiple bond rupture (Eq. 5). The second method (3-point method) tends to mimic the classical approach in which a statistically meaningful value of F^* depends on a statistically

representative value of r_f (see values in Table 1). The graph of F^* versus $\ln r_f$ obtained on a wide number of cells leads to a linear regression analysis ($R=0.6$ which is a rather low value due to intercellular variability). The slope of this relationship gives F_β and the intercept at $r_f=1$ pN/s enables to estimate K_{off}^0 from the vertical intercept. Practically, the large number method leads to $F_\beta = 280$ pN and $K_{off}^0=0.28$ s⁻¹ while the 3 point method ($R=0.99$) leads to $F_\beta = 379$ pN and $K_{off}^0=0.73$ s⁻¹. Because the energy landscape of the multiple bond is highly complex, it is difficult to conclude about the CyaA effect at the global level without considering the energetic characteristics of the individual rupture events. This is done below.

Local effects of CyaA at different indentation / retraction speeds

The effect of CyaA toxin (0.5 nM) on individual bonds can be characterized by the statistics of the local rupture force f^* obtained at the 3 different speeds tested. f^* values characterize the local rupture events observed on the force-distances curves already used to analyze the effect of CyaA toxin on global parameters. There are several rupture events appearing on each force-distance curve but the number of rupture event is susceptible to be affected by the exposure to CyaA toxin at 0.5 nM. Results presented in Fig. 7A reveal that, during CyaA exposure, f^* is significantly higher than control values. Note however that dispersion of data remains important likely due to the intercellular and molecular variability in the biological and mechanical cell properties. Histograms of local rupture forces for the different speeds tested (Fig. 7B) show that the values of f^* roughly follow a Gaussian distribution except at the highest speed tested. Nevertheless, the closeness between the median values and the mean values systematically observed at the 3 tested speeds (Fig. 7A) confirms that the values of f^* remain close to a Gaussian (or normal) distribution. The

effect of CyaA exposure on local rupture forces differs from the one observed at the global scale. One possible reason is that the local stiffness (and thus the local loading rate) measured during CyaA exposure is sensibly higher compared to that of control conditions (SM7). The analysis of local rupture events must therefore be directly performed as a function of the local loading rate.

The effect of CyaA on the local loading rate dependence of local cellular parameters

Individual rupture events can be characterized by the relationships between rupture force f^* and local loading rates r_f obtained from the slope on the final part of the retract force-displacement curve preceding the point of bond rupture (see Material and Method and insert in Fig. 1B). However, because the force rupture analysis requires an estimate of the loading rate, only values of rupture events preceded by a negative slope (local stiffness) are analyzed in the graph of Fig. 8. As for global data, we used the large number method and the 3-point method to analyze the local rupture events. Using the large number method, the relationship between f^* and $\ln r_f$ can be adequately fitted by a linear regression (R=0.77) (continuous line shown in Fig. 8) whose slope provides $f_\beta = 45$ pN and the intercept at $r_f = 1$ pN/s provides an estimate of $k_{off}^0 = 1.02$ s⁻¹ (Table 1). Using the 3-point method, the linear regression (R=0.99) leads to $f_\beta = 61$ pN and $k_{off}^0 = 1.94$ s⁻¹ showing that the two methods lead to consistent estimates. By comparison to the adhesion parameters obtained in control conditions (see dotted line in Fig. 8), the present evolutions of f_β and k_{off}^0 caused by CyaA toxin (see Table 1) suggest that the height of the energy barrier of the integrin-ligand is only moderately modified (lowered by CyaA toxin because k_{off}^0 is increased) while its position moves inward (x_β is decreased).

Discussion

The CyaA toxin is known to invade a wide range of eukaryotic target cells in which it produces high levels of cAMP thus altering the cellular physiology (Ladant and Ullmann, 1999; Vojtova *et al.*, 2006; Carbonetti, 2010; Eby *et al.*, 2010). Our results show that cAMP increase by order(s) of magnitude when alveolar epithelial cell line (A549) are exposed to CyaA toxin (Fig. 2C). CyaA toxin has been extensively studied for its molecular specificity and a unique mechanism of action (Ladant and Ullmann, 1999; Vojtova *et al.*, 2006; Carbonetti, 2010). Most particularly, due to a very specific mode of intoxication (direct translocation of the catalytic domain through the plasma membrane of the host cell), CyaA intoxication is rapid, producing increase in cAMP within seconds, letting CyaA toxin play a key role in the early phase of respiratory tract colonization (Ladant and Ullmann, 1999; Vojtova *et al.*, 2006; Carbonetti, 2010).

The first important information presently obtained by the MFS method used here is that alveolar epithelial cells shortly exposed to patho-physiological (small) dose of CyaA - not compromising cell viability - show that integrin-mediated cellular adhesion and to a lesser extent cell mechanics, are affected. This cytotoxic effect of CyaA occurs although A549 cells do not express the $\alpha_M\beta_2$ -integrin receptors which are classically targeted by CyaA for initial interactions with host (immune) cells. Interestingly, these alterations in cell adhesion and cell mechanics properties are measurable in conditions where no significant alteration of cell viability is detected.. In contrast, at high doses (supra-physiological) of CyaA toxin, the degradation of cell structure, cell viability, and cell function is clearly evident. Importantly an

enzymatically inactive form of CyaA (CyaAE5) has no effect on the measured adhesion and mechanical parameters indicating that the cAMP increase induced by CyaA is the cause of the measured cellular changes (Fig. 4).

The second interesting information brought by the present results concerns the type of modifications induced by CyaA in terms of failure of increase in the number of bonds associated, thus restraining the early phase of adhesion reinforcement. It appears that CyaA toxin - even at the smallest (patho-physiological) concentration of 0.5 nM - compromises this reinforcement by systematically decreasing the strength of newly formed collective adhesion system created at the early phase of integrin-ligand interactions. At the same time, we observe that cell height is increased, F-actin structure is affected and apical cell stiffness is increased. The advantage of the MFS approach compared to the more classical single bond force spectroscopy methods (Taubenberger *et al.*, 2007; Lehenkari and Horton, 1999) is to assess the CyaA-induced modifications in terms of bond structure and the number of non cooperative associated bond breaking stochastically (See Material and Methods). Indeed, the theoretical predictions described in this paper show that the number of bonds (here, the integrin receptors) implicated in the cell-probe adhesion can be estimated from the ratio $N = F_{\beta} / f_{\beta}$ for cells invaded by CyaA toxin and for control conditions. From this simple expression, we find: $N \approx 6$ after CyaA exposure and $N \approx 12$ (Table 1) in control conditions. It means that CyaA intoxication tends to markedly decrease the number of newly formed bonds generated at the apical cell surface by cell interaction with the coated probe. This decay in estimated adhesion bonds suggests that integrin clustering might be reduced in presence of CyaA toxin while analysis of individual rupture events show that adhesion energy of individual integrin-ligand bonds is only moderately affected (see Results and

values of f_β and k_{off}^0 in Table 1). In a consistent fashion, the range of local rupture forces found in cells exposed to CyaA (≈ 70 -150 pN, see SM7) is only slightly above that found in control conditions (≈ 60 -115 pN, see SM7) likely because loading rates during CyaA exposure tend to be higher (cells are becoming stiffer) (see SM7). Incidentally, the range of local force rupture found in A549 is in agreement with the range of rupture forces (40-100 pN) found with a method allowing to characterize single-integrin unbinding rupture events at the early phase of adhesion (Taubenberger *et al.*, 2007). In summary, the CyaA-induced adhesion weakening found at global cell scale would be essentially due to the decay in bond number association caused by CyaA intoxication and not really by a CyaA-induced change in chemical energy of the integrin-ligand binding.

Concerning the experimental conditions presently used with the MFS method, it should be indicated that to determine the bond number of the newly formed integrin adhesion systems, we needed to limit the range of loading rate to 2 decades. Using a 3 decade range of loading rates, Zhang *et al.* (Zhang *et al.*, 2004) found two activation potential energy barriers (a steep inner barrier and a more elevated outer barrier) for the integrin $\alpha_4\beta_1$ -vascular cell adhesion molecule-1 (VCAM-1) system. Indeed, in the case where the adhesion system must overcome a series of increasingly higher energy barriers before final dissociation, the dissociation kinetics of the complex at low loading rates is governed by the properties of the outmost barrier (Li *et al.*, 2003). Nevertheless, the outer barrier is the site of integrin activation and the most relevant energetic site for newly formed bonds implicated in adhesion reinforcement. By contrast to previous approaches on integrin receptors which could not provide the bond number per adhesive unit (Taubenberger *et al.*, 2007), the present MFS method offers this new possibility. Noteworthy, in our cellular model

of CyaA intoxication, it is precisely the number of newly formed bonds which is affected by cytotoxicity of CyaA.

A third important aspect highlighted by the present study is the ability of the CyaA toxin to efficiently invade A549 cells that are human alveolar basal epithelial cells. Although these cells do not express the $\alpha_M\beta_2$ integrin (Massin *et al.*, 2004; Merilahti *et al.*, 2016), the primary CyaA receptor on innate immune cells (Guermontprez *et al.*, 2001), we show here that CyaA, at concentrations (0.5 nM) that are expected to be reached *in vivo* at the surface of an epithelium colonized by *B. pertussis* (Eby *et al.*, 2013), elicits a substantial increase in intracellular cAMP and significantly alters the cell bio-mechanics and adhesion properties. These data thus suggest that in addition to the well-established and general role of CyaA in counteracting the innate immune responses thus facilitating the host colonization by *Bordetella pertussis*, CyaA may also have a local effect on the epithelium colonized by the bacteria. Eby *et al.*, 2010 previously reported that CyaA could invade polarized T84 cell monolayers or human airway epithelial cultures when applied to their basolateral membranes (Eby *et al.*, 2010). These polarized epithelial cells model the mucosal surface of the host, while the A549 cells are characterized by an alveolar epithelial phenotype and do not have a protective mucosal surface. Hence, various epithelial cell types could be targeted by CyaA during the course of bacterial infection. The direct, cAMP-mediated effects of CyaA on the mechanics (increase in Young's modulus) and adhesion properties of the intoxicated cells, as evidenced here, may thus contribute to the alteration of lung epithelia, a characteristic of the *B. pertussis* infection.

Acknowledgements

The authors thank Emilie Bequignon, André Coste, Estelle Escudier, Jean-François Papon, for their helpful advice and the fruitful discussions.

For this research, we acknowledge receipt of grants from Institut Pasteur, CNRS UMR 3528, Agence Nationale de la Recherche: ANR n° 2010-BLAN-1119-05 and from Fondation pour la Recherche Médicale: FRM programme Bio-Ingénierie pour la Santé 2014, DBS 20140930771.

Conflicts of interests

No competing interests declared.

References

- Ahmad, J. N., Cerny, O., Linhartova, I., Masin, J., Osicka, R. and Sebo, P. (2016) 'cAMP signalling of Bordetella adenylate cyclase toxin through the SHP-1 phosphatase activates the BimEL-Bax pro-apoptotic cascade in phagocytes', *18*, pp. 384-398. doi: 10.1111/cmi.12519.
- Bell, G. I. (1978) 'Models for the specific adhesion of cells to cells.', *Science (New York, N.Y.)*, 200(4342), pp. 618–27. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/347575>.
- Bulychev, S. I., Alekhin, V. P., Shorshorov, M. K., Ternovskii, A. P. and Shnyrev, G. D. (1975) 'Determining Young's modulus from the indenter penetration diagram', *Zavodskaya Laboratoriya*, 9, pp. 1137–1140.
- Butt, H. and Jaschke, M. (1995) 'Calculation of thermal noise in atomic force microscopy', *Nanotechnology*, 6, pp. 1–7. Available at: <http://iopscience.iop.org/0957-4484/6/1/001> (Accessed: 23 July 2015).
- Cannella, S. E., Yvette, V., Enguéné, N., Davi, M., Malosse, C., Cristina, A., Pérez, S., Chamot-rooke, J., Vachette, P., Durand, D., Ladant, D. and Chenal, A. (2017) 'Stability, structural and functional properties of a monomeric, calcium – loaded adenylate cyclase toxin, CyaA, from Bordetella pertussis', *Nature Publishing Group*. Nature Publishing Group, pp. 1–17. doi: 10.1038/srep42065.
- Carbonetti, N. H. (2010) 'Pertussis toxin and adenylate cyclase toxin: key virulence factors of Bordetella pertussis and cell biology tools.', *Future microbiology*, 5(3), pp. 455–69. doi: 10.2217/fmb.09.133.
- Domke, J. and Radmacher, M. (1998) 'Measuring the elastic properties of thin polymer films with the atomic force microscope', *Langmuir*, 14(18), pp. 3320–3325. Available at: <http://pubs.acs.org/doi/abs/10.1021/la9713006> (Accessed: 27 August 2013).
- Eby, J. C., Ciesla, W. P., Hamman, W., Donato, G. M., Pickles, R. J., Hewlett, E. L. and Lencer, W. I. (2010) 'Selective translocation of the Bordetella pertussis adenylate cyclase toxin across the basolateral membranes of polarized epithelial cells.', *The Journal of biological chemistry*, 285(14), pp. 10662–70. doi: 10.1074/jbc.M109.089219.
- Eby, J. C., Gray, M. C., Warfel, J. M., Paddock, C. D., Jones, T. F., Day, S. R., Bowden, J., Poulter, M. D., Donato, G. M., Merkel, T. J. and Hewlett, E. L. (2013) 'Quantification of the adenylate cyclase toxin of bordetella pertussis in vitro and during respiratory infection', *Infection and Immunity*, 81(5), pp. 1390–1398. doi: 10.1128/IAI.00110-13.
- Evans, E. (1998) 'Energy landscapes of biomolecular adhesion and receptor anchoring at interfaces explored with dynamic force spectroscopy.', *Faraday discussions*, (111), pp. 1–16.
- Evans, E. (2001) 'Probing the Relation Between Force—Lifetime—and Chemistry in Single Molecular Bonds', *Annual Review of Biophysics and Biomolecular Structure*. Annual Reviews 4139 El Camino Way, P.O. Box 10139, Palo Alto, CA 94303-0139, USA , 30(1), pp. 105–128. Available at: <http://www.annualreviews.org/doi/10.1146/annurev.biophys.30.1.105>.
- Evans, E. and Kinoshita, K. (2007) 'Using force to probe single-molecule receptor-cytoskeletal anchoring beneath the surface of a living cell.', *Methods in cell biology*, 83(7), pp. 373–96. doi: 10.1016/S0091-679X(07)83016-0.
- Evans, E. and Ritchie, K. (1997) 'Dynamic strength of molecular adhesion bonds.', *Biophysical journal*,

72(4), pp. 1541–55. doi: 10.1016/S0006-3495(97)78802-7.

Gray, M. C., Donato, G. M., Jones, F. R., Kim, T. and Hewlett, E. L. (2004) 'Newly secreted adenylate cyclase toxin is responsible for intoxication of target cells by *Bordetella pertussis*', *Molecular Microbiology*, 53(6), pp. 1709–1719.

Gueirard, P., Druilhe, A., Pretolani, M. and Guiso, N. (1998) 'Role of Adenylate Cyclase-Hemolysin in Alveolar Macrophage Apoptosis during *Bordetella pertussis* Infection In Vivo', 66(4), pp. 1718–1725.

Guermonprez, P., Khelef, N., Blouin, E., Rieu, P., Ricciardi-Castagnoli, P., Guiso, N., Ladant, D. and Leclerc, C. (2001) 'The Adenylate Cyclase Toxin of *Bordetella pertussis* Binds to Target Cells via the alphaMbeta2 Integrin (Cd11b/Cd18)', *Journal of Experimental Medicine*, 193(9), pp. 1035–1044. Available at: <http://jem.rupress.org/content/193/9/1035>.

Hersel, U., Dahmen, C. and Kessler, H. (2003) 'RGD modified polymers: biomaterials for stimulated cell adhesion and beyond', *Biomaterials*, 24(24), pp. 4385–4415. doi: 10.1016/S0142-9612(03)00343-0.

Humphries, J. D., Byron, A. and Humphries, M. J. (2006) 'Integrin ligands at a glance', *Journal of Cell Science*, 119(19), pp. 3901–3903. doi: 10.1242/jcs.03098.

Isabey, D., Féréol, S., Caluch, A., Fodil, R., Louis, B. and Pelle, G. (2013) 'Force distribution on multiple bonds controls the kinetics of adhesion in stretched cells.', *Journal of biomechanics*, 46(2), pp. 307–13. doi: 10.1016/j.jbiomech.2012.10.039.

Kamanova, J., Kofronova, O., Masin, J., Genth, H., Vojtova, J., Linhartova, I., Benada, O., Just, I. and Sebo, P. (2008) 'Adenylate Cyclase Toxin Subverts Phagocyte Function by RhoA Inhibition and Unproductive Ruffling', 181, pp. 5587–5597. doi: 10.4049/jimmunol.181.8.5587.

Karst, J. C., Enguéné, V. Y. N., Cannella, S. E., Subrini, O., Hessel, A., Debard, S., Ladant, D. and Chenal, A. (2014) 'Calcium, acylation, and molecular confinement favor folding of *Bordetella pertussis* adenylate cyclase cyaa toxin', *Journal of Biological Chemistry*. American Society for Biochemistry and Molecular Biology Inc., 289(44), pp. 30702–30716.

Khelef, N., Sakamoto, H. and Guiso, N. (1992) 'Both adenylate cyclase and hemolytic activities are required by *Bordetella pertussis* to initiate infection', *Microbial Pathogenesis*, 12(3), pp. 227–235.

Kuznetsova, T. G., Starodubtseva, M. N., Yegorenkov, N. I., Chizhik, S. a and Zhdanov, R. I. (2007) 'Atomic force microscopy probing of cell elasticity.', *Micron (Oxford, England : 1993)*, 38(8), pp. 824–33. doi: 10.1016/j.micron.2007.06.011.

Ladant, D., Glaser, P. and Ullmanns, A. (1992) 'Insertional Mutagenesis of *Bordetella pertussis* Adenylate Cyclase *', *The Journal of biological chemistry*, 267(4), pp. 2244–2250.

Ladant, D. and Ullmann, A. (1999) '*Bordetella pertussis* adenylate cyclase : a toxin with multiple talents', (99), pp. 172–176.

Lehenkari, P. P. and Horton, M. a (1999) 'Single integrin molecule adhesion forces in intact cells measured by atomic force microscopy.', *Biochemical and biophysical research communications*, 259(3), pp. 645–50. doi: 10.1006/bbrc.1999.0827.

Li, F., Redick, S. D., Erickson, H. P. and Moy, V. T. (2003) 'Force measurements of the alpha5beta1 integrin-fibronectin interaction.', *Biophysical journal*, 84(2 Pt 1), pp. 1252–62. doi: 10.1016/S0006-3495(03)74940-6.

- Massin, F., Rubinstein, E., Faure, G. C., Martinet, Y., Boucheix, C. and Béné, M. C. (2004) 'Tetraspan and beta-1 integrins expression pattern of the epithelial lung adenocarcinoma cell line A549 and its sensitivity to divalent cations.', *Cytometry. Part B, Clinical cytometry*. doi: 10.1002/cyto.b.20010.
- Merilahti, P., Tauriainen, S. and Susi, P. (2016) 'Human parechovirus 1 infection occurs via alphaVbeta1 integrin', *PLoS ONE*, 11(4). doi: 10.1371/journal.pone.0154769.
- Merkel, R., Nassoy, P., Leung, A., Ritchie, K. and Evans, E. (1999) 'Energy landscapes of receptor-ligand bonds explored with dynamic force spectroscopy.', *Nature*, 397(6714), pp. 50–3. doi: 10.1038/16219.
- Mouallem, M., Farfel, Z. and Hanski, E. (1990) 'Bordetella pertussis Adenylate Cyclase Toxin : Intoxication of Host Cells by Bacterial Invasion', 58(11), pp. 3759–3764.
- Nguyen, N. M., Angely, C., André Dias, S., Planus, E., Filoche, M., Pelle, G., Louis, B. and Isabey, D. (2017) 'Characterization of cellular adhesion reinforcement by multiple bond force spectroscopy in alveolar epithelial cells', *Biology of the Cell* (accepted) .
- Odorico, M., Teulon, J.-M., Bessou, T., Vidaud, C., Bellanger, L., Chen, S. W., Quéméneur, E., Parot, P. and Pellequer, J.-L. (2007) 'Energy landscape of chelated uranyl: antibody interactions by dynamic force spectroscopy.', *Biophysical journal*, 93(2), pp. 645–654. doi: 10.1529/biophysj.106.098129.
- Ohnishi, H., Miyake, M., Kamitani, S. and Horiguchi, Y. (2008) 'The morphological changes in cultured cells caused by Bordetella pertussis adenylate cyclase toxin', 279, pp. 174–179. doi: 10.1111/j.1574-6968.2007.01028.x.
- Otero, A. S., Yi, X. B., Gray, M. C., Szabo, G. and Hewlett, E. L. (1995) 'Membrane depolarization prevents cell invasion by Bordetella pertussis adenylate cyclase toxin', *The Journal of biological chemistry*, 270(17), pp. 9695–9697.
- Planus, E., Galiacy, S., Matthey, M., Laurent, V., Gavrilovic, J., Murphy, G., Clérici, C., Isabey, D., Lafuma, C. and d'Ortho, M.-P. (1999) 'Role of collagenase in mediating in vitro alveolar epithelial wound repair.', *Journal of cell science*, 112 (Pt 2, pp. 243–52. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9858477>.
- Qualmann, B., Kessels, M. M. and Kelly, R. B. (2000) 'Molecular Links between Endocytosis and the Actin Cytoskeleton', 150(5), pp. 111–116.
- Rankl, C., Kienberger, F., Wildling, L., Wruss, Jr., Gruber, H. J., Blaas, D. and Hinterdorfer, P. (2008) 'Multiple receptors involved in human rhinovirus attachment to live cells', *Proceedings of the National Academy of Sciences*, 105(46), pp. 17778–17783. doi: 10.1073/pnas.0806451105.
- Sean, G., Barnes, R., Hawdon, N., Shewchuk, L., Eibl, J., Lam, J. and Ulanova, M. (2010) 'Up-regulation of integrin expression in lung adenocarcinoma cells caused by bacterial infection: in vitro study', *Innate Immunity*, 16(1), pp. 14–26. doi: 10.1177/1753425909106170.
- Sulchek, T. a, Friddle, R. W., Langry, K., Lau, E. Y., Albrecht, H., Ratto, T. V, DeNardo, S. J., Colvin, M. E. and Noy, A. (2005) 'Dynamic force spectroscopy of parallel individual Mucin1-antibody bonds.', *Proceedings of the National Academy of Sciences of the United States of America*, 102(46), pp. 16638–43. doi: 10.1073/pnas.0505208102.
- Tamada, M., Sheetz, M. P. and Sawada, Y. (2004) 'Activation of a signaling cascade by cytoskeleton stretch', *Developmental Cell*, 7(5), pp. 709–718. doi: 10.1016/j.devcel.2004.08.021.
- Taubenberger, A., Cisneros, D. A., Friedrichs, J., Puech, P.-H., Muller, D. J. and Franz, C. M. (2007)

'Revealing early steps of $\alpha 2\beta 1$ integrin-mediated adhesion to collagen type I by using single-cell force spectroscopy', *Molecular biology of ...*, 18(May), pp. 1634–1644. doi: 10.1091/mbc.E06.

Tsukasaki, Y., Kitamura, K., Shimizu, K., Iwane, A. H., Takai, Y. and Yanagida, T. (2007) 'Role of Multiple Bonds Between the Single Cell Adhesion Molecules, Nectin and Cadherin, Revealed by High Sensitive Force Measurements', *Journal of Molecular Biology*, 367(4), pp. 996–1006.

Veneziano, R., Rossi, C., Chenal, A., Devoisselle, J.-M., Ladant, D. and Chopineau, J. (2013) 'Bordetella pertussis adenylate cyclase toxin translocation across a tethered lipid bilayer.', *Proceedings of the National Academy of Sciences of the United States of America*, 110(51), pp. 20473–8. Available at: <http://www.pnas.org/content/110/51/20473>.

Vojtova, J., Kamanova, J. and Sebo, P. (2006) 'Bordetella adenylate cyclase toxin: A swift saboteur of host defense', *Current Opinion in Microbiology*, 9, pp. 69–75. doi: 10.1016/j.mib.2005.12.011.

Williams, P. M. (2003) 'Analytical descriptions of dynamic force spectroscopy: Behaviour of multiple connections', *Analytica Chimica Acta*, 479(1), pp. 107–115.

Yaghi, A. and Dolovich, M. (2016) 'Airway Epithelial Cell Cilia and Obstructive Lung Disease', *Cells*, 40(5), pp. 1–19. doi: 10.3390/cells5040040.

Zhang, X., Chen, A., De Leon, D., Li, H., Noiri, E., Moy, V. T. and Goligorsky, M. S. (2004) 'Atomic force microscopy measurement of leukocyte-endothelial interaction.', *American journal of physiology. Heart and circulatory physiology*, 286(1), pp. H359-67. doi: 10.1152/ajpheart.00491.2003.

Legend of Figures

Figure 1: Sketch of the MFS method

Procedure for measuring adhesion and mechanical parameters by Multiple bond Force Spectroscopy (MFS) method using AFM.

(A) Typical time-dependent signal of force (in nN) recorded on AFM device showing the sequence of successive maneuvers (in relation to the probe position relatively to the cell sketched on top) used during an MFS experiment. 3 characteristic periods can be observed: (i) Approach and indentation of a RGD-coated spherical probe at predetermined constant speed ($v=5\mu\text{m/s}$ on this record) up to a maximal pressing force of 5nN (red part of the curve), (ii) Cell contact during 2 s and a contact force set at 5nN (green part of the curve), (iii) Retraction obtained at the same predetermined constant speed ($5\mu\text{m/s}$ on this record) (blue part of the curve).

(B) Typical force-distance curve obtained in a given A549 cell showing the indentation period (red curve) and the retraction period (blue curve). The Young modulus is obtained by Hertz model in the low range of indentation ($<0.5\mu\text{m}$) whereas global and local adhesion parameters are obtained on the retraction curve. The adhesion parameters are: (i) the maximal detachment force F^* in nN range (or global rupture force) which corresponds to peak (negative) value recorded on the retraction curve and the work of adhesion W_{adh} in $\text{nN}\times\mu\text{m}$ calculated from the striped area, (ii) the local rupture force f^* (step increase) which aims at characterizing (see insert) the single-bond rupture forces (in pN range) and the local cellular/molecular stiffness (k_{local}) from the negative slopes preceding the local bond ruptures. In case of a zero slope (plateau) which is attributed to tether extension with no

measurable cell stiffness, the local loading rate could not be estimated and the rupture force data did not enter in the statistics of the loading rate dependence of rupture force f^* .

Figure 2: Intracellular cAMP measurement and viability tests performed on A549 cells

Intracellular cAMP measurement (**A**) and viability test (**B** and **C**) performed on A549 cells for control conditions, and cells exposed to active (**B**) and enzymatically inactive variant of CyaA, CyaAE5 (**C**). Cell viability is measured as percentage of living cells after 15, 30 or 60 min of exposure.

(A) The cAMP intracellular concentrations in A549 cells for control conditions (n=6) or after exposure to the indicated concentrations of CyaA (n=6) for times corresponding to MFS experiment duration (15 min-60 min), or the enzymatically inactive variant CyaAE5 (n=6), were determined by an ELISA assay (see Material & Methods). This bar graph shows that the intracellular concentration of cAMP in cells exposed to CyaA toxin increases significantly with the increase in the concentration of CyaA toxin, while CyaAE5 has no detectable effect on cAMP as expected. **(B)** From left to right, control conditions (complete medium with the toxin dilution buffer) (n=6), and CyaA concentrated at 0.5, 5 and 10 nM (n=6). **(C)** From left to right, control conditions (complete medium with the toxin dilution buffer) (n=6), and CyaAE5, concentrated at 0.5, 5 and 10 nM (n=6).

The effect of CyaA on cell viability is dose dependent and not really time dependent. The cell viability decreases with increasing CyaA concentration up to significance at 10 nM ($p \leq 0.05$) but the time of exposure has a negligible effect within a maximal experimental duration of 60 min.) Error bars are \pm SEM * $p \leq 0.05$; ** $p \leq 0.01$; *** $p \leq 0.001$.

Figure 3: Cell imaging for F-actin structure and focal adhesion points.

Co-staining of F actin (phalloidin, in red) and of focal adhesion (phosphotyrosin PY99, in green) in fixed A549 cells after 30 min of exposure to control conditions or CyaA toxin. **(A, D)** F-actin staining, **(B, E)** focal adhesion staining, **(C, F)** merge. **(A-C)** Control conditions (n= 13), **(D-F)** Cell exposure to CyaA concentrated at 0.5 nM cells (n=13). **(G)** Quantification of F-actin fluorescence for control conditions and after exposure to CyaA concentrated at 0.5 nM. **(H)** Average number of focal adhesions per cell, control conditions and CyaA exposure. F-actin in control conditions is reticulated and exhibits a highly organized stress fibers network (white arrows in A). After CyaA exposure, F-actin structure is weakened which is confirmed by the significant decay ($p \leq 0.05$) in fluorescent F-actin between control conditions and CyaA-exposed cells (number of white arrows is decreased in D). Control cells also have numerous focal adhesions (white dotted arrows in B) linked to actin stress fibers near the cell periphery (grey arrows in C) and a smaller number around the cell centre. After CyaA exposure, focal adhesions at the periphery of the cell are less intense than in control (number of dotted and grey arrows is decreased in E and F respectively). There is a significant decay in the number of focal adhesions per cell after CyaA exposure (in H) ($p \leq 0.05$). Scale bar represents 20 μm . Error bars are $\pm\text{SEM}$. * $p \leq 0.05$; ** $p \leq 0.01$; *** $p \leq 0.001$.

Figure 4: MFS results in A549 cells exposed to various concentrations of CyaA toxin

(A) The maximal detachment force F^* (in nN), **(B)** the work of adhesion W_{adh} (in $\text{nN} \times \mu\text{m}$) and **(C)** the Young's modulus E_{cell} (in Pa) measured during MFS experiments. Each data point on A, B, C, corresponds to a different cell. Indentation and retraction speeds are fixed at $v=5\mu\text{m/s}$. Data are obtained in A549 cells for control condition, exposure to 3 concentrations of CyaA toxin (0.5, 5 and 10 nM) and 1 concentration (0.5nM) of CyaAE5 toxin. Compared to each control condition: F^* and W_{adh} , are significantly decreased for the 3 CyaA

concentration tested ($p < 0.001$). In the case of CyaAE5, there is no difference with the control condition. For E_{cell} , there is significant differences between control and exposure to CyaA toxin at the lowest CyaA concentration (0.5 nM) ($p < 0.001$) and the highest (10 nM) ($p < 0.05$) but no significant differences at 5nM. The squares in rectangular boxes mark the mean value, whisker bars the \pm SD values, and the horizontal line in each box gives the median value. * $p \leq 0.05$; ** $p \leq 0.01$; *** $p \leq 0.001$. Number of tested cells is detailed in SM 6. The lack of dose-dependent consistency in the cell mechanical behavior seems associated to a larger intercellular variability associated to CyaA exposure.

Figure 5: MFS results for global parameters in CyaA-exposed A549 cells, obtained at various indentation / retraction speeds

(A) The maximal detachment force F^* (in nN), (B) the work of adhesion W_{adh} (in nN $\times\mu$ m) and (C) the Young's modulus E_{cell} (in Pa) measured during MFS experiments and 3 different predetermined indentation/ retraction speeds: $v=2, 5$ and 10μ m/s. Each data point on A, B, C, corresponds to a different cell. Data are obtained in A549 cells for control condition, and exposure to the lowest concentrations of CyaA toxin (0.5 nM). Compared to their respective control condition, F^* and W_{adh} , are significantly decreased for the 3 tested speeds ($p < 0.001$). For E_{cell} , there is a systematic significant increase ($p < 0.001$) compared to the respective control condition. The squares in rectangular boxes mark the mean value, whisker bars the \pm SD values, and the horizontal line in each box gives the median value. . * $p \leq 0.05$; ** $p \leq 0.01$; *** $p \leq 0.001$. . Number of tested cells is detailed in SM 7.

Figure 6: Force-loading rate relationship at the global cellular level

The relationship between the maximal detachment force F^* (in pN) and the global loading rate r_f (in pN/s) plotted on a logarithmic scale for CyaA toxin at the smallest concentration (0.5 nM) and for the 3 predetermined speeds: $v=2\mu\text{m/s}$ (green points), $v=5\mu\text{m/s}$ (red points) and $v=10\mu\text{m/s}$ (purple points). The slope of the linear regression (continuous line) give the reference force F_β and the intercept with vertical axis at $r_f=1$ pN/s enables to calculate the natural dissociation rate K_{off}^0 (in s^{-1}) from the relation $\left[F^* \right]_{\ln r_f=0} = -f_\beta \ln(K_{off}^0 F_\beta)$. Note that each data point represents a different cell. The dotted line represents the linear regression obtained for control condition (data points of control condition are not plotted on this graph).

Figure 7: MFS results for local parameters obtained at various indentation / retraction speeds

In (A) the local rupture forces f^* (in pN) and in (B) the histograms of f^* (in pN) for the 3 different predetermined retraction speeds ($v=2, 5$ and $10 \mu\text{m/s}$) in control condition or for exposure to CyaA toxin concentrated at 0.5 nM. For each speed, the differences between cells exposed to CyaA toxin relatively to control conditions are significant ($p<0.001$). A Gaussian distribution is observed for CyaA toxin at $v=2$ and $5\mu\text{m/s}$. For (A), the squares in rectangular boxes mark the mean value, whisker bars the $\pm\text{SD}$ values, and the horizontal line in each box gives the median value. * $p\leq 0.05$; ** $p\leq 0.01$; *** $p\leq 0.001$.

Figure 8: Force-loading rate relationship at the local level

The relationship between the rupture force f^* (in pN) and the local loading rate r_f (in pN/s) plotted on natural logarithmic scale and calculated for CyaA toxin at 0.5 nM concentration and for the 3 predetermined speeds: $v=2\mu\text{m/s}$ (green points), $v=5\mu\text{m/s}$ (red points) and $v=10\mu\text{m/s}$ (purple points). The slope of the linear regression (continuous line) gives the reference force f_β and the intercept with vertical axis at $r_f = 1$ pN/s enables to calculate the natural dissociation rate k_{off}^0 (in s^{-1}) from the relation: $\left[f^* \right]_{\ln r_f = 0} = -f_\beta \ln(k_{off}^0 f_\beta)$. The dotted line represents the linear regression for control condition. Note that points of control condition are not plotted on this graph.

Table 1: The adhesion parameters obtained from the linear regressions analysis at global and local levels

Characterization of the relationships force-loading rate obtained globally at the cell scale and locally for each rupture event for CyaA toxin condition at 0.5 nM compared to control condition (values noted in parenthesis correspond to the dotted lines in Fig. 6 and 8). For the 3 first columns: values correspond to the slope and intercept parameters of the linear regression done on either the whole data set (continuous lines in Fig. 6 and 8) or the 3-points obtained by averaging of data corresponding to each constant speed. For the 3 last columns, adhesion parameters are deduced from the linear regression analysis. In control conditions, there are 332 force-curves (i.e., cells) analyzed for global rupture events and 171 ($\approx 50\%$) interpretable rupture events leading to the determination of the parameter values (in parenthesis) issued from the force-loading rate analysis. By contrast, with CyaA at 0.5 nM concentration, there are 187 force-curves (i.e., cells) analyzed for global rupture events and

only 50 ($\approx 27\%$) interpretable events, leading to the determination of the parameter values issued from the force-loading rate analysis presented in Fig. 8.