



**HAL**  
open science

## Future development of apricot blossom blight under climate change in Southern France

Paul Tresson, Laurent Brun, Iñaki Garcia de Cortazar-Atauri, Jean-Marc Audergon, Sophie Buléon, Hélène Chenevotot, Freddy Combe, Doriane Dam, Maxime Jacquot, Baptiste Labeyrie, et al.

### ► To cite this version:

Paul Tresson, Laurent Brun, Iñaki Garcia de Cortazar-Atauri, Jean-Marc Audergon, Sophie Buléon, et al.. Future development of apricot blossom blight under climate change in Southern France. *European Journal of Agronomy*, 2020, 112, pp.125960. 10.1016/j.eja.2019.125960 . hal-02361185

**HAL Id: hal-02361185**

**<https://hal.science/hal-02361185>**

Submitted on 15 Nov 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# 1Future development of apricot blossom blight under climate change in Southern France

2

3Paul Tresson<sup>1,2,7</sup>, Laurent Brun<sup>2</sup>, Iñaki Garcia de Cortazar-Atauri<sup>1</sup>, Jean-Marc Audergon<sup>3</sup>,

4Sophie Buléon<sup>4</sup>, Hélène Chenevotot<sup>2</sup>, Freddy Combe<sup>2</sup>, Doriane Dam<sup>2</sup>, Maxime Jacquot<sup>5</sup>,

5Baptiste Labeyrie<sup>6</sup>, Vincent Mercier<sup>2</sup>, Claude-Eric Parveaud<sup>5</sup>, Marie Launay<sup>1</sup>

7<sup>1</sup> INRA US1116 AgroClim, Domaine St. Paul, Site Agroparc, 84914 Avignon Cedex 9, France

8<sup>2</sup> INRA UERI Gotheron, 460 route de Gotheron, 26320 Saint-Marcel-Lès-Valence, France

9<sup>3</sup> INRA GAFL, Domaine Saint-Maurice, 67 Allée des Chênes CS60094 84143 MONTFAVET

10CEDEX

11<sup>4</sup> Chambre d'Agriculture Ardèche et Drôme, 145 Avenue Georges Brassens, 26500 Bourg-lès-

12Valence

13<sup>5</sup> GRAB, 460 route de Gotheron, 26320 Saint-Marcel-Lès-Valence, France

14<sup>6</sup> SEFRA, Quartier Marcellas - 26800 Etoile Sur Rhône

15<sup>7</sup> Present affiliations : CIRAD, UR GECO, Montpellier, France ; GECO, Univ. Montpellier,

16CIRAD, Montpellier, France ; LIRMM, Univ. Montpellier, CNRS, France.

17

18

19Corresponding author:

20Marie Launay

21Tel.: +33 4 32 72 23 67. E-mail address: marie.launay@inra.fr

22

## 23Abstract

24

25Climate change will have several consequences for agro-systems, one of which will concern

26changes to the development of pathogens. Because of the losses it causes, particularly in

27organic farming, *Monilinia laxa* is an important pathogen affecting apricot crops. This study  
28focuses on the consequences of climate change regarding blossom and twig blight (*Monilinia*  
29*laxa*) of apricot. To achieve this, a Climatic Index of cumulated Blight risk (CIB) was built, to  
30obtain the weighted sum of blossom blight incidence throughout the blooming period. An  
31epidemiological model to calculate the incidence of blossom blight during every potentially  
32infectious episode and based on biological parameters, was calibrated using a trap pot  
33experiment where trees were placed in orchards and subject to various meteorological  
34conditions. The CIB derived from this model was evaluated on field data, and was shown to  
35be a robust and useful tool to predict the effects of climate change on the development of  
36apricot blight. Then, using the CIB with a phenological model to predict blooming periods in  
37the future, we estimated the risks of apricot blight until 2100 on four contrasted apricot  
38cultivars and in three geographical zones under climate change scenarios RCP 4.5 and 8.5.  
39This study revealed different effects of climate change depending on the cultivar and altitude.  
40Apricot trees would bloom earlier (up to a difference of 50 days between 1950 and 2100)  
41under climate change. Under the combined effects of these shifts of blooming period and  
42changing climatic conditions, late cultivars such as Bergarouge might see a reduction in the  
43risk of blossom blight (down to 31%) because of warmer but dryer blooming periods. Other  
44varieties (e.g.: Bergeron) could experience an increase in this risk by up to 27% with a shift of  
45the blooming period towards rainier conditions at the highest altitudes. The results of this  
46study could be used to anticipate future changes as well as be used at present as a decision-  
47support tool for farmers.

48

#### 49**Key words**

50Blossom blight, *Monilinia laxa*, apricot orchards, *Prunus armeniaca* L., climate change,  
51modelling, phenology

52

### 531. Introduction

54

55

56 Climatic changes such as increase of air temperature and rainfall variability can directly  
57 and/or indirectly affect pathogens and the plant diseases they are causing, which has been  
58 recently reviewed (Trebicki and Finlay, 2019). All important life cycle stages of fungal  
59 pathogens are more or less directly influenced by the prevailing environmental conditions.  
60 The infection process is particularly dependent on the species specific temperature and  
61 humidity requirements of the pathogens. According to the review article by Juroszek and  
62 Tiedemann (2015) infection risk is the most frequently investigated plant disease parameter in  
63 disease risk simulation studies, where crop disease models have been linked to climate  
64 scenarios including, for example, downy mildew infection risk of grapevine (*Vitis vinifera* L.)  
65 published by Launay et al. (2014). As well as direct effects on pathogens, climate change will  
66 also affect plant phenology (Körner and Basler, 2010) and thus modify host-pathogen  
67 synchronism (Caubel et al., 2017). This indirect effect of climate change on disease  
68 development will be all the more crucial regarding pathogens that infect plants during a short  
69 and sensitive phenological stage, such as infectious diseases which attack during blooming  
70 periods.

71 Apricot (*Prunus armeniaca* L.) is an important crop in the Mediterranean region (14,000 ha  
72 planted and 180,000 T of fruit produced in 2014) and particularly in southern France (third  
73 most important fruit crop) (Lichou and Jay, 2012), but like other such crops, apricot requires  
74 repeated fungicide treatments to secure production. In 2012, an average apricot orchard in  
75 France received 11.8 treatments, including 8.1 against fungi (AGRESTE, 2014). Among the  
76 different fungi that affect apricot, *Monilinia spp.* causes the most losses (Hrustić et al., 2012;

77Oliveira Lino et al., 2016). Three species of *Monilinia* have a significant economic impact:  
78*Monilinia laxa*, *Monilinia fructicola* and *Monilinia fructigena* (Hrustić et al., 2012; Oliveira  
79Lino et al., 2016). The latter has been the subject of the most study because of the damages it  
80also causes to apple and pear crops. However, infections affecting stone fruits are mainly due  
81to *Monilinia laxa* and *Monilinia fructicola* (Oliveira Lino et al., 2016). *Monilinia laxa* can  
82infect apricot blossom, twigs and fruit. Blossom and twig blight are the principal concerns,  
83particularly for organic farmers, and can cause losses of up to 90% in Southern France  
84(Parveau et al., 2016).

85Apricot blossom is sensitive to blight (whether this is caused by *M. laxa* or *M. fructicola*) at  
86flowering. The flowers have been shown to be the most susceptible when they are wide open  
87(BBCH Stage 65; Hack et al., 1992) (Luo et al., 2001). In some cases, infected blossoms can  
88then convey infection to the twigs, generating a necrosis of internal tissues. Moreover, twig  
89blight can only be caused when transmitted via infected blossom (Agrios, 2005). During the  
90present study, we focused on twig and blossom blight, as there is a causal relationship  
91between the two, and did not address the problem of fruit rot.

92The development of *Monilinia* fungal infection on stone fruit blossom is linked to  
93meteorological conditions in several ways, as has been demonstrated by previous studies.  
94Experimenting in a growth chamber, Tamm et al. (1995) revealed the effects of the duration of  
95petal wetness and temperature on the development of blossom blight caused by *M. laxa* on  
96cherry (*Prunus avium* L.). Luo et al., (2001) produced similar results in 2001 in their study on  
97plums (*Prunus domestica* L.) infected by *M. fructicola*. Relative humidity and water activity  
98have also been shown to affect the development of *M. fructicola* on cherry blossom petals  
99(Koball et al., 1997). Casals et al., (2010) highlighted the importance of the same weather  
100factors to the germination of *M. fructicola* on Petri dishes (Casals et al., 2010). However,  
101although the effects of climate on the development of *Monilinia* have been studied before, to

102our knowledge this has only been done under controlled conditions. It may be difficult to  
103ensure reliable recordings of the variables used to describe incidence, mycelial development  
104or conidia germination in the field. For example, leaf wetness duration is widely used in crop-  
105disease epidemiological models but remains difficult to measure accurately (Gleason et al.,  
1062008).

107In the context of climate change, the increasing threat of apricot blight makes it necessary to  
108develop tools that will enable us to define/imagine crop systems adapted to future conditions.  
109There is therefore a need for bioclimatic modelling to anticipate the changes to come and  
110adapt our agro-systems (Jeschke and Strayer, 2008). Current epidemiological models that are  
111used as Decision Support Systems (DSS) are often of a purely statistical nature. They are  
112calibrated on current data for use at present but may lack robustness regarding any changes to  
113conditions which fall outside their domain of validity. On the other hand, bioclimatic and  
114physiological models are calibrated under controlled conditions (Petri dishes, growth  
115chambers, greenhouses) that may not reflect the reality in the field, so are once again  
116deficient. Faced with the future uncertainties inherent in climate change, our models need to  
117be adaptable and valid under different conditions (Maier et al., 2016).

118The study we present here only concerned *M. laxa*, as several identification tests performed  
119during the experiments based on the Lane identification key (Lane, 2002) had reported a great  
120majority of *M. laxa* cases (93.7%), rather than *M. fructicola* (6.3%) and *M. fructigena* (none).

121

122The aims of this study were to (1) produce a climatic index of blight risk thanks to an  
123epidemiological model simulating the incidence of blossom and twig blight caused by *M.*  
124*laxa* on apricot. This model, including biological parameters, was calibrated on field data  
125from a trap pot experiment and then evaluated on independent data from a network of  
126orchards;

127(2) determine the effects of climate change on apricot blossom blight. To do so, we first of all  
128applied a phenological process-based model to simulate flowering changes, and then  
129implemented our epidemiological model to calculate the incidence of blossom blight on four  
130contrasted apricot cultivars in three geographical ranges and under two climate change  
131scenarios.

132

## 1332. Materials and methods

134

### 135 2.1. Building a climatic index of cumulated blight risk

136The Climatic Index of cumulated Blight risk (CIB) that we propose is built in several sections.  
137A blossom incidence  $I$  is calculated using a modified version of the epidemiological model  
138proposed by Tamm et al. in 1995. This factor reflects primary blossom infection due to  
139inoculum dispersal and infection as a function of rainfall and temperature. This blossom  
140infection can then be transmitted to twigs. This transmission is not equal at all stages of  
141blossom development so incidence  $I$  is weighted according to the proportion of disease-  
142sensitive stages at the time of infection, to form a Weighted Incidence  $WI$ . The twig blight  
143observed at the end of blooming is thus the result of several infectious episodes and the  
144different  $WI$  experienced by an orchard are then summed to obtain a Climatic Index of  
145cumulated Blight risk (CIB) that reflects twig infection.

146

#### 147 2.1.1. Incidence of blossom blight

148The epidemiological model we used for this study was a modified version of that proposed by  
149Tamm et al., (1995) to describe the incidence of *M. laxa* blossom blight (*number of*  
150*symptomatic blossoms/total number of blossoms*) on sour cherry trees (Tamm et al., 1995).  
151This model describes a continuous response surface for any given temperature and wetness

152duration, making it usable under different conditions and appropriate for the study of climate  
153change. We nevertheless made several modifications to this model.

154First, Tamm's model takes wetness duration and temperature as input variables to generate  
155incidence data. During our study, we preferred to use rainfall rather than wetness duration to  
156explain blight incidence (see Supplementary Materials 1 for a comparison between wetness  
157duration and rainfall). One of the advantages of this approach is that rainfall data are easier to  
158measure accurately in the field.

159Second, in order to prevent the model from generating positive incidence values in the  
160absence of rainfall (as the trees displayed no symptoms under dry conditions), we added a  
161corrective term taking a zero-value for null rainfall and a one-value otherwise. In this context,  
162a supplementary factor built on precipitation ( $p$ ) and a constant  $\varepsilon$ ,  $p/(p + \varepsilon)$  with  $\varepsilon \neq 0$ , was  
163added to force the model to return no incidence if the rainfall is null.

164The equation of our modified version of the Tamm model is therefore as follows (eq.1):

165

$$166 I(p, T) = \frac{p}{p + \varepsilon} * i_{max} * \{1 - (1 - i_0(T)^{1-m}) * e^{-r(T)*p}\}^{\frac{1}{1-m}} \quad (1)$$

167with,

$$168 i_0(T) = \gamma_1 * \phi^{\gamma_2} * (1 - \phi) \quad (2)$$

$$169 r(T) = \rho_1 * \phi^{\rho_2} * (1 - \phi) \quad (3)$$

170where,

$$171 \phi = \frac{(T - T_{min})}{(T_{max} - T_{min})} \quad (4)$$

172

173The model returns the incidence  $I$  for a given rainfall  $p$  and temperature  $T$ . The model takes as  
174known parameters  $i_{max}$  (the maximum observed incidence), and  $T_{min}$  and  $T_{max}$  the minimal and  
175maximal cardinal temperatures for *M. laxa* development. The fit parameters  $\rho_1$ ,  $\rho_2$ ,  $\gamma_1$ ,  $\gamma_2$  and  $\varepsilon$



176 were estimated during model fitting. The shape parameter  $m$  was set at 0.9 as advised by  
177 Tamm (Tamm et al., 1995). The  $\phi$  factor describes a “bell curve” response to temperature. The  
178  $i_0(T)$  factor was originally designed to return an incidence in the absence of wetness duration.  
179 We retained it despite the supplementary factor that we added because we saw a rapid  
180 increase in the incidence observed on trees under low rainfall levels. This factor therefore  
181 describes incidence at low rainfall values. The  $r(T)$  factor is a rate parameter describing the  
182 response to rainfall.

183

184 In addition to this model, we also tested two others (see Supplementary Materials 2.): (i) a  
185 simple generic infection model based on epidemiological knowledge and proposed by  
186 Magarey and colleagues (Magarey et al., 2005, referred to below as “Magarey”), and (ii) a  
187 purely statistical model (linear regression, referred to as “LM”) established from our trap pot  
188 experiment dataset. We chose the modified Tamm model we because of its greater robustness  
189 and suitability for climate change studies (for details see Supplementary Materials 2.).

190

### 191 **2.1.2. Transmission to twigs**

192 As the proportion of disease-sensitive blossoms is evolving with phenology,  $I$  has to be  
193 weighted for rainy events that occur at different times during blooming.

194 Different blooming stages are susceptible in different ways to blossom blight (Luo et al.,  
195 2001). We observed that 58 to 65 BBCH stages (flower opening) displayed comparable  
196 sensitivity and the 57 stage (sepals opening) had less sensitivity; 57 stage infections were only  
197 possible in the context of a highly infectious event and were observed at lower proportions  
198 (data not shown). The Weighted Incidence  $WI$  was thus defined as:

199

$$200 WI = \frac{I \times (0,25 S_{57} + S_{5865})}{S_{57} + S_{5865}} \text{ if } I \geq 0,5 \quad (5)$$

$$WI = \frac{I \times S_{5865}}{S_{57} + S_{5865}} \text{ if } I < 0,5 \quad (6)$$

With  $S_{57}$  being the number of flowers at BBCH stage 57 and  $S_{5865}$  being the number of flowers at BBCH stages 58 to 65.

Because twig infection is caused by infected blossoms, use of these weightings was necessary to correctly describe transmission of the infection to twigs.

206

### 207 **2.1.3. Climatic index of cumulated blight risk**

The twig infection observed was the result of several blossom infections, each being caused by a rainy event during blooming. We chose to identify a single rainy event during the blooming period as a record of rainfall separated from another rainfall episode by at least 4 hours without rain or leaf wetness. We estimated that 4 hours was sufficiently long for the wetness caused by the rain to dry, so that the next event could be considered separately. Finally, by summing the  $WI$  associated with different rainy events, we were able to build a **Climatic Index of cumulated Blight risk** (CIB) reflecting the history of infectious events (eq. 7).

$$CIB = \sum_{i=1}^n WI_i \quad (7) \quad n \text{ being the number of recorded rainy events}$$

217

## 218 **2.2. Data**

219

### 220 **2.2.1. Trap pot experiment for model calibration**

CIB parameters corresponding to those enabling calculation of the blossom infection ( $I$ ) component were optimised with data from a trap pot experiment. This trap pot experiment was performed under semi-controlled conditions: we chose the weather events to which the pots would be exposed by taking them out into the orchard or returning them to the

225greenhouse. In the orchard, they continued to be exposed to field inoculums and current  
226weather conditions. The fact that we could choose the events to which the trees were exposed  
227enabled us to cover a broad range of weather conditions for model calibration, and at the same  
228time this contributed robustness to our model as the calibration was performed using field  
229data.

230The trap pot experiment for model calibration was carried out at the INRA Gotheron Research  
231Station (Southern France, 44° 58' 37" N, 4° 55' 48" E) over a two-year period, *i.e.* 2017 and  
2322018.

233Trap apricot trees in pots were set out in the orchard for a defined period of time (around 24  
234hours), during which they were exposed to recorded meteorological conditions (see 3.1.2) and  
235outdoor *Monilinia* inoculum. The pots were then moved to a greenhouse in which the  
236conditions were controlled and favourable to the expression of blossom blight.

237Apricot trees of the Bergarouge cultivar (Bergarouge® (A2914) Avirine (cov)) were used as  
238the trap pot trees. This cultivar is known to be very sensitive to blossom blight (Parveau et al.,  
2392016). The expression of symptoms in the event of exposure to *M. laxa* and favourable  
240climatic conditions for infection was therefore ensured. The orchard comprised three lines of  
24125 trees along a south-north axis.

242We used seven groups of six Bergarouge trees in pots during 2017 and nine groups of five  
243trees in 2018. Before exposure, the trees were maintained in a cold room (4°C) to keep their  
244phenology under control. Each tree within a group was exposed for the same period in the  
245orchard between the BBCH 57 and BBCH 65 stages (disease-sensitive stages) and during the  
246blooming period of the Bergeval orchard. The first group of trees was exposed on 27 February  
247and the last on 10 March in 2017, while in 2018 the trees were exposed for periods between 7  
248and 28 March. After exposure, pots were placed in a greenhouse under controlled conditions  
249(relative humidity >40%, ambient temperature between 5°C and 25°C).

250 With each group, a control tree was left in the greenhouse to prove the absence of any  
251 inoculum inside the greenhouse.

252

### 253 **2.2.2. Orchard network for model evaluation**

254 The model was tested on independent data by studying an orchard network in Southern  
255 France. The CIB was calculated for the orchards and compared versus a measured Twig  
256 Blight Incidence (TBI).

257

258 The network comprised 15 orchards located in the Drôme and Ardèche regions (Rhône Valley,  
259 France, ranging between 4°48'29"E - 4°58'54"E and between 45°14'52"N - 44°41'51"N).  
260 Thirteen orchards were studied in both 2017 and 2018 and two were studied in 2018, thus  
261 providing a total of 28 siteXyear measurements of Twig Blight Incidence. None of the  
262 orchards was treated against fungal diseases.

263 These 15 orchards were planted with two moderately sensitive cultivars: ten with the  
264 Bergeron cultivar (Bergeron (660)) and five with the Bergeval cultivar (Parveau et al., 2016).  
265 These cultivars are less sensitive to blossom blight than the Bergarouge trees used for the  
266 blossom blight model. However, this was not expected to alter the performance of the model  
267 (see Discussion).

268

269 Blighted twigs were counted on five random main branches per orchard to obtain one TBI  
270 notation per orchard:

$$271 TBI = \frac{\text{number of blighted twigs}}{\text{total number of twigs}} \quad (8)$$

272

273 The evolution of blossom phenological stages was recorded regularly (three times a week) on  
274 ten trees in each orchard. On several occasions, the proportion of blossom at each stage was

275estimated. The proportion of blossom at any time during the different stages was then  
276extrapolated linearly between two estimation dates.

277

### 278 **2.2.3. Recorded meteorological data**

279A weather station (IMT 200 Pessl Instruments, Weiz, Austria) was placed in the middle of  
280each orchard (trap pot experiment and each orchard in the orchard network) at a height of 1.80  
281m. Rainfall (mm), temperature (°C) and leaf wetness (min) were recorded at an hourly time  
282step using a rain gauge, temperature sensor and filter paper leaf sensor, respectively. Leaf  
283wetness was also measured with an electric resistance sensor to assess the reliability of the  
284measurements (this variable is used in Supplementary Materials 1).

285

### 286 **2.2.4. Future climate data**

287We performed the study using the predicted rainfall and temperature data of 46 DRIAS grids  
288(French climate change modelling project, <http://www.drias-climat.fr>), which are 8 x 8 km  
289wide. They were selected in the Rhône Valley at locations where apricot is currently being  
290cultivated.

291

292For more clarity we decided to group the DRIAS grids thus employed in several clusters. A  
293Hierarchical Ascending Classification was performed (mean temperature and mean rainfall as  
294entry variables) and reflected groups as a function of altitude (see Supplementary Materials  
2953). The following clusters were thus used (see *Fig. 1*):

296Cluster 1: altitude <100m

297Cluster 2: 100m ≤ altitude ≤ 400m

298Cluster 3: altitude > 400m

299

300 Present and future climatic conditions (between 1950 and 2100) were applied using the  
301 ALADIN-Climate regional climate model nested within the global ARPEGE model (Deque,  
302 2010). Three periods were simulated: the ‘recent past’ (RP, 1970–1999), ‘near future’ (NF,  
303 2020–2049) and ‘far future’ (FF, 2070–2099) according to two ‘representative concentration  
304 pathway’ emission scenarios, RCP4.5 (median) and RCP8.5 (pessimistic) (Pachauri et al.,  
305 2014).

306

## 307 **2.3. Application to climate change**

308

### 309 **2.3.1. Phenological model**

310 In order to apply our model to the context of climate change, it was necessary to forecast  
311 apricot blooming periods in the future. We therefore implemented a two-step phenological  
312 model on future climatic data.

313

#### 314 **2.3.1.1. Estimation of the mid-blooming date**

315 To simulate the blooming period for apricot we used the sequential phenological process-  
316 based model proposed by Andreini et al. (2014). The date of budbreak is simulated using the  
317 Smoothed Utah function (Bonhomme et al., 2010, advised by Andreini et al. 2014) and the  
318 time between budbreak and mid-blooming (F50) is simulated with a sigmoid model (Chuine  
319 et al., 2016). We estimated the mid-blooming dates for four cultivars chosen to have different  
320 precocities in terms of dormancy release and blooming: cv. Beliana (Beliana® Sayeb)  
321 (median to early blooming period), cv. Bergarouge (used during the trap pot experiment,  
322 median blooming period), cv. Bergeron (late blooming period, widely used in the Rhône  
323 Valley) and cv. Rouge du Roussillon (Rouge du Roussillon (A157)) (early blooming period).

324The models we used to estimate the F50 date had already been calibrated for the different  
325cultivars (Andreini et al., 2014, Chuine et al., 2016), (Garcia de Cortazar-Atauri et al., 2013).

326

### 327 **2.3.1.2. Estimation of the blooming period**

328The expand of the blooming period was estimated by studying the phenological data recorded  
329by the orchard network. No significant differences were found between the expands of the  
330blooming periods of Bergeron and Bergeval (T-test, p-value = 0.2196). Recordings on the two  
331cultivars were then grouped. We found that the proportion of disease sensitive stages during  
332blooming (degree-days) could be approximated using a Gaussian curve centred on the  
333maximum proportion of opened flowers (around 100% of opened flowers) and of a 28 degree-  
334day standard deviation. It was then possible to estimate the expand of the blooming period at  
335around F50, starting 122 degree-days before F50 and ending 133 degree-days after F50.

336

## 337 **2.4. Data analysis**

338Statistical analysis and computations were performed using R 3.4.3. Computations for  
339phenological modelling were performed using Phenology-Modeling-Platform 5.5  
340(<http://www.cefe.cnrs.fr/fr/recherche/ef/forecast/phenology-modelling-platform>) (Chuine et  
341al., 2013).

342

### 343 **2.4.1. Model calibration**

344Goodness-of-fit was assessed using the Root Mean Square Error (RMSE, eq.9) and Relative  
345Root Mean Square Error (RRMSE, eq.10).

346

$$347 RMSE = \sqrt{\frac{\sum (S_i - O_i)^2}{n}} \quad (9) \quad \text{and} \quad RRMSE = \frac{RMSE}{\bar{O}} \quad (10)$$

348

349 Where  $n$  is the number of observations,  $S_i$  the simulated value and  $\bar{O}$  the average of observed  
350 values  $O_i$ . RMSE can be broken down into two components representing systematic (bias or  
351 RMSEs) and unsystematic (dispersion or RMSEu) error (Willmott, 1981).

352

$$353 RMSE_s = \sqrt{\frac{\sum (\hat{S}_i - O_i)^2}{n}} \quad (11)$$

$$354 RMSE_u = \sqrt{\frac{\sum (\hat{S}_i - S_i)^2}{n}} \quad (12)$$

355

356 With  $\hat{S}_i$  being derived from the linear regression of observed versus simulated values:

357  $\hat{S}_i = a + b * O_i$  with  $a$  and  $b$  being the parameters of the regression.

358

## 359 **2.4.2. Model evaluation**

360 The performance of our model was then assessed from the correlation between the observed  
361 Twig Blight Incidence of orchards in the network and their respective calculated CIB.

362

## 363 **2.4.3. Algorithm for climate change study**

364 The method used for computation is described in Figure 2.

365

## 366 **3. Results**

367

### 368 **3.1. Model fitting and evaluation**

369 Our model revealed a RMSE of 6.12% incidence when comparing the simulated incidence on  
370 flowers with those observed in the calibration dataset. Given that the average incidence was  
371 27.34%, the RRMSE of our model was 22.43%, which could be considered to be a



372satisfactory performance. A comparison between simulated and observed incidences is  
373illustrated in Figure 3.

374

375

376Furthermore, the RMSEs was 1.86% whereas the RMSEu was 5.84%, meaning that the error  
377was mostly due to dispersal of the points (*i.e.* biological variability) and not to bias in the  
378chosen formalism. This can also be seen in Figure 4 which represents the residuals associated  
379with each point.

380

381The parameter values we obtained after optimisation are shown in Table 1.

382

383The CIB calculated using our model was correlated with Twig Blight Infection at  $R^2 = 0.46$   
384(Fig.5). Given the numerous parameters varying between the orchards (see Discussion), we  
385considered this performance to be satisfactory.

386

### 387 **3.2. Evolution of blooming dates**

388The predictions of the phenological model indicated a shift of all mid-blooming dates to an  
389earlier day in the year (DOY) for all cultivar and clusters. Under climate change scenario  
390RCP4.5, the median blooming date was 20 days earlier, from 89.9 DOY (*i.e.* 31 March in  
3911950) to 72.2 DOY (March 12 in 2100). This shift was more marked under scenario RCP8.5,  
392where the F50 date moved to 61.5 DOY (1<sup>st</sup> March) in 2100 (Fig. 6). Rouge du Roussillon and  
393Beliana displayed similar but moderated shifts toward earlier blooming dates (of around 20  
394days), but the Bergeron cultivar notably experienced a shift of almost 50 days in its mid-  
395blooming date in all clusters under the RCP8.5 scenario. Therefore, by 2100, the differences  
396in blooming periods between Bergeron and other cultivars would no longer be significant. On

397the contrary, Bergarouge experienced less variation in its blooming date and would thus  
398become the latest blooming cultivar by 2100 in both cluster 1 and cluster 2. We concluded  
399that under the RCP4.5 and RCP8.5 scenarios, but particularly with the latter, a shift towards  
400earlier blooming periods and a convergence between the blooming periods of different  
401cultivars would be observed, with consequences regarding the climatic conditions during  
402blooming and thus the risk of blossom blight.

403

### 404 **3.3. Evolution of climatic conditions during the blooming period**

405This future convergence of blooming periods caused a convergence of the climatic conditions  
406prevailing during the blooming of different cultivars. In particular, Bergeron and Bergarouge  
407experienced opposite and strong shifts. The blooming conditions for Bergarouge shifted  
408towards warmer but dryer conditions, and would be 2°C warmer in the far future than in the  
409past in all clusters and under both scenarios, the trees receiving 10 to 20 mm less rainfall  
410during blooming (Fig. 7). On the other hand, Bergeron saw a marked shift of its blooming  
411period towards earlier dates, so that this cultivar would experience cultivar conditions that  
412would be colder (2.8°C lower in cluster 3 and RCP8.5) and rainier (especially in cluster 1).  
413Rouge du Roussillon and Beliana saw more moderate shifts; those in cluster 3 shifted towards  
414dryer conditions while cluster 1 saw a temperature-related shift towards warmer conditions.

415

416From an epidemiological standpoint, the impacts of these changes to climatic conditions  
417during blooming could indeed be assessed by the CIB computation.

418

### 419 **3.4. Future risk of apricot blight**

420Significant differences in future CIB were only found in the eventuality of the RCP8.5  
421scenario, so this is the only one described in greater detail below.

422The CIB calculated by the model revealed different consequences of climate change that  
423varied according to the clusters and cultivars studied (Fig.8). Variations in the CIB between  
424past and far future ranged from +27% (Bergeron cluster 1) to -31% (Bergarouge, cluster 1).  
425However, cluster 1 (lowest altitudes) displayed greater inter-annual variations, although these  
426differences were not always significant.

427

428We noted a significant in the risk to which Bergarouge is exposed, in all clusters. This could  
429be linked to the shift of its blooming period to warmer but dryer conditions, the expected  
430positive effect from rising temperatures being counter-balanced by the negative effect of  
431lower rainfall (Fig. 7). On the other hand, an increase in risk may be possible in the near  
432future (nf) for Bergeron in cluster 3 (altitudes higher than 400 m) because of colder but rainier  
433blooming conditions. Beliana and Rouge du Roussillon experienced more diverse conditions  
434depending on the cluster, with a general trend regarding disease risk that stagnated or  
435diminished in the far future.

436

#### 4374. Discussion

438This study generated a climatic index of cumulated blight risk based on an epidemiological  
439model, describing the blossom and twig blight caused by *M. laxa* on apricot. The model we  
440used was built using biological parameters driving development of the fungus (optimal  
441growth temperatures, response to rain). This gave the model robustness, enabling its use  
442within the framework of a changing climate. Furthermore, by comparison to the original  
443Tamm model, this model was calibrated and evaluated on easily measurable variables and in  
444the field, so that it is more applicable and closer to observed incidence. To our knowledge,  
445this approach – integrating the incidence of both blossom and twig blight in the same CIB

446index, based on biological parameters and easily accessible weather inputs (daily temperature  
447and rainfall), and valuable under field conditions – is entirely new.

448To estimate the effects of climate change on blight risks, we first of all determined the  
449sensitivity periods of several cultivars. By estimating the present and future CIB at different  
450altitudes (and even latitudes to a lesser extent), we were able to reveal different changes as a  
451function of the precocities of the trees and their altitudes. The late blooming cultivar Bergeron  
452might bloom earlier and face rainier conditions, leading to an increase in the incidence of twig  
453blight; on the other hand, the median cultivar Bergarouge might shift towards later blooming  
454and dryer conditions, accompanied by a lower blight incidence at all altitudes.

455

#### 456 **4.1. Variabilities affecting evaluation**

457The results of evaluating this model using data from the orchard network could be considered  
458as satisfactory ( $R^2 = 0.44$ ). Indeed, numerous parameters varied between orchards in the  
459network. For example, the amount of *Monilinia* inoculum could vary at the landscape or  
460regional levels. Furthermore, the fitting performances of the epidemiological model  
461(RRMSE = 0.22) were equal to or better than the fitting performances regarded as satisfactory  
462by recent comparable studies (e.g.: Gouache et al., 2015; Morales et al., 2018).

463We set blossom-twig transmission according to our empirical observations. However,  
464variations in twig infection via blossom could impact the performance of the model. For  
465instance, blossom-twig transmission could also depend on climatic conditions. A large part of  
466the correlation between the calculated CIB and observed TBI was due to the choice of  
467weightings. More observations might help to improve our understanding of blossom-twig  
468transmission. For example, orchards that experienced rain at the end of the blooming period  
469displayed greater sensitivity to twig blight infection. This factor could be determined by  
470means of other trap pot experiments or the analysis of a larger orchard network experiencing a

471 variety of conditions, with a broader range of latitudes; for example, combining  
472 Mediterranean and more continental climatic conditions.

473 The biological construction of our epidemiological model means that it could be used to  
474 address future conditions at a single location (in the present case, the Rhône Valley) but  
475 should also make the model usable at various geographical locations. For example, testing our  
476 model in other apricot growing regions such as Spain or Turkey would further assess the  
477 robustness of the model we propose here.

478 It should be noted that the model used to calculate blossom blight infection was calibrated on  
479 Bergarouge trees, which are more sensitive to blossom blight than Bergeval and Bergeron  
480 (Parveau et al., 2016). However, because Bergeron and Bergeval display comparable  
481 sensitivity (and thus comparable differences in sensitivity versus Bergarouge), the correlation  
482 between CIB and TBI should not be affected mathematically by a difference in sensitivity  
483 between Bergarouge and Bergeron or Bergeval.

484

#### 485 **4.2. Working hypotheses**

486 We only studied climate-related factors during this study, in order to assess changes to the  
487 infection risk in line with climate change. Nevertheless, factors of a genetic (e.g. resistance),  
488 physiological (e.g. water stress) or epidemiological (e.g. inoculum repartition) nature should  
489 also be taken into account.

490 Here, the apricot cultivars were only compared through the lens of precocities, but a factor  
491 reflecting varietal sensitivity could be added. Such study would enable assessment of the  
492 respective roles of phenology and genetics: is it better for a tree to avoid blight or resist it?

493 As for physiological factors, two types of interactions have been documented to date: "cross-  
494 protection" for plants whose resistance to biotic stresses is increased by the onset of abiotic  
495 stress, and "cross-vulnerability" for plants whose susceptibility to biotic stress is increased

496under abiotic stress (Fones and Gurr, 2017). These processes might modify the host response  
497to increased biotic and abiotic stresses under climate change, and should therefore be  
498considered under a more integrative modelling approach.

499Further, inoculum levels may vary at a landscape or regional level. For example, they may be  
500affected by the provenance of air masses, as has already been shown for *Botrytis cinerea*  
501(Leyronas and Nicot, 2013). Moreover, amounts of primary inoculum may also vary as a  
502function of previous disease levels or orchard management practices (Lichou and Jay, 2012).

503The *Monilinia* inoculum may differ in terms of both quantity and quality involving a change  
504to the predominant *Monilinia* species. For instance, because it is better suited to warmer  
505temperatures (Casals et al., 2010) and displays pesticide resistance (Lichou and Jay, 2012),  
506*Monilinia fructicola* may become more important than *Monilinia laxa* in the European  
507inoculum landscape as climate change progresses. In addition, the sexual reproduction of  
508*Monilinia laxa* has not yet been observed under natural conditions in Europe (Hrustić et al.,  
5092012) but warmer temperatures could trigger this development cycle (Agrios, 2005). Such a  
510change would render obsolete the model we present here, as this aspect of development  
511remains a limiting factor. These features could also affect epidemiological concerns over the  
512coming decades. Epidemiological studies on *M. fructicola* under semi-controlled conditions  
513would then be of value in the context of future research.

514

#### 515 **4.3. Uncertainty of climatic variables**

516Variations affecting the risk predicted by the model were mainly due to rainfall. However, this  
517variable is hard to predict in climate change scenarios, and climatic models can generate  
518markedly different predictions (Jouzel et al., 2014). For instance, the DRIAS data we used are  
519based on the ALADIN model proposed by the French National Weather Research Centre  
520(Centre National de Recherches Météorologiques) and used to forecast weather in the context

521of climate change; it predicts an overall reduction of rainfall in France between [1976-2005]  
522and [2071-2100] under the RCP8.5 scenario in summer (-0.38 mm.day<sup>-1</sup>). On the other hand,  
523the WRF (Weather Research and Forecasting) model proposed by the US National Center for  
524Atmospheric Research predicts an overall increase of rainfall of the same scope (+0.32  
525mm.day<sup>-1</sup>) (Jouzel et al., 2014), hence the uncertainty attached to ongoing modelling  
526approaches used to study future impacts of climate change.

527

### 5285. Conclusion

529The Climatic Index of cumulated Blight risk we propose here offers an efficient reflection of  
530twig blight infection calculated from the weighted sum of blossom blight infection episodes  
531( $R^2 = 0.44$  with independent evaluation). Blossom blight infections were estimated with a  
532good fit to an epidemiological model (RRMSE = 0.22, largely due to unsystematic error).  
533This model, calibrated and evaluated on field data (using easily measurable variables such as  
534rainfall) and based on biological parameters, was shown to be a robust and useful tool to  
535predict the consequences of climate change regarding the development of apricot blight.  
536Because of a shift of the blooming period, the Bergarouge apricot cultivar could experience a  
537long term reduction in the climatic risk to which it is exposed (in the far future). By contrast,  
538cultivars such as Beliana or Bergeron could see a medium-term (near future) increase of this  
539risk of blossom blight at altitudes above 100 m. The various conditions under which our  
540model can be employed means that it could be applied to conditions other than those in the  
541Rhône Valley, and possibly worldwide.

542

### 543Acknowledgements

544This study was funded by the GIS Fruits Network, FAN de Bio and FEADER ClimArbo  
545projects. The authors gratefully acknowledge the assistance received from Emilie Freyssinet

546and William Chambeyron during field and lab work. The authors are also grateful to Benoît  
547Saunier and Tom Fillot for their careful reviews of the paper.

548

#### 549References

550

551AGRESTE, 2014. Enquête pratiques phytosanitaires en arboriculture 2012.

552

553Agrios, G., 2005. Plant Pathology, 5th Edition. ed. ELSEVIER.

554

555

556

557Andreini, L., de Cortázar-Atauri, I.G., Chuine, I., Viti, R., Bartolini, S., Ruiz, D., Campoy,

558J.A., Legave, J.M., Audergon, J.-M., Bertuzzi, P., 2014. Understanding dormancy release in

559apricot flower buds (*Prunus armeniaca* L.) using several process-based phenological models.

560Agricultural and Forest Meteorology 184, 210–219.

561<https://doi.org/10.1016/j.agrformet.2013.10.005>

562

563

564

565

566

567Bonhomme, M., Rageau, R. et Lacoïnte, A., 2010. Optimization of endodormancy release

568models, using series of endodormancy release data collected in France. Acta Hort. 2010.

569Vol. 872, pp. 51-60. <https://doi.org/10.17660/ActaHortic.2010.872.4>

570



571Casals, C., Viñas, I., Torres, R., Griera, C., Usall, J., 2010. Effect of temperature and water  
572activity on in vitro germination of *Monilinia* spp.: Effect of temperature and water activity on  
573*Monilinia* spp. *Journal of Applied Microbiology* 108, 47–54. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-5742672.2009.04402.x)

575

576

577

578Chuine, I., Garcia de Cortazar-Atauri, I., Kramer, k., Hänninen, H., 2013. *Plant Development*  
579Models, in: *Phenology: An Integrative Environmental Science*. Springer, Dordrechts,  
580Netherlands, pp. 275–293.

581

582Chuine, I. Bonhomme, M., legave,J-M., Garcia de Cortazar-Atauri, I., Charrier, G., Lacoite,  
583A., Améglio, T., 2016. Can phenological models predict tree phenology accurately in the  
584future? The unrevealed hurdle of endodormancy break. *Global Change Biology* 22, 3444-  
5853460.

586

587Deque, M., 2010. Regional climate simulation with a mosaic of RCMs. *Meteorol. Z.* 19,  
588259–266.

589

590

591

592Fones, H.N., Gurr, S.J., 2017. NO(X)ious gases and the unpredictability of emerging plant  
593pathogens under climate change. *BMC Biol.* 15.

594

595 Garcia de Cortazar-Atauri, I., Bertuzzi, P., Andreini, L., Madon, F., Audergon, J-M., 2013.  
596 Adaptation de l'abricotier aux changements climatiques : Estimation des risques génétiques et  
597 géographiques liés aux avortements floraux et à l'époque de floraison. Le réseau  
598 agroclimatique de l'INRA et la base associée de données. Rapport CTPS Abricot, 54 p.

599

600

601

602 Gleason, M.L., Duttweiler, K.B., Batzer, J.C., Taylor, S.E., Sentelhas, P.C., Monteiro,  
603 J.E.B.A., Gillespie, T.J., 2008. Obtaining weather data for input to crop disease-warning  
604 systems: leaf wetness duration as a case study. *Scientia Agricola* 65, 76–87.  
605 <https://doi.org/10.1590/S0103-90162008000700013>

606

607 Gouache, D., Léon, M. S., Duyme, F., Braun, P., 2015. A novel solution to the variable  
608 selection problem in Window Pane approaches of plant pathogen – Climate models:  
609 Development, evaluation and application of a climatological model for brown rust of wheat.  
610 *Agricultural and Forest Meteorology* 205 pp. 51-59.  
611 <http://dx.doi.org/10.1016/j.agrformet.2015.02.013>

612

613 Hack, H., Bleiholder, H., Buhr, L., Meier, U., Schnock-Fricke, U., Weber, E., Witzemberger,  
614 A., 1992. Einheitliche Codierung der phänologischen Entwicklungsstadien mono- und  
615 dikotyler Pflanzen – Erweiterte BBCH-Skala, *AllgemeinNachrichtenbl. Deut.*  
616 *Pflanzenschutzd.*, 44 pp. 265-270.

617

618

619

620Hrustić, J., Mihajlović, M., Grahovac, M., Delibašić, G., Bulajić, A., Krstić, B., Tanović, B.,  
6212012. Genus *Monilinia* on pome and stone fruit species. *Pesticidi i fitomedicina* 27, 283–297.

622

623Jeschke, J.M., Strayer, D.L., 2008. Usefulness of Bioclimatic Models for Studying Climate  
624Change and Invasive Species. *Annals of the New York Academy of Sciences* 1134, 1–24.  
625<https://doi.org/10.1196/annals.1439.002>

626

627Jouzel, J., Déqué, M., Jouini, M., Planton, S., Vautard, R., 2014. Le climat de la France au  
628XXIe siècle (No. 4). Ministère de l'Ecologie, du Développement durable et de l'Energie.

629

630Juroszek P, von Tiedemann A, 2015. Linking plant disease models to climate change scenarios  
631to project future risks of crop diseases: a review. *Journal of Plant Diseases and Protection* 122,  
6323-15.

633

634Koball, D.C., Wilcox, W.F., Seem, R.C., 1997. Influence of incubation-period humidity on the  
635development of brown rot blossom blight of sour cherry. *Phytopathology* 87, 42–49.

636

637

638

639Lane, C.R., 2002. A synoptic key for differentiation of *Monilinia fructicola*, *M. fructigena* and  
640*M. laxa*, based on examination of cultural characters. *EPPO Bulletin* 32, 489–493.

641

642Launay M, Caubel J, Bourgeois G, Huard F, Cortazar-Atauri IG, Bancal MO, Brisson N,  
6432014. Climatic indicators for crop infection risk: Application to climate change impacts on

644 five major foliar diseases in Northern France. *Agriculture, Ecosystems, and Environment* 197,  
645 147-158.

646

647 Leyronas, C., Nicot, P.C., 2013. Monitoring viable airborne inoculum of *Botrytis cinerea* in  
648 the South-East of France over 3 years: relation with climatic parameters and the origin of air  
649 masses. *Aerobiologia* 29. <https://doi.org/10.1007/s10453-012-9280-0>

650

651 Lichou, J., Jay, M., 2012. Monographie abricot. CTIFL.

652

653 Luo, Y., Morgan, D.P., Michailides, T.J., 2001. Risk analysis of brown rot blossom blight of  
654 prune caused by *Monilinia fructicola*. *Phytopathology* 91, 759–768.

655

656 Magarey, R.D., Sutton, T.B., Thayer, C.L., 2005. A Simple Generic Infection Model for Foliar  
657 Fungal Plant Pathogens. *Phytopathology* 95, 92–100. [https://doi.org/10.1094/PHYTO-95-](https://doi.org/10.1094/PHYTO-95-6580092)  
658 0092

659

660 Maier, H.R., Guillaume, J.H.A., van Delden, H., Ridell, G.A., Haasnoot, M., Kwakkel, J.H.,  
661 2016. An uncertain future, deep uncertainty, scenarios, robustness and adaptation: How do  
662 they fit together? *Environmental Modelling & Software* 81, 154-164.  
663 <https://doi.org/10.1016/j.envsoft.2016.03.014>

664

665 Morales G., Moragrena, C., Montesinos, E., Llorente, I., 2018. Environmental and inoculum  
666 effects on epidemiology of bacterial spot disease of stone fruits and development of a disease  
667 forecasting system. *Eur J Plant Pathol* 152 pp. 635-651. [https://doi.org/10.1007/s10658-018-](https://doi.org/10.1007/s10658-018-6681507-7)  
668 1507-7

669

670

671

672

673

674Oliveira Lino, L., Pacheco, I., Mercier, V., Faoro, F., Bassi, D., Bornard, I., Quilot-Turion, B.,

6752016. Brown Rot Strikes *Prunus* Fruit: An Ancient Fight Almost Always Lost. Journal of

676Agricultural and Food Chemistry 64, 4029–4047. <https://doi.org/10.1021/acs.jafc.6b00104>

677

678Pachauri, R.K., Meyer, L., Equipe de rédaction principale, 2014. IPCC, 2014: Climate Change

6792014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment

680Report of the Intergovernmental Panel on Climate Change (No. 5). GIEC, Genève.

681

682Parveau, C.-É., Gomez, C., Mercier, V., Brun, L., Guillermin, A., Clauzel, G., Broquaire, J.-

683M., Audergon, J.-M., 2016. Guide des sensibilités variétales aux bio-agresseurs - Fiche n°3 :

684Monilioses sur fleur. L'Arboriculture Fruitière 5.

685

686

687

688

689

690

691

692

693

694

695

696Tamm, L., Minder, C.E., Flückiger, W., 1995. Phenological analysis of brown rot blossom

697blight of sweet cherry caused by *Monilinia laxa*. *Phytopathology* 85, 401–408.

698

699Trebicki P, Finlay K, 2019. Pests and diseases under climate change; it's threats to food

700security, 229-248. In: Yadav et al (eds). *Food Security and Climate Change*. Wiley Blackwell.

701

702Willmott, C.J., 1981. ON THE VALIDATION OF MODELS. *Physical Geography* 2, 184–

703194. <https://doi.org/10.1080/02723646.1981.10642213>

704

Fig.1: Localisation of the grids studied and their associated clusters (Best viewed in colour).




Fig. 2: Method used to calculate climate change for a given year  $n$  and cultivar. (1) Computations for year  $n$  start on the 15<sup>th</sup> of October of year  $n-1$ . For the dormancy model, it is assumed that the cold needs of apricot have not started to be fulfilled prior to this date. (2) Phenological models compute the F50 date given the daily average temperature. (3) The expand of the blooming period around F50 is estimated given the distribution of disease-sensitive stages. (4) Daily rainfall and temperature during the blooming period are extracted and (5) CIB is calculated for year  $n$






Fig 3: Simulated vs. Observed values for blossom blight incidence ( $I$ , dimensionless).  $R^2 = 0.941$ ,

Intercept = 0.009; slope = 0.935.




Fig 4: 3D representation of our model as a surface response of blossom blight incidence to rainfall (mm) and temperature (°C) (left) and residuals associated with each point of the dataset as a function of rainfall and temperature (right). The size of each point represents the value of the residual and the colour the sign of the residual. Lines correspond to different predicted levels of incidence.




Fig 5: Predicted Climatic Index of cumulated Blight risk (CIB, cumulated %) versus observed Twig Blight Incidence (TBI, dimensionless).





Fig 6: Evolution of the mid-blooming date (DOY) for the different cultivars and clusters under scenario RCP8.5. Linear regression lines are shown for clarity.



Cultivar ··· Beliana — Bergarouge ·-· Bergeron - - Rouge

Fig 7: Mean temperature (°C) and mean cumulated rainfall (mm) experienced during blooming for the different cultivars and clusters. The start of the line represents the situation during the [1970-1999] period and the head of the arrow the situation forecasted for the [2070-2099] period.





Fig 8: Mean simulated CIB ( $\pm$ one standard deviation) by period (p: past, nf: near future, pf: far future) for each cluster and cultivar under RCP 8.5. The colours are red for a significant increase in the risk and green for a significant reduction in the risk. Letters are the result of the Kruskal-Wallis comparison between periods within one Cultivar-Cluster panel (best viewed in colour).



*Table.1: Parameter values of the fitted model. Bolt parameters were chosen priori to optimisation according to the method described by Tamm et al. 1995*

Parameter	Value
$T_{\min}$	0
$T_{\max}$	31
E	0.029
$i_{\max}$	0.79
$m$	0.9
$\rho_1$	1.575
$\rho_2$	1.965
$\gamma_1$	1.709
$\gamma_2$	1.472