



Wolbachia can enhance Plasmodium infection in mosquitoes: implications for malaria control?

Grant Hughes, Ana R Rivero, Jason Rasgon

► To cite this version:

Grant Hughes, Ana R Rivero, Jason Rasgon. Wolbachia can enhance Plasmodium infection in mosquitoes: implications for malaria control?. PLoS Pathogens, 2014, 10 (9), pp.e1004182. 10.1371/journal.ppat.1004182 . hal-02361640

HAL Id: hal-02361640

<https://hal.science/hal-02361640>

Submitted on 21 Sep 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Opinion

Wolbachia Can Enhance Plasmodium Infection in Mosquitoes: Implications for Malaria Control?

Grant L. Hughes¹, Ana Rivero², Jason L. Rasgon^{1*}

1 The Huck Institutes of The Life Sciences, The Center for Infectious Disease Dynamics and the Department of Entomology, Pennsylvania State University, State College, Pennsylvania, United States of America, **2** Maladies Infectieuses et Vecteurs: Écologie, Génétique, Évolution et Contrôle, MIVEGEC (UMR CNRS-UM1-UM2 5290, IRD 224), Montpellier, France

The symbiotic bacterium *Wolbachia* is an attractive agent for vector-borne pathogen control. It has long been studied for its ability to manipulate host reproduction and spread into arthropod populations [1]. These properties, coupled with the recently identified ability to inhibit diverse pathogens [2–6], open avenues for its use in controlling vector-borne disease. Numerous *Wolbachia*-based control strategies are being investigated (reviewed in [7–9]), with some studies having progressed to field trials [10,11]. However, a worrying trend is emerging whereby *Wolbachia* infections have been demonstrated to enhance rather than suppress pathogens in some systems [12–18]. *Plasmodium* parasites, which are the causal agent of malaria, seem particularly prone to *Wolbachia*-mediated pathogen enhancement [13–16].

Wolbachia-based strategies have been proposed to control malaria [19]. *Anopheles* mosquitoes (the vectors of human malaria parasites) are not naturally infected by *Wolbachia* [20,21], but artificial transfer of this bacterium between species can be accomplished in the laboratory (reviewed in [22]). Pathogen interference phenotypes appear to be most prominent when *Wolbachia* is transferred into a novel host [16,23]. Given that *Anopheles* are for the most part naturally uninfected by *Wolbachia* (but see [24]), they can be considered an open niche for infection and a prime mosquito genus for *Wolbachia*-based control strategies. However, the main impediment for developing a control strategy is the difficulty in creating a stable artificial infection in *Anopheles* [19]. While examining *Plasmodium* interference in a stably infected host is the gold standard, a more convenient system is to transiently infect mosquitoes by intrathoracic microinjection. Using this system, the infection persists during the lifetime of the transinfected individual but is not transmitted to its offspring. Transient infection allows the rapid assessment of *Wolbachia*-host interactions without the need for generating stable artificial infections [5]. It is uncertain how representative transient infec-

tions are of stable inherited associations; however, similarities in tissues tropism and fitness costs incurred upon the host between stable and transiently infected *Anopheles* mosquitoes are evident [5,14,25]. Furthermore, both types of infection have been shown to inhibit the human malaria parasite *Plasmodium falciparum* [5,25]. However, studies using transient infection models have found that *Wolbachia* can enhance certain *Plasmodium* species [13,14].

The *Plasmodium* interference phenotype is therefore not universal, but context dependent. While *P. falciparum* is suppressed by the *wAlbB* strain of *Wolbachia* from *Aedes albopictus* [5,25], transient infections have shown the opposite effect on rodent malaria parasites. *Anopheles gambiae* transiently infected with *wAlbB* exhibited enhanced *P. berghei* development at the oocyst stage [14]. Similarly, *wAlbB* increased the number of *P. yoelii* oocysts in *An. stephensi*, although the phenotype was modulated by temperature [13]. At a temperature optimal for parasite development, *Wolbachia* increased parasite intensity compared to uninfected controls, but at warmer temperatures, *Wolbachia* inhibited *Plasmodium* development [13].

While *P. falciparum* is a major parasite in sub-Saharan Africa, four other parasites also cause human malaria worldwide: *P. malariae*, *P. ovale*, *P. knowlesi*, and *P. vivax* (the etiological agent of the most prevalent form of relapsing malaria). To our knowledge, the effect of *Wolbachia* on these other human *Plasmodium* parasites

is unknown. The question is relevant for two reasons. First, the precedent that a particular *Wolbachia* strain can inhibit one parasite yet enhance another has already been documented [5,14], indicating that effects on parasites can be species-specific. Troublingly, *P. malariae*, *P. ovale*, *P. knowlesi*, and *P. vivax* are phylogenetically more closely related to rodent malaria parasites, which are enhanced by *Wolbachia* infections [13,14], than they are to *P. falciparum* (Figure 1) [26,27]. Second, many human *Plasmodium* parasites occur in sympatry and are transmitted by the same vectors. A case in point is *P. falciparum* and *P. vivax*, both of which occur in sympatry over large stretches of the Asian continent where they are both transmitted by *An. stephensi* [28,29]. Any potential control strategy devised in regions where more than one parasite species occurs needs to thoroughly investigate the effect of *Wolbachia* on all parasite species transmitted by the vector, as well as other pathogens such as filarial worms or arboviruses (both as single infections and in the context of coinfections) to ensure that *Wolbachia*-infected mosquitoes do not inadvertently enhance transmission of secondary pathogens.

While difficult, forecasting the long-term evolutionary response in this tripartite relationship between *Wolbachia*, *Plasmodium*, and *Anopheles* is very important. Natural *Wolbachia*-mosquito associations in which the symbiont and the host have tightly coevolved exist and may provide powerful models for studying the long-term evolutionary effects of *Wolbachia*

Citation: Hughes GL, Rivero A, Rasgon JL (2014) *Wolbachia* Can Enhance *Plasmodium* Infection in Mosquitoes: Implications for Malaria Control? PLoS Pathog 10(9): e1004182. doi:10.1371/journal.ppat.1004182

Editor: Glenn F. Rall, The Fox Chase Cancer Center, United States of America

Published: September 4, 2014

Copyright: © 2014 Hughes et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This research was funded by NIH grant R21AI070178 to JLR. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* Email: jlr54@psu.edu

Wolbachia association with host

- ⊙ Natural infection
- Stable artificial infection
- ◆ Transient infection




Figure 1. Representative phylogenetic dendrogram of *Plasmodium* parasites, their vertebrate hosts, and the influence of *Wolbachia* infection on parasite development within the mosquito vector. The protective effect of *Wolbachia* is variable and dependent on the *Wolbachia* strain and the insect host background, suggesting that complex tripartite interactions influence the effect on *Plasmodium*. The type of association between *Wolbachia* with the vector may also influence *Plasmodium*. Only one human malaria parasite (*P. falciparum*) has been assessed, while the effect of *Wolbachia* infection on the other four human parasites is unknown. Arrows indicate suppression (green), enhancement (red), or no effect (grey) of *Plasmodium*. The type of association within the host is depicted by symbols (target: natural infection, square: stable artificial infection, diamond: transient artificial infection). Numbers indicate: (1) the phenotype is temperature sensitive, (2) *Wolbachia* infection also increases insect life span [31], which has implications for pathogen transmission. Phylogeny was reconstructed based on work from Carlton et al. [26] and Martinsen et al. [27].

doi:10.1371/journal.ppat.1004182.g001

infections. The evidence currently available suggests that natural *Wolbachia* infections can also enhance malaria parasite development within the mosquito. *Aedes fluviatilis* naturally infected with the wFlu *Wolbachia* strain had a significantly higher number of *P. gallinaceum* oocysts compared to an *Ae. fluviatilis* line which had been cleared of the *Wolbachia* infection [15]. *Ae. fluviatilis* is not,

however, a natural vector of *P. gallinaceum*, and it is well known that the outcome of experiments using such laboratory models can differ significantly from those of natural mosquito-*Plasmodium* combinations (e.g., Boete [30]). Recent studies carried out in *Culex pipiens* mosquitoes, which are naturally infected with the wPip *Wolbachia* strain and transmit the avian malaria parasite *P.*

relictum, have also demonstrated *Plasmodium* enhancement. In this natural system, *Wolbachia* protects the mosquito host against the detrimental fitness effects incurred by *Plasmodium* infection [31] and increases the susceptibility of *C. pipiens* to *P. relictum*, with wPip-infected mosquitoes having a higher prevalence of *Plasmodium* sporozoites in the salivary glands [16]. These studies show that the

Plasmodium-inhibiting properties of *Wolbachia* are far from universal; certain mosquito–*Wolbachia*–*Plasmodium* combinations and experimental conditions transform *Wolbachia*-infected mosquitoes into better vectors of malaria parasites. This is worrisome for the general implementation of *Wolbachia*-based control strategies.

Given that *Wolbachia*-based control strategies will use stable transinfected mosquitoes, the key question is whether stable and natural infections will behave in the same way. The stable transfer of *Wolbachia* into the host likely alters many aspects of host homeostasis, as evidenced by the novel phenotypes induced by infection [32–34], and as such, these associations likely differ from natural associations where *Wolbachia* and its host have coevolved. Another question is whether stable artificial infections will evolve over time. Theory and empirical studies show that these maternally transmitted bacteria will tend to evolve towards mutualistic associations with their host [35–38]. However, the evolutionary outcomes of pathogen interference or enhancement are harder to predict. A more complete mechanistic understanding of how *Wolbachia* infection modulates *Plasmodium* parasites is critical to address these important evolutionary questions and to evaluate if they are likely to occur in timescales relevant for disease control.

To date, two stable artificial *Wolbachia* transinfections have been assessed for their

effect on *Plasmodium*. First, an *Aedes aegypti* line infected with *wMelPop* had inhibited *P. gallinaceum* infection [4]; *Ae. aegypti* is not, however, the natural vector of this parasite. Second, and more recently, the *wAlbB* strain was stably transferred into *An. stephensi*, one of the main vectors of human malaria in Asia [25]. This groundbreaking work demonstrated that stable artificial infections in epidemiologically relevant malaria vectors are feasible, and that *P. falciparum* can be inhibited by *Wolbachia* within its natural vector. If the severe fitness effects induced by *Wolbachia* in *Anopheles* can be overcome [25], then this approach is highly promising.

The work by Bian and colleagues [25] dramatically enhances the prospect for the use of *Wolbachia* in a malaria control strategy, but many questions still remain. What are the effects of *Wolbachia* on the other four species of *Plasmodium* parasites that infect humans? How relevant are transient infection models? Do some strains of *Wolbachia* enhance pathogens in a field context? What are the long-term evolutionary consequences of novel *Wolbachia*-host associations on *Plasmodium* development within the insect host? What are the mechanisms behind pathogen interference and enhancement of *Wolbachia* on *Plasmodium* parasites, and are the mechanisms of enhancement seen in rodent and avian model systems relevant to human malaria parasites? How influential are environmental variables on

pathogen inhibition phenotypes? While many of these questions may be difficult to answer in the short term, assessing the relevance of transient infections would seem within the grasp of the scientific community. Although challenging, understanding the evolutionary consequences of novel *Wolbachia* associations on pathogen transmission and identifying the mechanisms behind *Wolbachia* modulation of *Plasmodium* is critical for developing effective control strategies and assessing their long-term feasibility. Insights from non-*Anopheline* systems where *Wolbachia* naturally infects the vector may be useful in this regard [16,31,39].

In conclusion, *Wolbachia*-based control of vector-borne pathogens is a promising novel strategy that has many advantages over other conventional and contemporary control methods. The development of a stable infection in *Anopheles* means the prospect of *Wolbachia*-based control of malaria can now be entertained [25], but many important questions need to be resolved before this idea can become a reality. While the concerns raised here focus on *Plasmodium*, these issues are relevant for *Wolbachia* control of any vector-borne pathogen [18]; we suggest that transinfected mosquitoes intended for release into nature should be assessed for inhibition (or lack thereof) of all relevant pathogens circulating in the system.

References

- Werren JH, Baldo L, Clark ME (2008) *Wolbachia*: master manipulators of invertebrate biology. *Nat Rev Microbiol* 6: 741–751.
- Hedges LM, Brownlie JC, O'Neill SL, Johnson KN (2008) *Wolbachia* and virus protection in insects. *Science* 322: 702.
- Kambris Z, Cook PE, Phuc HK, Sinkins SP (2009) Immune activation by life-shortening *Wolbachia* and reduced filarial competence in mosquitoes. *Science* 326: 134–136.
- Moreira LLA, Iturbe-Ormaetxe I, Jeffery JA, Lu G, Pyke AAT, et al. (2009) A *Wolbachia* symbiont in *Aedes aegypti* limits infection with Dengue, Chikungunya, and *Plasmodium*. *Cell* 139: 1268–1278.
- Hughes GL, Koga R, Xue P, Fukatsu T, Rasgon JL (2011) *Wolbachia* infections are virulent and inhibit the human malaria parasite *Plasmodium falciparum* in *Anopheles gambiae*. *PLoS Pathog* 7: e1002043.
- Kambris Z, Blagborough AM, Pinto SB, Blagrove MSC, Godfray HCJ, et al. (2010) *Wolbachia* stimulates immune gene expression and inhibits *Plasmodium* development in *Anopheles gambiae*. *PLoS Pathog* 6: e1001143.
- McGraw EA, O'Neill SL (2013) Beyond insecticides: new thinking on an ancient problem. *Nat Rev Microbiol* 11: 181–193.
- Bourtzis K, Dobson SL, Xi Z, Rasgon JL, Calvitti M, et al. (2014) Harnessing mosquito–*Wolbachia* symbiosis for vector and disease control. *Acta Trop* 132S: S150–S163.
- Iturbe-Ormaetxe I, Walker T, O'Neill SL (2011) *Wolbachia* and the biological control of mosquito-borne disease. *EMBO Rep* 12: 508–518.
- Hoffmann AA, Montgomery BL, Popovici J, Iturbe-Ormaetxe I, Johnson PH, et al. (2011) Successful establishment of *Wolbachia* in *Aedes* populations to suppress dengue transmission. *Nature* 476: 454–457.
- Walker T, Johnson PH, Moreira LA, Iturbe-Ormaetxe I, Freni FD, et al. (2011) The *wMel* *Wolbachia* strain blocks dengue and invades caged *Aedes aegypti* populations. *Nature* 476: 450–453.
- Graham RI, Grzywacz D, Mushobizi WL, Wilson K (2012) *Wolbachia* in a major African crop pest increases susceptibility to viral disease rather than protects. *Ecol Lett* 15: 993–1000.
- Murdock CC, Blanford S, Hughes GL, Rasgon JL, Thomas MB (2013) Temperature alters malaria transmission blocking by *Wolbachia*. *Sci Rep* 4: 3932.
- Hughes GL, Vega-Rodríguez J, Xue P, Rasgon JL (2012) *Wolbachia* strain *wAlbB* enhances infection by the rodent malaria parasite *Plasmodium berghei* in *Anopheles gambiae* mosquitoes. *Appl Environ Microbiol* 78: 1491–1495.
- Baton LA, Pacidônio EC, Gonçalves DDS, Moreira LA (2013) *wFlu*: characterization and evaluation of a native *Wolbachia* from the mosquito *Aedes fluviatilis* as a potential vector control agent. *PLoS ONE* 8: e59619.
- Zélé F, Nicot A, Berthomieu A, Weill M, Duron O, et al. (2013) *Wolbachia* increases susceptibility to *Plasmodium* infection in a natural system. *Proc Biol Sci* 281: 20132837.
- Hussain M, Lu G, Torres S, Edmonds JH, Kay BH, et al. (2013) Effect of *Wolbachia* on replication of West Nile virus in a mosquito cell line and adult mosquitoes. *J Virol* 87: 851–858.
- Dodson BL, Hughes GL, Paul O, Matarachio AC, Kramer LD, et al. (2014) *Wolbachia* Enhances West Nile Virus (WNV) Infection in the Mosquito *Culex tarsalis*. *PLoS Negl Trop Dis* 8: e2965.
- Walker T, Moreira LA (2011) Can *Wolbachia* be used to control malaria? *Mem Inst Oswaldo Cruz* 106 (Suppl. 1): 212–217.
- Ricci I, Cancrini G, Gabrielli S, D'Amelio S, Favi G (2002) Searching for *Wolbachia* (Rickettsiales: Rickettsiaceae) in mosquitoes (Diptera: Culicidae): large polymerase chain reaction survey and new identifications. *J Med Entomol* 39: 562–567.
- Rasgon JL, Scott TW (2004) An initial survey for *Wolbachia* (Rickettsiales: Rickettsiaceae) infections in selected California mosquitoes (Diptera: Culicidae). *J Med Entomol* 41: 255–257.
- Hughes GL, Rasgon JL (2014) Transinfection: a method to investigate *Wolbachia*-host interactions and control arthropod-borne disease. *Insect Mol Biol* 23: 141–151.
- Bian G, Xu Y, Lu P, Xie Y, Xi Z (2010) The endosymbiotic bacterium *Wolbachia* induces resistance to Dengue virus in *Aedes aegypti*. *PLoS Pathog* 6: e1000833.
- Baldini F, Segata N, Pompon J, Marcenac P, Robert Shaw W, et al. (2014) Evidence of natural

- Wolbachia* infections in field populations of *Anopheles gambiae*. Nat Commun 6: 3985.
25. Bian G, Joshi D, Dong Y, Lu P, Zhou G, et al. (2013) *Wolbachia* invades *Anopheles stephensi* populations and induces refractoriness to *Plasmodium* infection. Science 340: 748–751.
 26. Carlton JM, Escalante AA, Neafsey D, Volkman SK (2008) Comparative evolutionary genomics of human malaria parasites. Trends Parasitol 24: 545–550.
 27. Martinsen ES, Perkins SL, Schall JJ (2008) A three-genome phylogeny of malaria parasites (*Plasmodium* and closely related genera): evolution of life-history traits and host switches. Mol Phylogenet Evol 47: 261–273.
 28. Korgaonkar NS, Kumar A, Yadav RS, Kabadi D, Dash AP (2012) Mosquito biting activity on humans & detection of *Plasmodium falciparum* infection in *Anopheles stephensi* in Goa, India. Indian J Med Res 135: 120–126.
 29. Adak T, Singh OP, Das MK, Wattal S, Nanda N (2005) Comparative susceptibility of three important malaria vectors *Anopheles stephensi*, *Anopheles fluviatilis*, and *Anopheles sudaicus* to *Plasmodium vivax*. J Parasitol 91: 79–82.
 30. Boëte C (2005) Malaria parasites in mosquitoes: laboratory models, evolutionary temptation and the real world. Trends Parasitol 21: 445–447.
 31. Zélé F, Nicot A, Duron O, Rivero A (2012) Infection with *Wolbachia* protects mosquitoes against *Plasmodium*-induced mortality in a natural system. J Evol Biol 25: 1243–1252.
 32. Clancy DJ, Hoffmann AA (1997) Behavior of *Wolbachia* endosymbionts from *Drosophila simulans* in *Drosophila serrata*, a novel host. Am Nat 149: 975–988.
 33. Suh E, Mercer D, Fu Y, Dobson SL (2009) Pathogenicity of life-shortening *Wolbachia* in *Aedes albopictus* after transfer from *Drosophila melanogaster*. Appl Env Microbiol 75: 7783–7788.
 34. Bouchon D, Rigaud T, Juchault P (1998) Evidence for widespread *Wolbachia* infection in isopod crustaceans: molecular identification and host feminization. Proc Biol Sci 265: 1081–1090.
 35. McGraw EA, Merritt DJ, Droller JN, O'Neill SL (2002) *Wolbachia* density and virulence attenuation after transfer into a novel host. Proc Natl Acad Sci U S A 99: 2918–2923.
 36. Weeks AR, Turelli M, Harcombe WR, Reynolds KT, Hoffmann AA (2007) From parasite to mutualist: rapid evolution of *Wolbachia* in natural populations of *Drosophila*. PLoS Biol 5: E114.
 37. Turelli M (1994) Evolution of incompatibility-inducing microbes and their hosts. Evolution 48: 1500–1513.
 38. Carrington LB, Hoffmann AA, Weeks AR (2010) Monitoring long-term evolutionary changes following *Wolbachia* introduction into a novel host: the *Wolbachia* popcorn infection in *Drosophila simulans*. Proc Biol Sci. 277: 2059–2068.
 39. Hughes GL, Samuels SK, Shaikh K, Rasgon JL, Vardo-Zalik AM (2014) Discrimination of the *Plasmodium mexicanum* vectors *Lutzomyia stewarti* and *Lutzomyia vexator* by a PCR-RFLP assay and *Wolbachia* infection. J Vector Ecol 39: 224–227.