


## Celtic Sea glaciation : open questions

- How far have ice sheets advanced SW into the Celtic Sea? Surprisingly, we are still not sure.
- Subglacial and glacimarine sediments cored in the 1970s on and between a system of shelf-crossing seafloor ridges<sup>1,2</sup> indicated ice extent to at least the mid-shelf of the Irish and UK sectors<sup>3,4</sup> (Fig. 1).
- Recent coring of subglacial sediments on and adjacent to ridges indicates ice extent >100 km farther seaward to the Irish & UK shelf edge<sup>5-7</sup> (Fig. 1, red line).
- Ice extent in the French sector remains unconstrained.
- What are the ridges? Integration of cores with seismic data suggests they could be a) post-glacial tidal sand banks, or b) relict glaciofluvial landforms<sup>5,6</sup>.



## Mid-shelf landforms from multibeam bathymetry

### Rectilinear network of seafloor features :

- Ridge segments**: en échelon elements up to 55 m high, 10 km wide and 10s km long
- Transverse ribs**: up to 15 m high, 1 km wide and 10 km long; arcuate to linear, no consistent spacing or asymmetry; occur both lateral to ridges and as axial extensions forming low ridges (see A-B vs C-D)
- Sand waves**: 3-7 m high, up to 0.6 km wide; consistent spacing, asymmetry, widths; occur in south of area, on and between ridges (see I-J)




Fig. 3 – perspective view of sand waves vs ribs and ridges




Fig. 2 – multibeam coverage (GLAMAR 2009) of a 25 x 100 km area of Irish-UK mid-shelf (see Fig. 1)



## Regional submarine landforms

### North Celtic Sea : incisions<sup>8</sup> (tunnel-valleys)

- up to 250 m in relief, 5 km wide, 20 km long
- at least 3 stratigraphic levels (mainly infilled)
- meltwater drainage during at least 3 glaciations

### Celtic Sea shelf : seafloor ridges

- up to 60 m high, 10 km wide, 300 km long
- fan-shaped, apex points to North Celtic Sea
- axial rotation of up to 80° (W-E to almost N-S)
- axial convergences seaward, some landward
- up & down axial profiles (segments 10s km long)



## Glacigenic drape of ridges and ribs ?

- Ridges shown to be truncated by a regional unconformity overlain by a fining-upward succession<sup>6</sup> (layers A & B, Fig. 5)
- Sample & sonar data show layer B to include gravel (to >1 m size) of presumed glacial origin (ice-raftered or reworked till)<sup>1,2</sup>




Fig. 5 – Stratigraphic model of Irish-UK ridges including coarse layer B'

- Comparison to multibeam features suggests gravel also present in areas of ribs (Fig. 6)




Fig. 6 – BGS seafloor sediment map<sup>1,2</sup> compared with multibeam features

## Are we looking at tidal or glaciofluvial landforms ?

### Tidal banks and sand waves : too big + wrong morphologies ?

- Up to 50% higher than largest known tidal banks, and at upper limit of sand wave heights → low slopes imply that, if degraded, both were originally far bigger than any known features
- Lack morphologies consistent with the cyclonic flow systems typical of tidal systems (curvature into parallelism with bank crests, opposite symmetry on opposite sides of banks...)

### Giant eskerine ridges and glaciofluvial (De Geer) moraines ?

- Comparable to rectilinear networks of giant eskers (up to 10 km wide, 40 m high) and De Geer moraines in Canada and Finland<sup>9,10</sup>, formed by subaqueous outwash during ice margin retreat
- Consistent with evidence of subglacial and glacimarine sediments in ridges and ribs in Irish and UK sectors
- Broad implications for ice sheet extent and dynamics – notably in the French sector




Fig. 8 – Giant eskers (black) and De Geer moraines (red) in Finland; IIa = Salpausselkä glaciofluvial moraines<sup>10</sup>

➤ These hypotheses invite testing through targeted acquisition of data from ridges and ribs in the Irish, UK and French sectors

### References

- Pantin HM, Evans CDR (1984) Quaternary history of the central & southwestern Celtic Sea. *Marine Geology* **57**: 259–293
- Evans CDR et al. (1990) *Geology of the Western English Channel...* UK Offshore Regional Report, BGS; HMSO: 93 pp.
- Scourse JD et al. (1990) Sedimentology... of glacimarine sediments... Celtic Sea. *GSL Special Publication* **53**: 329–347
- Sejrup H-P et al. (2005) Pleistocene glacial history of the NW European continental margin. *Mar Pet Geol* **22**: 1111–1129
- Praeg D, McCarron S, Dove D et al. (2015) Ice sheet extension to the Celtic Sea shelf edge... *Quat Sci Rev* **111**, 107–112
- Lockhart EA, Scourse JD, Praeg D et al. (2018) Stratigraphic investigation... Celtic Sea megaridges... *Quat Sci Rev* **198**: 156–170
- Scourse JD et al. (2019) Advance and retreat of the marine-terminating Irish Sea Ice Stream... *Marine Geology* **412**: 53–68
- Tappin D et al. (1994) *Geology of Cardigan Bay and the Bristol Channel...* UK Offshore Regional Report, BGS; HMSO : 107 pp.
- Vaillette JJ (1986) Former southwesterly ice flows in the Abitibi - Timiskaming region... *Can. J. Earth Sci.* **23** : 1724–1741
- Makinen J (2003) Time-transgressive deposits... glaciofluvial (esker) sediments in Köyliö, SW Finland. *Sedimentology* **50** : 327–360




Fig. 7 – Seismic reflection profiles (top pair, Chirp; bottom, sparker) along and across ridges and ribs, tied to BGS vibrocores (locations on Fig. 2)