



**HAL**  
open science

## Network inference for truncated gaussian data

Anne Gégout-Petit, Aurélie Gueudin, Clémence Karmann

► **To cite this version:**

Anne Gégout-Petit, Aurélie Gueudin, Clémence Karmann. Network inference for truncated gaussian data. European Meeting of Statisticians, Jul 2019, Palermo, Italy. hal-02369239

**HAL Id: hal-02369239**

**<https://hal.science/hal-02369239>**

Submitted on 18 Nov 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Network inference for truncated gaussian data

Anne Gégout, Aurélie Gueudin and Clémence Karmann

Université de Lorraine, Inria Nancy Grand Est, FRANCE

*anne.gegout-petit@univ-lorraine.fr*

EMS 2019

Palermo

# Introduction

- Infer network defined by conditional dependencies

# Introduction

- Infer network defined by conditional dependencies
- Gaussian variables zero inflated by double truncation

# Introduction

- Infer network defined by conditional dependencies
- Gaussian variables zero inflated by double truncation
- Applications for the modelling of interactions between microbacterial populations
  - left truncation: replication phenomena
  - right truncation: non restrictive hypothesis for theoretical results !

# Content

- 1 Modelling
- 2 Estimation process
  - Covariance matrix estimation
  - Precision matrix estimation
- 3 Theoretical results
- 4 Simulation studies

- 1 Modelling
- 2 Estimation process
  - Covariance matrix estimation
  - Precision matrix estimation
- 3 Theoretical results
- 4 Simulation studies

# Modelling

## Modelling

Let  $\mathbf{X} \sim \mathcal{N}_p(0, \Sigma^*)$  be a gaussian  $p$ -vector with  $\Sigma_{jj}^* = 1$  for all  $j$ .  $\mathbf{X}$  is not observed but we observe  $\mathbf{Y}$  defined from  $\mathbf{X}$  by:

$$Y_j = \mathbb{1}_{a_j \leq X_j \leq b_j} X_j \text{ where } a_j < b_j.$$

A non directed graph structure is given by the inverse of the covariance matrix  $\Theta^* := (\Sigma^*)^{-1}$  :

$$\begin{aligned} X_j \longleftrightarrow X_k &\iff X_j \not\perp\!\!\!\perp X_k \mid (X_l)_{l \neq j, k} \\ &\iff \text{cor}(X_j, X_k \mid (X_l)_{l \neq j, k}) \neq 0 \\ &\iff \Theta_{jk}^* \neq 0. \end{aligned}$$



# Aim

## Aim

Infer the latent graph structure given by the precision matrix  $\Theta^*$  of the gaussian vector  $\mathbf{X}$  from the observations of the truncated vector  $\mathbf{Y}$ .

## Theoretical tools - Pairs likelihood

Let us look at the likelihood  $\mathcal{L}_{jk}(\Sigma_{jk}^*, y)$  of the pairs  $(Y_j, Y_k)$ ,  $j < k$  where :

$$\mathcal{L}_{jk}(\sigma, y) = \sum_{a,b=0}^1 \phi_{ab,jk}(\sigma, y_j, y_k) n_{ab}(y_j, y_k),$$

with:

## Theoretical tools - Pairs likelihood

Let us look at the likelihood  $\mathcal{L}_{jk}(\Sigma_{jk}^*, y)$  of the pairs  $(Y_j, Y_k)$ ,  $j < k$  where :

$$\mathcal{L}_{jk}(\sigma, y) = \sum_{a,b=0}^1 \phi_{ab,jk}(\sigma, y_j, y_k) n_{ab}(y_j, y_k),$$

with:

- $n_{ab}(y_j, y_k) = \mathbb{1}_{\zeta_j=a, \zeta_k=b}$  où  $\zeta_l = \begin{cases} 1 & \text{si } y_l \in [a_l, b_l] \setminus \{0\}, \\ 0 & \text{si } y_l = 0. \end{cases}$

## Theoretical tools - Pairs likelihood

Let us look at the likelihood  $\mathcal{L}_{jk}(\Sigma_{jk}^*, y)$  of the pairs  $(Y_j, Y_k)$ ,  $j < k$  where :

$$\mathcal{L}_{jk}(\sigma, y) = \sum_{a,b=0}^1 \phi_{ab,jk}(\sigma, y_j, y_k) n_{ab}(y_j, y_k),$$

with:

- $n_{ab}(y_j, y_k) = \mathbb{1}_{\zeta_j=a, \zeta_k=b}$  où  $\zeta_l = \begin{cases} 1 & \text{si } y_l \in [a_l, b_l] \setminus \{0\}, \\ 0 & \text{si } y_l = 0. \end{cases}$
- $\phi_{11,jk}(\sigma, y_j, y_k) = f(y_j, y_k, \sigma)$

## Theoretical tools - Pairs likelihood

Let us look at the likelihood  $\mathcal{L}_{jk}(\Sigma_{jk}^*, y)$  of the pairs  $(Y_j, Y_k)$ ,  $j < k$  where :

$$\mathcal{L}_{jk}(\sigma, y) = \sum_{a,b=0}^1 \phi_{ab,jk}(\sigma, y_j, y_k) n_{ab}(y_j, y_k),$$

with:

- $n_{ab}(y_j, y_k) = \mathbb{1}_{\zeta_j=a, \zeta_k=b}$  où  $\zeta_l = \begin{cases} 1 & \text{si } y_l \in [a_l, b_l] \setminus \{0\}, \\ 0 & \text{si } y_l = 0. \end{cases}$
- $\phi_{11,jk}(\sigma, y_j, y_k) = f(y_j, y_k, \sigma)$
- $\phi_{01,jk}(\sigma, y_j, y_k) = \phi_{01,jk}(\sigma, y_k) = \int_{[a_j, b_j]^c} f(x, y_k, \sigma) dx$
- $\phi_{10,jk}(\sigma, y_j, y_k) = \phi_{10,jk}(\sigma, y_j) = \int_{[a_k, b_k]^c} f(y_j, y, \sigma) dy$

## Theoretical tools - Pairs likelihood

Let us look at the likelihood  $\mathcal{L}_{jk}(\Sigma_{jk}^*, y)$  of the pairs  $(Y_j, Y_k)$ ,  $j < k$  where :

$$\mathcal{L}_{jk}(\sigma, y) = \sum_{a,b=0}^1 \phi_{ab,jk}(\sigma, y_j, y_k) n_{ab}(y_j, y_k),$$

with:

- $n_{ab}(y_j, y_k) = \mathbb{1}_{\zeta_j=a, \zeta_k=b}$  où  $\zeta_l = \begin{cases} 1 & \text{si } y_l \in [a_l, b_l] \setminus \{0\}, \\ 0 & \text{si } y_l = 0. \end{cases}$
- $\phi_{11,jk}(\sigma, y_j, y_k) = f(y_j, y_k, \sigma)$
- $\phi_{01,jk}(\sigma, y_j, y_k) = \phi_{01,jk}(\sigma, y_k) = \int_{[a_j, b_j]^c} f(x, y_k, \sigma) dx$
- $\phi_{10,jk}(\sigma, y_j, y_k) = \phi_{10,jk}(\sigma, y_j) = \int_{[a_k, b_k]^c} f(y_j, y, \sigma) dy$
- $\phi_{00,jk}(\sigma, y_j, y_k) = \phi_{00,jk}(\sigma) = \iint_{[a_j, b_j]^c \times [a_k, b_k]^c} f(x, y, \sigma) dx dy.$

- 1 Modelling
- 2 Estimation process
  - Covariance matrix estimation
  - Precision matrix estimation
- 3 Theoretical results
- 4 Simulation studies

# Context

Let  $\mathbf{Y} := (Y^{(1)}, \dots, Y^{(n)})$  be an i.i.d.  $n$ -sample du vecteur  $Y$ . For every  $j < k$ , let us denote:

- $\mathcal{L}_{jk}^{(n)}(\Sigma_{jk}^*, \mathbf{y})$  the likelihood of the pairs sample  $((Y_j^{(i)}, Y_k^{(i)}))_{1 \leq i \leq n}$  given by:

$$\mathcal{L}_{jk}^{(n)}(\sigma, \mathbf{y}) = \prod_{i=1}^n \mathcal{L}_{jk}(\sigma, y^{(i)}).$$

- $L_{jk}^{(n)}(\Sigma_{jk}^*, \mathbf{y})$  the log-likelihood  $L_{jk}^{(n)}(\Sigma_{jk}^*, \mathbf{y}) = \log(\mathcal{L}_{jk}^{(n)}(\sigma, \mathbf{y}))$



## Step 1: covariance matrix $\Sigma^*$

- empirical covariance matrix from  $\mathbf{Y}$ : poor performance

## Step 1: covariance matrix $\Sigma^*$

- empirical covariance matrix from  $\mathbf{Y}$ : poor performance
- Likelihood Maximisation of  $\mathbf{Y}$  :  $2^p$  terms...

## Step 1: covariance matrix $\Sigma^*$

- empirical covariance matrix from  $\mathbf{Y}$ : poor performance
- Likelihood Maximisation of  $\mathbf{Y}$  :  $2^p$  terms...

$\rightsquigarrow$  estimator of  $\Sigma^*$  term to term from the pairs  $(Y_j, Y_k)$ . We estimate each coefficients  $\tilde{\Sigma}_{jk}^{(n)}$  by maximising the log-likelihood of the sample  $((Y_j^{(i)}, Y_k^{(i)}))_{i=1, \dots, n}$ .

## Step 1: covariance matrix $\Sigma^*$

- empirical covariance matrix from  $\mathbf{Y}$ : poor performance
- Likelihood Maximisation of  $\mathbf{Y}$  :  $2^p$  terms...

$\rightsquigarrow$  estimator of  $\Sigma^*$  term to term from the pairs  $(Y_j, Y_k)$ . We estimate each coefficients  $\tilde{\Sigma}_{jk}^{(n)}$  by maximising the log-likelihood of the sample  $((Y_j^{(i)}, Y_k^{(i)}))_{i=1, \dots, n}$ .

### Estimator of the covariance matrix

Estimator  $\tilde{\Sigma}^{(n)}$  of  $\Sigma^*$  is defined by:

$$\tilde{\Sigma}_{jk}^{(n)} = \operatorname{argmax}_{|\sigma| \leq 1} L_{jk}^{(n)}(\sigma, \mathbf{y}). \quad (1)$$

## Step 2: precision matrix $\Theta^*$

Idea : estimate  $\Theta^*$  thanks to the graphical Lasso (Friedman et al. (2007)) for the GGM

$\rightsquigarrow$  is to maximize the penalized log-likelihood of the Gaussian model.

$$\hat{\Theta}^{(n)} = \operatorname{argmax}_{\Theta \succ 0} \log \det(\Theta) - \operatorname{trace}(\Theta \hat{\Sigma}^{(n)}) - \lambda_n \|\Theta\|_{1, \text{off}}. \quad (2)$$

with  $\hat{\Sigma}^{(n)}$  being the empirical covariance matrix of  $\mathbf{X}$

## Step 2: precision matrix $\Theta^*$

Idea : estimate  $\Theta^*$  thanks to the graphical Lasso (Friedman et al. (2007)) for the GGM

$\rightsquigarrow$  is to maximize the penalized log-likelihood of the Gaussian model.

$$\hat{\Theta}^{(n)} = \underset{\Theta \succ 0}{\operatorname{argmax}} \log \det(\Theta) - \operatorname{trace}(\Theta \hat{\Sigma}^{(n)}) - \lambda_n \|\Theta\|_{1,\text{off}}. \quad (2)$$

with  $\hat{\Sigma}^{(n)}$  being the empirical covariance matrix of  $\mathbf{X}$  Problem:

$\Sigma^{(n)}$  unachievable!  $\rightsquigarrow$  replaced by  $\tilde{\Sigma}^{(n)}$

## Step 2: precision matrix $\Theta^*$

Idea : estimate  $\Theta^*$  thanks to the graphical Lasso (Friedman et al. (2007)) for the GGM

$\rightsquigarrow$  is to maximize the penalized log-likelihood of the Gaussian model.

$$\hat{\Theta}^{(n)} = \underset{\Theta \succ 0}{\operatorname{argmax}} \log \det(\Theta) - \operatorname{trace}(\Theta \hat{\Sigma}^{(n)}) - \lambda_n \|\Theta\|_{1,\text{off}}. \quad (2)$$

with  $\hat{\Sigma}^{(n)}$  being the empirical covariance matrix of  $\mathbf{X}$  Problem:

$\Sigma^{(n)}$  unachievable!  $\rightsquigarrow$  replaced by  $\tilde{\Sigma}^{(n)}$

### Estimator of the precision matrix

Estimator  $\tilde{\Theta}^{(n)}$  of  $\Theta^*$  is defined by:

$$\tilde{\Theta}^{(n)} = \underset{\Theta \succ 0}{\operatorname{argmax}} \log \det(\Theta) - \operatorname{trace}(\Theta \tilde{\Sigma}^{(n)}) - \lambda_n \|\Theta\|_{1,\text{off}}. \quad (3)$$

- 1 Modelling
- 2 Estimation process
  - Covariance matrix estimation
  - Precision matrix estimation
- 3 **Theoretical results**
- 4 Simulation studies



# Tools

**Step 1:** Mei & al (2017), *under suitable assumptions, good properties of the population risk can be carried to the empirical risk, even in a non-convex case*

$\rightsquigarrow$  control of  $|\tilde{\Sigma}_{j,k}^{(n)} - \Sigma_{j,k}^*|$

# Tools

**Step 1:** Mei & al (2017),  
 $\rightsquigarrow$  control of  $|\tilde{\Sigma}_{j,k}^{(n)} - \Sigma_{j,k}^*|$

**Step 2:** Ravikumar & al (2011), *estimating the concentration matrix under sparsity conditions without specific distributional assumptions, but rather analyze the estimator in terms of the tail behavior of  $\max_{j,k} |\tilde{\Sigma}_{j,k}^{(n)} - \Sigma_{j,k}^*|$*

# Hypotheses

**(H1)** For all  $j < k$ ,  $|\Sigma_{jk}^*| \neq 1$ .

# Hypotheses

**(H1)** For all  $j < k$ ,  $|\Sigma_{jk}^*| \neq 1$ .

**(H2)** Let  $j < k$ . Let  $g$  be a function:

$$\sigma \in [-1 + \delta, 1 - \delta] \mapsto \mathbb{E}\left(L_{jk}^{(n)}(\sigma, \mathbf{y})\right).$$

- $(-1 + \delta)$  et  $(1 - \delta)$  are not critical points of  $g$ ,
- $g$  admits a finite number of critical points,
- all the critical points of  $g$ , different from  $\Sigma_{jk}^*$ , are non-degenerated, that is:

$$\sigma \neq \Sigma_{jk}^*, \quad g'(\sigma) = 0 \Rightarrow g''(\sigma) \neq 0.$$

# Hypotheses

**(H1)** For all  $j < k$ ,  $|\Sigma_{jk}^*| \neq 1$ .

**(H2)** Let  $j < k$ . Let  $g$  be a function:

$$\sigma \in [-1 + \delta, 1 - \delta] \mapsto \mathbb{E}\left(L_{jk}^{(n)}(\sigma, \mathbf{y})\right).$$

- $(-1 + \delta)$  et  $(1 - \delta)$  are not critical points of  $g$ ,
- $g$  admits a finite number of critical points,
- all the critical points of  $g$ , different from  $\Sigma_{jk}^*$ , are non-degenerated, that is:

$$\sigma \neq \Sigma_{jk}^*, \quad g'(\sigma) = 0 \Rightarrow g''(\sigma) \neq 0.$$

**(H3)** Technical (not written). The underlying intuition is to limit the influence of "non-edged" terms on edge terms.

Intermediate result on the estimated covariance matrix  $\tilde{\Sigma}^{(n)}$ 

## Proposition

Under **(H1)** and, **(H2)**, let  $0 < \rho < 1$ . There exist known constants  $B$ ,  $C$  and  $D$  such that if  $n$  satisfies  $\frac{n}{\log n} \geq C \log\left(\frac{B}{\rho}\right)$ , then the estimated covariance matrix  $\tilde{\Sigma}^{(n)}$  satisfies:

$$\mathbb{P}\left(\|\tilde{\Sigma}^{(n)} - \Sigma^*\|_{\infty} \geq D \sqrt{\frac{\log n}{n} \log\left(\frac{B}{\rho}\right)}\right) \leq \frac{\rho(\rho - 1)}{2} \rho,$$

where  $\|A\|_{\infty} = \max_{j,k \in \{1, \dots, p\}} |A_{jk}|$  is the infinite norm of matrix  $A \in \mathbb{R}^{p^2}$ .

$\rightsquigarrow$  Mei *et al.* (2017)

# Final result about $\tilde{\Theta}^{(n)}$

## Theorem

Under **(H1)**, **(H2)** et **(H3)**. Let  $c > 2$ . There exist known constants  $B$ ,  $C$  and  $D$  such that for all  $n$  satisfying  $\frac{n}{\log n} > f(B, C, D, c, \Sigma^*)$  and

$\lambda_n \propto \sqrt{\frac{\log n}{n} \log(Bp^c)}$ , we have with probability  $1 - \frac{1}{p^{c-2}}$  :

$$(a) \quad \|\tilde{\Theta}^{(n)} - \Theta^*\|_\infty \leq D \sqrt{\frac{\log n}{n} \log(Bp^c)}.$$

(b)  $E(\tilde{\Theta}^{(n)}) \subset E(\Theta^*)$  and vertices  $(j, k)$  is correctly detected as soon as:  $|\Theta_{jk}^*| > D \sqrt{\frac{\log n}{n} \log(Bp^c)}$ .

↪ Ravikumar *et al.* (2011)

- 1 Modelling
- 2 Estimation process
  - Covariance matrix estimation
  - Precision matrix estimation
- 3 Theoretical results
- 4 Simulation studies



## Simulations design

- $p = 100$  covariates,  $n = 500$  observations

## Simulations design

- $p = 100$  covariates,  $n = 500$  observations
- Gaussian  $\mathbf{X}$  with a chain structure (99 edges) :  
 $X_1 \longleftrightarrow X_2 \longleftrightarrow \dots \longleftrightarrow X_{100}$

## Simulations design

- $p = 100$  covariates,  $n = 500$  observations
- Gaussian  $\mathbf{X}$  with a chain structure (99 edges) :  
 $X_1 \longleftrightarrow X_2 \longleftrightarrow \dots \longleftrightarrow X_{100}$
- Different truncation thresholds:
  - Identical thresholds :  $a = -0.5, b = 2$
  - Non identical thresholds  $a = \text{seq}(-1, 0), b = \text{seq}(0.5, 3)$



## Simulations design

- $p = 100$  covariates,  $n = 500$  observations
- Gaussian  $\mathbf{X}$  with a chain structure (99 edges) :  
 $X_1 \longleftrightarrow X_2 \longleftrightarrow \dots \longleftrightarrow X_{100}$
- Different truncation thresholds:
  - Identical thresholds :  $a = -0.5, b = 2$
  - Non identical thresholds  $a = \text{seq}(-1, 0), b = \text{seq}(0.5, 3)$
- Detection rate over 50 repetitions and comparison with naive graphical Lasso directly on truncated data  $\mathbf{Y}$ .

$a = -0.5 ; b = 2$

Nous



Graphical Lasso




$a = \text{seq}(-1, 0)$ ,  $b = \text{seq}(0.5, 3)$


Nous

Graphical Lasso








(a) Structure “cluster”.



(b) Structure “random”.





(c) Structure “hub”.





**Figure:** Detection rates obtained by our method (estimated truncation points) and by the graphical Lasso applied directly to the truncated data. The configurations of the truncation points are "identical only". Detection rates are obtained over 50 independent repetitions for  $n = 500$  observations of  $p = 100$  variables. "cluster" structure; red = true edges.







**Figure:** Detection rates obtained by our method (estimated truncation points) and by the graphical Lasso applied directly to the truncated data. The configurations of the truncation points are "identical only". Detection rates are obtained over 50 independent repetitions for  $n = 500$  observations of  $p = 100$  variables. "random" structure; red = true edges.



**Figure:** Detection rates obtained by our method (estimated truncation points) and by the graphical Lasso applied directly to the truncated data. The configurations of the truncation points are "identical only". Detection rates are obtained over 50 independent repetitions for  $n = 500$  observations of  $p = 100$  variables. "hub" structure; red = true edges.

# Grazie mille !

-  Mei S., Bei Y., and Montanari A. The landscape of empirical risk for non-convex losses. *arXiv preprint arXiv:1607.06534*, 2017.
-  Ravikumar P., Wainwright M. J., Raskutti G., and Yu B. High-dimensional covariance estimation by minimizing  $\ell_1$ -penalized log-determinant divergence. *Electronic Journal of Statistics*, 5:935-980, 2011.