



HAL
open science

Spatial risk assessment in case of multiple nuclear release scenarios

Oussama Raboun, Eric Chojnacki, Céline Duffa, David Rios Insua, Alexis Tsoukiàs

► **To cite this version:**

Oussama Raboun, Eric Chojnacki, Céline Duffa, David Rios Insua, Alexis Tsoukiàs. Spatial risk assessment in case of multiple nuclear release scenarios. *Socio-Economic Planning Sciences*, 2020, 70, pp.100721. 10.1016/j.seps.2019.06.006 . hal-02369416

HAL Id: hal-02369416

<https://hal.science/hal-02369416v1>

Submitted on 29 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Spatial risk assessment in case of multiple nuclear release scenarios

Oussama Raboun ^{a,b}, Eric Chojnacki ^b, Céline Duffa ^b, David Rios Insua ^c, Alexis Tsoukiàs ^a

^a Univ. Paris-Dauphine, PSL Research University, CNRS, UMR [7243], LAMSADE, 75016 PARIS, FRANCE

^b Institut de Radioprotection et de Sureté Nucléaire (IRSN), CADARACHE, FRANCE

^c Instituto de Ciencias Matemáticas (ICMAT) / CSIC, MADRID, SPAIN

Abstract

This paper presents an approach aimed at assessing multiple criteria spatial risk, where several methods are used for decision aiding purposes, with an application on marine nuclear releases. The case study simulates a post-accident analysis evaluating the impact upon the bay of Toulon of an accidental nuclear release. The problem is characterized by the presence of spatial features, multiple criteria describing the involved assets and uncertainties represented through multiple release scenarios and their corresponding probabilities.

Keywords: Multiple criteria decision support, Marine pollution, Environmental risk assessment, Risk rating.

1. Introduction

This work is part of a larger project aimed at developing theoretical and practical tools aiding to synthesize multiple criteria spatial risks in case of multiple nuclear release scenarios. A literature review with relevant papers on the integration of multiple criteria decision analysis tools in spatial decision problems until 2006 can be viewed at <http://publish.uwo.ca/~jmalczew/list.htm>. Despite the existing literature aiming to understand the processes governing the fate of radionuclides in the environment, [1], [2], [3], we note that the concentration of a given isotope is a necessary but not sufficient information for making informed decisions. Let us consider the example of two geographic zones: the first one is characterised by an average concentration level and very important economic and environmental assets while the second one is highly contaminated but does not present any economic or environmental relevance. Clearly, the involved stakeholders will be more sensitive to the impact in the first geographic plot.

Our case study deals with simulated releases from a nuclear submarine at the bay of Toulon, where one of the most important bases of the French Navy is located. In case of a nuclear accident, the incumbent prefect needs synthetic information to support decisions, such as banning certain economic activities, setting a new water management policy at each relevant zone or impeding the access to specific areas. The IRSN¹ is in charge of a project aimed at improving models predicting dispersion and assessing the impact of radionuclides in the environment, see www.irsn.fr/FR/Larecherche/Organisation/Programmes/Amorad/Pages/projet-Amorad.aspx#.Wl4GxiN7TOQ. In order to provide supplementary post-accident management tools allowing to evaluate environment and economic impacts, we have developed an approach in which data associated to assets involved in the bay are paired with maps displaying the concentration level of a given isotope generating criteria maps. Each map describes the impact of a release concentration for a given criterion. We then use a multiple criteria aggregation procedure generating impact maps taking into account all assets. The final step consists of aggregating uncertain information over release scenarios (release positions, sea conditions,...) through an outranking approach. Our case study serves as a template that can be extended to other release events and geographical areas.

¹Institut de Radioprotection et de Sureté Nucléaire is a French center of expertise and research in radioprotection and safety of nuclear installations. More information can be found at <http://www.irsn.fr>.

26 The originality of our work stands on the way we structured and modeled a practical issue, starting from
27 the raw question “How can we evaluate the impact of a nuclear accident, similar to that of Fukushima, in the
28 marine area?” The practical case was offered by the bay of Toulon, due to the presence of nuclear submarines
29 in its port, characterised by the presence of multiple assets and two levels of spatial decomposition. In this
30 paper, we propose the models used to assess the impact of a nuclear release on each asset involved, in case
31 we are interested in identifying the most impacted assets or areas with respect to each asset, as well as to
32 evaluate the global impact taking into account all considered assets.

33 The paper is organised as follows. Section 2 describes the case study including different decompositions
34 of the area of interest and the associated data. In Section 3, we introduce the main theoretical concepts used
35 in this work. We present in Section 4 the construction procedure of the criteria functions characterizing and
36 evaluating the Bay. In Section 5, we show the results of the multiple criteria aggregation and the aggregation
37 of release scenarios. We end up with a discussion. Several appendices provide additional details about this
38 work and its results.

39 2. Case study

40 The area of interest is the Bay of Toulon (in what follows, we will use the Bay to refer to it), where
41 a major basis of the French naval force is located, including nuclear submarines, besides being a densely
42 inhabited area with important economic activities. Thus, there is a possibility of major negative impacts in
43 case an accidental nuclear release takes place. Two features are identified in this study:

- 44 • Multiple impacts over different assets characterising the Bay.
- 45 • Uncertainties relative to accident parameters, to be modeled through scenarios.

46 In a radioactive release several isotopes may be present such as cesium-137, cesium-134, silver-110 or
47 iodine-131. In our case, we will focus on cesium-137 characterised by a half-life of 30.17 years. However, the
48 developed methodology does not depend on the considered radionuclide.

49 In our problem context, our objective is to set a decision aiding model based on consequences induced
50 by an accident. The available information includes:

- 51 • scientific facts and results: The dispersion model of radionuclides in the marine environment;
- 52 • geographic features: Each geographic zone has special characteristics such as the income associated
53 with tourism or fishing;
- 54 • norms: Including the maximum allowable levels of concentration for fishing or forbidding an activity.

55 2.1. Assets data

56 A decomposition of the Bay was carried out within the “Bay contract” by the “Syndicat Intercommunal
57 de l’Aire Toulonnaise” (SIAT, 1998 and 2002). This decomposition was based on the following criteria:

- 58 • A physical criterion, relying mainly on the geomorphology and local hydrodynamics of water bodies.
- 59 • A biological criterion, taking into account the presence of particular ecosystems.
- 60 • A socioeconomic criterion, based upon the presence of certain special activities such as ports and
61 military activities.

62 In what follows, we adopt the above mentioned division, with seven homogeneous zones illustrated in Figure
63 1:

- 64 1. The north of the small bay, characterised by maritime and military activities. It includes a military
65 port, freight, passenger transport, boaters and professional fishers.
- 66 2. The bay of Lazaret, characterised by aquaculture and tourism activities.

- 67 3. From the beaches Mourillon, Saint-Mandrier, until Cape Brown. The entrance to the small harbour
 68 is also characterised by military activity, a port, boaters and maritime transport. Its particularity lies
 69 in the fact that it represents a natural area of ecological faunistic and floristic interest, due to the
 70 presence of seagrass Posidonia.
- 71 4. From zone 3 to “Commune le Pradet”. This part is characterised by an important fishing activity,
 72 tourism activities and a high presence of seagrass Posidonia.
- 73 5. From Cap Sicié to Saint-Elme, characterised by several seaside activities. There is mainly swimming,
 74 boating, diving and professional fishing activities. This area is characterised by ecological richness,
 75 particularly a high presence of seagrass Posidonia. Moreover, there are three protected zones at “Anse
 76 des Sablettes”, the “Islands of the Two Brothers” and Cape Sicié.
- 77 6. From Marégau Point to Cape Cepet. This area is mainly dedicated to military activities. There is
 78 also tourism activities and seagrass Posidonia. This last is an important asset for sea life.
- 79 7. The rest of the bay with no land boundary is mainly characterised by professional fishing.




Figure 1: Decomposition of the bay into seven homogeneous zones.

80 In order to evaluate the consequences of accidents, we distinguished two types of attributes: economic and
 81 environmental.

82 2.1.1. Economic attributes

83 Two types of activities are present in the Bay:

- 84 • commercial activities linked with water quality: fishing, water sports, diving, professional fishing and
 85 aquaculture;
- 86 • Non-commercial activities such as swimming and leisure fishing.

87 Non-commercial activities seem to be not directly linked with economic assets. However, they have a strong
 88 influence over the touristic attractiveness of each zone which might induce an economic impact.

89 As far as the economic axis is concerned, we shall evaluate the impact of a released cesium-137 concen-
 90 tration based on three attributes:

- 91 • Professional fishing (F), based on an estimation of the annual economic impact of the fish caught. The
 92 data comes from the “Système d’Informations Halieutiques” (SIH-2007). Table 1 provides the annual
 93 turnover associated with professional fishing at each of the zones.

Zones	1	2	3	4	5	6	7
Annual turnover (k€)	300	300	965	965	934	1286	1000

Table 1: Annual turnover of professional fishing in 2007.

- Fish farming (FF), supports raising fish and shellfish. The area characterised by this activity is zone 2, representing an important economic asset for Toulon. The main characteristic is that the fish are more impacted by water quality as they cannot swim outside the breeding areas. The turnover generated by this activity in 2007 was 2129 (k€).
- Tourist attractiveness (T), refers mainly to swimming, diving and water sports. The economic value of swimming is assessed based on the income of restaurants located at sea shore and accommodations at each municipality. Thus, the value associated with the commune of Toulon will be assigned to zones 1 and 3; that of Seyne-Sur-Mer to zone 2; Pradet to zone 4; and, finally the commune of Saint-Mandrier to zone 6 (zone 7 has no coastline). Data associated with this attribute come from INSEE-Sirene 2007 for the catering sector, Chambre de Commerce et d’Industrie CCI-PACA 2007 for water sports and boaters and BVA-Ifremer 2007 for non professional fishing. Table 2 summarises the turnover associated with touristic activities in the Bay.

Zones	1	2	3	4	5	6	7
Annual turnover (k€)	34 839.5	29 593	20 828.5	13 591.5	23 113	24 483	1 131

Table 2: Annual turnover of Tourism in 2007.

2.1.2. Environmental axis

As far as the environmental axis is concerned, we shall focus our attention on the presence of seagrass Posidonia. This is one of the most important ecosystems in Mediterranean coastal zones, playing the same importance as forests in terrestrial areas: It is essential for the preservation of the balance of sea-life, [5], [11], as it:

1. Influences coastal water quality, through significant oxygen production and sediment trapping.
2. Is at the base of many trophic networks, for the production of plant and animal biomass.
3. Plays a fundamental role in the hydrodynamic protection of the coastline and beaches.
4. Fixes sediments and reduces the turbidity of the water, preventing their resuspension during storms.

Data on the mapping of seagrass Posidonia are rare, mostly very old, and its evolutionary dynamics are poorly known. Nevertheless, we have qualitative information on its presence at each geographic zone. Table 3 summarises its presence in the Bay.

Zones	1	2	3	4	5	6	7
Degree of the presence of seagrass Posidonia	Absent	Absent	Average	High	High	High	Absent

Table 3: Presence of seagrass Posidonia in the Bay (2002).

Cesium concentration might be included as a relevant environmental indicator representing water quality. However, we will not consider it independently, since we use it to assess criteria and we are interested in its impact on assets characterising the bay.

121 *2.2. Generating concentration data*

122 Many studies have been conducted to model the physical dispersion process of radioactive substances in
 123 the marine environment, e.g. [1], [2], [3], [9], [8], [13]. These have led to the development of simulation tools,
 124 such as STERNE², which we have used in our case study. The input parameters required by this tool are
 125 the type of sea currents, the release position and the quantity initially released.

126 *2.2.1. Sources of uncertainty*

127 Since accidental nuclear releases are related with the routes undertaken by the submarines, there will be
 128 two main sources of uncertainty in our case study³:

- 129 • The sea conditions (wind, currents, ...), at the time of the release, identified by a parameter β . In
 130 the case of Toulon, they are dominated by wind [13], and their probabilities can be estimated using a
 131 meteorologic database.
- 132 • The position $RP = (x_{RP}, y_{RP})$ where the release takes place, being, respectively, the latitude and
 133 longitude. We identify the main typical routes for submarines with some uncertainty around them.

134 We modeled uncertainty about the accident parameters through representative scenarios. We shall consider
 three sea conditions with their associated probabilities, as described in Table 4: and four initial release

Scenario	Prevailing wind	Probability
β_1	Mistral	q_1
β_2	East	q_2
β_3	Steady	q_3

Table 4: Discretisation of sea conditions.

135 positions with their associated probabilities, as specified in Table 5, displayed by red crosses in Figure 1.

Position in the map	Scenarios	Probability
Zone 5	$RP_1 = (43.053, 5.89)$	r_1
Zone 6	$RP_2 = (43.053, 5.96)$	r_2
Zone 4	$RP_3 = (43.079, 5.975)$	r_3
Zone 1	$RP_4 = (43.103, 5.918)$	r_4

Table 5: Discretisation of initial release positions.

136 The corresponding probabilities will be assessed in Section 5.2, where we shall synthesise the twelve scenarios.
 137

138 *2.2.2. Assessing cesium concentration*

139 The approach proposed here is driven by the contaminant concentration at each plot of the bay. This,
 140 in turn, will be driven by the amount initially released as well as the release position RP and sea conditions
 141 β . Based on a hydrodynamic model [8] sketched in Appendix A, we may estimate the concentration of the
 142 radioactive substance in water (respectively in a marine organism) at any point $z = (x, y)$ in the map, which
 143 we designate $c_w(z, RP, \beta)$ (respectively $c_o(z, RP, \beta)$).

144 STERNE offers the possibility of using tracking points to simulate the concentration evolution of a given
 145 isotope. We discretised the bay into several geographic units, represented by their tracking points in the
 146 center. The previous decomposition of the Bay in 7 homogeneous zones was too rough to be applied for

²Simulation du Transport et du transfert d'Eléments Radioactifs dans l'environNement marin, translated as Simulation of radionuclide transport and transfer in marine environments

³The amount initially released can be also considered as a source of uncertainty. However, in this work we shall fix it to $10^{15}Bq$, i.e. a very important release.

147 the estimation of the contaminant concentration, as it may lead to missing significant concentrations. We
 148 instead defined 97 geographical units adjusted to the map of the Bay. Each geographic unit is defined by
 149 two representative points: a tracking point at the bottom of the sea and another one at 1m depth. The
 150 reasons for choosing two depth levels are related with the nature of the chosen attributes. In what follows,
 151 we shall use “geographic zone” to refer to the first decomposition in 7 zones and “geographic units” to the
 152 second decomposition, in 97 units. Figure 2 displays 10 evolution curves of cesium concentration at 1m
 153 depth at the 10 most contaminated zones based on the maximum concentration attained.




Figure 2: 10 highest evolutions of cesium concentration in water over time.

154 We summarize the concentration evolution curves through their mean and maximum values. Figures 3
 155 and 4 display, respectively, the maximum and mean values corresponding to tracking points at 1m depth
 156 for the release scenario (*mistral*; RP_2).

	5,86_5,87	5,87_5,88	5,88_5,89	5,89_5,9	5,9_5,91	5,91_5,92	5,92_5,93	5,93_5,94	5,94_5,95	5,95_5,96	5,96_5,97	5,97_5,98	5,98_5,99	5,99_6	6_6,01	6,01_6,02
43,11_43,12			3,09E-10	1,09E-09	3,53E-08	4,87E-07	1,44E-04									
43,1_43,11			2,09E-09	1,09E-08	2,63E-06	7,85E-05	1,00E+00	3,24E+01	4,52E+01	6,54E+01	7,77E+01	5,84E+01	9,02E+01	1,55E+03	2,89E+03	
43,09_43,1						3,23E-03	4,62E+00	3,35E+01	6,63E+01	4,34E+01	5,29E+01	2,24E+02	1,02E+03	1,80E+03	7,72E+03	1,51E+04
43,08_43,09					2,66E-03	6,13E-02	1,14E+00	3,46E+01	2,50E+01	8,36E+01	6,50E+02	4,14E+03	8,56E+03	3,58E+04	7,31E+04	8,42E+04
43,07_43,08			1,45E+03	1,25E+03						2,39E+04	1,07E+04	2,95E+04	1,54E+05	2,55E+05	2,45E+05	1,70E+05
43,06_43,07		3,42E+03	2,00E+03	5,21E+03	2,82E+04	6,96E+04	8,04E+04	3,96E+05	9,64E+05	6,84E+04	1,10E+05	1,55E+05	3,07E+05	4,11E+05	4,05E+05	3,37E+05
43,05_43,06	2,75E+03	8,86E+02	1,71E+03	4,38E+03	1,20E+04	2,09E+04	1,30E+05	3,62E+05	3,30E+05	1,87E+05	1,73E+05	2,59E+05	1,09E+05	1,85E+05	3,31E+05	1,85E+05
43,04_43,05	2,01E+03	1,64E+03	4,06E+03	3,53E+03	2,06E+04	2,48E+04	4,51E+04	9,23E+04	2,56E+05	2,61E+05	7,11E+04	4,92E+04	5,74E+04	1,39E+05	1,08E+05	1,61E+07

Figure 3: Maximum concentration, 1m depth, at the 97 geographic units for (*mistral*; RP_2).

157 Empty cells in both figures correspond to land space. In all simulations we face a factor of 10 between
 158 the average and maximum values. We aggregate both values and move from a cardinal to an ordinal scale by
 159 assigning each zone to a corresponding concentration level. This can be achieved in several ways depending
 160 on the eventual compensation between both values. In our work we considered a geometric mean between
 161 them as their is a scale factor between both values. Figure 5 illustrates their aggregation considering the
 162 same level of importance for both evaluations at each zone.

163 A colour coding will reflect the contamination level at each geographic unit. We consider 5 levels from less
 164 to more contaminated. The cutting levels are fixed based on expert judgment. Level 1 is displayed in blue,
 165 2 in green, 3 in yellow, 4 in orange and 5 in red. We shall use this grading colour in the rest of the paper. As
 166 a first way to display the information, we could present the map $(z, c_w(z, RP, \beta))$, which provides, for each
 167 geographic unit z , the estimated contamination level, in an ordinal scale, given specific initial conditions.

	5,86_5,87	5,87_5,88	5,88_5,89	5,89_5,9	5,9_5,91	5,91_5,92	5,92_5,93	5,93_5,94	5,94_5,95	5,95_5,96	5,96_5,97	5,97_5,98	5,98_5,99	5,99_6	6_6,01	6,01_6,02
43,11_43,12			1,27E-12	3,75E-11	1,35E-09	2,08E-08	4,21E-06									
43,1_43,11			3,24E-11	4,24E-10	1,82E-07	6,21E-06	3,05E-02	3,00E+00	5,59E+00	8,53E+00	9,65E+00	7,28E+00	1,31E+01	1,31E+02	3,19E+02	
43,09_43,1						2,35E-04	1,67E-01	5,13E+00	1,18E+01	7,32E+00	9,17E+00	1,98E+01	7,05E+01	1,92E+02	4,07E+02	1,15E+03
43,08_43,09					1,47E-04	1,88E-03	5,47E-02	5,37E+00	4,46E+00	6,85E+00	5,45E+01	2,81E+02	8,48E+02	1,99E+03	3,42E+03	5,99E+03
43,07_43,08			2,10E+02	1,91E+02						1,22E+03	8,22E+02	2,41E+03	1,25E+04	2,77E+04	2,55E+04	1,90E+04
43,06_43,07		4,56E+02	2,79E+02	6,90E+02	3,80E+03	7,25E+03	4,50E+03	1,98E+04	4,37E+04	5,73E+03	9,49E+03	1,72E+04	3,85E+04	4,05E+04	3,79E+04	2,74E+04
43,05_43,06	4,32E+02	1,42E+02	2,52E+02	5,93E+02	1,76E+03	2,58E+03	6,92E+03	2,11E+04	1,56E+04	1,35E+04	1,64E+04	2,50E+04	1,08E+04	1,95E+04	2,70E+04	1,36E+04
43,04_43,05	3,76E+02	3,69E+02	5,58E+02	7,99E+02	1,85E+03	2,95E+03	4,19E+03	5,82E+03	1,15E+04	1,31E+04	4,72E+03	3,45E+03	3,16E+03	8,44E+03	9,61E+03	1,57E+05

Figure 4: Average concentration, 1m depth, at the 97 geographic units for (*mistral*; RP_2).

Csw	5,86_5,87	5,87_5,88	5,88_5,89	5,89_5,9	5,90_5,91	5,91_5,92	5,92_5,93	5,93_5,94	5,94_5,95	5,95_5,96	5,95_5,97	5,97_5,98	5,98_5,99	5,99_6	6_6,01	6,01_6,02
43,11_43,12			3,44E-10	1,13E-09	3,67E-08	5,08E-07	1,48E-04									
43,1_43,11			2,79E-09	1,13E-08	2,82E-06	8,47E-05	1,03E+00	3,54E+01	5,08E+01	7,39E+01	8,73E+01	6,57E+01	1,03E+02	1,68E+03	3,21E+03	
43,09_43,1						3,47E-03	4,78E+00	3,87E+01	7,82E+01	5,08E+01	6,21E+01	2,44E+02	1,09E+03	1,99E+03	8,13E+03	1,62E+04
43,08_43,09					2,81E-03	6,32E-02	1,20E+00	4,00E+01	2,94E+01	9,04E+01	7,05E+02	4,42E+03	9,40E+03	3,78E+04	7,65E+04	9,01E+04
43,07_43,08			1,66E+04	1,45E+04						2,51E+04	1,15E+04	3,19E+04	1,67E+05	2,83E+05	2,70E+05	1,89E+05
43,06_43,07		3,87E+03	2,28E+03	5,90E+03	3,20E+04	7,69E+04	8,49E+04	4,16E+05	1,01E+06	7,41E+04	1,20E+05	1,72E+05	3,45E+05	4,51E+05	4,43E+05	3,64E+05
43,05_43,06	3,18E+03	1,03E+03	1,97E+03	4,98E+03	1,38E+04	2,35E+04	1,37E+05	3,83E+05	3,45E+05	2,01E+05	1,90E+05	2,84E+05	1,20E+05	2,04E+05	3,58E+05	1,99E+05
43,04_43,05	2,39E+03	2,01E+03	4,62E+03	4,33E+03	2,24E+04	2,78E+04	4,93E+04	9,81E+04	2,67E+05	2,74E+05	7,59E+04	5,26E+04	6,05E+04	1,47E+05	1,18E+05	1,63E+07

Figure 5: Contamination level for 97 geographic units for (*mistral*; RP_2).

168 Figure 6 displays the contamination level induced by the release scenario (*mistral*; RP_2) using the previous
169 colour code.




Figure 6: Map displaying the contamination level corresponding to (*mistral*; RP_2).

170 3. Multiple criteria decision analysis

171 In Section 2.1, we described the Bay of Toulon as a rich area where several assets are involved and can be
172 impacted in case of a nuclear release. Our first objective is to define functions, which we shall call criteria,
173 allowing us to assess the impact on each asset at each geographic unit. Each function evaluates a geographic
174 unit from a single perspective. In consequence, we shall associate with each criterion a map evaluating the
175 impact on the corresponding asset ⁴. We shall consider the four criteria expressed on an ordinal scale, see
176 section 4, all of which need to be taken into account in an appropriate multiple criteria formulation.

⁴Considering each criterion function separately, we can either identify the most impacted geographic units or compute the expected impact.

177 The field of multiple criteria decision analysis (MCDA) offers a set of operational tools and methodologies
 178 to incorporate the decision maker’s preferences as well as any information allowing the decision analyst to
 179 evaluate a set of actions described by multiple attributes. In real-world cases, several problem statements
 180 can be considered referring to the way in which decision aiding is envisaged, see [18]:

- 181 • clustering (partition the set of alternatives into unordered not pre-defined equivalence classes; the
 182 clusters).
- 183 • assignment (partition the set of alternatives into unordered pre-defined equivalence classes).
- 184 • rating (partition the set of alternatives into ordered pre-defined equivalence classes).
- 185 • ranking (partition the set of alternatives into ordered not pre-defined equivalence classes).

186 Modeling a MCDA problem requires representing preferences either measuring their values, as in the case
 187 of multi-attribute value theory, or directly using binary relations, as in the case of social choice theory and
 188 outranking based methods, see [4].

189 In our case, we aim at assigning each geographic unit to the corresponding impact level. We consider five
 190 predefined and ordered impact categories C_1, \dots, C_5 , ranked from best to worst $C_h \gg C_{h+1} \forall h \in \{1, \dots, 4\}$
 191 where \gg refers to a complete order on the set of categories. Hence, the type of decision aid required here
 192 is a rating problem statement.

193 Two main methods corresponding to two different approaches deal with rating problems: UTADIS and
 194 ELECTRE-TRI. The UTADIS method was first presented in [6], being a variant of the well-known UTA
 195 method [12]. UTADIS consists of defining a marginal utility function over criteria, taking respectively the
 196 value 0 and 1 for the least and most preferred values of each criterion, and evaluating each action with an
 197 additive utility function. Such methods are suitable in multiple criteria problems where trade-offs among
 198 criteria are possible and meaningful. Alternatively, the ELECTRE-TRI method is an outranking based
 199 procedure first introduced in [19]. This method uses a majority rule, while respecting a minority using a
 200 veto rule, to compare the actions to the profiles characterizing categories; ELECTRE-TRI method is detailed
 201 in Appendix B. The MCDA procedure used in this work is based on ELECTRE-TRI, as trade-offs among
 202 the criteria were not interpretable.

203 4. Construction of criteria

204 The multiple criteria problem at hand adopts a rating formulation in which we consider the four criteria
 205 reflected in Table 6, with scales referring to the raw impact of a nuclear accident at each geographic unit.
 206 All criteria considered to evaluate the Bay are based on water quality through the concentration of cesium
 207 in water. Hence, the criteria will measure the impact of a given concentration on the assets involved at each
 208 geographic zone.

	Criteria	scale
1	Fishing	impact level
2	Fish Farming	impact level
3	Seagrass Posidonia	Impact level
4	Tourism	impact level

Table 6: Criteria and scales.

209 We start by presenting the typology of impact functions, allowing to associate with each concentration
 210 level an impact on an asset. For example, given a concentration level, the impact function will assess
 211 the proportion of tourists giving up visiting a geographic unit, the proportion of fishes not allowed to be
 212 commercialised or the impact on seagrass Posidonia. In the second part of this section, we construct the
 213 criteria functions, taking into account the impact function and the data associated with the assets. For
 214 instance, the tourism criterion is evaluated based on the income in a geographic zone, when there is no

215 accident, multiplied by the proportion of tourists giving up visiting the such unit given a concentration level
 216 (impact function).

217 4.1. Typology of impact functions

218 We aim now at evaluating the impact of a given level of contamination on each asset. The considered
 219 impact functions are based on two hypotheses:

- 220 • independent geographic units. As units are small, we do not consider mutual influences between
 221 neighbouring units. Thus, the impact on a geographic unit will only depend on its concentration level.
- 222 • The impact function does not depend on geographic units, as it depends on the characteristics of the
 223 assets.

224 Three types of impact functions will be considered. The choice of them will depend on the characteristics
 225 of attributes and the decision maker’s preferences:

- 226 • Heaviside function: We consider that a given asset is impacted from a certain level of concentration.
 227 This function is used in evaluating the impact on seagrass Posidonia.
- 228 • Linear function: no impact is considered before a first threshold is met while an important impact is
 229 assumed after the second one. Between both thresholds, the impact is linear. This type of function
 230 can be chosen when the population response is linearly proportional to pollution levels.
- 231 • Cumulative function: It is more suitable for modeling social phenomena for which the number of
 232 people influencing the evaluation of areas is important. We will use this function to assess the impact
 233 on tourism and fishing.



Figure 7: Heaviside impact function.




Figure 8: Linear impact function.




Figure 9: Cumulative impact function.

234 The cumulative impact function requires calibration reflecting the impact of different levels of concen-
 235 tration on a given asset. For example, qualitatively, the higher the concentration, the less tourists will visit
 236 the corresponding polluted area. This function can be derived through a weighted sum of linear functions,
 237 of type 2, representing each the impact assessment by a pool of experts, assessing a “tolerance threshold”
 238 and a “reaction threshold”. Alternatively, we can calibrate the median for each contamination level as we
 239 do here. This approach is inspired by the probability equivalent method for assessing utilities [10]. Let
 240 us call the cumulative impact function $prop_i(c_k)$, where i refers to a geographic unit i and c_k is the level
 241 of contamination in the marine organism $k = o$ or in seawater $k = w$. Our objective is to find for a few
 242 concentrations c_{k_1}, \dots, c_{k_5} ⁵, the corresponding $prop_i(c_{k_1}), \dots, prop_i(c_{k_5})$, through expert judgment, and then
 243 adjust a curve. Note that $prop_i(c_k)$ will essentially be uncertain and we shall focus on assessing its median
 244 using lottery comparison.

245 In what follows we apply this approach to the attribute Tourism and thus $k = w$. For this we compare
 246 two lotteries:

- 247 • Lottery A , represents throwing a fair coin in which the expert wins 100 € if he obtains Head and 0 €
 248 is he obtains "Tail". This serves as reference.

⁵ 5 represents the number of contamination levels introduced in Section 2.2.2

- 249 • Lottery B , represents the calibrated event and gives the expert 100 € if $prop \geq q$ and 0 € otherwise,
 250 where $prop = prop_i(c_w)$ is the proportion of tourists giving up visiting a geographic unit in case c_w is
 251 high enough and q is the calibrating value.

252 We ask the expert whether he prefers A to B ($A \succeq B$). In such case, we have: $100 \times \frac{1}{2} + 0 \times \frac{1}{2} \geq$
 253 $100 \times P(prop \geq q) + 0 \times (1 - P(prop \geq q))$; we need to adjust q to approximate the median. For this, we can
 254 design an iterative procedure to converge to it, bounding it from below and above. Initially, the bounding
 255 interval is $[0, 1]$ and we iteratively split it depending on the responses of the expert. Specifically, we use
 256 $q = \frac{Y+X}{2}$, for $[X, Y]$ and adjust X and Y according to expert responses, with $X = 0, Y = 1$ initially. For
 257 a large number of iterations this will converge to the median. Figure 10 displays the calibration for a few
 258 concentration levels using the above procedure. The same approach remains valid for the fishing attribute,
 for which $k = o$.




Figure 10: Calibration of proportion of tourists giving up visiting a cell.

259

260 4.2. Tourism

261 We construct first the criterion function for tourism, referring to the level of economic loss related with
 262 the tourism sector. This last is assessed as $prop_i(c_w(z_i, s), T)Inc_i(T)$ where $prop_i(c_w(z_i, s), T)$ represents
 263 the proportion of tourists refraining from visiting the geographic unit i under the incumbent release scenario
 264 and $Inc_i(T)$ represents the income associated with the geographic unit i . The function $prop_i(c_w(z_i, s), T)$
 265 has been assessed in section 4.1, Figure 10.

266 In order to evaluate the economic importance of each geographic unit, an issue with the spatial decom-
 267 position arises:

- 268 • Data associated with tourism revenues are available just for the seven geographic zones. We parti-
 269 tioned the annual turnover proportionally between all geographic units constituting each of the seven
 270 geographic zones.
- 271 • Some geographic units are shared between several homogeneous zones. The solution adopted is to
 272 evaluate the geographic units by considering the turnover proportionally to the surface occupied by
 273 geographic zones at the geographic unit. This entails the use of the same decomposition as for cesium
 274 concentration simulations.

275 Thus, the estimated annual turnover at each geographic unit is

$$Inc_i(T) = \sum_{j \in Z, st: Z \cap \{i\} \neq \emptyset} \frac{S_{ij}}{\sum_{i \in U \cup Z} S_{ij}} Tur_j(T),$$

276 where U and Z represent, respectively, the set of geographic units (decomposition of the Bay adopted
 277 to forecast cesium concentration) and the set of geographic zones (decomposition made to describe the
 278 attributes); T refers to the asset Tourism; S_{ij} the maritime surface (land excluded) belonging both to the
 279 geographic unit i of U and the zone j of Z ; $Tur_j(T)$ the turnover associated with geographic zone j .

We denote by $g_T(i, s)$, the function of the tourism criterion rating the geographical unit i , given a scenario s . Such function would be ⁶:

$$g_T(i, s) = \begin{cases} 1, & \text{if } prop_i(c_w(z_i, s), T)Inc_i(T) \times 97 < 10^4 \\ 2, & \text{if } 10^4 \leq prop_i(c_w(z_i, s), T)Inc_i(T) \times 97 < 10^6 \\ 3, & \text{if } 10^6 \leq prop_i(c_w(z_i, s), T)Inc_i(T) \times 97 < 10^7 \\ 4, & \text{if } 10^7 \leq prop_i(c_w(z_i, s), T)Inc_i(T) \times 97 < 10^8 \\ 5, & \text{if } 10^8 \leq prop_i(c_w(z_i, s), T)Inc_i(T) \times 97 \end{cases}$$

280 where $10^4, 10^6, 10^7, 10^8$ represent the economic losses delimiting each impact category. The cutting thresh-
 281 olds used in the different criteria are assessed based on expert judgment. Figure 11 shows the assessment of
 282 the tourism criterion for the mistral-type marine currents and release point RP_2 .

Tourism	5,86_5,87	5,87_5,88	5,88_5,89	5,89_5,9	5,90_5,91	5,91_5,92	5,92_5,93	5,93_5,94	5,94_5,95	5,95_5,96	5,95_5,97	5,97_5,98	5,98_5,99	5,99_6	6_6,01	6,01_6,02
43,11_43,12			1	1	1	1	1									
43,1_43,11			1	1	1	1	1		3	3	3	3	4	4	4	
43,09_43,1						1	1	3	3	3	3	4	4	4	4	5
43,08_43,09					1	1	1	3	3	3	3	4	4	4	5	5
43,07_43,08			4	4						5	5	5	5	5	5	5
43,06_43,07			4	4	4	5	5	5	5	5	5	5	5	5	5	5
43,05_43,06	4	4	4	4	4	5	5	5	5	5	5	5	5	5	5	5
43,04_43,05	4	4	4	4	4	5	5	5	5	5	5	5	5	5	5	5

Figure 11: evaluation of the tourism criterion under the scenario ($Mistral, RP_2$)

283 We can display the above results through maps ⁷. As an example, the map in Table 7 represents the impact
 284 on the tourism criterion under scenario ($Mistral, RP_2$). Results corresponding to the other scenarios can
 285 be found in the supplementary material.




Table 7: Tourism criterion maps for the twelve scenarios

286 Some relevant information can be assessed in this way. For example we can identify areas which are most
 287 at risk from the perspective of tourism. (e.g. the red ones)

⁶97 in the criterion function refers to the number of geographic units

⁷We used the same colour coding as in Figure 6

The economic loss in the bay associated with scenario $s = (\beta_h, RP_k)$ can be obtained through spatial aggregation, without considering interactions between neighbouring geographic units based on:

$$\sum_i prop_i(c_w(z_i, \beta_h, RP_k), T) Inc_i(T).$$

288 The expected economic loss in the whole area, through aggregating uncertainties over initial conditions,
289 $s = (\beta_h, RP_k)$, would be: e

$$\varphi_{TA} = \sum_i Inc_i(T) \sum_{h=1}^3 \sum_{k=1}^4 prop_i(c_w(z_i, (\beta_h, RP_k), T) q_h r_k,$$

290 which we denote $\varphi_{TA} = \sum_i Inc_i(T) prop_i(c_w, T)$.

The expected income in tourism sector when there is no accident would be:

$$\varphi_T = \sum_i Inc_i(T),$$

Then, the expected income on the whole area after an accidental release would be:

$$\varphi_T - \varphi_{TA}.$$

We could also use relative losses. For example, for the income from tourism, it would be:

$$\frac{\varphi_{TA}}{\varphi_T}.$$

291 All these indices, derived from the process of the construction of the tourism criterion may help the decision
292 maker assessing the impact of an eventual accident over the tourism sector.

293 4.3. Fishing

294 We assess now the fishing criterion function, focusing on the economic loss on the fishing sector at
295 each geographic unit. Such loss is evaluated by coupling the proportion of fish not authorised for sale
296 and the economic income before the accident in a geographic unit. Thus, the economic loss would be
297 $prop_i(c_o(z_i, s), P_e) Inc_i(P_e)$, where $prop_i(c_o(z_i, s), P_e)$ represents the impact function associated with the
298 fishing sector, $c_o(z_i, s)$ denotes the contamination level in fish and $Inc_i(P_e)$ represents the income from the
299 fishing sector at the geographic unit i . It should be mentioned that, for this criterion, we will consider
300 tracking points both at 1m depth and at the bottom of the sea. This is justified by the presence of fish at
301 all sea levels in this region.

302 The impact function $prop_i(c_o(z_i, s), P_e)$, is characterised by two thresholds:

- 303 • The first one reflects the level at which responsible authorities begin to control the cesium concentration
304 in fish before selling.
- 305 • The second one represents the level at which authorities prohibit consumption of fish caught at a
306 given geographic unit. We shall consider the second threshold to be $500Bq/kg$ equal to the maximum
307 allowable level of contamination for authorising fish consumption.

308 Between both thresholds, the impact is considered non-linear. The calibration process in section 4.1 is
309 applicable. The only modification would be to use $prop = prop_i(c_o)$ in lottery B , reflecting the proportion
310 of fish not allowed for sale given the level of cesium concentration c_o in fish.

In order to evaluate the economic importance of a geographic unit, $Inc_i(P_e)$, we use the same solution
for the two spatial decompositions as for tourism. Thus, the annual turnover at each geographic unit is
defined as:

$$Inc_i(P_e) = \sum_{j \in Z, st: Z \cap \{i\} \neq \emptyset} \frac{S_{ij}}{\sum_{i \in U \cup Z} S_{ij}} Tur_j(P_e),$$

311 where $Tur_j(P_e)$ refers to the turnover of fishing associated with geographic zone j .

We denote by $g_{P_e}(i, s)$, the fishing criterion rating the geographic unit i under scenario s . Such function would be

$$g_{P_e}(i, s) = \begin{cases} 1, & \text{if } prop_i(c_o(z_i, s), P_e)Inc_i(P_e) \times 97 < 10^3 \\ 2, & \text{if } 10^3 \leq prop_i(c_o(z_i, s), P_e)Inc_i(P_e) \times 97 < 10^5 \\ 3, & \text{if } 10^5 \leq prop_i(c_o(z_i, s), P_e)Inc_i(P_e) \times 97 < 10^6 \\ 4, & \text{if } 10^6 \leq prop_i(c_o(z_i, s), P_e)Inc_i(P_e) \times 97 < 5.10^6 \\ 5, & \text{if } 5.10^6 \leq prop_i(c_o(z_i, s), P_e)Inc_i(P_e) \times 97 \end{cases}$$

312 Table 8 shows the assessment of the fishing criterion maps for the twelve scenarios. As for tourism, we can
 313 derive the economic loss in the bay, the expected loss, the relative loss and the expected income in relation
 314 with the fishing asset.




Table 8: Fishing criterion maps for the twelve scenarios

315 4.4. Fish Farming

316 For this criterion, as for fishing, we use the concentration level of cesium in organisms (fish and shellfish
 317 farming). Because of the special characteristics of the fish farming activity, the impact on this sector will
 318 not be assessed at a geographic unit but at the whole geographic zone 2:

- 319 • Unlike the fishing indicator, where fish can swim through many geographic units, fish in aqua-farms
 320 cannot leave geographic zone 2 and, thus, they are just impacted by the water quality of this zone.
- 321 • The economic relevance of all geographic units in zone 2 is the same.

322 To assess this criterion, we consider $\overline{c_o(s)} = \max_i(c_o(z_i, s))$, where z_i is a geographic unit in zone 2. The
 323 economic income associated with the fish farming sector will not be considered on the criterion evaluation,
 324 as it is the same in all geographic units of zone 2. However, this last will represent a relevant information
 325 to assess the criterion's importance during the multicriteria aggregation procedure.

326 We denote by $g_{F_f}(s)$, the fishing criterion evaluating the geographic zone 2 under scenario s . Such
 327 function can be interpreted as a rate representing the impact on the fish farming sector.

$$g_{F_f}(s) = \begin{cases} 1, & \text{if } \overline{c_o(s)} < 100 \\ 2, & \text{if } 100 \leq \overline{c_o(s)} < 200 \\ 3, & \text{if } 200 \leq \overline{c_o(s)} < 300 \\ 4, & \text{if } 300 \leq \overline{c_o(s)} < 400 \\ 5, & \text{if } 500 \leq \overline{c_o(s)} \end{cases}$$

328 $500Bq/kg$ and $100Bq/kg$ are respectively the maximum allowable level to consume fishes from Fukushima
 329 before and after the accident. Table 9 shows the assessment of the fish-farming criterion maps for the twelve
 330 scenarios.




Table 9: Fish-Farming criterion maps for the twelve scenarios

331 4.5. Seagrass "*Posidonia Oceanica*"

We assess now a criterion function in relation with the impact of a radioactive release scenario on seagrass for each geographic unit. Unlike the previous ones, the seagrass *Posidonia* criterion rates the impact of a concentration level at a geographic unit level. Such impact represents a coupling between contamination levels, through the corresponding impact function, and scores associated with the presence of seagrass *Posidonia* at each geographic unit. To assess the corresponding impact function, we consider a Heaviside function, Figure 7, defined through

$$Imp_i(c_w) = \begin{cases} 0, & \text{if } c_w \in \{1, 2\} \\ 1, & \text{if } c_w \in \{3, 4, 5\} \end{cases}$$

332 where c_w refers to the level of cesium concentration in seawater. Degrees of the presence of seagrass *Posidonia*
 333 are described on an ordinal scale in Table 3. We denote by $LHp(j)$ the score associated with the degree of
 334 presence of seagrass *Posidonia* in geographic zone j , with the following scores:

- 335 • 0: Absence;
- 336 • 1: Weak presence;
- 337 • 2: Average presence;
- 338 • 3: Strong presence.

339 At this level, we need to solve the problem of both spatial decompositions in our problem. This asset is
 340 characterised by the lack of information about the exact distribution of seagrass *Posidonia* in the geographic
 341 units. Thus, we shall assume that its presence is uniform in all of them. This generates the following cases:

- 342 • For each geographic unit entirely included in a geographic zone, we consider that it has the same
 343 degree of presence of seagrass *Posidonia* as for the geographic zone;
- 344 • For geographic units shared between several geographic zones, we consider a weighted sum of the
 345 different degrees of presence of the seagrass in geographic zones. Weights in this work represent the
 346 relative surface at each geographic unit belonging to a given geographic zone.

The function describing these two cases would be

$$Sc(i) = \sum_{j \in Z, st: Z \cap \{i\} \neq \emptyset} \frac{S_{ij}}{S_i} LH_p(j)$$

where $Sc(i)$ represents the score associated with the presence of Posidonia at zone i , S_{ij} the surface (land excluded) of the geographic zone j and geographic unit i and S_i the surface of geographic unit i . We denote by $RSc(i)$, the rounded value of $Sc(i)$. We denote by $g_{Sp}(i, s)$, the seagrass Posidonia criterion rating the geographic unit i , under scenario s

$$g_{Sp}(i, (\beta_k, z_{RP_j})) = Imp_i(z_i, c_w((\beta_k, RP_j), Sp)) RSc(i) + 1.$$

347 Table 10 shows the assessment of the seagrass Posidonia criterion for the twelve scenarios.




Table 10: Seagrass-Posidonia criterion maps for the twelve scenarios

348 Again, we could compute various aggregated indices.

349 5. Multiple impacts

350 We finally consider the case with multiple criteria. Recall that for each scenario (β_i, RP_j) , which occurs
 351 with probability $p_{ij} = q_i \times r_j$, we obtain four criterion maps:

- 352 • Fishing: $g_{P_e}(\beta_i, RP_j)$
- 353 • Fish farming: $g_{F_f}(\beta_i, RP_j)$
- 354 • Seagrass Posidonia: $g_{S_p}(\beta_i, RP_j)$
- 355 • Tourism: $g_T(\beta_i, RP_j)$

356 The aim of this section is to aggregate effects due to:

- 357 • multiple criteria.
- 358 • uncertainty.

359 In the first part of this section, we shall solve the multiple criteria problem. In the second part, we
 360 aggregate uncertainties by considering scenarios as criteria evaluating the geographic units in the aggregated
 361 maps with respect to their corresponding importance (probabilities).

362 5.1. ELECTRE-TRI for multiple criteria aggregation

The problem at hand is a rating one. To solve it we use the ELECTRE TRI method. The first step consists of rating each geographic unit X for each scenario (β_i, RP_j) :

$$X(\beta_i, RP_j) = (g_{P_e}(\beta_i, RP_j), g_{F_f}(\beta_i, RP_j), g_{S_p}(\beta_i, RP_j), g_T(\beta_i, RP_j)).$$

363 We consider the following notation:

- 364 • the set of criteria \mathcal{F} , with criteria \mathcal{F}_j characterised by an importance (weight) w_j .
- 365 • the set \mathcal{C} of predefined impact categories. Each category \mathcal{C}_k is characterised by a lower bound, called
- 366 limiting profile, which we denote $r^k = (r_j^k)_{k \in \mathcal{F}}$.

367 The idea is, then, to compare the performance of each geographic unit with the limiting profiles to assign it
 368 to the corresponding category. Figure 12 illustrates the issue where the axes represent the criteria and we
 369 aim to assign x , a geographic unit, to one of the five predefined categories by comparing it with the limiting
 370 profiles.




Figure 12: Illustration of the multiple criteria problem

371 5.1.1. Assessing criteria weights

372 The literature reports several methods for assessing ELECTRE-TRI parameters from assignment exam-
 373 ples, [7], [14], [15], [17], [20]. We use a simplified version of the optimisation model in [15], by assuming that
 374 we are able to assess, with the aid of the decision maker, the limiting profiles. We denote by $A = \{A_1, \dots, A_5\}$
 375 the learning set where the assignments are previously known, with $A_k = \{a_{ki}; a_{ki} \in \mathcal{C}_k\}$. The learning set
 376 consists on assignment examples: examples of performance vectors, for which the rating is previously known.

Under the previous assumption, and based on the majority rule, an alternative $a_k \in A_k$ from the learning set is assigned to category \mathcal{C}_k if there is a weighted-majority of criteria in favour of “ a_k is at least as good as the limiting profile r^k ” and there is no weighted-majority in favour of “ a_k is at least as good as the limiting profile r^{k+1} ”. This can be written as

$$\sum_{j \in \mathcal{F}, g_j(a_{ki}) \geq r_j^k} w_j \geq c,$$

and

$$\sum_{j \in \mathcal{F}, g_j(a_{ki}) \geq r_j^{k+1}} w_j < c,$$

where c is the concordance threshold. Such inequalities are equivalent to the following equalities, introducing the slack variables x_{ki} and y_{ki} :

$$\sum_{j \in \mathcal{F}, g_j(a_{ki}) \geq r_j^k} w_j - x_{ki} = c$$

and

$$\sum_{j \in \mathcal{F}, g_j(a_{ki}) \geq r_j^{k+1}} w_j + y_k = c.$$

377 If the slack variables x_{ki} and y_{ki} are positive, the assignment made by the decision maker corresponds to
 378 the assignment done through the pessimistic procedure of ELECTRE TRI: the lower the minimum of these
 379 values, the less adapted is the model. In case one of these slack variables is negative, the concordance
 380 principle is not sufficient to justify the assignment and we need to assess the veto threshold. Thus, we
 381 need to maximise the minimum of both slack variables to take into account the worst assignment from the
 382 decision maker through

$$\max \min_{\substack{a_{ki} \in A_k \\ A_k \in A}} (x_{ki}, y_{ki})$$

383 and we also need to maximise the ability of the model to assign alternatives correctly through

$$\max \sum_{\substack{a_{ki} \in A_k \\ A_k \in A}} (x_{ki} + y_{ki}).$$

384 We then consider the following decision variables:

- 385 • Weight vector. $w_j, \forall j \in \mathcal{F}$
- 386 • Concordance threshold c
- 387 • Slack variables $x_{ki}, y_{ki}, \forall a_{ki} \in A_k, \forall k$

and the following objective function to be maximised

$$\text{maximise} \left(\min_{\substack{a_{ki} \in A_k \\ A_k \in A}} (x_{ki}, y_{ki}) + \epsilon \sum_{\substack{a_{ki} \in A_k \\ A_k \in A}} (x_{ki} + y_{ki}) \right) \quad (1)$$

Problem (1) is equivalent to

$$\begin{aligned} & \text{maximise} \quad \delta + \epsilon \sum_{\substack{a_{ki} \in A_k \\ A_k \in A}} (x_{ki} + y_{ki}) \\ & \text{s.t} \quad \delta \leq x_{ki}, \forall a_{ki} \in A_k, \forall A_k \in A \\ & \quad \delta \leq y_{ki}, \forall a_{ki} \in A_k, \forall A_k \in A. \end{aligned}$$

388 In order to assess criteria weights, we add to the previous model the following constraints:

- 389 • Two constraints related with the slack variables $\forall a_{ki} \in A_k, \forall A_k \in A$, $\sum_{\substack{j \in \mathcal{F} \\ g_j(a_{ki}) \geq r_j^k}} w_j - x_{ki} = c$ and

$$390 \quad \sum_{\substack{j \in \mathcal{F} \\ g_j(a_{ki}) \geq r_j^{k+1}}} w_j + y_{ki} = c.$$

- 391 • The majority constraint related to the concordance principle $c > 0.5$.
- 392 • We assume that all criteria are relevant, $w_j < c, \forall j \in \mathcal{F}$.
- 393 • The strict positivity and normalisation of weights: we respectively have $\forall j \in \mathcal{F}, w_j > 0$ and $\sum_j w_j = 1$.

We finally use the following model:

$$\begin{aligned}
& \text{maximise} && \delta + \epsilon \sum_{\substack{a_{ki} \in A_k \\ A_k \in A}} (x_{ki} + y_{ki}) \\
& \text{s.t} && \delta \leq x_{ki}, \forall a_{ki} \in A_k, \forall A_k \in A, \\
& && \delta \leq y_{ki}, \forall a_{ki} \in A_k, \forall A_k \in A, \\
& && \sum_{\substack{j \in \mathcal{F} \\ g_j(a_{ki}) \geq r_j^k}} w_j - x_{ki} = c, \forall a_{ki} \in A_k, \forall A_k \in A, \\
& && \sum_{\substack{j \in \mathcal{F} \\ g_j(a_{ki}) \geq r_j^{k+1}}} w_j + y_{ki} = c, \forall a_{ki} \in A_k, \forall A_k \in A, \\
& && \sum_j w_j = 1, \\
& && w_j < c, \forall j \in \mathcal{F}, \\
& && w_j > 0, \forall j \in \mathcal{F}, \\
& && 0.5 < c < 1.
\end{aligned} \tag{2}$$

Example: We consider the following learning sets:

$$A_2 = \{(1, 3, 2, 1), (1, 2, 3, 1), (3, 1, 1, 1)\}$$

$$A_3 = \{(2, 4, 3, 3), (4, 2, 3, 2), (3, 3, 2, 4)\}$$

$$A_4 = \{(2, 4, 4, 5), (4, 4, 4, 3), (5, 5, 3, 3)\}$$

$$A_5 = \{(5, 4, 4, 5), (5, 4, 5, 3), (3, 3, 5, 5)\}$$

394 The limiting profile of a category \mathcal{C}_k is the vector (k, k, k, k) . Therefore, we will not consider a learning
395 set associated with category \mathcal{C}_1 (no impact), since it does not provide us with any relevant information. A
396 profile in A_1 will always dominate $(1, 1, 1, 1)$; based on the majority principle it will always be outranked by
397 $(2, 2, 2, 2)$, otherwise it will not be assigned to \mathcal{C}_1 . Hence, both x_1 and y_1 are positive.

398 The tourism and fishing sectors are more sensitive than that of fish farming since they are present in
399 most of the geographic units. Hence, we consider two additional constraints, $w_1 \geq w_2$ and $w_4 \geq w_2$.

400 The solution of the model (2) is:

- 401 • weights: $w_1 = 0.33$; $w_2 = 0.1$; $w_3 = 0.23$; $w_4 = 0.34$;
- 402 • concordance threshold: $c = 0.54$;
- 403 • the slack variables:
 - 404 – slacks associated with $A_2 = \{(1, 3, 2, 1), (1, 2, 3, 1), (3, 1, 1, 1)\}$ are $(x_{21} = -0.17, y_{21} = 0.41)$,
 - 405 $(x_{22} = -0.07, y_{22} = 0.17)$, $(x_{23} = -0.17, y_{23} = 0.17)$;
 - 406 – slacks associated with $A_3 = \{(2, 4, 3, 3), (4, 2, 3, 2), (3, 3, 2, 4)\}$ are $(x_{31} = 0.15, y_{31} = 0.41)$, $(x_{32} =$
 - 407 $0.35, y_{32} = 0.17)$, $(x_{33} = 0.25, y_{33} = 0.17)$;
 - 408 – slacks associated with $A_4 = \{(2, 4, 4, 5), (4, 4, 4, 3), (5, 5, 3, 3)\}$ are $(x_{41} = 0, y_{41} = 0.17)$, $(x_{42} =$
 - 409 $0.15, y_{42} = 0.51)$, $(x_{43} = -0.7, y_{43} = 0.07)$;
 - 410 – slacks associated with $A_5 = \{(5, 4, 4, 5), (5, 4, 5, 3), (3, 3, 5, 5)\}$ are $x_{51} = 0.15$, $x_{52} = 0.05$, $x_{53} =$
 - 411 0.05 .

412 Negative slack variables can be justified due to the non consideration of the veto threshold in our linear
 413 model. For example, vector $(1, 3, 2, 1)$, using the majority principle, should be assigned to category \mathcal{C}_1 , since
 414 $w_1 + w_4 > c$. However, $(1, 3, 2, 1)$ is assigned to 2 because of its performance (a rate 3) under the fish farming
 415 criterion, and thus we cannot consider that there is no considerable impact. A similar remark is valid for
 416 $(1, 2, 3, 1)$, $(3, 1, 1, 1)$ and $(5, 5, 3, 3)$. Based on an observation over assignment examples with negative slack
 417 variables, a threshold value equal to 2 is the minimum value justifying the assignments.

418 *5.1.2. Assignment zones to the predefined categories*

419 We show now the results of the multiple criteria aggregation procedure using ELECTRE-TRI. The
 420 parameters we use are derived from the example in Section 5.1.1:

- 421 • criteria weights: $w_1 = 0.33$; $w_2 = 0.1$; $w_3 = 0.23$; $w_4 = 0.34$;
- 422 • concordance threshold: $c = 0.54$;
- 423 • veto threshold $v = 2$;

424 In what follows, we display the criteria-maps associated with the scenarios $(Mistral, RP_3)$, and the corre-
 425 sponding aggregated map.

426 **Scenario $(Mistral, RP_3)$**




Figure 13: Criteria maps for the $(mistral, RP_3)$

427 These maps display the criteria for fishing, fish farming, seagrass Posidonia and tourism, respectively. They
 428 are assessed based on the level of cesium concentration, from 1 to 5, where level 1 refers to low concentration
 429 and level 5 to a high concentration, and the vulnerability of each geographic unit from a given asset point
 430 of view. For example, zones 1 and 2 are not very impacted because of a low level of concentration; however,
 431 zone 3 is characterised by a level 5 of cesium concentration, crossed with important tourist and fishing
 432 activities, an average presence of seagrass Posidonia and no activity of fish farming. Thus, the outcome of
 433 the multiple criteria aggregation mostly associate a rate 1 to geographic units in zones 1 and 2 and a rate
 434 5 in zone 3 (recall that $w_1 + w_4 = 0.67 > 0.54$ and there is no discordance). The result of the aggregation
 435 is displayed in Figure 14.




Figure 14: The aggregated map for $(mistral, RP_3)$ scenario

436 *5.2. Uncertainty aggregation*

437 The aim of this section is to model uncertainties represented through different accident scenarios. We
438 need to establish a global rate for geographic units. We can aggregate the impact induced by different sce-
439 narios either before aggregating criteria or after the aggregation. The most common technique synthesising
440 uncertainties is to compute expected values. We used the expected impact in section 4, before the multiple
441 criteria aggregation procedure, in order to evaluate the sensitivity of each geographic unit from a single
442 criterion point of view. In this section, we deal with the case in which we want to synthesize uncertainties
443 related with the accident scenarios after the multiple criteria aggregation, section 5.1.

444 Computing the expected impact at each geographic unit allows for compensation between rates with
445 respect to the probabilities over scenarios. In our context, such compensation is not desirable since the
446 performance of geographic units under each scenario is modeled through rates. The aggregation procedure
447 proposed in this section is based on the concordance and discordance principles, reflected in ELECTRE TRI,
448 and can be solved as any multiple criteria rating problem, by considering scenarios as criteria, probabilities
449 as weights and geographic units as alternatives to be evaluated.

450 *5.2.1. Probabilities and ELECTRE TRI parameters*

451 In this section, we assess uncertainties over the initial conditions and the ELECTRE TRI parameters
452 to rate the geographic units. In section 2.2.1, we defined three sea conditions, corresponding to different
453 types of wind. In what follows, we associate to the types of wind the following probabilities [8]: For mistral
454 $q_1 = 0.4$, for east wind $q_2 = 0.4$ and for steady wind $q_3 = 0.2$.

455 To assess probabilities over the four release positions, we assume that the closer we are to the naval
456 base, the greater the probability of a release. Such hypothesis can be transcribed through the following
457 inequalities $r_i > r_j$ where $i > j$, with $r_i > 0$, $\sum_{i=1}^4 r_i = 1$. One possible assessment would be $r_1 = 0.5$, $r_2 =$
458 0.25 , $r_3 = 0.15$ and $r_4 = 0.1$, which we use in our initial analysis.

459 In what follows, we shall assume such values. A sensitivity analysis with respect to them, based on
460 intervals, would be necessary, but we shall not include it in this paper. Observe now that this “multiple
461 criteria decision making problem” is characterised by:

- 462 • The criteria evaluating the geographic units: the release scenarios;
- 463 • The weights of criteria: the probabilities $p_{ij} = q_i \times r_j$;
- 464 • Under each scenario, impacts on geographic units are rated from 1 to 5. We shall consider the same
465 scale for the aggregated rate;
- 466 • The veto threshold: $v = 2$.

467 *5.2.2. Results of the aggregation*

468 We represent now the results of the aggregation over the 12 scenarios using ELECTRE TRI and the
469 parameters in section 5.2.1.

470 From Table 11, we notice that for the release position RP_1 , the geographic zone 5 is highly impacted. This
471 is justified by the simulated release position in Zone 5, and the high importance of economic environmental
472 assets in this area. This remark is still valid for Zones 3 and 6 for RP_2 and 3 and 4 for RP_3 . depending
473 on the direction of wind, other zones might be highly impacted. For instance, considering the scenarios
474 characterised by a Steady wind, the impacted zones are those close to the release position. We also note
475 from Table 11 that the most impacting scenarios are those corresponding to East wind. The main reason
476 is the high dispersion of radionuclides in the majority of geographic zones due to the sea currents, which
477 impact many assets.

478 Zone 1 where the simulated RP_4 took place is highly impacted, rated 4, but less impacted than other
479 neighbouring zones, even if the contamination level is the highest. This is due to several reasons, such as
480 the non presence of seagrass Posidonia and fish farming activity, representing a total weight $w_2 + w_3 = 0.33$,
481 the low income from fishing activity compared to the other geographic zones




Table 11: The aggregated maps for the twelve scenarios




Figure 15: The aggregated map for the 12 release scenarios

482 Aggregating the release scenarios, we note from Figure 15 that geographic zone 1 seems the less impacted.
 483 The reason of such level of impact in zone 1 is justified by the low presence or absence of the majority of
 484 assets and the low level of concentration at several release scenarios. The other geographic zones are either
 485 rated 4 or 5, since scenarios corresponding to East wind, occurring with a probability of 0.4, impact highly
 486 the majority of the Bay and scenarios RP_2 , RP_3 and RP_4 in the case of Mistral type of wind, occurring
 487 with a total probability of 0.2, impact highly zones 3, 4, 5, 6 and 7.

488 6. Discussion

489 For the typical reasons related to a real world application we were induced to make a number of sim-
 490 plification hypotheses which we discuss here. Clearly these also indicate relevant research directions to
 491 explore.

492 6.1. Analysis of the multiple criteria aggregation procedures

493 The multiple criteria aggregation procedure used in this paper is based upon the concordance and non
 494 discordance principles. The obtained results are coherent. Nevertheless, the use of ELECTRE TRI method
 495 might lead to inconsistent results. For instance, let us consider the impact vector $(5, 1, 1, 3)$ characterizing
 496 the geographic units in zone 7 in case of scenario (*Mistral*, RP_3). Because of the discordance principle
 497 $(5, 1, 1, 3)$ will be rated 4. Let us consider a fictitious geographic unit characterized by $(4, 3, 3, 3)$, using the

498 same parameters, this last will be rated 3. However, (5, 1, 1, 3) is strictly preferred to (4, 3, 3, 3). Other
499 inconsistencies, might come from Condorcet Paradox due to the concordance principle.

500 6.2. Evaluating a map

501 A relevant question for the decision maker can be, how can we rate a geographic area? The answer to
502 this question is not simple. The rate of a geographic space depends upon:

- 503 • the characteristics of the problem, e.g. we may have interactions between geographic units (or not)
504 [16];
- 505 • the aggregation path, e.g. one possible path is aggregating the multiple criteria problem, then syn-
506 thesising uncertainties before rating the global map. Changing this order may lead to a different
507 result.

508 In this work, the interaction effects between geographic units is not taken into account because, in all
509 simulations, geographic units belonging to the same category of impact are grouped together.

510 7. Conclusion

511 We have presented an approach to assess spatial risks, in cases characterised by the presence of several
512 assets, spatial characteristics and uncertainties over the accident parameters (mainly the release position
513 and sea currents). The developed approach is illustrated through an application of nuclear releases in the
514 marine environment. The methodology aims to assess the impact of a nuclear accident at a geographic space
515 (in our case the Bay of Toulon) as part of a post-accident analysis. In order to evaluate the impact of a
516 nuclear release on a geographic space, several methods were used for decision aiding purposes. The procedure
517 developed consists of representing uncertainties through accident scenarios, structuring impact indices for
518 each asset and under each scenario, and synthesising these indices using a multiple criteria aggregation
519 procedure, describing the general impact over the studied area. We then aggregated uncertainties to evaluate
520 the vulnerability of the studied area regarding the accident scenarios. At a next step, we shall establish a
521 robustness analysis and study the possible recommendations to one or several decision markers, depending
522 on their risk aversion.

523 Acknowledgment

524 The authors would like to thank the “Agence Nationale de la Recherche” (ANR) project AMORAD
525 for supporting this work and the COST-action IS1304 for supporting a short term scientific mission at the
526 “Instituto de Ciencias Matematicas” (ICMAT) on expert judgment.

527 Appendix 1. The hydrodynamic model

528 1.1. The advection diffusion equation

To assess the radiological impact of an accidental release in seawater, the IRSN has developed a hydro-
dynamic model tool called STERNE (*Simulation du Transport et du transfert d'Éléments Radioactifs dans
l'environnement marin*, translated as *Simulation of radionuclide transport and transfer in marine envi-
ronments*) to simulate the dispersion of radionuclides in the marine area. This tool is based on the tracer
advection diffusion equation estimating the dispersion of radionuclides:

$$\frac{\partial C}{\partial t} - \frac{\partial}{\partial x} \left[D_x \frac{\partial c}{\partial x} - u_x c \right] - \frac{\partial}{\partial y} \left[D_y \frac{\partial c}{\partial y} - u_y c \right] - \frac{\partial}{\partial z} \left[D_z \frac{\partial c}{\partial z} - u_z c \right] = F(c, t)$$

529 where C is the radionuclide concentration; u , the advection current; and, finally, D is the turbulent diffusion
530 tensor. The model is illustrated with Figure 16:

531 Since it is difficult to solve this equation analytically, the most common procedure consists of discretising
532 time, the choice of time step depending on the mesh size and maximum sea current velocity for the area
533 considered and sigma-coordinates and calculates this concentration at each grid point and time step. This
534 model takes into account the half life of each radionuclide considered.




Figure 16: Schematic diagram of STERNE implementation principle

535 *1.2. Input data*

536 For each time step and mesh, the hydrodynamic data required as input to dispersion calculations includes

- 537 • The cumulative water fluxes in x , y and z directions; free surface elevation and diffusion coefficients
- 538 (set to calculate the exact quantity of water passing through the grid meshes at each instant and
- 539 should satisfy the continuity equation)
- 540 • The free surface elevation and diffusion coefficients.

541 Hydrodynamic models are generated based on hindcasts and forecasts of meteorological and tidal forcing.
 542 Source terms are characterised by:

- 543 • known quantities of radionuclide releases.
- 544 • known localisations (Release point coordinates).
- 545 • instants of releases.

546 **Appendix 2. Brief introduction of ELECTRE TRI**

547 ELECTRE TRI is a rating method, aiming to assign elements of a set A to one of predefined ordered
 548 categories C_1, \dots, C_p . Such categories are ranked from the worst to the best: $C_{h+1} \gg C_h \forall h \in \{1, \dots, p-1\}$
 549 where \gg refers to a complete order on the set of categories, [19]. This method uses a majority rule while
 550 respecting a minority using a veto rule, in order to compare elements of a set A (representing actions) to the
 551 profiles characterising categories. Let us denote r_1, \dots, r_p the limiting profiles characterising the p categories,
 552 r_k refers to the upper limit of category C_k and the lower limit of category C_{k+1} , $k = 1, 2, \dots, p$ and R the
 553 set of the associate indices. Let F denote the set of the indices of the criteria g_1, g_2, \dots, g_m . Without loss of
 554 generality, we make the assumption that preferences increase with the value on each criterion. ELECTRE
 555 TRI is based on an outranking relation S . Roughly speaking, an outranking relation can be interpreted as,
 556 "at least as good as". In a first step, we aim at constructing an outranking relation S characterising how
 557 actions compare to each limiting profile. Thus, we use S to assign each action to a specific category. The
 558 procedure can be described as follows:

- Partial concordance index $c_j(a, r_h) \in [0, 1], \forall j \in F, h \in R$: IT represents a weight of the proposition
a is at least as good as a certain r_h from the criterion j point of view. The formulation of partial
 concordance index is:

$$c_j(a, r_h) = \begin{cases} 1, & \text{if } g_j(r_h) - g_j(a) \leq 0 \\ 0, & \text{if } g_j(r_h) - g_j(a) > 0 \end{cases}$$

559 This index takes 1 to denote a full approval of the proposition " a is at least as good as r_h " from the
 560 criterion j point of view.

- Global concordance index $c(a, r_h) \in [0, 1], \forall h \in R$: represents the majority rule, i.e. the global weight of all criteria approving the proposition "a is at least as good as r_h ".

$$c(a, r_h) = \frac{\sum_{j \in F} w_j c_j(a, r_h)}{\sum_{j \in F} w_j}$$

561 where $w_j, j \in F$ refers to the weight associated to the criterion j .

- Discordance index $d_j(a, r_h) \in [0, 1], \forall j \in F, h \in R$: represent the respect of minority rule, i.e. when the difference between a certain r_h and a for a given criterion j is greater than a threshold, called veto threshold, the outranking relation between a and r_h is vetoed.

$$d_j(a, r_h) = \begin{cases} 1, & \text{if } g_j(r_h) - v_j(r_h) \geq g_j(a) \\ 0, & \text{otherwise} \end{cases}$$

562 where $v_j(r_h), j \in F, h \in R$ refers to the veto threshold associated with the criterion j .

- Credibility index or the outranking relation $\sigma(a, r_h)$ aggregating the concordance and the discordance.

564 In the ELECTRE TRI method, the assignment of a depends on the values of $\sigma(a, r_h)$, $\sigma(r_h, a)$ and a
565 cutting threshold λ . When $\sigma(a, r_h) \geq \lambda$, a outranks r_h , denoted aSr_h . Four possible situation may occur:

- $\sigma(a, r_h) \geq \lambda, \sigma(r_h, a) \geq \lambda \implies aIr_h$, i.e. a is indifferent to r_h
- $\sigma(a, r_h) < \lambda, \sigma(r_h, a) < \lambda \implies aRr_h$, i.e. a is incomparable to r_h
- $\sigma(a, r_h) \geq \lambda, \sigma(r_h, a) < \lambda \implies aPr_h$, i.e. a is preferred to r_h
- $\sigma(a, r_h) < \lambda, \sigma(r_h, a) \geq \lambda \implies r_hPa$, i.e. r_h is preferred to a

570 The assignment is done using two procedures:

- Pessimistic (conjunctive) procedure. It consists on the pairwise comparaison between each action a and the limiting profil r_h starting from $h = p$ to $h = 0$. We stop this procedure when aSr_h , and potentially a will be assigned to C_{h+1} .
- Optimistic (disjunctive) procedure. We compare each action a and the limiting profil r_h starting from $h = 1$ to $h = p$. We stop this procedure when r_hSa , and potentially a will be assigned to C_h .

576 The imperfection of knowledge about evaluations of criteria can be taken into account when defining the
577 thresholds of the aggregation model. However, it is not easy for the decision maker to provide precise and
578 complete information about weights and thresholds. Numerous technics were proposed in the literature to
579 elicit these parameters, [7], [14], [15], [17], [20].

580 References

- 581 [1] M. Aoyama and K. Hirose. Temporal variation of 137 cs water column inventory in the North Pacific since the 1960s.
582 *Journal of environmental radioactivity*, 69(1):107–117, 2003.
- 583 [2] M. Aoyama, K. Hirose, T. Miyao, Y. Igarashi, and P. Povinec. 137cs activity in surface water in the western North Pacific.
584 *Journal of Radioanalytical and Nuclear Chemistry*, 248(3):789–793, 2001.
- 585 [3] M. Aoyama, D. Tsumune, and Y. Hamajima. Distribution of 137cs and 134cs in the North Pacific Ocean: impacts of the
586 TEPCO Fukushima-Daiichi NPP accident. *Journal of Radioanalytical and Nuclear Chemistry*, 296(1):535–539, 2013.
- 587 [4] D. Bouyssou, Th. Marchant, M. Pirlot, A. Tsoukiàs, and Ph. Vincke. *Evaluation and decision models with multiple*
588 *criteria: Stepping stones for the analyst*. Springer Verlag, Boston, 1st edition, 2006.
- 589 [5] S. Bussotti, P. Guidetti, and G. Matricardi. Morphological analysis of Posidonia oceanica flowers from a reef formation
590 and a shallow water meadow of the Ligurian Sea (north-western Mediterranean). *Vie et milieu*, 48(1):55–62, 1998.
- 591 [6] J.M. Devaud, G. Groussaud, and E. Jacquet-Lagrece. UTADIS: Une methode de construction de fonctions d'utilite
592 additives rendant compte de jugements globaux. In *European working group on MCDA, Bochum, Germany*, 1980.

- 593 [7] L. Dias, V. Mousseau, J. Figueira, and J. Clímaco. An aggregation/disaggregation approach to obtain robust conclusions
594 with ELECTRE TRI. *European Journal of Operational Research*, 138(2):332–348, 2002.
- 595 [8] C. Dufresne, C. Duffa, and V. Rey. Wind-forced circulation model and water exchanges through the channel in the Bay
596 of Toulon. *Ocean Dynamics*, 64(2):209–224, 2014.
- 597 [9] C. Estournel, E. Bosc, M. Bocquet, C. Ulses, P. Marsaleix, V. Winiarek, I. Osvath, C. Nguyen, T. Duhaut, F. Lyard,
598 et al. Assessment of the amount of cesium-137 released into the Pacific Ocean after the Fukushima accident and analysis
599 of its dispersion in Japanese coastal waters. *Journal of Geophysical Research: Oceans*, 117(C11), 2012.
- 600 [10] P. H. Farquhar. State of the art: utility assessment methods. *Management science*, 30(11):1283–1300, 1984.
- 601 [11] M. C. Gambi, M. Lorenti, G. F. Russo, M. B. Scipione, and V. Zupo. Depth and seasonal distribution of some groups
602 of the vagile fauna of the *Posidonia oceanica* leaf stratum: structural and trophic analyses. *Marine Ecology*, 13(1):17–39,
603 1992.
- 604 [12] E. Jacquet-Lagrèze and Y. Siskos. Assessing a set of additive utility functions for multicriteria decision making: the UTA
605 method. *European Journal of Operational Research*, 10:151–164, 1982.
- 606 [13] P. Le Hir, R. S. Jacinto, B. Thouvenin, L. Guézennec, Ph. Bassoullet, Ph. Cugier, F. Leboulenger, R. Hocdé, P. Lesueur,
607 and L. Romaña. Courants, vagues et marées(les mouvements de l’eau). *Programme scientifique Seine-aval*, 2001.
- 608 [14] V. Mousseau. Eliciting information concerning the relative importance of criteria. In *Advances in multicriteria analysis*,
609 pages 17–43. Springer, 1995.
- 610 [15] V. Mousseau and R. Slowinski. Inferring an ELECTRE TRI model from assignment examples. *Journal of global opti-
611 mization*, 12(2):157–174, 1998.
- 612 [16] S. A. M. Takougang, V. Brison, and M. Pirlot. Two models for comparing decisional maps. *International Journal of
613 Multicriteria Decision Making* 73-74, 3(2-3):129–156, 2013.
- 614 [17] T. Tervonen, J. Figueira, R. Lahdelma, and P. Salminen. An inverse approach for ELECTRE III. *Instituto de Engenharia
615 de Sistemas e Computadores de Coimbra*, 2004.
- 616 [18] A. Tsoukiàs. On the concept of decision aiding process. *Annals of Operations Research*, 154:3 – 27, 2007.
- 617 [19] W. Yu. *Aide multicritère à la décision dans le cadre de la problématique du tri: Concepts, méthodes et applications*. PhD
618 thesis, Paris 9, 1992.
- 619 [20] J. Zheng, S. A. M. Takougang, V. Mousseau, and M. Pirlot. Learning criteria weights of an optimistic Electre Tri sorting
620 rule. *Computers & Operations Research*, 49:28–40, 2014.