

1 **Accepted: Pest Management Science**

2

3 **Azadirachtin effects on reproductive traits and progeny survival in**
4 ***Drosophila melanogaster* (Diptera)**

5

6 **Running title:** Azadirachtin reproductive toxicity

7

8 Chemseddine M. Oulhaci¹, Béatrice Denis², Samira Kilani-Morakchi¹, Jean-Christophe.
9 Sandoz², Laure Kaiser-Arnaud², Dominique Joly² and Nadia Aribi¹

10

11

12 ¹*Laboratoire de Biologie Animale Appliquée. Faculté des Sciences. Université Badji Mokhtar*
13 *Annaba. BP12, 23000, Annaba, Algeria.*

14 ²*Laboratoire Evolution, Génomes, Comportements, Ecologie, UMR 9191, CNRS, IRD,*
15 *Université Paris-Sud et Université Paris-Saclay, 91198 avenue de la Terrasse, Gif-sur-Yvette*
16 *Cedex, France.*

17

18

19 **Corresponding author**

20

21 Nadia Aribi

22 nadia.aribi@univ-annaba.dz

23

24 Tel : +213551273853

25

26 *Laboratoire de Biologie Animale Appliquée. Faculté des Sciences. Université Badji Mokhtar Annaba.*
27 *BP12, 23000, Annaba, Algeria.*

28

29

30

31

32

33

1 **ABSTRACT**

2 BACKGROUND - Azadirachtin is a prominent natural pesticide and represents an alternative
3 to conventional insecticides. It has been successfully used against insect pests but has also
4 various effects on non-target organisms, that still are poorly known. Here, we investigated
5 lethal and sublethal effects of Azadirachtin on treated adults of *Drosophila melanogaster*
6 (Meigen) and examined determined and several parameters of reproduction (fertility,
7 spermatogenesis, and oogenesis) ..

8 RESULTS – Azadirachtin, applied topically on newly emerged adults, increased mortality
9 with a positive dose-dependent relationship. The LD₅₀ (0.63 µg) was determined 24 hours
10 after treatment using a non-linear regression. Azadirachtin induced a significant reduction in
11 reproductive potential in treated flies, particularly when females were treated. In addition, the
12 number of cysts and the apical nuclei positions within the cysts decreased by 29.7% and 20%
13 respectively in males. In females, the number of oocytes per ovary and the volume of basal
14 oocytes also decreased by 16.1% and 32.4% respectively.

15 CONCLUSION - Azadirachtin causes significant toxic effects in both sexes and decreases the
16 fecundity in non-targeted species. Females are more sensitive to Azadirachtin.

17

18 **Key words:** natural pesticide, Azadirachtin, toxicity, spermatogenesis, oogenesis, *Drosophila*

19

1 I. INTRODUCTION

2 Crop losses caused by pests may represent the equivalent of the food needed to feed over 1
3 billion people.¹ The use of pesticides is therefore required to protect agrosystems. However,
4 their side effects induce important health^{2,3} and environmental problems.^{4,5} Consequently,
5 natural compounds with pesticidal properties (biopesticides) have been investigated for
6 decades and offer a more sustainable solution to pest control than synthetic or conventional
7 products.⁸

8 Azadirachtin, a tetranortriterpenoid derived from the seeds of the Indian neem tree
9 (*Azadirachta indica* A. Juss, Meliaceae), is one of the main commercialized biopesticides and
10 remains the most successful botanical pesticide in agricultural use worldwide.^{9,10} It possesses
11 a strong toxicity against insect pests of different orders.^{9,11-14} Besides its insecticidal action,
12 Azadirachtin is also used in traditional medicine, in Asia and in Africa as an antidiabetic,
13 immunostimulant, antimicrobial, antiviral, contraceptive and anticancer remedy.¹⁵⁻¹⁷

14 Azadirachtin is an insecticide with rapid biodegradability and without resistance problems
15 due to its chemical complexity.⁹ This pesticide is also reported to be relatively safer than most
16 conventional insecticides.^{18,19} It is non-toxic to humans and warm-blooded vertebrates²⁰ and
17 without genotoxicity for mammals.^{21,22} However, several studies have questioned the safety
18 of neem products on non-target species, but the results remain contradictory.²³⁻²⁸ For example,
19 behavior and morphological alterations have been highlighted in bees,²⁸ while Bernades *et*
20 *al.*²⁶ did not note any effect on worker bee mortality, flight, or respiration rate; furthermore,
21 in aquatic organisms, Azadirachtin is used as a potential immunostimulant and a promising
22 agent to control fish parasites.²⁷ Because Azadirachtin acts as an insect growth disruptor²⁹
23 negative side effects on beneficial arthropods are expected.

24 Azadirachtin interferes with the endocrine system and interacts with juvenile hormone and
25 20-hydroxyecdysone. It induces growth and moulting abnormalities and affects ovarian

1 development, oocyte structure, fecundity, oviposition and egg viability.^{9, 12,30-32} Azadirachtin
2 also shows moderate to strong cytotoxicity, neurotoxicity and antimitotic effects.^{9,33-36} It acts
3 on the expression of genes related to development in *Drosophila melanogaster*.¹⁸ In addition,
4 varied effects have been noted according to the tested Azadirachtin formulation.^{12,37}
5 Nevertheless, the mechanism of action of this pesticide remains elusive.^{18,19}
6 Direct effects of Azadirachtin on various developmental processes, related to the signaling of
7 20-hydroxyecdysone, have been previously reported.⁹ However, the mechanisms of action of
8 Azadirachtin on reproduction are still poorly known. Azadirachtin has been shown to
9 influence oviposition behavior in several insects,^{11,38} but its precise effects on reproductive
10 parameters would benefit from being more deeply investigated. Therefore, the aim of this
11 study was double: on the one hand, it quantified the side effects of Azadirachtin on the non-
12 target insect *D. melanogaster* using doses allowing 25 and 50% of survivors, *i.e.* mimicking
13 possible exposure to the insecticide residues; on the other hand, it filled some gaps in
14 evaluating Azadirachtin reprotoxic effects in this biological model species. Effects were
15 assessed (1) following treatment on newly emerged adults, and (2) on various reproductive
16 traits of adult males and females. Previous studies have already shown that Azadirachtin
17 impacts the reproduction of *D. melanogaster* treated at the pupae³⁹ and larvae stages.³⁸ Thus,
18 we considered its impact on the survival of progeny at various development stages and on
19 fertility in both sexes. We quantified the number of descendants produced by surviving adults
20 (numbers of eggs, larvae, pupae and adults obtained at the F₁ generation) and also the
21 fecundity of males and females (number of cysts and potential anomalies of spermatocyte
22 nucleus positions; number of oocytes and size of basal oocyte). A toxicity study was first
23 carried out to determine lethal doses for *D. melanogaster* adults in our experimental
24 conditions.

25

1 **2. MATERIAL AND METHODS**

2 **2.1 Insect rearing**

3 *Drosophila melanogaster* (Canton-S) was reared on standard cornmeal medium at $25 \pm 2^\circ\text{C}$,
4 70% relative humidity under a 12-hour light/dark photoperiod. Flies were transferred every
5 three days to avoid larval competition and to regularly provide abundant progeny for testing.

6 **2.2 Insecticide and toxicity tests**

7 The Neem Azal (1% Azadirachtin; TrifolioM GmbH; Lahnau, Germany) commercial
8 formulation, was dissolved in acetone and topically applied on adults (sexes combined after
9 screening and showing any difference between male and female) less than 6 hours after
10 ecdysis (1 μl per insect according to Di Prisco *et al.*).⁴⁰ Control insects were treated with
11 solvent alone. Five doses (0.1, 0.2, 0.4, 0.6 and 1.2 μg of Neem Azal acetone solution) were
12 tested,³⁹ considering 3 replicates per dose, each consisting of 300 insects. Adult mortality
13 percentages were calculated between 24 and 96 hours after treatment and corrected in
14 accordance with Abbott.⁴¹ A non-linear regression was used to determine the lethal doses
15 (LD) LD₂₅ and LD₅₀, corresponding to 25% and 50% of adult mortality respectively, with
16 their corresponding 95% fiducial limits (95% FL) and Hill slope.

17 **2.3 Progeny output**

18 Newly moulted (< 6 hours) *D. melanogaster* adults were treated topically with Azadirachtin at
19 the LD₂₅ and LD₅₀. Subsequently, one male and one female from the control and treated series
20 were placed in a Petri dish (90 x 14.2 mm) containing standard cornmeal medium to obtain
21 their progeny. Eight repetitions of different types of couples were realized (control male +
22 control female; control male + female LD₂₅; male LD₂₅ + control female; male LD₂₅, + female
23 LD₂₅; control male + female LD₅₀; male LD₅₀ + control female; male LD₅₀ + female LD₅₀).
24 After 48 hours, the adults were removed and generation 1 (F₁) was followed daily until all

1 adults had emerged. The numbers of eggs, larvae, pupae and adults that emerged from each
2 series of experiments were counted.

3 **2.4 Cyst parameters in males**

4 Developmental factors leading to male infertility were investigated by analyzing post-meiotic
5 abnormalities, *i.e.* cyst numbers and spermatocyte nucleus localization, both parameters
6 influencing the number of sperm produced.⁴² Newly moulted male adults (< 6 hours) were
7 treated topically with Azadirachtin at the LD₅₀ (0.63 µg). Individuals from the control and
8 treated series were placed in vials containing standard cornmeal medium. After 48 hours,
9 testes were dissected in a phosphate-buffered saline (PBS) solution with 4', 6-diamidino-2-
10 phenylindole (DAPI) allowing nuclei staining.⁴³ Cysts were gently spread on the blade and
11 covered with a cover slip. Cysts were then observed under a fluorescence microscope and the
12 number of cysts per male and the position of spermatocyte nuclei within the cyst were
13 counted. Thirty repetitions per condition were performed.

14 **2.5 Ovarian parameters in females**

15 To detect a possible impact of Azadirachtin on female fecundity, classical ovarian parameters
16 were investigated (number of oocytes and size of basal oocyte). Newly emerged female adults
17 (< 6 hours) were treated topically with the Neem Azal at the LD₅₀ (0.63 µg). Individuals from
18 the control and treated series (LD₅₀: 0.63 µg) were placed in vials containing standard
19 cornmeal medium. After 48 hours, ovaries were dissected out. After removal of the
20 circumovarian fat body, the numbers of oocytes were scored together with the volume of the
21 basal oocyte.⁴⁴

22 **2.6 Statistical analysis**

23 The mean ± standard deviations (SD) were calculated for each experimental group. Data from
24 the toxicity assay were analyzed using non-linear sigmoid curve fitting. The goodness of fit to
25 the curve model was evaluated on the basis of R² values. The homogeneity of variances was

1 checked using Bartlett's and Shapiro Wilk Tests. Analyses of variance (one-way and two-
2 way) were performed and followed by Tukey's HSD test for multiple comparisons when
3 significant. Repeated-measure Anova was also used for progeny output between stages for the
4 same couple. Numbers of cysts and ovarian parameters were compared between the treated
5 and control groups using Student's t test. All calculations were performed using GraphPad
6 Prism (v6.01 for Windows).

7

8 **3. RESULTS**

9 **3.1 Insecticidal activity**

10 Azadirachtin, applied topically on newly emerged adults of *D. melanogaster*, induces an
11 insect mortality, 24 hours after treatment. Corrected mortality percentages were 13.92 ± 1.43
12 % for the lowest dose (0.1 μg) and 71.80 ± 3.11 % for the highest dose (1.2 μg). The mortality
13 percentage recorded in untreated animals was 5.48 ± 0.53 % (control mortality, Fig.1).
14 Statistical analysis revealed a significant dose effect ($F_{4, 10} = 62.43$; $p < 0.0001$) and Tukey's
15 HSD test showed a significant increase in mortality with increasing doses (Fig. 1). The lethal
16 doses (LD) recorded with 95% fiducial limits (95% FL) were $\text{LD}_{25} = 0.23$ (0.13-0.38), $\text{LD}_{50} =$
17 0.63 (0.44-0.91) and $\text{LD}_{90} = 4.85$ (1.47-15.98) μg (Table 1). The non-linear regression fitted
18 on these results indicated a Hill slope of 1.08 (0.54-1.62). No significant effect of the time
19 elapsed after treatment (24 to 96 hours) was observed ($F_{5, 60} = 1.05$; $p = 0.39$; Table 2).

20 **3.2 Progeny output**

21 To evaluate the progeny from surviving treated adults, we monitored the different
22 developmental stages (eggs, larvae, pupae and adults) in the progeny of couples formed with
23 control animals (female, F_C ; male, M_C) and/or animals treated with the LD_{50} (female, F_T , male
24 M_T). Statistical analysis showed a significant decrease in the numbers of eggs, larvae, pupae
25 and adults in the F_1 generation for all treated series compared to couples from control

1 individuals (Repeated-measure ANOVA: treatment effect: $F_{6, 42} = 241.90$, $p < 0.0001$; stage
2 effect: $F_{3, 21} = 15.94$, $p < 0.0001$; interaction: $F_{18, 126} = 0.60$, $p = 0.89$; Fig. 2). The comparison
3 of the different series revealed a significant dose effect. Irrespective of the developmental
4 stage considered, Tukey's HSD test detailed three groups ($M_C + F_C > M_T + F_C > M_C + F_T =$
5 $M_T + F_T$). This result reveals a greater sensitivity of females to Azadirachtin.

6 In addition, for all stages evaluated, a dose-dependent effect was noted and a more significant
7 reduction was recorded at the LD_{50} than at LD_{25} (supplementary Fig. 1). For homogenization
8 of the results along the core of the paper, the values of treated series with LD_{25} are not
9 presented in figure 2.

10 **3.3 Cyst parameters**

11 The effect of Azadirachtin treatment was tested on male fecundity. Azadirachtin was applied
12 topically at LD_{50} ($0.63 \mu\text{g}$) on male *D. melanogaster* on the day of emergence. The number of
13 cysts per testis was counted 48 hours after emergence. Control adults displayed 28.47 ± 2.16
14 cysts per testis. This number decreased significantly after topical application of Azadirachtin
15 and reached 20 ± 1.69 ($t_{58} = 3.08$; $p = 0.003$) (Fig. 3A). In addition, the percentage of
16 spermatocyte nuclei in abnormal position significant increased in LD_{50} males ($\chi^2 = 5.30$, 1; z
17 $= 2.30$, $p = 0.021$) (Fig. 3B).

18 **3.4 Ovarian parameters**

19 The effect of Azadirachtin treatment was tested on female fecundity. Azadirachtin was
20 applied topically at LD_{50} ($0.63 \mu\text{g}$) on female *D. melanogaster* on the day of emergence. The
21 number of oocytes was counted 48 hours after emergence. We found a significant decrease in
22 the number of oocytes in the treated series (13.90 ± 0.55) compared to controls (16.57 ± 0.53 ;
23 $t_{58} = 3.43$; $p = 0.0011$, Fig. 4A). Furthermore, the volume of the basal oocyte showed a
24 reduction in treated females ($0.0048 \pm 0.0003 \text{ mm}^3$) compared to controls (0.0071 ± 0.0005
25 mm^3) ($t_{58} = 3.87$; $p = 0.0003$, Fig. 4B).

1

2 **4. DISCUSSION**

3 In this study, we investigated Azadirachtin effects on adult toxicity and reproduction in
4 *D. melanogaster*. Azadirachtin, applied topically on newly emerged adults, induced
5 comparable mortality when evaluated at different times (24 – 96 hours) after treatment. This
6 result is in agreement with those of Andreazza *et al.*⁴⁵ on two other Drosophilidae, *Drosophila*
7 *suzukii* (Matsumara) and *Zaprionus indianus* (Gupta). Azadirachtin acts with a dose-response
8 relationship and the LD₅₀ we obtained (0.63 µg or 630 ppm, evaluated 24 h after treatment) is
9 close to that found by Bezzar-Bendjazia *et al.*,³⁸ despite the fact that these authors used
10 topical application at the last larval stage (0.67 µg). Azadirachtin seems less toxic (1.17 µg)
11 when applied on newly formed pupae of *D. melanogaster*.³⁹ Neem-oil, another Azadirachtin
12 formulation applied topically at the same stage, induces similar toxicity in *Drosophila*³⁹ in
13 spite of two antifeedants (nimbine and salanine) in its composition. This can be explained by
14 the mode of application; indeed, oral administration could have provided different results. The
15 most likely explanation for this effect is a lower susceptibility due to the slow metabolism at
16 this stage of development. Nevertheless, the pupal stage, a critical phase for adult formation,
17 remains very responsive to Azadirachtin,³⁹ which inherently induces a drastic remodeling of
18 most tissues and organs.⁴⁶

19 Azadirachtin presents a stronger toxicity in *D. suzukii* and *Z. indianus* adults comparatively to
20 *D. melanogaster*.⁴⁵ Indeed, a concentration of only 12 ppm causes 40% mortality in those
21 species. Similar observations were reported by De Andrade-Coelho *et al.*⁴⁷ in female adults of
22 another Diptera *Lutzomyia longipalpis* (Lutz and Neiva). Previously, Azadirachtin was
23 reported to impair, with variable effects, the survival of different insect species.^{11-14,48} Strong
24 variations in insects' susceptibility to Azadirachtin were noted depending on insect order,
25 species, formulation, or the method of application.^{12,26,28,49-51} Sensitivity of insects to

1 insecticides varies also with the regulation of receptors,⁵² penetration rate through the cuticle,
2 absorption by insects, transport in tissues of the body and metabolism.⁵³

3 Several mechanisms may be at play in causing acute Azadirachtin toxicity. Azadirachtin
4 effects on adult mortality may be linked to the cytotoxicity and induction of apoptosis causing
5 general disruption in the organs of *D. melanogaster*,^{54,55} and in others species, like
6 *Spodoptera frugiperda*.^{9,56} As supposed by Lai *et al.*,¹⁸ Shao *et al.*⁵⁴ revealed that Azadirachtin
7 induced apoptotic cell death *via* insulin receptors and its signaling pathways.

8 Mortality can increase in *D. melanogaster* if the nutritional balance is impacted⁵⁷ and this
9 parameter might be affected by Azadirachtin. Indeed, in *Drosophila*, rhythmic activity of
10 feeding muscles is controlled by specific motor neurons⁵⁸ and cytotoxicity can be explained
11 by a blockage of voltage-gated calcium channels and their impact on cholinergic synapses.³⁴

12 Azadirachtin could also induce morphological and physiological damages on the salivary
13 glands, as was noted by Remedio *et al.*³⁶ in the mite (*Rhipicephalus sanguineus*, Latreille). In
14 addition, the metabolic fight against the insecticide's toxicity^{13,39} may involve a high energy
15 requirement.⁵⁹ These processes affecting the flies' nutritional state, immediately following
16 metamorphosis, could be potentially fatal and may contribute to the mortality observed in this
17 study.

18 Quantitative and qualitative variations in nutrition significantly influence reproductive
19 physiology and behavior as well as overall reproductive success.⁶⁰ In *Drosophila*, disruptions
20 in energy homeostasis severely affect reproduction⁶¹⁻⁶³ and a correct lipid composition is
21 crucial for spermatogenesis.⁶⁴ Effects of Azadirachtin on fecundity and oviposition have been
22 reported in *D. melanogaster*,^{38,39} other Diptera^{31,32} and several insects orders.^{9,12,28,51}

23 Observations made in the current study indicate possible causes for these reprotoxic effects.
24 Indeed, the fecundity and fertility of *D. melanogaster* that survived Azadirachtin treatment
25 was adversely affected, resulting in a decreased progeny, and these depressive effects were

1 positively correlated with a reduction of cyst and oocyte numbers, an increase in spermatocyte
2 nucleus anomalies and a decrease in the size of basal oocytes, all of which can explain the
3 reduced progeny of treated adults. Females of *Drosophila* seem more sensitive than males in
4 spite of a similar toxicity; this may be explained by a distinct physiology between the sexes. It
5 is known that Azadirachtin causes significant effects on the reproductive system of both male
6 and female insects by causing alterations and morphological changes in oocytes and
7 spermatocytes.^{9,35} In addition, the reduction of the egg layer can be explained by the
8 neurotoxic action of Azadirachtin *via* the role of follicular adrenergic signaling in ovulation,
9 demonstrated recently in *D. melanogaster*.⁶⁵

10 The diminution of the size of basal oocytes in females, in agreement with several
11 authors,^{9,12,28,48} may be linked to a vitellogenesis impairment already noted in *D.*
12 *melanogaster* after Azadirachtin treatment.³⁹ Azadirachtin alters or prevents the formation of
13 a new actin cytoskeleton, resulting in disruption of cell division and blocked transport,⁹ which
14 may affect the dumping of cytoplasmic contents of nurse cells to the oocyte.⁶⁶ Azadirachtin
15 can interfere with vitellogenin synthesis and/or its uptake, affecting oocyte development *via*
16 juvenile hormone (JH). For the normal progress of oogenesis, vitellogenesis and
17 spermatogenesis, a proper balance between JH and 20-hydroxyecdysone (20E) is of
18 paramount importance in *D. melanogaster*.^{66,67}

19 In recent literature, ecdysone or JH signaling pathways seem implicated in the regulation of
20 microvilli formation in follicular epithelial cells⁶⁸ or in intestine changes (enterocyte
21 remodeling) to yield a larger organ to support increased lipid metabolism.⁶⁹ Consequently, the
22 impairment of flies' reproductive capacity could be explained by an antagonist action of
23 Azadirachtin on these two essential hormones. The reproductive process is linked to
24 nutritional signals controlled by the insulin signaling pathways, which are situated within a
25 complex physiological regulatory network that also encompasses the lipophilic hormones, 20-

1 hydroxyecdysone and JH.^{70,71} Furthermore, direct action of Azadirachtin on the nervous
2 system³⁴ can also interfere indirectly on different endocrinological processes in insects.⁷²
3 In conclusion, our study shows strong toxic effects of Azadirachtin, from 24 hours after
4 treatment, on the non-target species *D. melanogaster*; this species, however, is less sensitive
5 than the targeted species.^{11,45,47} This study also highlights a more drastic impact of
6 Azadirachtin on progeny output in females compared to males. Further investigations are
7 needed to achieve a better understanding of these effects.

8

9 **ACKNOWLEDGEMENTS**

10 We are very grateful to M. Bonneau and S. Nortier for their help in insect media preparation.
11 We thank Claude Wicker-Thomas for fruitful discussions and M. Eden for English editing.
12 This work was supported by grants from the National Fund for Scientific Research of Algeria
13 (Laboratory of Applied Animal Biology) to N. Soltani, and from IDEEV (Institut Diversité,
14 Ecologie et Evolution du Vivant, FR3284, CNRS, AAP-2014) to D. Joly.

15

16 **REFERENCES**

- 17 1. Birch A N E, Begg G S and Squire G R, How agro-ecological research helps to address
18 food security issues under new IPM and pesticide reduction policies for global crop
19 production systems. *J Exp Bot* **62**:3251-3261(2011).
- 20 2. Nicolopoulou-Stamati P, Maipas S, Kotampasi C, Stamatis P and Hens L, Chemical
21 pesticides and human health: the urgent need for a new concept in agriculture. *Front Public*
22 *Health* **4**:148 (2016).
- 23 3. Vrijheid M, Casas M, Gascon M, Valvi D and Nieuwenhuijsen M, Environmental
24 pollutants and child health, A review of recent concerns. *Int J Hyg Environ Health* **219**:331-
25 342 (2016).

- 1 4. Foley J A, DeFries R, Asner G P, Barford C, Bonan G, Carpenter S R, Chapin F S, Coe M
2 T, Daily G C, Gibbs H K, Helkowski J H, Holloway T, Howard E A, Kucharik C J, Monfreda
3 C, Patz J A, Prentice I C, Ramankutty N and Snyder P K, Global consequences of land
4 use. *Science* **309**:570-574 (2005).
- 5 5. Köhler H R and Triebkorn R, Wildlife ecotoxicology of pesticides: can we track effects to
6 the population level and beyond? *Science* **341**:759-765 (2013).
- 7 6. Villaverde J J, Sevilla-Morán B, Sandín-España P, López-Goti C and Alonso-Prados J L,
8 Biopesticides in the framework of the European pesticide regulation (EC) No.
9 1107/2009. *Pest Manag Sci* **70**:2-5 (2014).
- 10 7. Mkenda P, Mwanauta R, Stevenson P C, Ndakidemi P, Mtei K and Belmain S R, Extracts
11 from field margin weeds provide economically viable and environmentally benign pest
12 control compared to synthetic pesticides. *PloS One* **10**:e0143530 (2015).
- 13 8. Olson S, An analysis of the biopesticide market now and where it is going. *Outlooks Pest*
14 *Manag* **26**:203-206 (2015).
- 15 9. Mordue L A J, Morgan E D and Nisbet A J, Azadirachtin, a natural product in insect
16 control, in *Comprehensive Molecular Insect Science*, ed. by Gilbert L I, Iatrou K and Gill SS,
17 Elsevier Oxford, UK, pp. 117-135 (2005).
- 18 10. Isman M B and Grieneisen M L, Botanical insecticide research: many publications,
19 limited useful data. *Trends Plant Sci* **19**:140-145 (2014).
- 20 11. Tomé H V, Martins J C, Corrêa A S, Galdino T V S, Picanço M C and Guedes R N C,
21 Azadirachtin avoidance by larvae and adult females of the tomato leafminer *Tuta*
22 *absoluta*. *Crop Prot* **46**:63-69 (2013).
- 23 12. Abedi Z, Saber M, Vojoudi S, Mahdavi V and Parsaeyan E, Acute, sublethal and
24 combination effects of azadirachtin and *Bacillus thuringiensis* on the cotton bollworm,
25 *Helicoverpa armigera*. *J Insect Sci* **14**:1-9 (2014).

- 1 13. Dere B, Altuntas H and Nurullohoğlu Z U, Insecticidal and oxidative effects of
2 azadirachtin on the model organism *Galleria mellonella* L. (Lepidoptera: Pyralidae). *Arch*
3 *Insect BiochemPhysiol* **89**:138-152 (2015).
- 4 14. Poland T M, Ciaramitaro T M and McCullough D G, Laboratory evaluation of the toxicity
5 of systemic insecticides to emerald ash borer larvae. *J Econ Entomol* **109**:705-716 (2016).
- 6 15. Koriem K M, Review on pharmacological and toxicological effects of oleum
7 azadirachtin oil. *Asian Pac J Trop Biomed* **3**:834-840 (2013).
- 8 16. Hao F, Kumar S, Yadav N and Chandra D, Neem components as potential agents for
9 cancer prevention and treatment. *Biochim Biophys Acta* **1846**:247-257(2014).
- 10 17. Hummel H E, Langner S , Hein D F, Sanguanpong U and Schmutterer H, Unusually
11 versatile plant genus *Azadirachta* with many useful and so far incompletely exploited
12 properties for agriculture, medicine and industry. *Acta Fytotechn Zootechn* **18**:169-175
13 (2016).
- 14 18. Lai D, Jin X, Wang H, Yuan M and Xu H, Gene expression profile change and growth
15 inhibition in *Drosophila* larvae treated with azadirachtin. *J Biotechnol* **185**:51-56 (2014).
- 16 19. Wang H, Lai D, Yuan M and Xu H, Growth inhibition and differences in protein profiles
17 in azadirachtin-treated *Drosophila melanogaster* larvae. *Electrophoresis* **35**:1122-1129
18 (2014).
- 19 20. Salehzadeh A, Jabbar A, Jennens L, Ley S V, Annadurai R S, Adams Rand Strang R H C,
20 The effects of phytochemical pesticides on the growth of cultured invertebrate and vertebrate
21 cells. *Pest Manag Sci* **58**:268-276 (2002).
- 22 21. Celestino D, Braoios G I, Ramos R S, Gontijo L M and Guedes R N C, Azadirachtin-
23 mediated reproductive response of the predatory pirate bug *Blaptostethus pallescens*. *Biol*
24 *Control* **59**:697-705 (2014).

- 1 22. Saxena A and Kesari V P, Lack of genotoxic potential of pesticides, spinosad,
2 imidacloprid and neem oil in mice (*Mus musculus*). *J Environ Biol* **37**:291-295 (2016).
- 3 23. Lima D B, Melo J W S, Guedes N M P, Gontijo L M, Guedes R N C and Gondim Jr M G
4 C, Bioinsecticide-predator interactions: Azadirachtin behavioral and reproductive impairment
5 of the coconut mite predator *Neoseiulus baraki*. *PloS One* **10**:e0118343 (2015).
- 6 24. Luna-Cruz A, Rodríguez-Leyva E, Lomeli-Flores J R, Ortega-Arenas L D, Bautista-
7 Martínez N and Pineda S, Toxicity and residual activity of insecticides against *Tamarixia*
8 *triozae* (Hymenoptera: Eulophidae), a parasitoid of *Bactericera cockerelli* (Hemiptera:
9 Triozidae). *J Econ Entomol* **108**:2289-2295 (2015).
- 10 25. Turchen L M, Golin V, Butnariu A R, Guedes R N C and Pereira M J B, Lethal and
11 sublethal effects of insecticides on the egg parasitoid *Telenomus podisi* (Hymenoptera:
12 Platygasteridae). *J Econ Entomol* **109**:84-92 (2016).
- 13 26. Bernardes R C, Tomé H V, Barbosa W F, Guedes R N C, and Lima M A P, Azadirachtin
14 induced antifeeding in Neotropical stingless bees. *Apidologie* 1-11. (2016).
- 15 27. Banerjee A, Manna S, and Saha S K, Effect of aqueous extract of *Azadirachta indica* A.
16 Juss (neem) leaf on oocyte maturation, oviposition, reproductive potentials and embryonic
17 development of a freshwater fish ectoparasite *Argulus bengalensis* Ramakrishna, 1951
18 (Crustacea: Branchiura). *Parasitol Res* **113**:4641-4650 (2014).
- 19 28. Barbosa W F, Tomé H V, Bernardes R C, Siqueira M A L, Smagghe G and Guedes R N
20 C, Biopesticide-induced behavioral and morphological alterations in the stingless bee
21 *Melipona quadrifasciata*. *Environ Toxicol Chem* **34**:2149-2158 (2015).
- 22 29. Pener M P and Dhadialla T S, An overview of insect growth disruptors; applied
23 aspects. *Adv In Insect Phys* **43**:1-162 (2012).

- 1 30. Lucantoni L, Giusti F, Cristofaro M, Pasqualini L, Esposito F, Lupetti P and Habluetzel
2 A, Effects of a neem extract on blood feeding, oviposition and oocyte ultrastructure in
3 *Anopheles stephensi* Liston (Diptera: Culicidae). *Tissue and Cell* **38**:361-371 (2006).
- 4 31. Dembo E G, Abay S M, Dahiya N, Ogboi J S, Christophides G K, Lupidi G, Chianese G,
5 Lucantoni L and Habluetzel A, Impact of repeated NeemAzal®-treated blood meals on the
6 fitness of *Anopheles stephensi* mosquitoes. *Parasit Vectors* **8**:94 (2015).
- 7 32. Evans B G and Hallett R H, Efficacy of biopesticides for management of the swede midge
8 (Diptera: Cecidomyiidae). *J Econ Entomol* **109**:2159-2167 (2016).
- 9 33. Huang J F, Shui K J, Li H Y, Hu M Y and Zhong G H, Antiproliferative effect of
10 azadirachtin A on *Spodoptera litura* SI-1 cell line through cell cycle arrest and apoptosis
11 induced by up-regulation of p53. *Pestic Biochem Physiol* **99**:16-24 (2011).
- 12 34. Qiao J, Zou X, Lai D, Yan Y, Wang Q, Li W, Deng S, Xu H and Gu H, Azadirachtin
13 blocks the calcium channel and modulates the cholinergic miniature synaptic current in the
14 central nervous system of *Drosophila*. *Pest Manag Sci* **70**:1041-1047 (2014).
- 15 35. Remedio R N, Nunes P H, Anholeto L A, Oliveira P R and Camargo-Mathias M I,
16 Morphological effects of neem (*Azadirachta indica* A. Juss) seed oil with known azadirachtin
17 concentrations on the oocytes of semi-engorged *Rhipicephalus sanguineus* ticks (Acari:
18 Ixodidae). *Parasitol Res* **114**:431-444 (2015).
- 19 36. Remedio R N, Nunes P H, Anholeto L A, Oliveira P R, Sá I C G and Camargo-Mathias M
20 I, Morphological alterations in salivary glands of *Rhipicephalus sanguineus* ticks (Acari:
21 Ixodidae) exposed to neem seed oil with known azadirachtin concentration. *Micron* **83**:19-31
22 (2016).
- 23 37. Da Costa J T, Forim M R, Costa E S, De Souza J R, Mondego J M and Junior A L B,
24 Effects of different formulations of neem oil-based products on control *Zabrotes subfasciatus*
25 Boheman, 1833 (Coleoptera: Bruchidae) on beans. *J Stored Prod Res* **56**:49-53 (2014).

- 1 38. Bezzar-Bendjazia R, Kilani-Morakchi S and Aribi N, Larval exposure to azadirachtin
2 affects fitness and oviposition site preference of *Drosophila melanogaster*. *Pestic Biochem*
3 *Physiol* **133**:85-90 (2016).
- 4 39. Boulahbel B, Aribi N, Kilani-Morakchi S and Soltani N, Insecticidal activity of
5 azadirachtin on *Drosophila melanogaster* and recovery of normal status by exogenous 20-
6 hydroxyecdysone. *Afr Entomol* **23**:224-233(2015).
- 7 40. Di Prisco G, Cavaliere V, Annoscia D, Varricchio P, Caprio E, Nazzi F, Gargiulo G and
8 Pennacchio F, Neonicotinoid clothianidin adversely affects insect immunity and promotes
9 replication of a viral pathogen in honey bees. *Proc Natl Acad Sci* **110**:18466-18471 (2013).
- 10 41. Abbott W S, A method of computing the effectiveness of an insecticide. *J Econ*
11 *Entomol* **18**:265-267 (1925).
- 12 42. Lindsley D L and Tokuyasu K. T, Spermatogenesis, in *Genetics and Biology in*
13 *Drosophila*, ed. by Ashburner M and Wright T. Academic Press 2nd edition, New York, pp.
14 225-294 (1980).
- 15 43. Montchamp-Moreau C, Joly D, Abnormal spermiogenesis is associated with the X-linked
16 sex-ratio trait in *Drosophila simulans*. *Heredity* **79**:24-30 (1997).
- 17 44. Lumbreras C J, Galante E and Mena J, Ovarian condition as an indicator of the phenology
18 of *Bubas bubalus* (Coleoptera: Scarabaeidae). *Ann Entomol Soc Am* **84**:190-194 (1991).
- 19 45. Andrezza F, Bernardi D, Baronio C A, Pasinato J, Nava D E and Botton M, Toxicities
20 and effects of insecticidal toxic baits to control *Drosophila suzukii* and *Zaprionus indianus*
21 (Diptera: Drosophilidae). *Pest Manag Sci* **73**:146-152 (2017).
- 22 46. Ureña E, Manjón C, Franch-Marro X and Martín D, Transcription factor E93 specifies
23 adult metamorphosis in hemimetabolous and holometabolous insects. *Proc Natl Acad Sci*
24 **111**:7024-7029. (2014).

- 1 47. De Andrade-Coelho C A, De Souza N A, Silva V C, Souza A A, Gonzalez M S and
2 Rangel E F, Effects of azadirachtin on the biology of *Lutzomyia longipalpis* (Diptera:
3 Psychodidae: Phlebotominae) adult female, the main vector of American visceral
4 leishmaniasis. *J Med Entomol* **51**:891-895 (2014).
- 5 48. Ahmad S, Ansari M S and Muslim M, Toxic effects of neem based insecticides on the
6 fitness of *Helicoverpa armigera* (Hübner). *Crop Prot* **68**:72-78 (2015).
- 7 49. Efrom C F S, Redaelli L R, Meirelles R N and Ourique C B, Side-effects of pesticides
8 used in the organic system of production on *Apis mellifera* Linnaeus, 1758. *Braz Arch Biol*
9 *Technol* **55**:47-53 (2012).
- 10 50. Asaduzzaman M, Shim J K, Lee S and Lee K Y, Azadirachtin ingestion is lethal and
11 inhibits expression of ferritin and thioredoxin peroxidase genes of the sweetpotato whitefly
12 *Bemisia tabaci*. *J Asia Pac Entomol* **19**:1-4 (2016).
- 13 51. Tofel K H, Nukenine E N, Stähler M and Adler C, Degradation of azadirachtin A on
14 treated maize and cowpea and the persistence of *Azadirachta indica* seed oil on
15 *Callosobruchus maculatus* and *Sitophilus zeamais*. *J Stored Prod Res* **69**:207-212 (2016).
- 16 52. Lavialle-Defaix C, Moignot B, Legros C and Lapied B, How does calcium-dependent
17 intracellular regulation of voltage-dependent sodium current increase the sensitivity to the
18 oxadiazine insecticide indoxacarb metabolite decarbomethoxylated JW062 (DCJW) in insect
19 pacemaker neurons? *J Pharmacol Exp Ther* **333**:264-272 (2010).
- 20 53. Besard L, Mommaerts V, Abdu-Alla G and Smagghe G, Lethal and sublethal side-effect
21 assessment supports a more benign profile of spinetoram compared with spinosad in the
22 bumblebee *Bombus terrestris*. *Pest Manag Sci* **67**:541-547 (2011).
- 23 54. Shao X, Lai D, Zhang L and Xu H, Induction of autophagy and apoptosis via
24 PI3K/AKT/TOR pathways by azadirachtin A in *Spodoptera litura* Cells. *Sci Rep* **6**:35482
25 (2016).

- 1 55. Xu L, Li S, Ran X, Liu C, Lin R and Wang J, Apoptotic activity and gene responses in
2 *Drosophila melanogaster* S2 cells, induced by azadirachtin A. *Pest ManagSci* **72**:1710-
3 1717(2016).
- 4 56. Wang Z, Cheng X, Meng Q, Wang P, Shu B, Hu Q, Hu M and Zhong G, Azadirachtin-
5 induced apoptosis involves lysosomal membrane permeabilization and cathepsin L release in
6 *Spodoptera frugiperda* Sf9 cells. *Int J Biochem Cell Biol* **64**:126-135 (2015).
- 7 57. Lee K P, Dietary protein: carbohydrate balance is a critical modulator of lifespan and
8 reproduction in *Drosophila melanogaster*: a test using a chemically defined diet. *J Insect*
9 *Physiol* **75**:12-19 (2015).
- 10 58. Friedrich J, Sorge S, Bujupi F, Eichenlaub M P, Schulz N G, Wittbrodt J and Lohmann I,
11 Hox function is required for the development and maintenance of the *Drosophila* feeding
12 motor unit. *Cell Rep* **14**:850-860 (2016).
- 13 59. Piri F, Sahragard A and Ghadamyari M, Sublethal effects of spinosad on some
14 biochemical and biological parameters of *Glyphodes pyloalis* Walker. *Plant Prot Sci* **50**:135-
15 144(2014).
- 16 60. Thompson J N and Pellmyr O, Evolution of oviposition behavior and host preference in
17 Lepidoptera. *Annu Rev Entomol* **36**:65-89 (1991).
- 18 61. Sieber M H and Spradling A C, Steroid signaling establishes a female metabolic state and
19 regulates SREBP to control oocyte lipid accumulation. *Curr Biol* **25**:993-1004 (2015).
- 20 62. Moghadam N N, Holmstrup M, Manenti T, Mouridsen M B, Pertoldi C and Loeschke V,
21 The role of storage lipids in the relation between fecundity, locomotor activity, and lifespan of
22 *Drosophila melanogaster* longevity-selected and control lines. *PloS One* **10**:e0130334 (2015).
- 23 63. Zheng H, Yang X and Xi Y, Fat body remodeling and homeostasis control in
24 *Drosophila*. *Life Sci* **167**:22-31 (2016).

- 1 64. Laurinyecz B, Péter M, Vedelek V, Kovács A L, Juhász G, Maróy P, Vigh L, Balogh G
2 and Sinka R, Reduced expression of CDP-DAG synthase changes lipid composition and leads
3 to male sterility in *Drosophila*. *Open Biol*, **6**:150-169 (2016).
- 4 65. Deady L D and Sun J, A follicle rupture assay reveals an essential role for follicular
5 adrenergic signaling in *Drosophila* ovulation. *PLoS Genet* **11**:e1005604 (2015).
- 6 66. Swevers L, Raikhel A S, Sappington T W, Shirk P and Iatrou K, Vitellogenesis and post-
7 vitellogenic maturation of the insect ovarian follicle, in *Comprehensive Molecular Insect*
8 *Science*, ed. by Gilbert L I, Iatrou K and Gill S S, Elsevier Oxford,UK, pp. 87-156 (2005).
- 9 67. Wu Z, Guo W, Xie Y and Zhou S, Juvenile hormone activates the transcription of cell-
10 division-cycle 6 (Cdc6) for polyploidy-dependent insect vitellogenesis and oogenesis. *J Biol*
11 *Chem* **291**:5418-5427 (2016).
- 12 68. Romani P, Gargiulo G and Cavaliere V, The ecdysone receptor signalling regulates
13 microvilli formation in follicular epithelial cells. *Cell Mol Life Sci* **73**:409-425 (2016).
- 14 69. Shingleton A W, Physiology: Female flies have the guts for reproduction. *Curr*
15 *Biol* **25**:R716-R718 (2015).
- 16 70. Toivonen J M and Partridge L, Endocrine regulation of aging and reproduction in
17 *Drosophila*. *Mol Cell Endocrinol* **299**:39-50 (2009).
- 18 71. Rauschenbach I Y, Karpova E K, Burdina E V, Adonyeva N V, Bykov R A, Ilinsky Y Y,
19 Menshanov N E and Gruntenko N E, Insulin-like peptide DILP6 regulates juvenile hormone
20 and dopamine metabolism in *Drosophila* females. *Gen Comp Endocrinol* **243**:1-9 (2017).
- 21 72. Nässel D R and Broeck J V, Insulin/IGF signaling in *Drosophila* and other insects: factors
22 that regulate production, release and post-release action of the insulin-like peptides. *Cell Mol*
23 *Life Sci* **73**:271-290 (2016).
- 24
25

1 **Table 1.** Toxicity of Azadirachtin tested by topical application, at different doses, on the day
2 of *Drosophila melanogaster* adult emergence: lethal doses and their fiducial limits.

3

	Doses (μg)	Fiducial limits
LD 90	4.85	[1.47– 15.98]
LD 50	0.63	[0.44 – 0.91]
LD 25	0.23	[0.13 – 0.38]
Hill slope	1.08 [0.54–1.61]	
R square	0.96	

4


1 **Table 2.** Toxicity of Azadirachtin tested by topical application, at different doses, on the day
2 of *Drosophila melanogaster* adult emergence: corrected mortality (%) from 24 to 96 hours
3 (mean \pm SD; n = 3 replicates of 300 insects). Means followed by the same uppercase letter do
4 not differ significantly according to Tukey's HSD test at level $p \leq 0.05$.

5

Times (h) Doses (μg)	24	48	72	96
0.1	13.92 \pm 1.43 A	15.96 \pm 2.19 A	14.72 \pm 2.24 A	14.65 \pm 0.97 A
0.2	26.16 \pm 3.58 B	23.67 \pm 1.13 B	28.12 \pm 1.22 B	27.77 \pm 5.84 B
0.4	34.84 \pm 0.44 C	33.85 \pm 4.61 C	35.02 \pm 5.92 C	34.89 \pm 4.57 C
0.6	43.16 \pm 0.74 D	42.46 \pm 2.49 D	43.42 \pm 2.47 D	43.87 \pm 6.90 D
1.2	71.80 \pm 3.11 E	69.88 \pm 1.13 E	70.45 \pm 6.20 E	69.88 \pm 1.16 E

6


7



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

Figure 1. Toxicity of Azadirachtin tested by topical application, at different doses, on the day of *Drosophila melanogaster* adult emergence: corrected mortality (%) at 24 hours (mean \pm SD; n = 3 replicates of 300 insects). Control mortality: 5.48 ± 0.53 %. Different letters indicate a significant difference between control and treated series according to Tukey's HSD test.


1
2
3
4



5
6
7
8
9

10 **Figure 2.** Effect of Azadirachtin tested by topical application (0.63 μg), on the day of
11 *Drosophila melanogaster* adult emergence, on the number of progeny (eggs, larvae L3,
12 pupae, adults; M_C: male control; F_C: female control; M_T: male treated; F_T: female treated)
13 (mean \pm SD; n = 8 by couple). Different letters indicate a significant difference between the
14 control and treated series of the same stage of development at level $p \leq 0.05$.

15




1

2

3 **Figure 3.** Effects of Azadirachtin applied by topical application (0.63 μg) on day of
 4 *Drosophila melanogaster* adult emergence on male fecundity parameters (mean \pm SD). **A:**
 5 Number of cysts per testis 48h after emergence. **B:** Percentage of spermatocyte nuclei in
 6 abnormal position. Numbers within each bar indicate the number of repetitions;* above bars
 7 indicate significant difference at $p \leq 0.05$;** above bars indicate significant differences at $p \leq$
 8 0.01.

1



2

3


4 **Figure 4.** Effects of Azadirachtin applied by topical application (0.63 μg) on day of
5 *Drosophila melanogaster* adult emergence on female fecundity parameters (mean \pm SD). **A:**
6 Number of oocytes per ovary 48h after emergence. **B:** Volume of the basal oocyte 48h after
7 emergence. Numbers within bar indicate the number of repetitions; ** above bars indicate
8 significant differences $p \leq 0.01$. *** Above bars indicate significant differences $p \leq 0.001$.

9

1 **Supplementary figure 1**

2

3



4

5

6 Effect of Azadirachtin tested by topical application (LD₂₅: 0.23 μg and LD₅₀: 0.63 μg), on the
 7 day of *Drosophila melanogaster* adult emergence, on the number of progeny (eggs, larvae L3,
 8 pupae, adults; M_C: male control; F_C: female control; M₂₅ and M₅₀: male treated at LD₂₅ and
 9 LD₅₀; F₂₅ and F₅₀: female treated at LD₂₅ and LD₅₀) (mean ± SD; n = 8 by couple). Different
 10 letters indicate a significant difference between control and treated series at the same stage of
 11 development at level p ≤ 0.05.

12