



**HAL**  
open science

# ON THE RADIUS OF SPATIAL ANALYTICITY FOR THE INVISCID BOUSSINESQ EQUATIONS

Feng Cheng, Chao-Jiang Xu

► **To cite this version:**

Feng Cheng, Chao-Jiang Xu. ON THE RADIUS OF SPATIAL ANALYTICITY FOR THE INVISCID BOUSSINESQ EQUATIONS. 2019. hal-02373432

**HAL Id: hal-02373432**

**<https://hal.science/hal-02373432>**

Preprint submitted on 21 Nov 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ON THE RADIUS OF SPATIAL ANALYTICITY FOR THE INVISCID BOUSSINESQ EQUATIONS

FENG CHENG, CHAO-JIANG XU

ABSTRACT. In this paper, we study the problem of analyticity of smooth solutions of the inviscid Boussinesq equations. If the initial datum is real-analytic, the solution remains real-analytic on the existence interval. By an inductive method we can obtain lower bounds on the radius of spatial analyticity of the smooth solution.

## 1. INTRODUCTION

In this paper, we consider the following multi-dimensional inviscid Boussinesq equation on the torus  $\mathbb{T}^d$ ,

$$\begin{cases} \partial_t u + (u \cdot \nabla)u + \nabla p = \theta e_d, \\ \partial_t \theta + (u \cdot \nabla)\theta = 0, \\ \operatorname{div} u = 0, \\ u(x, 0) = u_0(x), \theta(x, 0) = \theta_0(x), \end{cases} \quad (1.1)$$

with  $\operatorname{div} u_0 = 0$ . Here,  $u = (u_1, \dots, u_d)$  is the velocity field,  $p$  the scalar pressure, and  $\theta$  the scalar density.  $e_d$  denotes the vertical unit vector  $(0, \dots, 0, 1)$ . The Boussinesq systems play an important role in geophysical fluids such as atmospheric fronts and oceanic circulation (see, e.g., [15, 22, 24]). Moreover, the Boussinesq systems are important for the study of the Rayleigh-Benard convection, see [10, 12].

Besides the physical importance, the inviscid Boussinesq equations can also be viewed as simplified model compared with the Euler equation. In the case  $d = 2$ , the 2D inviscid Boussinesq equations share some key features with the 3D Euler equations such as the vortex stretching mechanism. It was also pointed out in [23] that the 2D inviscid Boussinesq equations are identical to the Euler equations for the 3D axisymmetric swirling flows outside the symmetric axis.

The inviscid Boussinesq equations have been studied by many authors through the years, for instance [6, 7, 13, 15, 21, 26, 28, 30]. Specially, Chae and Nam [7] studied local existence and uniqueness of the inviscid Boussinesq equation and some blow-up criterion in the Sobolev space, Yuan [30] and Liu et al. [21] in the Besov space, Chae and Kim [6] and Cui et al. [11] in the Hölder spaces, Xiang and Yan [29] in the Triebel-Lizorkin-Lorentz spaces. It was remarked that the global regularity for the inviscid Boussinesq equations even in two dimensions is a challenging open problem in mathematical fluid mechanics.

---

2010 *Mathematics Subject Classification.* 35Q35, 76B03, 76W05.

*Key words and phrases.* Boussinesq equations, Analyticity, Radius of analyticity.

In this paper, we are concerned with the analyticity of smooth solutions of the inviscid Boussinesq equations (1.1). The analyticity of the solution for Euler equations in the space variables, for analytic initial data is an important issue, studied in [1, 2, 3, 4, 16, 17, 18]. In particular, Kukavica and Vicol [16] studied the analyticity of solutions for the Euler equations and obtained that the radius of analyticity  $\tau(t)$  of any analytic solution  $u(t, x)$  has a lower bound

$$\tau(t) \geq C(1+t)^{-2} \exp\left(-C_0 \int_0^t \|\nabla u(s, \cdot)\|_{L^\infty} ds\right) \quad (1.2)$$

for a constant  $C_0 > 0$  depending on the dimension and  $C > 0$  depending on the norm of the initial datum in some finite order Sobolev space. The same authors in [17] obtained a better lower bound for  $\tau(t)$  for the Euler equations in a half space replacing  $(1+t)^{-2}$  by  $(1+t)^{-1}$  in (1.2). In [9], we have investigated the Gevrey analyticity of the smooth solution for the ideal MHD equations following the method of [16]. The approach used in [16, 17, 18, 9] relies on the energy method in infinite order Gevrey-Sobolev spaces. Recently, Capiello and Nicola [5] developed a new inductive method to simplify the proof of [16, 17]. In this paper, we shall apply this inductive method to study the analyticity of smooth solution for the inviscid Boussinesq equations. The main additional difficulty arise from the estimate of the weak coupling term  $u \cdot \nabla \theta$ .

The paper is organized as follows. In Section 2, we will give some notations and state our main results. In Section 3, we first recall some known results and then give some lemmas which are needed to prove the main Theorem. In Section 4, we finish the proof of Theorem 2.1.

## 2. NOTATIONS AND MAIN THEOREM

In this section we will give some notations and function spaces which will be used throughout the following arguments. Throughout the paper,  $C$  denotes a generic constant which may vary from line to line. Since we work on the torus  $\mathbb{T}^d$  throughout the paper, we shall write the function space  $L^2$  or  $H^k$  to represent the functions that are square integrable or square integrable up to  $k$ -th derivative without mentioning the domain  $\mathbb{T}^d$ .

Let  $v = (v_1, \dots, v_d)$  be a vector function, we say that  $v \in L^2$  which means  $v_i \in L^2$  for each  $1 \leq i \leq d$ . We denote the  $L^2$  norm of  $v$  by  $\|v\|_{L^2} = \sqrt{\sum_{1 \leq i \leq d} \|v_i\|_{L^2}^2}$ . Let  $\rho$  be a scalar function, we say the pair  $(v, \rho) \in L^2$  if  $v, \rho \in L^2$ . We denote the  $L^2$  norm of the pair  $(v, \rho)$  to be

$$\|(v, \rho)\|_{L^2} = \sqrt{\|v\|_{L^2}^2 + \|\rho\|_{L^2}^2}.$$

Denote  $\langle \cdot, \cdot \rangle$  to be the inner product in  $L^2$  either for vector function or scalar function.

In [20], it is stated that a smooth function  $f$  is uniformly analytic in  $\mathbb{T}^d$  if there exist  $M, \tau > 0$  such that

$$\|\partial^\alpha f\|_{L^\infty} \leq M \frac{|\alpha|!}{\tau^{|\alpha|}}, \quad (2.1)$$

for all multi-indices  $\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{N}_0^d$ , where  $|\alpha| = \alpha_1 + \dots + \alpha_d$ . The supremum of the constant  $\tau > 0$  in (2.1) is called the radius of the analyticity of  $f$ . Notice that we can also replace the  $L^\infty$  norm with a Sobolev norm  $H^k$ ,  $k \geq 0$ .

Let  $n \geq 0$  be an integer and  $\alpha, \beta \in \mathbb{N}^d$  be multi-indices, then the sequence

$$M_n = \frac{n!}{(n+1)^2}$$

satisfies

$$\sum_{\beta < \alpha} \binom{\alpha}{\beta} M_{|\alpha-\beta|} M_{|\beta|+1} \leq C |\alpha| M_{|\alpha|}, \quad (2.2)$$

for all multi-index  $\alpha, \beta \in \mathbb{N}^d$  and some universal constant  $C$ , for proof please refer to [1].

With these notations, we can state our main results.

**Theorem 2.1.** *Let  $k > \frac{d}{2} + 1$ ,  $(u_0, \theta_0)$  be analytic in  $\mathbb{T}^d$ , satisfying  $\operatorname{div} u_0 = 0$  and*

$$\|\partial^\alpha (u_0, \theta_0)\|_{H^k} \leq BA^{|\alpha|-1} |\alpha|! / (|\alpha| + 1)^2, \quad \alpha \in \mathbb{N}^d, \quad (2.3)$$

for some  $B \geq \frac{9}{4} \|(u_0, \theta_0)\|_{H^{2k+1}}$  and  $A \geq 1$ . Let  $(u(t, x), \theta(t, x))$  be the corresponding  $H^k$  maximal solution of the inviscid Boussinesq equations (1.1), with the initial datum  $(u_0, \theta_0)$ . Then  $(u(t, x), \theta(t, x))$  is analytic, and there exists constants  $C_0, C_1 > 0$ , depending only on  $k$  and  $d$ , such that the radius of analyticity satisfying

$$\tau(t) \geq \frac{1}{A(1 + C_1 B t)} \exp\left(-C_0 \int_0^t (1 + \|\nabla u(s, \cdot)\|_{L^\infty} + \|\nabla \theta(s, \cdot)\|_{L^\infty}) ds\right). \quad (2.4)$$

**Remark 2.1.** In the case  $\theta = 0$ , Theorem 2.1 recovers the result of Kukavica and Vicol [16] and Capiello and Nicola [5] for the incompressible Euler equation.

**Remark 2.2.** When the dimension  $d = 2$ , the blow-up criterion proved by Chae and Nam in [7] stated that the solution remains smooth up to  $T$  as long as  $\int_0^T \|\nabla \theta(\cdot, s)\|_{L^\infty} ds < \infty$ . So, it will be very interesting if the quantity  $\int_0^t (1 + \|\nabla u(\cdot, s)\|_{L^\infty} + \|\nabla \theta(\cdot, s)\|_{L^\infty}) ds$  in the lower bound of the radius of analytic solution in (2.4) can be replaced by  $\int_0^t \|\nabla \theta(\cdot, s)\|_{L^\infty} ds$ .

### 3. THE ESTIMATE OF THE SOBOLEV NORM

In order to prove the main Theorem 2.1, we recall the following results about the local existence and uniqueness of  $H^k$ -solution of the inviscid Boussinesq equations (1.1) which is a proposition in [27] for  $d = 3$ .

**Theorem 3.1** (Wang-Xie, Proposition 1.2 of [27]). *If  $(u_0(x), \theta_0(x)) \in H^3(\Omega)$  and  $u_0(x)$  satisfies the divergence-free condition, then there exists  $T_2 > 0$  such that the inviscid problem (1.1) admits a unique solution*

$$(u, \theta) \in C(0, T_2; H^3(\Omega)) \cap C^1(0, T_2; H^2(\Omega)).$$

The domain considered in [27] is a bounded domain with smooth boundary conditions and this case can be naturally extended to periodic domain with periodic boundary conditions. The proof is due to the argument in [8] and [19]. Since it is standard, we omit the details here. When the dimension  $d = 2$ , Chae and Nam [7] also proved the local existence and blow-up criterion.

In order to prove the main Theorem, we will need the following Lemma.

**Lemma 3.2.** *Let  $d = 2, 3$ , and  $k > \frac{d}{2} + 1$  be fixed. Let  $(u, \theta) \in C(0, T_1; H^k)$  be the corresponding maximal  $H^k$ -solution of (1.1) with initial data  $(u_0, \theta_0) \in H^k$*

and  $u_0$  satisfies the divergence-free condition and periodic boundary condition, then  $\forall 0 \leq t < T_1$ ,

$$\begin{aligned} \|(u(t, \cdot), \theta(t, \cdot))\|_{H^k} &\leq \|(u_0, \theta_0)\|_{H^k} \\ &\times \exp\left(C_0 \int_0^t (1 + \|\nabla u(s, \cdot)\|_{L^\infty} + \|\nabla \theta(s, \cdot)\|_{L^\infty}) ds\right), \end{aligned} \quad (3.1)$$

where the constant  $C_0$  depending on the dimension and  $k$ .

*Proof.* Since the initial data  $(u_0, \theta_0) \in H^k$  satisfies  $\nabla \cdot u_0 = 0$  and the periodic boundary conditions, the local existence of the  $H^k$ -solution  $(u, \theta)$  is already known. We here only need to show the  $H^k$  energy estimate (3.1).

Let  $\alpha \in \mathbb{N}^d$  satisfies  $0 \leq |\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_d \leq k$ . We first apply the  $\partial^\alpha$  on both sides of the first equation of (1.1) and then take the  $L^2$ -inner product with  $\partial^\alpha u$  with both sides, which gives

$$\frac{1}{2} \frac{d}{dt} \|\partial^\alpha u\|_{L^2}^2 + \langle \partial^\alpha (u \cdot \nabla u), \partial^\alpha u \rangle = \langle \partial^\alpha (\theta e_d), \partial^\alpha u \rangle, \quad (3.2)$$

where the pressure term  $\langle \nabla \partial^\alpha p, \partial^\alpha u \rangle = 0$  is due to the fact  $u$  is divergence free and the domain considered here is a periodic domain.

We then apply the  $\partial^\alpha$  on both sides of the second equation of (1.1) and take the  $L^2$  inner product with  $\partial^\alpha \theta$  on both sides, which gives

$$\frac{1}{2} \frac{d}{dt} \|\partial^\alpha \theta\|_{L^2}^2 + \langle \partial^\alpha (u \cdot \nabla \theta), \partial^\alpha \theta \rangle = 0. \quad (3.3)$$

Now adding (3.2) with (3.3) and taking summation over  $0 \leq |\alpha| \leq k$ , we can obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|(u, \theta)\|_{H^k}^2 &= \sum_{0 \leq |\alpha| \leq k} \langle \partial^\alpha (\theta e_d), \partial^\alpha u \rangle \\ &- \sum_{0 \leq |\alpha| \leq k} \langle \partial^\alpha (u \cdot \nabla u), \partial^\alpha u \rangle - \sum_{0 \leq |\alpha| \leq k} \langle \partial^\alpha (u \cdot \nabla \theta), \partial^\alpha \theta \rangle. \end{aligned} \quad (3.4)$$

The Hölder inequality implies that

$$\left| \sum_{0 \leq |\alpha| \leq k} \langle \partial^\alpha (\theta e_d), \partial^\alpha u \rangle \right| \leq \|\theta\|_{H^k} \|u\|_{H^k}. \quad (3.5)$$

Notice that  $u$  is divergence free, by use of the Sobolev inequality which can be found in [23] we can obtain

$$\left| \sum_{0 \leq |\alpha| \leq k} \langle \partial^\alpha (u \cdot \nabla u), \partial^\alpha u \rangle \right| \leq C \|\nabla u\|_{L^\infty} \|u\|_{H^k}^2, \quad (3.6)$$

where the constant  $C$  depends on  $k$  and the dimension  $d$ . In the same way, we can obtain

$$\left| \sum_{0 \leq |\alpha| \leq k} \langle \partial^\alpha (u \cdot \nabla \theta), \partial^\alpha \theta \rangle \right| \leq C (\|\nabla u\|_{L^\infty} \|u\|_{H^k} + \|\nabla \theta\|_{L^\infty} \|u\|_{H^k}) \|\theta\|_{H^k}, \quad (3.7)$$

where the constant  $C$  also depends on  $k$  and the space dimension  $d$ . Substituting (3.5), (3.6) and (3.7) into (3.4), we obtain

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|(u, \theta)\|_{H^k}^2 &\leq C (\|\nabla u\|_{L^\infty} + \|\nabla \theta\|_{L^\infty}) \|(u, \theta)\|_{H^k}^2 + C \|(u, \theta)\|_{H^k}^2 \\ &\leq C (1 + \|\nabla u\|_{L^\infty} + \|\nabla \theta\|_{L^\infty}) \|(u, \theta)\|_{H^k}^2, \end{aligned} \quad (3.8)$$

where  $C$  is some constant depending on  $k, d$ . By the Gronwall inequality, (3.1) is then proved.  $\square$

Lemma 3.2 tells us that the solution for the inviscid Boussinesq equation has the same Sobolev regularity as the initial data. In the following Lemma we will show that if the initial data  $(u_0, \theta_0) \in H^k$  for arbitrary  $k \geq 3$ , there exists an interval  $[0, T]$  uniformly with respect to  $k$  such that the unique smooth solution  $(u, \theta) \in L^\infty([0, T]; H^k)$ .

**Lemma 3.3.** *Let  $(u, \theta)$  be the  $H^3$ -solution of the inviscid Boussinesq equation (1.1) on the time interval  $[0, T]$ , with initial data  $(u_0, \theta_0) \in H^3$ . Then for all  $k \geq 3$ , if  $(u_0, \theta_0) \in H^k$ , the corresponding solution  $(u, \theta)$  satisfies*

$$(u, \theta) \in L^\infty([0, T], H^k) \quad \text{for all } k \geq 3.$$

*Proof.* We claim that for every  $3 \leq m \leq k$  there exist a constant  $C_m$  such that

$$\sup_{0 \leq s \leq T} \|(u(\cdot, s), \theta(\cdot, s))\|_{H^m} \leq C_m.$$

For  $m = 3$  the statement follows from our assumption. Take  $4 \leq m \leq k$  and suppose that the statement is true for  $m - 1$ , i. e.

$$\sup_{0 \leq s \leq T} \|(u(\cdot, s), \theta(\cdot, s))\|_{H^{m-1}} \leq C_{m-1}.$$

We take the  $H^m$  inner product of the first equation of (1.1) with  $u$  and take the  $H^m$  inner product of the second equation of (1.1) with  $\theta$ , which gives

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|(u, \theta)\|_{H^m}^2 &= \sum_{0 \leq |\alpha| \leq m} \langle \partial^\alpha (\theta e_d), \partial^\alpha u \rangle \\ &\quad - \sum_{0 \leq |\alpha| \leq m} \langle \partial^\alpha (u \cdot \nabla u), \partial^\alpha u \rangle - \sum_{0 \leq |\alpha| \leq m} \langle \partial^\alpha (u \cdot \nabla \theta), \partial^\alpha \theta \rangle. \end{aligned}$$

For  $4 \leq m \leq k$ , by (3.8) we obtain

$$\frac{d}{dt} \|(u, \theta)\|_{H^m} \leq C(1 + \|\nabla u\|_{L^\infty} + \|\nabla \theta\|_{L^\infty}) \|(u, \theta)\|_{H^m},$$

where  $C$  is a constant depending on  $m, d$ . Then the Gronwall inequality yields

$$\begin{aligned} \|(u(\cdot, t), \theta(\cdot, t))\|_{H^m} &\leq \|(u_0, \theta_0)\|_{H^m} \\ &\quad \times \exp \left( C \int_0^t (1 + \|\nabla u(\cdot, s)\|_{L^\infty} + \|\nabla \theta(\cdot, s)\|_{L^\infty}) ds \right). \end{aligned} \quad (3.9)$$

By Sobolev embedding inequality, we have

$$\|\nabla u\|_{L^\infty} \leq C' \|u\|_{H^3}, \quad \|\nabla \theta\|_{L^\infty} \leq C' \|\theta\|_{H^3},$$

for some constant  $C'$ . Then from (3.9) and the assumption, we have

$$\begin{aligned} \|(u(\cdot, t), \theta(\cdot, t))\|_{H^m} &\leq \|(u_0, \theta_0)\|_{H^m} \\ &\quad \times \exp \left( C \int_0^t (1 + C' \|u(\cdot, s)\|_{H^3} + C' \|\theta(\cdot, s)\|_{H^3}) ds \right) \\ &\leq \|(u_0, \theta_0)\|_{H^m} \exp \left( C(1 + 2C'C_3)t \right). \end{aligned}$$

So it easily follows that

$$\sup_{0 \leq s \leq T} \|(u(\cdot, s), \theta(\cdot, s))\|_{H^m} \leq C_m,$$

which proves the Lemma by induction.  $\square$

**Remark 3.1.** In Lemma 3.3, the uniform lifespan  $[0, T]$  of the Sobolev solution is independent of the Sobolev order  $k$  which allows us to take  $k \rightarrow \infty$ . In other words, if the initial datum  $(u_0, \theta_0)$  is  $C^\infty$ , then solution  $(u(t, x), \theta(t, x))$  is also  $C^\infty$  for almost every  $t \in [0, T]$ .

#### 4. PROOF OF THEOREM 2.1

In this Section, we will give the proof of the main theorem.

*Proof of Theorem 2.1.* By Lemma 3.3, we know that the solution  $(u, \theta)$  is smooth, since  $(u_0, \theta_0)$  is. Now we claim that for all  $|\alpha| = N > 2$  we have

$$\begin{aligned} \frac{\|(\partial^\alpha u(t, \cdot), \partial^\alpha \theta(t, \cdot))\|_{H^k}}{M_{|\alpha|}} &\leq 2BA^{N-1}(1 + C_1 Bt)^{N-2} \\ &\times \exp\left(C_0(N-1) \int_0^t (1 + \|\nabla u(s)\|_{L^\infty} + \|\nabla \theta(s)\|_{L^\infty}) ds\right), \end{aligned} \quad (4.1)$$

where  $C_0, C_1$  are positive constants depending only on  $k$  and  $d$ . We set

$$\mathcal{E}_N[(u, \theta)(t)] = \sup_{|\alpha|=N} \frac{\|(\partial^\alpha u(t, \cdot), \theta(t, \cdot))\|_{H^k}}{M_{|\alpha|}}.$$

To prove the claim, we proceed by induction on  $N$ . The result is true for  $N = 2$  by (3.1) with notice that  $k + 2 < 2k + 1$  and  $B \geq \frac{9}{4} \|(u_0, \theta_0)\|_{H^{2k+1}}, A \geq 1$ . Hence, let  $N \geq 3$  and assume (4.1) holds for multi-indices  $\alpha$  of length  $2 \leq |\alpha| \leq N - 1$  and prove it for  $|\alpha| = N$ .

For  $|\alpha| = N, |\gamma| \leq k$ , we first apply  $\partial^{\alpha+\gamma}$  on both sides of the first and the second equation of (1.1) and then take the  $L^2$ -inner product with  $\partial^{\alpha+\gamma}u$  and  $\partial^{\alpha+\gamma}\theta$  respectively, which gives

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\partial^{\alpha+\gamma}(u, \theta)\|_{L^2}^2 + \langle \partial^{\alpha+\gamma}(u \cdot \nabla u), \partial^{\alpha+\gamma}u \rangle + \langle \partial^{\alpha+\gamma}(u \cdot \nabla \theta), \partial^{\alpha+\gamma}\theta \rangle \\ = \langle \partial^{\alpha+\gamma}(\theta e_d), \partial^{\alpha+\gamma}u \rangle. \end{aligned} \quad (4.2)$$

Denote  $Lw = L_u w := u \cdot \nabla w$  and

$$\begin{aligned} \mathcal{I}_1 &= [\partial^{\alpha+\gamma}, L]u := \partial^{\alpha+\gamma}(u \cdot \nabla u) - u \cdot \nabla \partial^{\alpha+\gamma}u, \\ \mathcal{I}_2 &= [\partial^{\alpha+\gamma}, L]\theta := \partial^{\alpha+\gamma}(u \cdot \nabla \theta) - u \cdot \nabla \partial^{\alpha+\gamma}\theta, \\ \mathcal{I}_3 &= (\partial^{\alpha+\gamma}\theta)e_d. \end{aligned}$$

Taking summation with  $0 \leq |\gamma| \leq k$  in (4.2), we have by the Cauchy-Schwartz inequality

$$\frac{d}{dt} \|\partial^\alpha(u(t, \cdot), \theta(t, \cdot))\|_{H^k} \leq \sum_{0 \leq |\gamma| \leq k} \|\mathcal{I}_1\|_{L^2} + \sum_{0 \leq |\gamma| \leq k} \|\mathcal{I}_2\|_{L^2} + \|\partial^\alpha(u, \theta)\|_{H^k}, \quad (4.3)$$

where we used the the standard argument in [pp. 47-48, [25]]. It remains to estimate  $\mathcal{I}_1$  and  $\mathcal{I}_2$ . Note that by the Leibniz rule, we can expand the expression of  $\mathcal{I}_1$  as follows

$$\begin{aligned}\mathcal{I}_1 &= \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \partial^\gamma (\partial^{\alpha-\beta} u \cdot \nabla \partial^\beta u) \\ &= \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \sum_{\substack{\delta \leq \gamma, \\ |\beta|+|\delta| < |\alpha|+|\gamma|}} \binom{\gamma}{\delta} \partial^{\alpha-\beta+\gamma-\delta} u \cdot \nabla \partial^{\beta+\delta} u,\end{aligned}$$

where the restriction  $|\beta|+|\delta| < |\alpha|+|\gamma|$  is due to the fact that  $\langle u \cdot \nabla \partial^{\alpha+\gamma} u, \partial^{\alpha+\gamma} u \rangle = 0$  because  $u$  is divergence free. By [5], we have

$$\begin{aligned}\|\mathcal{I}_1\|_{L^2} &\leq \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \sum_{\substack{\delta \leq \gamma, \\ |\beta|+|\delta| < |\alpha|+|\gamma|}} \binom{\gamma}{\delta} \|\partial^{\alpha-\beta+\gamma-\delta} u \cdot \nabla \partial^{\beta+\delta} u\|_{L^2} \\ &\leq C \left\{ NM_N \|\nabla u\|_{L^\infty} \mathcal{E}_N[(u, \theta)] + NM_N B^2 A^{N-1} \right. \\ &\quad \left. \times \exp \left[ C_0(N-1) \int_0^t (1 + \|\nabla u(s)\|_{L^\infty} + \|\nabla \theta(s)\|_{L^\infty}) ds \right] (1 + C_1 B t)^{N-3} \right\}.\end{aligned}$$

We then follow the ideal of [5] to estimate  $\mathcal{I}_2$ . Similarly, we can expand  $\mathcal{I}_2$  as

$$\begin{aligned}\mathcal{I}_2 &= \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \partial^\gamma (\partial^{\alpha-\beta} u \cdot \nabla \partial^\beta \theta) \\ &= \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \sum_{\substack{\delta \leq \gamma, \\ |\beta|+|\delta| < |\alpha|+|\gamma|}} \binom{\gamma}{\delta} \partial^{\alpha-\beta+\gamma-\delta} u \cdot \nabla \partial^{\beta+\delta} \theta.\end{aligned}\tag{4.4}$$

Then we divide the summation of the right of (4.4) into three parts

$$\mathcal{I}_2 = \mathcal{I}_{21} + \mathcal{I}_{22} + \mathcal{I}_{23},$$

where

$$\begin{aligned}\mathcal{I}_{21} &= \sum_{\substack{\beta \leq \alpha \\ 0 \neq |\beta| \leq |\alpha|-2}} \binom{\alpha}{\beta} \sum_{\substack{\delta \leq \gamma, \\ |\beta|+|\delta| < |\alpha|+|\gamma|}} \binom{\gamma}{\delta} \partial^{\alpha-\beta+\gamma-\delta} u \cdot \nabla \partial^{\beta+\delta} \theta, \\ \mathcal{I}_{22} &= \sum_{\beta=\alpha} \binom{\alpha}{\alpha} \sum_{|\delta|=|\gamma|-1} \binom{\gamma}{\delta} \partial^{\gamma-\delta} u \cdot \nabla \partial^{\alpha+\delta} \theta \\ &\quad + \sum_{|\beta|=|\alpha|-1} \binom{\alpha}{\beta} \sum_{\delta=\gamma} \binom{\gamma}{\gamma} \partial^{\alpha-\beta} u \cdot \nabla \partial^{\beta+\gamma} \theta \\ &\quad + \sum_{\beta=0} \binom{\alpha}{\beta} \sum_{\delta=0} \binom{\gamma}{\delta} \partial^{\alpha+\gamma} u \cdot \nabla \theta,\end{aligned}$$



and

$$\begin{aligned}\mathcal{I}_{23} &= \sum_{\beta=\alpha} \binom{\alpha}{\alpha} \sum_{|\delta|\leq|\gamma|-2} \binom{\gamma}{\delta} \partial^{\gamma-\delta} u \cdot \nabla \partial^{\alpha+\delta} \theta \\ &\quad + \sum_{|\beta|=|\alpha|-1} \binom{\alpha}{\beta} \sum_{|\delta|\leq|\gamma|-1} \binom{\gamma}{\gamma} \partial^{\alpha-\beta} u \cdot \nabla \partial^{\beta+\gamma} \theta \\ &\quad + \sum_{\beta=0} \binom{\alpha}{\beta} \sum_{\delta \neq 0} \binom{\gamma}{\delta} \partial^{\alpha+\gamma} u \cdot \nabla \theta.\end{aligned}$$

Estimation of  $\mathcal{I}_{21}$ : With the fact that  $H^k$  is an algebra if  $k > \frac{d}{2} + 1$  and  $|\gamma| \leq k$ , we have

$$\begin{aligned}\|\mathcal{I}_{21}\|_{L^2} &\leq \sum_{\substack{\beta \leq \alpha \\ 0 \neq |\beta| \leq |\alpha|-2}} \binom{\alpha}{\beta} \sum_{\substack{\delta \leq \gamma, \\ |\beta|+|\delta| < |\alpha|+|\gamma|}} \binom{\gamma}{\delta} \|\partial^{\alpha-\beta+\gamma-\delta} u \cdot \nabla \partial^{\beta+\delta} \theta\|_{L^2} \\ &\leq C \sum_{\substack{\beta \leq \alpha \\ 0 \neq |\beta| \leq |\alpha|-2}} \binom{\alpha}{\beta} \sum_{\substack{\delta \leq \gamma, \\ |\beta|+|\delta| < |\alpha|+|\gamma|}} \binom{\gamma}{\delta} \|\partial^{\alpha-\beta} u\|_{H^k} \|\nabla \partial^{\beta} \theta\|_{H^k}.\end{aligned}\tag{4.5}$$

Noting that  $2 \leq |\alpha - \beta| \leq N - 1$  and  $2 \leq |\beta| + 1 \leq N - 1$ , the hypothesis (4.1) for  $2 \leq |\alpha| \leq N - 1$  indicates that

$$\begin{aligned}\|\partial^{\alpha-\beta} u\|_{H^k} &\leq M_{|\alpha-\beta|} 2BA^{|\alpha-\beta|-1} \\ &\quad \times \exp \left[ C_0(|\alpha - \beta| - 1) \int_0^t (1 + \|\nabla u(s)\|_{L^\infty} + \|\nabla \theta(s)\|_{L^\infty}) ds \right] (1 + C_1 Bt)^{|\alpha-\beta|-2}\end{aligned}\tag{4.6}$$

and

$$\begin{aligned}\|\nabla \partial^{\beta} \theta\|_{H^k} &\leq M_{|\beta|+1} 2BA^{|\beta|} \\ &\quad \times \exp \left[ C_0(|\beta|) \int_0^t (1 + \|\nabla u(s)\|_{L^\infty} + \|\nabla \theta(s)\|_{L^\infty}) ds \right] (1 + C_1 Bt)^{|\beta|-1}.\end{aligned}\tag{4.7}$$

Substituting (4.6) and (4.7) into (4.5) and employing (2.2), we obtain

$$\begin{aligned}\|\mathcal{I}_{21}\|_{L^2} &\leq C \sum_{\substack{\beta \leq \alpha \\ 0 \neq |\beta| \leq |\alpha|-2}} \binom{\alpha}{\beta} M_{|\alpha-\beta|} M_{|\beta|+1} B^2 A^{N-1} \\ &\quad \times \exp \left[ C_0(N-1) \int_0^t (1 + \|\nabla u(s)\|_{L^\infty} + \|\nabla \theta(s)\|_{L^\infty}) ds \right] (1 + C_1 Bt)^{N-3} \\ &\leq CNM_N B^2 A^{N-1} \\ &\quad \times \exp \left[ C_0(N-1) \int_0^t (1 + \|\nabla u(s)\|_{L^\infty} + \|\nabla \theta(s)\|_{L^\infty}) ds \right] (1 + C_1 Bt)^{N-3}.\end{aligned}$$

Estimation of  $\mathcal{I}_{22}$ : In a similar way, we rewrite  $\mathcal{I}_{22}$  as

$$\begin{aligned}\mathcal{I}_{22} &= \sum_{\beta=\alpha} \binom{\alpha}{\alpha} \sum_{|\delta|=|\gamma|-1} \binom{\gamma}{\delta} \partial^{\gamma-\delta} u \cdot \nabla \partial^{\alpha+\delta} \theta \\ &\quad + \sum_{|\beta|=|\alpha|-1} \binom{\alpha}{\beta} \sum_{\delta=\gamma} \binom{\gamma}{\gamma} \partial^{\alpha-\beta} u \cdot \nabla \partial^{\beta+\gamma} \theta \\ &\quad + \sum_{\beta=0} \binom{\alpha}{\beta} \sum_{\delta=0} \binom{\gamma}{\delta} \partial^{\alpha+\gamma} u \cdot \nabla \theta \\ &= \mathcal{R}_{21} + \mathcal{R}_{23} + \mathcal{R}_{23}.\end{aligned}$$

For  $\mathcal{R}_{21}$ , we obtain

$$\begin{aligned}\|\mathcal{R}_{21}\|_{L^2} &\leq \sum_{\beta=\alpha} \binom{\alpha}{\alpha} \sum_{|\delta|=|\gamma|-1} \binom{\gamma}{\delta} \|\partial^{\gamma-\delta} u \cdot \nabla \partial^{\alpha+\delta} \theta\|_{L^2} \\ &\leq \sum_{\beta=\alpha} \binom{\alpha}{\alpha} \sum_{|\delta|=|\gamma|-1} \binom{\gamma}{\delta} \|\partial^{\gamma-\delta} u \cdot \nabla \partial^{\alpha+\delta} \theta\|_{L^2} \\ &\leq C \|\nabla u\|_{L^\infty} \|\partial^\alpha \theta\|_{H^k}.\end{aligned}\tag{4.8}$$

for some constant  $C$  depending on  $k$ .

For  $\mathcal{R}_{22}$ , we have the following estimate

$$\begin{aligned}\mathcal{R}_{22} &\leq \sum_{|\beta|=|\alpha|-1} \binom{\alpha}{\beta} \sum_{\delta=\gamma} \binom{\gamma}{\gamma} \|\partial^{\alpha-\beta} u \cdot \nabla \partial^{\beta+\gamma} \theta\|_{L^2} \\ &\leq \sum_{|\beta|=|\alpha|-1} N \sum_{\delta=\gamma} \binom{\gamma}{\gamma} \|\partial^{\alpha-\beta} u \cdot \nabla \partial^{\beta+\gamma} \theta\|_{L^2} \\ &\leq CdNM_N \|\nabla u\|_{L^\infty} \mathcal{E}_N[\theta].\end{aligned}\tag{4.9}$$

Then for  $\mathcal{R}_{23}$ , we have

$$\begin{aligned}\|\mathcal{R}_{23}\|_{L^2} &\leq \sum_{\beta=0} \binom{\alpha}{\beta} \sum_{\delta=0} \binom{\gamma}{\delta} \|\partial^{\alpha+\gamma} u \cdot \nabla \theta\|_{L^2} \\ &\leq \|\nabla \theta\|_{L^\infty} \|\partial^\alpha u\|_{H^k}.\end{aligned}\tag{4.10}$$

Summing up the estimates of (4.8), (4.9) and (4.10), we obtain

$$\|\mathcal{I}_{22}\|_{L^2} \leq C(\|\nabla u\|_{L^\infty} \|\partial^\alpha \theta\|_{H^k} + \|\nabla \theta\|_{L^\infty} \|\partial^\alpha u\|_{H^k} + dNM_N \|\nabla u\|_{L^\infty} \mathcal{E}_N(\theta)).$$

Estimate of  $\mathcal{I}_{23}$ : We divide  $\mathcal{I}_{23}$  as follows

$$\begin{aligned}\mathcal{I}_{23} &= \sum_{\beta=\alpha} \binom{\alpha}{\alpha} \sum_{|\delta|\leq|\gamma|-2} \binom{\gamma}{\delta} \partial^{\gamma-\delta} u \cdot \nabla \partial^{\alpha+\delta} \theta \\ &\quad + \sum_{|\beta|=|\alpha|-1} \binom{\alpha}{\beta} \sum_{|\delta|\leq|\gamma|-1} \binom{\gamma}{\gamma} \partial^{\alpha-\beta+\gamma-\delta} u \cdot \nabla \partial^{\beta+\gamma} \theta \\ &\quad + \sum_{\beta=0} \binom{\alpha}{\beta} \sum_{\delta\neq 0} \binom{\gamma}{\delta} \partial^{\alpha+\gamma-\delta} u \cdot \nabla \partial^\delta \theta \\ &= \mathcal{R}_{31} + \mathcal{R}_{32} + \mathcal{R}_{33}.\end{aligned}$$

For  $\mathcal{R}_{31}$ , we have

$$\begin{aligned}
\|\mathcal{R}_{31}\|_{L^2} &\leq C \sum_{\beta=\alpha} \binom{\alpha}{\beta} \sum_{|\delta|\leq|\gamma|-2} \binom{\gamma}{\delta} \|\partial^{\gamma-\delta} u \cdot \nabla \partial^{\alpha+\delta} \theta\|_{L^2} \\
&\leq C \|\partial^{\gamma-\delta} u\|_{L^\infty} \|\nabla \partial^{\alpha+\delta} \theta\|_{L^2} \\
&\leq C \|\partial^{\gamma-\delta} u\|_{H^k} \|\partial^\alpha \theta\|_{H^{k-1}} \\
&\leq C \|\partial^{\gamma-\delta}(u, \theta)\|_{H^k} \|\partial^\alpha(u, \theta)\|_{H^{k-1}} \\
&\leq C \|(u_0, \theta_0)\|_{H^{2k}} \exp \left[ C_0 \int_0^t (1 + \|\nabla u\|_{L^\infty} + \|\nabla \theta\|_{L^\infty}) ds \right] \sup_{j:\alpha_j \geq 1} \|\partial^{\alpha-e_j}(u, \theta)\|_{H^k} \\
&\leq C \|(u_0, \theta_0)\|_{H^{2k}} \exp \left[ C_0 \int_0^t (1 + \|\nabla u\|_{L^\infty} + \|\nabla \theta\|_{L^\infty}) ds \right] \\
&\quad \times M_{N-1} B A^{N-2} \exp \left[ C_0(N-2) \int_0^t (1 + \|\nabla u\|_{L^\infty} + \|\nabla \theta\|_{L^\infty}) ds \right] (1 + C_1 B t)^{N-3} \\
&\leq C M_{N-1} B^2 A^{N-2} \exp \left[ C_0(N-1) \int_0^t (1 + \|\nabla u\|_{L^\infty} + \|\nabla \theta\|_{L^\infty}) ds \right] (1 + C_1 B t)^{N-3},
\end{aligned} \tag{4.11}$$

where we used the fact  $|\gamma - \delta| + k \leq 2k$  in (4.11), the inductive hypothesis (4.1), and the fact that  $B > \|(u_0, \theta_0)\|_{H^{2k+1}}$ . For  $\mathcal{R}_{32}$ , we have

$$\begin{aligned}
\|\mathcal{R}_{32}\|_{L^2} &\leq \sum_{|\beta|=|\alpha|-1} \binom{\alpha}{\beta} \sum_{|\delta|\leq|\gamma|-1} \binom{\gamma}{\delta} \|\partial^{\alpha-\beta+\gamma-\delta} u \cdot \nabla \partial^{\beta+\gamma} \theta\|_{L^2} \\
&\leq |\alpha| \|\partial^{\alpha-\beta+\gamma-\delta} u\|_{L^\infty} \|\nabla \partial^{\beta+\gamma} \theta\|_{L^2} \\
&\leq C |\alpha| \|\partial^{\gamma-\delta} u\|_{H^k} \|\partial^\beta \theta\|_{H^k} \\
&\leq C |\alpha| \|\partial^{\gamma-\delta}(u, \theta)\|_{H^k} \|\partial^\beta(u, \theta)\|_{H^k} \\
&\leq C N M_{N-1} \|(u_0, \theta_0)\|_{H^{2k}} \exp \left[ C_0 \int_0^t (1 + \|\nabla u\|_{L^\infty} + \|\nabla \theta\|_{L^\infty}) ds \right] \\
&\quad \times B A^{N-2} \exp \left[ C_0(N-2) \int_0^t (1 + \|\nabla u\|_{L^\infty} + \|\nabla \theta\|_{L^\infty}) ds \right] (1 + C_1 B t)^{N-3} \\
&\leq C M_{N-1} B^2 A^{N-2} \exp \left[ C_0(N-1) \int_0^t (1 + \|\nabla u\|_{L^\infty} + \|\nabla \theta\|_{L^\infty}) ds \right] (1 + C_1 B t)^{N-3}.
\end{aligned} \tag{4.12}$$

For  $\mathcal{R}_{33}$ , in a similar way we have

$$\begin{aligned}
\|\mathcal{R}_{33}\|_{L^2} &\leq \sum_{\beta=0} \binom{\alpha}{\beta} \sum_{\delta \neq 0} \binom{\gamma}{\delta} \|\partial^{\alpha+\gamma-\delta} u \cdot \nabla \partial^\delta \theta\|_{L^2} \\
&\leq C \|\partial^\alpha u\|_{H^{k-1}} \|\nabla \partial^\delta \theta\|_{H^k} \\
&\leq C M_{N-1} B^2 A^{N-2} \exp \left[ C_0(N-1) \int_0^t (1 + \|\nabla u\|_{L^\infty} + \|\nabla \theta\|_{L^\infty}) ds \right] (1 + C_1 B t)^{N-3}.
\end{aligned} \tag{4.13}$$

Summing up (4.11), (4.12) and (4.13), we obtain

$$\begin{aligned} \|\mathcal{I}_2\|_{L^2} &\leq CNM_N(\|\nabla u\|_{L^\infty} + \|\nabla\theta\|_{L^\infty})\mathcal{E}_N[(u, \theta)] \\ &\quad + CNM_NB^2A^{N-1} \exp\left(C_0(N-1) \int_0^t [1 + \|\nabla u(s)\|_{L^\infty} + \|\nabla\theta(s)\|_{L^\infty}] ds\right). \end{aligned}$$

Combining the estimates  $\mathcal{I}_1$  and  $\mathcal{I}_2$ , we have from (4.3)

$$\begin{aligned} \frac{d}{dt} \frac{\|\partial^\alpha(u, \theta)(t)\|_{H^k}}{M_{|\alpha|}} &\leq C(N-1)(1 + \|\nabla u(t)\|_{L^\infty} + \|\nabla\theta(t)\|_{L^\infty})\mathcal{E}_N[(u, \theta)(t)] \\ &\quad + CNA^{N-1}B^2 \exp\left(C_0(N-1) \int_0^t (1 + \|\nabla u(s)\|_{L^\infty} + \|\nabla\theta(s)\|_{L^\infty}) ds\right) \\ &\quad \times (1 + C_1Bt)^{N-3}, \end{aligned} \tag{4.14}$$

where we used the Cauchy-Schwartz inequality and the fact  $\frac{N}{N-1} \leq \frac{3}{2}$  if  $N \geq 3$  and the constant  $C > 0$  depending only on the dimension  $d$  and  $k$ .

Now we integrate (4.14) from 0 to  $t$  and take the supremum on  $|\alpha| = N$ . We obtain

$$\begin{aligned} \mathcal{E}_N[(u, \theta)(t)] &\leq \mathcal{E}_N[(u_0, \theta_0)] \\ &\quad + \int_0^t C(N-1)(1 + \|\nabla u(t)\|_{L^\infty} + \|\nabla\theta(t)\|_{L^\infty})\mathcal{E}_N[(u, \theta)(s)] ds \\ &\quad + \int_0^t CNA^{N-1}B^2(1 + C_1Bs)^{N-3} \\ &\quad \times \exp\left(C_0(N-1) \int_0^s (1 + \|\nabla u(\ell)\|_{L^\infty} + \|\nabla\theta(\ell)\|_{L^\infty}) d\ell\right) ds. \end{aligned} \tag{4.15}$$

We can take  $C_0 \geq C$  in (4.15), so that Grownwall inequality [Lemma 2.1 in [5]] gives

$$\begin{aligned} \mathcal{E}_N[(u, \theta)(t)] &\leq \exp\left(C_0(N-1) \int_0^t (1 + \|\nabla u\|_{L^\infty} + \|\nabla\theta\|_{L^\infty}) ds\right) \\ &\quad \times [\mathcal{E}_N[(u, \theta)(0)] + CNB^2A^{N-1} \int_0^t (1 + C_1Bs)^{N-3} ds] \\ &\leq \exp\left(C_0(N-1) \int_0^t (1 + \|\nabla u\|_{L^\infty} + \|\nabla\theta\|_{L^\infty}) ds\right) \\ &\quad \times [\mathcal{E}_N[(u, \theta)(0)] + \frac{CN}{C_1(N-2)}BA^{N-1}(1 + C_1Bt)^{N-2}]. \end{aligned}$$

Note that we have  $\mathcal{E}_N[(u, \theta)(0)] \leq BA^{N-1}$  by the assumption (2.3). If we choose  $C_1 = 3C$ , so that  $\frac{3C}{C_1} = 1$ , since  $A \geq 1, N \geq 3$  we have

$$\begin{aligned} \mathcal{E}_N[(u, \theta)(0)] + \frac{CN}{C_1(N-2)}BA^{N-1}(1 + C_1Bt)^{N-2} \\ \leq [BA^{N-1} + \frac{3C}{C_1}BA^{N-1}(1 + C_1Bt)^{N-2}] \\ \leq 2BA^{N-1}(1 + C_1Bt)^{N-2}, \end{aligned}$$

and we obtain exactly (4.1) for  $|\alpha| = N$ . Then the theorem is proved. □

**Acknowledgements.** The research of the second author is supported partially by “The Fundamental Research Funds for Central Universities of China”.

## REFERENCES

- [1] S. Alinhac and G. Métivier, Propagation de l’analyticité des solutions de systèmes hyperboliques non-linéaires, *Invent. Math.*, **75** (1984), 189-204.
- [2] S. Benachour, Analyticité des solutions périodiques de l’équation d’Euler en trois dimensions, *C. R. Acad. Sci. Paris Sér. A-B* **283** (1976), A107-A110.
- [3] C. Bardos and S. Benachour, Domaine d’analyticité des solutions de l’équation d’Euler dans un ouvert de  $\mathbb{R}^n$ , *Ann. Scuola Norm. Sup. Pisa Cl. Sci.*, (4) **4**(4) (1977), 647-687.
- [4] C. Bardos, S. Benachour and M. Zerner, Analyticité des solutions périodiques de l’équation d’Euler en deux dimensions, *C. R. Acad. Sci. Paris Sér. A-B* **282**(17) (1976), A995-A998.
- [5] M. Cappelletto and F. Nicola, Some remarks on the radius of spatial analyticity for the Euler equations. *Asymptot. Anal.* **91** (2015), no. 2, 103-110.
- [6] D. Chae, S. Kim and H. Nam, Local existence and blow-up criterion of Hölder continuous solutions of the Boussinesq equations. *Nagoya Math. J.*, **155** (1999), 55-80.
- [7] D. Chae and H. Nam, Local existence and blow-up criterion for the Boussinesq equations. *Proc. Roy. Soc. Edinburgh Sect. A*, **127** (1997), no. 5, 935-946.
- [8] D. Chae and O. Imanuvilov, Generic solvability of the axisymmetric 3D Euler equations and the 2D Boussinesq equations. *J. Diff. Eqns.*, **156** (1999), 1-17.
- [9] F. Cheng and C.-J. Xu, On the Gevrey regularity of solutions to the 3D ideal MHD equations, *Discrete Contin. Dyn. Syst.*, **39** (2019), 6485-6506.
- [10] P. Constantin and C. Doering, Infinite Prandtl number convection. *J. Stat. Phys.*, **94** (1999), 159-172.
- [11] X. Cui, C. Dou and Q. Jiu, Local well-posedness and blow up criterion for the inviscid Boussinesq system in Hölder spaces, *J. Partial Differ. Equ.*, **25** (2012), 220-238.
- [12] C. Doering and J. Gibbon, Applied Analysis of the Navier-Stokes Equations, Cambridge Texts in Applied Mathematics, Cambridge University Press, Cambridge, 1995.
- [13] W. E and C. Shu, Small scale structures on Boussinesq convection. *Phys. Fluids*, **6** (1994), 49-58.
- [14] A. Gill, Atmosphere-Ocean Dynamics. *Academic Press*, London, 1982.
- [15] B. Guo, Spectral method for solving two-dimensional Newton-Boussinesq equation. *Acta Math. Appl. Sinica (English Ser.)*, **5** (1989), 208-218.
- [16] I. Kukavica and V. Vicol, On the radius of analyticity of solutions to the three-dimensional Euler equations. *Proc. Amer. Math. Soc.* **137** (2009), 669-677.
- [17] I. Kukavica and V. Vicol, The domain of analyticity of solutions to the three dimensional Euler equations in half space. *Discrete Contin. Dyn. Syst.*, **29** (2011), no. 1, 285-303.
- [18] C. Levermore and M. Oliver, Analyticity of solutions for a generalized Euler equation. *J. Differential Equations*, **133** (1997), 321-339.
- [19] T. T. Li and W. C. Yu. Boundary value problems for quasilinear hyperbolic system. *Duke University Mathematics Series*, Vol. V (Durham, NC: Duke University Mathematics Department, 1985).
- [20] J. Lions and E. Magenes, “Problèmes aux limites non homogènes et applications”, Vol. 3, 1970, Dunod, Paris.
- [21] X. Liu, M. Wang and Z. Zhang, Local well-posedness and blowup criterion of the Boussinesq equations in critical Besov spaces, *J. Math. Fluid Mech.*, **12** (2010), 280-292.
- [22] A. Majda, Introduction to PDEs and Waves for the Atmosphere and Ocean, Courant Lecture Notes in Mathematics, vol. 9, AMS/CIMS, 2003.
- [23] A. Majda and A. Bertozzi, Vorticity and incompressible flow. Cambridge University Press, 2002.
- [24] J. Pedlosky, Geophysical Fluid Dynamics, Springer-Verlag, New York, 1987.
- [25] J. Rauch, Hyperbolic Partial Differential Equations and Geometric Optics, *GSM Series*, **133**, Amer. Math. Soc., Providence, 2012
- [26] Y. Taniuchi, A note on the blow-up criterion for the inviscid 2D Boussinesq equations, *Lecture Notes in Pure and Appl. Math.*, **223**, (2002), 131-140.

- [27] J. Wang and F. Xie, Zero dissipation limit and stability of boundary layers for the heat conductive Boussinesq equations in a bounded domain. *Proc. Roy. Soc. Edinburgh Sect. A*, **145** (2015), no. 3, 611-637.
- [28] X. Xu, Local existence and blow-up criterion of the 2-D compressible Boussinesq equations without dissipation terms, *Discrete Contin. Dyn. Syst.*, **25** (2009), 1333-1347.
- [29] Z. Xiang and W. Yan, On the well-posedness of the Boussinesq equation in the Triebel-Lizorkin-Lorentz spaces, *Abstr. Appl. Anal.*, **2012** (2012), Art. ID 573087, 17 pp.
- [30] B. Yuan, Local existence and continuity conditions of solutions to the Boussinesq equations in Besov spaces, *Acta Math. Sinica (Chin. Ser.)*, **53** (2010), 455-468.

FENG CHENG, HUBEI KEY LABORATORY OF APPLIED MATHEMATICS, FACULTY OF MATHEMATICS AND STATISTICS, HUBEI UNIVERSITY, WUHAN 430062, P.R. CHINA

*E-mail address:* [fengcheng@hubu.edu.cn](mailto:fengcheng@hubu.edu.cn)

CHAO-JIANG XU, DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY OF AERONAUTICS AND ASTRONAUTICS, 211106 NANJING, CHINA, AND, UNIVERSITÉ DE ROUEN, CNRS UMR 6085, LABORATOIRE DE MATHÉMATIQUES, 76801 SAINT-ETIENNE DU ROUVRAY, FRANCE

*E-mail address:* [xuchaojiang@nuaa.edu.cn](mailto:xuchaojiang@nuaa.edu.cn)