



HAL
open science

Involvement of ion channels and transporters in carcinoma angiogenesis and metastasis

Sonia Martial

► **To cite this version:**

Sonia Martial. Involvement of ion channels and transporters in carcinoma angiogenesis and metastasis. *American Journal of Physiology - Cell Physiology*, 2016, 310, pp.710 - 727. 10.1152/ajp-cell.00218.2015 . hal-02375861

HAL Id: hal-02375861

<https://hal.science/hal-02375861v1>

Submitted on 26 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Involvement of ion channels and transporters in carcinoma angiogenesis**
2 **and metastasis**

3

4 **Author:** Sonia MARTIAL

5

6 **Affiliation:**

7 IRCAN, CNRS UMR 7284, Inserm U1081, Université Nice-Sophia Antipolis, NICE, FRANCE

8

9 **Running title:** Ion transport modulates tumor angiogenesis and metastasis

10

11 **Corresponding author:**

12 Dr. Sonia MARTIAL

13 CNRS UMR 7284 – INSERM U 1081 – UNS

14 Institute for Research on Cancer and Aging, Nice

15 33 avenue Valombrose

16 06189 NICE CEDEX 2

17 FRANCE

18 **e-mail :** sonia.martial@unice.fr

19 **Tel :** +33 4 92 03 12 27 / +33 4 92 03 12 28

20 **Fax :** +33 4 92 03 12 35

21

22 **ABSTRACT**

23

24 Angiogenesis is a finely tuned process, which is the result of the equilibrium between pro- and anti-angiogenic
25 factors. In solid tumor angiogenesis, the balance is highly in favor of the production of new, but poorly
26 functional blood vessels, initially intended to provide growing tumors with nutrients and oxygen. Among the
27 numerous proteins involved in tumor development, several types of ion channels are overexpressed in tumor
28 cells, as well as in stromal and endothelial cells. Ion channels thus actively participate in the different hallmarks
29 of cancer, especially in tumor angiogenesis and metastasis. Indeed, from their strategic localization in the
30 plasma membrane, ion channels are key operators of cell signaling, as they sense and respond to environmental
31 changes. This review aims to decipher how ion channels of different families are intricately involved in the
32 fundamental angiogenesis and metastasis hallmarks, which lead from a nascent tumor to systemic
33 dissemination. An overview of the possible use of ion channels as therapeutic targets will also be given,
34 showing that ion channel inhibitors or specific antibodies may provide effective tools, in a near future, in the
35 treatment of carcinomas.

36

37

38 **KEYWORDS:** ion channel, tumor vascularization, cell signaling, ion channel-targeted therapy

39

40

41 **ABBREVIATIONS:** HIF, hypoxia-inducible factor; CRC, colorectal carcinoma; ccRCC, clear cell renal cell
42 carcinoma; CAIX, carbonic anhydrase IX; RVD, regulatory volume decrease; PDAC, pancreatic ductal
43 adenocarcinoma; VEGFR, VEGF receptor; TSP, thrombospondine; MMP, matrix metalloproteinase; SOCE,
44 store-operated calcium entry; CRAC, Ca²⁺-release activated channel; ROCE, receptor-operated calcium entry;
45 IP3, inositol 1,4,5-trisphosphate; EMT, epithelial-mesenchymal transition; TME, tumor microenvironment;
46 STIM, stromal interaction molecule; MAM, mitochondrion-associated ER membrane; NMDAR, N-methyl, D-
47 aspartate Receptor; CML, chronic myeloid leukemia; ER, endoplasmic reticulum; BiP, binding immunoglobulin
48 protein; NSAID, non-steroid anti-inflammatory drug; GPCR, G protein-coupled receptor.

49

50 **I. Introduction**

51 Carcinoma is a solid cancer developing from epithelial cells and is the most widely spread form of human
52 cancer. Among the phenomena leading to the development and progression of this type of cancer, one of the
53 most important is angiogenesis. Indeed, tumor vascularization is essential for a growing tumor to obtain oxygen
54 and nutrients needed for its ongoing growth, *in situ*. Tumor neovascularization is also a means for the tumor to
55 grow away from its primary site towards a remote neo-colonized structure (metastasis).

56 In the last 10 to 15 years, many efforts have been made on the elucidation of which genes might be directly
57 involved or affected during the different steps of tumorigenesis, described by Hanahan and Weinberg as the
58 hallmarks of cancer (95, 96). Multiple studies demonstrated that, among these genes, plasma membrane ion
59 channels and transporters are clearly modulated during carcinogenesis, in different types of carcinomas (9, 20,
60 31, 75, 78, 93, 133, 147, 201, 207). Several different roles have been ascribed to ion channels during
61 carcinogenesis, depending on the step of the tumorization process and on the organ in which it takes place.

62 In this review, we do not intend to make an exhaustive list of channels involved in cancer development. Instead,
63 we choose to focus on the involvement and/or alteration of some ion channels in angiogenesis, from the
64 vascularization of recently formed tumors to the metastasis cascade. Indeed, neo-formation of tumors involves
65 pH changes, which are associated with proton exchange through the plasma membrane (127, 209, 217) and
66 water transport (17, 162). The subsequent tumor growth needs neovascularization, associated with Ca^{2+} , K^+ and
67 Na^+ channels (5, 61, 201). Finally, Na^+ , K^+ , Ca^+ and Cl^- channels are implicated in the transition to metastasis
68 (32, 73, 75).

69 The main ion channels involved in tumor development processes and mentioned in this review are listed in
70 Table 1.

71

72 II. Neangiogenesis

73 A. The fate of new tumors

74 In the early steps of its formation, a tumor depends solely on the host blood vessels (104). However, this initial
75 peripheral vasculature quickly disappears, leaving the tumor and its microenvironment in a hypoxic state (84,
76 97). Folkman (77) was the first to show that, under these conditions, the tumor growth is limited to a few cubic
77 millimeters, unless reorganization of a new blood supply is promoted (neoangiogenesis).

78 The lowered local O_2 pressure (pO_2) observed near cancer cells and the resulting tumor microenvironment
79 acidification activate hypoxia-inducible factors, especially HIF-1 (44, 58, 115). HIF-1 α is stabilized by
80 phosphorylation (172) and dimerizes with the constitutively expressed HIF-1 β to form the transcription factor
81 HIF-1, which in turn stimulates a set of genes involved in tumor development. Conversely, recent reports
82 present HIF-2 as the tumorigenic form of HIF, whereas HIF-1 would be a tumor suppressor (227, 263).

83 Initially, the cancer cells undergo metabolic changes, among which increased glucose consumption and lactate
84 production (the Warburg's effect) lead to acidosis of the cells. In conjunction with the higher need for glucose,
85 several authors report a HIF-dependent expression of the GLUT-1 glucose transporter in breast or colorectal
86 carcinoma cell (CRC) models (25, 84, 132) and a repressed mitochondrial function (175). In clear cell renal cell
87 carcinomas (ccRCC), some isoforms of glucose transporters are equally increased, including the GLUT-5
88 fructose transporter (240). Moreover, in order to prevent acidosis induced by the accumulation of lactate, cancer
89 cells increase the amount and/or activity of several families of acid-extruding plasma membrane proteins. In
90 several carcinoma models, these HIF-1 increased proteins include: NHE1, the Na^+/H^+ exchanger (79, 170, 231),
91 the MCT1 and MCT4 forms of monocarboxylate transporters, dedicated to lactate extrusion (62, 184, 246), the
92 SLC4A4 (NBC) Na^+/HCO_3^- cotransporter (1, 180), involved in the bicarbonate reabsorption which contributes
93 to intracellular pH recovery. Acid-sensing ion channels (ASIC1, 2, 3) are also up-regulated in adenoid cystic
94 carcinomas, following the acidification of the extracellular space (257). Finally, carbonic anhydrases (especially
95 CAIX, the main carbonic anhydrase found in hypoxic tumors (64)), dedicated to microenvironment pH
96 modulation are also upregulated (132, 171, 180). The different ion channels involved at this stage are
97 summarized in Figure 1. Activation of all these transport proteins eventually contribute to a modest elevation of
98 the tumor cell intracellular pH, while the extracellular medium is acidified.

99 Subsequently, the newly constituted tumor will only grow larger if it manages to regenerate a constant blood
100 supply to provide O₂ and nutrients to the tumor cells and to rid the cells of metabolites. Two main families of
101 plasma membrane ion channels have been proposed to be involved in this tumor developmental stage and
102 regulated by HIF. Transient receptor potential (TRP) channels contribute to calcium homeostasis and are
103 upregulated in several cancer cell types, thus playing an important part in Ca²⁺ influx, regulatory volume
104 decrease (RVD) and cell cycle progression (63, 254). A variety of K⁺ channels are also involved in the HIF-
105 induced regulation of proliferation and revascularization, namely the voltage-dependent “ether-à-gogo”
106 potassium channel (EAG1, KCNH1, Kv10.1) (31, 177), and the two-pore domain potassium channels TASK
107 (30) and TREK (243). In pancreatic ductal adenocarcinoma (PDAC), the Ca²⁺-dependent potassium channel of
108 intermediate conductance, IK_{Ca} (K_{Ca}3.1) is up-regulated and contributes to cell proliferation (114).

109

110 B. Angiogenic switch

111 Tumor angiogenesis is the key step for tumor growth, invasion and metastasis. Tumor angiogenesis mostly
112 relies on the same processes as those involved in physiological angiogenesis (27, 45, 176). Particularly,
113 ischemia and hypoxia are in both cases, major initiators of angiogenesis processes (“on” switches). Moreover,
114 tumor angiogenesis and physiological angiogenesis heavily depend on VEGF secretion, as was described
115 elsewhere and as we will describe here for tumor-dependending angiogenesis. The main differences are: 1)
116 Pathological angiogenesis is more dependent upon VEGF than physiological angiogenesis (45); and 2)
117 Physiological angiogenesis events recede when vessel perfusion starts and upon disappearance of the stimuli
118 that gave rise to neoangiogenesis (“off” switch), whereas tumor angiogenesis is a continuous process, fueled by
119 the steady secretion of tumor-induced angiogenesis factors (45).

120 The concept of “angiogenic switch” was formulated in order to characterize angiogenesis as a balance between
121 processes that tend to favor and processes that tend to prevent angiogenesis (94). When tumor cells enter a
122 hypoxic state, activation of HIF leads to the subsequent activation of a multiple array of pro-angiogenic factors;
123 thus, the regular balance of angiogenesis is lost: the angiogenic switch has been turned “ON”. The most studied
124 HIF-regulated pro-angiogenesis factor is probably VEGF, the angiogenesis cytokine that regulates endothelial
125 cell proliferation and blood vessel formation. Both VEGF and its receptors VEGFR1 (FLT-1) and VEGFR2

126 (KDR/FLK1), which are located principally on the surface of endothelial cells, are upregulated by HIF (123,
127 185). Fibroblast Growth Factor (FGF) is also regulated by hypoxia and may act synergistically with VEGF to
128 amplify angiogenesis (125).

129 The “angiogenic switch” is turned “ON”, in several types of cancer cells, including mouse colon carcinoma,
130 mouse melanoma and human neuroblastoma, when several types of ion channels are modulated. P2X₇R, the
131 non-selective cationic pore, has been shown to promote tumor growth and VEGF release (4, 5, 7). TRPC
132 channels are functionally coupled to VEGF or FGF, playing important roles in cancer development steps (76,
133 156). Moreover, in renal cell carcinoma (RCC), the TRPC4 Ca²⁺ channel expression is decreased, which impairs
134 Ca²⁺ metabolism, hence preventing the production of thrombospondin-1 (TSP1), an angiogenesis inhibitor
135 (238). Numerous studies report overexpression of most families of K⁺ channels in human cancers. In particular,
136 the closely related voltage-dependent hERG and EAG1 K⁺ channels are both involved in the processes leading
137 to angiogenesis. Indeed, hERG channels regulate angiogenesis in CRC by increasing the HIF-1 activated VEGF
138 expression and secretion in a β1-integrin-dependent manner and via a PI3K/Akt pathway (46). This regulation
139 has been confirmed in transgenic mice expressing hERG channels and treated with azoxymethane. These mice
140 present increased staining for VEGF in immunohistochemistry (IHC) studies and a higher total number of blood
141 vessels, consistent with neoangiogenesis, as well as colorectal lesions, on the whole showing acceleration of the
142 colorectal cancer phenotype (74). EAG1, but not EAG2 contributes to tumor progression, as demonstrated by
143 xenograft tumor formation resulting from implanted MDA-MB-435S breast cancer cells, EAG1-transfected
144 CHO cells and EAG1- or EAG2-transfected NIH-3T3 cells. Furthermore, EAG1 stimulates VEGF secretion,
145 measured in CHO- or 3T3-transfected cells. Finally, EAG1 increases HIF-1 amount in EAG1-expressing cells,
146 thus demonstrating that EAG1 is involved in the angiogenic switch and the subsequent neoangiogenesis (61).

147 The role of Ca²⁺ channels in conjunction with angiogenic switch is worth to mention. Indeed, it has been
148 demonstrated that VEGF utilizes Ca²⁺ signaling to promote endothelial cell proliferation and subsequent
149 neoangiogenesis, which we largely discuss in the next section. However, it is noteworthy that some Ca²⁺
150 channels, mostly of the TRP family, are upregulated by hypoxia in cancer cells (76). The subsequent increase in
151 intracellular Ca²⁺ concentration is at the basis of cancer cell hallmarks such as cell proliferation or migration
152 (76).

153

154 C. Neovascularization

155 Tumor vasculature is a crucial feature of carcinoma development. New blood vessels develop by sprouting
156 (angiogenesis) or intussusception from pre-existing vessels (144, 199). They can also emerge from the assembly
157 of endothelial precursors derived from the bone marrow (vasculogenesis) (189). Sprouting is a very well studied
158 process, which is involved in tumor blood vessel formation. Angiogenesis is a multistep process that begins
159 when endothelial cell VEGF receptors are stimulated by VEGF (19, 34) to give rise to new blood vessels
160 growing towards the avascular tumor that initially secreted the stimulating cytokine. The different steps of the
161 process are detailed below:

162

163 1. VEGF-induced blood vessel dilatation

164 Most research on tumor angiogenesis has focused on VEGF-A. VEGF-A signaling occurs through two VEGF
165 receptors, VEGFR1 and VEGFR2 (19, 144). Although VEGF-A has the highest affinity for VEGFR1, the
166 higher tyrosine kinase activity of VEGFR2 makes it the primary receptor involved in angiogenesis processes.
167 VEGFR1, conversely, is involved in macrophage chemotaxis, as well as in tumor cell survival and invasion
168 (136, 205, 255). Additionally, VEGFR1 may also have a negative role on angiogenesis and tumorigenesis
169 processes, only acting as a decoy receptor (102, 142). VEGF-A binds to its receptor and this first results in
170 blood vessel dilatation, as recently shown by *in vivo* studies of breast tumors, by sonographic examination (41).
171 Like most mammalian cells, endothelial cells express cell-swelling activated Cl⁻ channels, called the Volume-
172 Regulated Anion Channel (VRAC) (166, 167). Interestingly, receptor tyrosine kinases such as VEGFRs can
173 serve as mechanosensors (40). As such, they are involved in the VRAC signaling pathway. Thus, one might
174 infer that VEGF signaling in endothelial cell is mediated by VRAC Cl⁻ channels, as shown by the suppression of
175 angiogenesis by VRAC inhibitors (146).

176

177 2. Disorganization of the vascular bed

178 Angiogenesis partly proceeds by the VEGF stimulation of endothelial cell matrix metalloproteinases (MMPs)
179 (188). Secretion of these MMPs, more precisely MMP-2 and MMP-9 (18, 250), triggers the dissolution of the
180 basement membrane surrounding these endothelial cells, thus provoking the destabilization of the entire blood

181 vessel wall and leading to vessel leak (34). Among the multiple mechanisms that cooperate for the close
182 regulation of MMP activity, Ca^{2+} signaling is one of the most important (121). The work, led by Kato,
183 demonstrates that under extracellular acidification, B16 mouse melanoma cell line displays increased voltage-
184 gated Ca^{2+} channel expression and activity, resulting in an increase in intracellular Ca^{2+} concentration and in
185 MMP-9 induction. Thus, Ca^{2+} signals and channels are potentially the most important points in the regulation of
186 the VEGF-triggered blood vessel reorganization. However, the work conducted by Kato and colleagues
187 addresses the other types of Ca^{2+} channels that may be involved in the complex processes going from VEGF
188 upregulation to basement membrane disorganization. Indeed, under a hypoxic stimulus, the formation of a
189 VEGF-VEGFR complex at the endothelial cell plasma membrane can also result in intracellular Ca^{2+} increase,
190 due to the release of intracellular Ca^{2+} stores (Fig. 2). The depletion of these stores in turn results in Ca^{2+} influx,
191 via plasma membrane Ca^{2+} channels which results in store-operated calcium entry or SOCE (for review, see
192 (75)). Several Ca^{2+} channels of the transient receptor potential (TRP) superfamily appear to be involved in
193 SOCE and help explain how SOCE contributes to VEGF-induced basement membrane disruption. Among the
194 TRPCs (canonical), TRPC1 activation magnifies the VEGF-induced increase in endothelial cell monolayer
195 transepithelial permeability (111, 178). Conversely, inhibition of the TRPC1 activity reduces this transepithelial
196 permeability increase (111). TRPC4 is a necessary intermediate in the response of human pulmonary artery
197 endothelial cells to VEGF stimulation (71). TRPC6 is also presented as a good candidate in the process of
198 VEGF-induced vascular permeability increase. Indeed, as shown in individually perfused frog microvessels,
199 stimulation of TRPC6 generates an increase in the hydraulic conductivity (L_p) of the vessels (186). The Ca^{2+} -
200 release activated Ca^{2+} (CRAC) channel (composed of the plasma membrane Orai1 subunit coupled with the
201 endoplasmic reticulum STIM subunit) is also involved in VEGF-dependent SOCE (139). Indeed, in human
202 umbilical vein endothelial cells, inhibition or disruption of Orai1 reduces VEGF-induced calcium entry, cell
203 migration and *in vitro* angiogenesis. The increase in transepithelial permeability may result from both in-cell
204 and in-between-cell pore creation (43, 98). Finally, VEGF effects can also occur via receptor-operated-calcium
205 entry (ROCE), as shown on human microvascular endothelial cells (42); TRPC3 and TRPC6 are two examples
206 of channels involved in calcium-mediated vascular permeabilization.

207 In conclusion, VEGF-induced calcium-dependent events leading to the basement membrane disorganization,
208 though resulting in the same final outcome, appear to be the consequence of several calcium entry pathways.

210 3. Endothelial cell proliferation, migration and organization

211 Breakdown of the blood vessels (i.e. dissolution of the basement membrane) leads endothelial cells to be
212 released into the extracellular matrix (ECM) where they proliferate and organize to generate new blood vessels,
213 growing towards the tumor that initially emitted stimulating signals. Angiogenic sprouting is a guided process in
214 which the vasculature expands outwards from the preexisting vessel, following a gradient of angiogenic factors
215 such as VEGF-A (85) or soluble VEGFR1 (38). In this regard, Gerhardt and colleagues demonstrated that the
216 first cell of the column, the so-called “endothelial tip cell,” consists of a single, non-proliferative, highly
217 polarized endothelial cell, which is the sensor of the emitted elongation signal. The cell migration along this
218 signal (chemotaxis) paves a path for the cells that follow, the “stack cells”, which undergo high rate mitosis to
219 form the newly elongating blood vessel. At the endothelial cell level, it is now recognized that VEGFR2 is the
220 main actor of the angiogenic cascade. VEGF, when binding to VEGFR2, activates a signaling cascade starting
221 from auto-phosphorylation of VEGFR2, subsequent phosphorylation and activation of phospholipase C- γ (PLC-
222 γ), increased production of inositol 1,4,5-trisphosphate (IP3) and resulting increase in the Ca^{2+} release from
223 endoplasmic reticulum. The subsequent rise in calcium influx by the aforementioned SOCE process involves
224 TRPC channels, as already discussed (see chapter II.C.2) and promotes endothelial cell migration via a
225 Ca^{2+} /calmodulin/calcineurin pathway (72). The role of SOCE was also established in bovine artery endothelial
226 cells for the different mechanisms of angiogenesis (i.e., cell sprouting, cell proliferation and cell migration) and
227 involves the Ca^{2+} /calmodulin/protein kinase cascade (15).

228 One of the fundamental roles of ion channels in endothelial Ca^{2+} signaling is the fine-tuning of the
229 electrochemical gradient for Ca^{2+} (118). Several types of ion channels cooperate in elaborating the cell
230 membrane potential, which regulates calcium entry into the endothelial cells (118). Among those, K^+ channels,
231 especially Ca^{2+} -activated K^+ channels (251), inwardly rectifying K^+ channels (K_{ir}) and voltage-dependent K^+
232 channels, are the major classes of ion channels involved in setting the membrane potential (165), which in turn
233 influences the driving force for the Ca^{2+} via SOCE. Indeed, blockers of K^+ channels, as well as plasma
234 membrane depolarization, stop the cell cycle in G1 phase, thus reducing cell proliferation (153, 165). However,
235 among the Ca^{2+} -dependent K^+ channels, the Ca^{2+} -dependent intermediate conductance K^+ channel IK1 (SK4,

236 KCNN4) increases cell proliferation with a completely different mechanism, in a HEK overexpressing system.
237 Indeed, the potassium current itself does not seem to mediate the effect, since a non-transporting mutant of the
238 channel induces the same proliferation. So it seems that the IK1-induced proliferative effect is not SOCE-
239 mediated (152). The cell membrane potential is partly regulated by Cl⁻ channels (247). Hence, volume-regulated
240 anion channels (VRAC), Ca²⁺-activated Cl⁻ channels (CaCC), as well as cystic fibrosis transmembrane regulator
241 (CFTR) (145) might be important regulators of SOCE, which drives the proliferation step of angiogenesis.

242

243 4. Maturation of the new blood vessel

244 Two steps are essential for completion of angiogenesis:

245 (1) A new vessel is formed when the elongating sprout lumenizes to create a tube, which will eventually be able
246 to transport fluid and the multiple blood components: this step is called tubulogenesis (117, 143). At least two
247 models of tubulogenesis have been described. The “cell hollowing” model (currently the most studied) evolves
248 from a chain of single endothelial cells (117). This model involves intracellular vesicle formation, then fusion of
249 these vesicles into vacuoles, first intracellularly, and then from cell to cell, all along the sprout, when the
250 vacuoles get bigger. The future lumen vacuole membrane finally acquires hallmarks of an apical membrane,
251 thus defining the apical-basolateral polarity of the neo-formed vessel. In the less studied “cord hollowing”
252 model, which happens in a multi-cellular elongated stack, the cells first lose their initial polarity, then
253 basolateral membrane markers are accumulated at the membrane facing the extracellular matrix, while vesicles
254 are accumulated and get fused on the opposite membrane, hence forming an initial lumen that expands
255 afterwards (179, 248). In the “cell hollowing” model, the formation of vacuoles is linked to the activation of
256 intracellular chloride channels of the CLIC family. The chloride intracellular channel (CLIC) family of proteins
257 contains six mammalian members (CLIC1-6), from which only CLIC1 and CLIC4 are expressed in endothelial
258 cells (230). CLIC proteins can change from a soluble state to a membrane-linked state. In the latter case, when
259 inserted in such organelles as mitochondria, Golgi membranes, nuclear membranes, etc., CLICs function as
260 chloride channels, although very different in structure from the other known Cl⁻ channels (65, 233). In the “cell
261 hollowing” model of tubulogenesis, CLIC4 is present in human umbilical vein endothelial cells and promotes
262 several processes involved in angiogenesis, including luminal formation (229). Particularly, endothelial cells

263 derived from transgenic mice, in which the *Clc4* gene has been invalidated, fail to proceed to the vacuole
264 creation and extension process. Tung and colleagues show that the enlarging vacuoles acidify during their
265 formation, involving an electrogenic vacuolar proton-ATPase (vH^+ -ATPase), and propose that CLIC4 supports
266 this acidification by creating an electrical gradient for the proton transfer. The same article proposes that CLIC1
267 may partially compensate for the absence of CLIC4 in the process of tubulogenesis, although CLIC1 seems
268 mainly committed to proliferation and migration processes rather than to morphogenesis of endothelial cells
269 (230).

270 (2) The new hollow tube thus created is finally stabilized by the formation of a new basement membrane.
271 Indeed, in physiological angiogenesis, accessory cells, such as pericytes and smooth muscle cells are recruited
272 to the newly developed vessel, to form a new basement membrane (54). In tumor angiogenesis, however, the
273 newly formed blood vessels are very heterogeneous in structure and function (54). Their architecture is very
274 chaotic and disorganized, with abnormal branching, inflated cells and tortuous courses. Subsequently, the
275 association between endothelial and mural cells is altered, becoming loose and leading to highly permeable
276 vessels, which, paradoxically, are not adapted to a fluent and regular irrigation of the tumor, hence favoring its
277 dissemination to other, supposedly more welcoming, locations (35). The leaky vessels allow the tumor to escape
278 the noxious environment in which it grows (metastasis). Again, Ca^{2+} channels of the TRP family, namely
279 TRPV4 Ca^{2+} channels, are key determinants in the balance between normal and defective vasculature (2).
280 Indeed, tumoral endothelial cells naturally contain low levels of TRPV4. Conversely, overexpression or
281 pharmacological activation of TRPV4 normalizes the vasculature and restores the endothelial cell layer physical
282 and biophysical properties.

283

284 **III. Metastasis**

285 **A. The tumor microenvironment (TME): role and ion channel profile**

286 The extracellular environment of tumor cells is viewed as a complex network of stromal cells, deregulated
287 vasculature, extracellular membrane (ECM) proteins, growth factors and cytokines, all of which generated or
288 activated by the tumor cells, acting together to influence the growth, behavior and malignancy of the tumor

289 (236), and constantly evolving while the tumor grows and becomes more aggressive. The physical and chemical
290 characteristics of the TME is cancer-type dependent, but there are a number of typical, cancer type-independent
291 features, among which hypoxia, acidic pH, high levels of lactate and low levels of glucose (21, 82). In this
292 context, ion channels and transporters are important specialized proteins involved in the regulation of signaling
293 pathways associated with these properties. All the cells composing the tumor microenvironment (TME) are
294 laden with several types of misexpressed/overexpressed ion channels, which participate in the crosstalk
295 described by Arcangeli, between the tumor and its TME (8). These ion channels also contribute to maintain the
296 harsh conditions that allow tumor growth (33). Concerning the vasculature, we have already discussed the
297 channels involved in neoangiogenesis and mentioned the importance of endothelial cell Ca^{2+} homeostasis
298 throughout this process. Mesenchymal cells (fibroblasts, myofibroblasts, immune cells) also express ion
299 channels, particularly lymphocytes that infiltrate the tumors express K^+ channels of the $\text{K}_v1.3$ and $\text{K}_{Ca}3.1$ types,
300 which are involved in T cell activation via Ca^{2+} signaling (124). Interestingly, tumor-associated macrophages
301 express $\text{P2X}_7\text{R}$ non-selective cation channels and inwardly-rectifying K_{IR} K^+ channels; both families are
302 involved in the Ca^{2+} -dependent activation of macrophages (8).

303

304 B. Role of the epithelial-mesenchymal transition (EMT) in the metastasis process

305 Metastasis of epithelial cancer cells is a multistep process, which involves mobilization of the cells from their
306 original location, migration of the cells, invasion of surrounding locations, intravasation of cancer cells into
307 newly formed tumoral blood vessels, cell survival in the blood flow and seeding to more distant new niches,
308 where the cells start proliferating again, thus creating secondary tumors. Multiple oncogenic signals originating
309 from the TME convert the cells into their new metastatic and invasive phenotype. These signals lead the tumor
310 cells to acquire several biological capabilities described as hallmarks of cancer (95, 96) and mediated by an
311 epithelial-mesenchymal transition (EMT) process, otherwise called the metastatic cascade. During this cascade,
312 cells, which originally display epithelial cell features (i.e., adhesion to each other and to the basement
313 membrane, high level of E-cadherin involved in the epithelial integrity (16)) are progressively transformed into
314 mesenchymal cells (116). Briefly, during this process, polarized epithelial cells acquire a new phenotype, which
315 includes increased resistance to death, enhanced migratory capacities and invasiveness. The complete spectrum

316 of signaling agents that contribute to EMT remains an open question. EMT is a complicated process and ion
317 channels from several different families play a particularly important role in this transition (249).

318

319 1. Mobilization of the cells: involvement of ion channels

320 Loss of cell-cell contacts is the first step of EMT. This phenotype is associated with repression of cell junction
321 proteins, such as E-cadherin, and correlated with stimulation of both vimentin (an intermediate filament protein)
322 and N-cadherin (neural cadherin, normally expressed in migrating non-tumoral cells) synthesis (107, 164, 226).

323 This process has been shown in breast cancer cells to be Ca^{2+} -dependent (52). In fact, induction of EMT is
324 clearly related to the expression of TRPM7, the melastatin-like TRP Ca^{2+} channel. Indeed, upregulation of
325 TRPM7 regulates the expression of the EMT marker vimentin. Conversely, downregulation of the channel
326 prevents EMT induction in breast cancer cells. In colorectal carcinoma cells, the Ca^{2+} -activated KCNN4
327 ($\text{K}_{\text{Ca}3.1}$; SK4) channels have been shown to be involved in EMT promotion, probably through participation in
328 the cell hyperpolarization, subsequent induction of Ca^{2+} entry via voltage-dependent Ca^{2+} channels and an
329 increase in intracellular Ca^{2+} concentration (129).

330 Tumor cell mobilization is the next step in the metastatic cascade. This step involves migration of the cells from
331 their initial site and invasion of the surrounding stroma and vasculature. This is a complex process requiring the
332 coordination of multiple macromolecules involved in adhesion, cytoskeletal dynamics and digestion of the
333 ECM. Among the extracellular elements favoring EMT, hypoxia and microenvironment acidification have been
334 the most studied and are involved in tumor cell mobilization. As discussed earlier in this review, tumor acidity
335 is a direct result of hypoxia (83, 237). Indeed, low pO_2 raises the rate of glycolysis, which produces elevated
336 levels of lactate and protons in tumor cells. In order to maintain pH_i within a tight range ($\text{pH } 7.2 - 7.4$), the
337 tumor cells upregulate H^+ extrusion, which, in turn, results in microenvironmental acidosis, down to $\text{pH } 6.5$ (50)
338 or even to $\text{pH } 5.8$ as shown in highly metastatic MDA-MB-435S breast tumor cells (208). Extracellular acidosis
339 increases even more due to hypoxia-induced activation of carbonic anhydrases, especially CAIX (223).
340 Consequently, tumor cells are equipped with mechanisms that allow them to sense and react to the low pH_e to
341 which they have been exposed. pH sensors of several types are present in the plasma membrane of mammalian
342 tumor cells and these pH sensors are categorized into G-protein coupled receptors (GPCRs) and non-GPCR pH

343 sensors (50). GPCR signaling pathways are manifold: ERK pathway/ Ca^{2+} release (105), G_s -protein/cAMP,
344 $\text{G}_{12/13}$ -protein/Rho, and G_q -protein/phospholipase C pathways (140). pH sensing results in reorganization of the
345 cytoskeleton (e.g. actin rearrangement, stress fiber synthesis, etc.), thus contributing to tumor migration and
346 eventually to tumor progression (39). Non-GPCR proteins include ion channels of several families and
347 participate in migration and invasion phenotypes. Indeed, expression of the tumor cell cation-permeable Acid
348 Sensing Ion Channels (ASICs) is pH-dependent. ASIC1a is associated with tumor cell migration and invasion in
349 hepatocellular carcinoma cells (112); ASIC2a and ASIC3 have been identified in the plasma membrane of
350 adenoid cystic carcinoma cells and proposed to be involved in the acquired invasive capabilities of the cells
351 (257). The Ca^{2+} channel TRPV1 is also activated by low pH (259) and is correlated with tumor cell proliferation
352 in human prostate cancer cell lines (159). Both families of channels are involved in cancer-associated pain (57,
353 259) and plasma membrane phospholipids are proposed to be modulators of ASIC and TRPV1 activities (128).
354 Other acid-sensitive channels comprise several members of the Ca^{2+} channel TRP family, ionotropic
355 purinoceptors (P2X), inward rectifier K^+ channels (K_{ir}), voltage-activated K^+ channels (K_v family), L-type Ca^{2+}
356 channels (Ca_v , CACN), hyperpolarization-activated and cyclic nucleotide-gated cation channels (HCN), gap
357 junction channels, and Cl^- channels. They are involved in the process of tumor cell migration (217).
358 Interestingly, the Ca^{2+} -activated, high conductance, BKCa K^+ channels are not involved in breast cancer
359 progression, as shown by the absence of effect of iberiotoxin (an inhibitor of BKCa) in the processes of
360 proliferation, survival, migration or invasion in several breast cancer cell lines (200).
361 Migratory cells are polarized along their longitudinal axis, with hypoxia- and TME acidosis-dependent
362 protrusion of lamellipodia at the cell front edge and coordinated retraction of the rear edge (213). This process is
363 intracellular Ca^{2+} concentration-sensitive, thus involving Ca^{2+} release from intracellular stores and SOCE
364 channels (228). It creates the amoeboid movement characteristic of migrating tumor cells which is highly
365 dependent on ion and water flux (210). Indeed, the formation of lamellipodia is accompanied by relocalization
366 at the leading edge of a number of plasma membrane transport proteins, among which CAIX, NHE, MCT (all
367 involved in pH modulation), anion exchangers (such as NBC transporters or the anion exchanger AE2) and
368 water channels (aquaporins; AQP). It is proposed that the coordinated action of all these channels and
369 transporters makes it possible for the cells to establish the osmotic gradient that subsequently induces AQP-
370 mediated water flux at the leading edge of the cell. Water extrusion by AQP, at the rear end of the cell, is

371 mediated by the combined activity of Ca^{2+} -activated K^+ channels ($\text{K}_{\text{Ca}3.1}$) and Volume-Regulated Anion
372 Channels (VRACs) and leads to regulatory volume decrease (RVD) and shrinkage of the cells (206).

373

374 2. Invasion of surrounding TME and vasculature

375 Invasion of healthy tissues by tumor cells is critically dependent on the local degradation of ECM and basement
376 membrane, which otherwise would constitute a barrier to tumor metastasis. The degradation process thus creates
377 the escape pathway by which single tumor cells leave their primary location and colonize surrounding areas.
378 This degradation results from the low extracellular pH produced by tumor cells and relies on the secretion or
379 membrane expression of acid-dependent proteases (MMPs and cathepsins). Soluble (MMP2, MMP9) or
380 membrane-bound (MMP14) MMPs are key enzymes in tumor cell invasiveness, involved in both ECM and
381 basement membrane degradation (187, 193, 253) and involved as well in the angiogenic sprouting (18, 53).
382 MMP activity is promoted by low pH_e localized at the front part of migrating cells. As stated above, NHE1 is
383 one of the main actors of this extracellular acidification. While the acidification process appears as a crucial step
384 to weaken and break the ECM barrier, the physical progression of aggressive tumor cells relies on the formation
385 of ECM-degrading protrusions, called invadopodia, at the leading edge of the cells. NHE1 has been localized
386 within these invadopodia (29) and this localization has been correlated with extracellular acidification and MMP
387 expression and activity (88), thus suggesting the invadopodia as the major sites of ECM digestion and NHE1 as
388 the main channel involved in this process.

389 Voltage-gated sodium channels are highly expressed in epithelial tumors. In the MDA-MB-231 breast cancer
390 cell line, the $\text{Na}_\text{v}1.5$ Na^+ channel, by driving Na^+ influx and depolarizing the cells, greatly increases NHE1
391 potency to promote extracellular acidification and tumor invasion (23, 24, 86).

392 Calcium channels are also involved in the migration and invasion processes. Indeed, TRP channels, which
393 mediate SOCE Ca^{2+} entry and regulate intracellular $[\text{Ca}^{2+}]$, especially TRPV2 (158) and TRPM8 (169),
394 positively influence carcinoma invasion by upregulating MMP2 and MMP9 synthesis and production. Thus,
395 MMP production may be an intracellular $[\text{Ca}^{2+}]$ -dependent process. Interestingly, high concentrations of ATP
396 are observed in the TME of most hypoxic solid tumors (181). ATP acts as a signaling agent by activating
397 plasma membrane ligand-gated receptors of the P2X family, namely P2X₇R cation channels promote migration

398 and invasion, as shown in T47D breast cancer cells (253). P2X₇R cation channels mediate Ca²⁺ influx in tumor
399 cells thus contributing to the increase in intracellular [Ca²⁺]. In the breast cancer cell line MDA-MB-435S, ATP
400 activation of P2X₇R cation channels activates Ca²⁺-dependent SK3-mediated potassium currents, thus favoring
401 the elongation of cell protuberances, which, in turn, promote cell migration (110). In this model, the P2X₇R-
402 promoted invasive phenotype also relies on the activation of proteolytic enzymes, essentially cathepsin B.
403 Correlatively, pharmacological inhibition or downregulation of P2X₇R channels in PC-3M human prostate
404 carcinoma-derived cell lines inhibit ATP-driven migration and invasion (193). This Ca²⁺-dependent process
405 requires activation of either PI3/AKT or ERK1/2 signaling pathway. It reduces the expression of proteins
406 involved in the maintenance of cell-cell contacts (e.g. E-cadherin and claudin-1) and correlatively increases the
407 expression of Snail (an inhibitor of E-cadherin), interleukin 8 (IL-8, a promoter of migration and angiogenesis)
408 and MMPs, thus contributing to the invasion phenomenon. The metastasis- and malignancy-promoting status of
409 P2X₇R, however, is still an important field in cancer research. Indeed, recent studies have been designed to
410 explore the effects of P2X₇R on tumor development and metastatic outcome *in vivo*. Hofman and collaborators
411 demonstrate that colitis-associated cancers (colon adenocarcinoma, colon carcinoma) are favored by P2X₇R
412 silencing or pharmacological inhibition (103). A different study reveals that P2X₇R restricts tumor progression
413 by developing antitumoral immunity (4).

414

415 3. Metastasis: transit in blood vessels and extravasation

416 Tumor cell metastasis occurs when the cells successfully escape from their primary location, overcome the step
417 of transit in the circulation as unattached cells and finally disseminate to their secondary sites by extravasation
418 from the vessels and reattachment to a new ECM support.

419 Circulating tumor cells do not always survive long enough to metastasize. It is estimated that only 0.01% of the
420 cells that enter the vasculature will eventually survive the harsh conditions to which they are submitted (87).

421 Tumor cells must resist anoikis whereby apoptosis is triggered by inadequate or loss of cell anchorage. Among
422 the multiple signals involved in anoikis resistance, several types of ion channels have been described (137).

423 Indeed, to effectively resist detachment-induced apoptosis, tumor cells utilize mechanisms that eventually limit
424 the rise in intracellular [Ca²⁺]. Ca²⁺ channels are thus downregulated, especially those acting via SOCE (130).

425 Depletion in other Ca^{2+} channels (TRPM2, TRPV6) also tends to reduce cell apoptosis (211). Activity of
426 caspases and other DNA-degrading enzymes, as well as a decrease in membrane potential have been linked to
427 loss of intracellular $[\text{K}^+]$ by K^+ channels. Cell shrinkage involved in apoptotic volume decrease (AVD) results
428 from a loss of KCl by K^+ and Cl^- channels, with a concomitant loss of water. Thus, downregulation of K^+ (K_v ,
429 K_{Ca} , K_{ATP} , K_{IR} , $\text{K}_{2\text{P}}$) and Cl^- channels also prevents apoptosis (26). Alternatively, anoikis-resistant cells can
430 develop regulatory volume increase (RVI) whereby Cl^- channels of the volume-regulated (VRAC) family,
431 especially prostate cancer cell ClC-3 (138), which are involved in the maintenance of cell volume under
432 hyperosmotic stress, are upregulated in cells that resist apoptosis. Interestingly, Ca^{2+} -activated Cl^- channels
433 (CLCA) may be either activated or not activated in anoikis-resistant cells, as shown in mammary gland cells
434 (67, 244), thus suggesting a different function of these channels in cancer cells.

435 Tumor cells are subjected to mechanical forces resulting from the blood stream, especially in narrow capillaries.
436 Mechanical stress is the main destruction pathway that tumor cells must avoid in order to be able to form
437 metastases. To prevent destruction, they develop integrin-involving adhesion processes and eventually get
438 attached to the endothelial bed of the vessel in which they are traveling (81). This interaction between
439 circulating tumor cells and endothelial cells is fundamental for the rest of the metastasis process, as it
440 determines the site where the cells will eventually exit from the vessel, hence initiating the extravasation
441 process. Gout and Huot have described the whole process of extravasation (87). Briefly, this process involves a
442 first step of tumor cell adhesion to endothelial cells, utilizing specific receptors. Tumor cells rolling along the
443 endothelial layer characterize this “light adhesion” step. A second step of firmer cell attachment then takes
444 place, mediated by cytokines and cell adhesion molecules. Extravasation (or diapedesis) finally occurs when the
445 cells cross the endothelial barrier through cell-cell junctions. As for several steps of the tumorigenesis process,
446 the importance of intracellular $[\text{Ca}^{2+}]$ and of the associated Ca^{2+} channels is also demonstrated for metastasis of
447 tumor cells. Indeed Orai1 , the pore-forming component of the store-operated Ca^{2+} channel and its associated
448 STIM1 Ca^{2+} sensor are associated with cell migration, invasion and metastasis in human breast carcinoma cells
449 (256). *In vivo* studies in a zebrafish model corroborate the importance of the $\text{Orai1}/\text{STIM1}$ partners and of
450 SOCE Ca^{2+} entry in the process of cell extravasation (262). Indeed, inhibition of the store-operated Ca^{2+} entry or
451 knockdown of Orai1 in nasopharyngeal carcinoma cells prevents extravasation of the cells from vasculature in a
452 zebrafish hematogenous metastasis model.

453 Finally, diapedesis is the result of tumor cells passing in between insufficiently tight endothelial cell junctions
454 (due to damaged vasculature). The alterations in cell shape necessary for diapedesis are most likely explained by
455 K^+ and Cl^- fluxes and by the following water fluxes. Indeed, it has been already shown that glioma cell
456 migration and invasion are supported by K^+ and Cl^- channels, which, by governing transmembrane water
457 movements and the resulting cell shape, are important for glioma invasion through narrow spaces (215).
458 Colonization of the secondary site by the tumor is not simply a matter of gaining access to this site. Tumor cells
459 will only be able to colonize compatible target tissues. The host tissue must then adapt to be able to foster the
460 colonizing tumor cells; this is the concept of “pre-metastatic niche” (212). Tumor cells must also adapt to their
461 new stroma and become able to secrete all the molecules necessary for their efficient attachment and ulterior
462 vascularization (91). Hence, there are many barriers to overcome before the tumor actually metastasizes.

463

464 **IV. Sigma 1 receptor: the main organizer?**

465 In the present review, we will skip the long history of sigma-1 receptor (Sig-1R), which has been widely studied
466 on the functional, pharmacological, structural and mechanistic points of view for the last 30 – 40 years (11, 148,
467 245). Similarly, we will not open this discussion to the sigma-2 receptor, which has not been cloned yet,
468 although it was shown to be involved in cancer development (234). Instead, we will focus on the specific role of
469 chaperone of Sig-1R that is of major interest in the context of cancer development.

470 Sig-1R is a 25 kD, endoplasmic reticulum (ER)-resident, ligand-regulated protein, which is involved in many
471 diseases, ranging from toxic substance addiction to stroke and cancer (10, 11, 148). It is mainly present in
472 normal brain, heart and liver, but is highly expressed in tumor cell lines and human cancer tissues of various
473 origins, including lung, colon, prostate carcinomas, sarcomas, breast tumors, brain tumors, and melanomas (10,
474 11, 113, 173, 241), in which it interferes with cell cycle and proliferation (10, 197). Interestingly, in breast
475 cancer cells, Sig-1R positively correlates with the metastatic grade of the cells, i.e. highly metastatic cells
476 contain higher levels of Sig-1R mRNA and protein than their less aggressive counterparts (10).

477 Sig-1R is anchored in the ER membrane, in a specific region called mitochondrion-associated ER membrane
478 (MAM). At rest, it is in a “dormant” state, associated with a chaperone protein, called GRP78 or BiP (100). BiP

479 is a main regulator of ER function: it has a high Ca^{2+} -buffering capacity and serves to store Ca^{2+} , which is
480 essential to many signaling pathways. Under specific ligand stimulation or under prolonged ER stress such as
481 ER Ca^{2+} depletion, Sig-1R is up-regulated, dissociates from BiP and becomes able to act as a chaperone (100)
482 after translocation to other ER areas or to the plasma membrane (101). In the plasma membrane, Sig-1R
483 interacts with – and regulates the activity of – several ion channels (Ca^{2+} , K^+ , Na^+ , Cl^-), receptors (NMDA
484 receptor) or kinases (Src) (148).

485 In the context of cancer development, as we discussed earlier in this review, many ion channels of diverse
486 classes (voltage-gated, calcium-gated, volume activated) are involved in a number of tumor cell processes.
487 Voltage-gated ion channels are involved in cell processes including proliferation, apoptosis, angiogenesis,
488 adhesion to ECM and migration. Interestingly, Sig-1R interacts with most of these channels, as demonstrated in
489 several works (12, 13, 126, 197). Patch-clamp studies and western blot assays in the presence of Sig-1R ligands
490 (e.g. (+)-pentazocine, igmesine, and 1,3-di(2-tolyl)guanidine [DTG]) were conducted in small cell lung cancer
491 (SCLC) derived cell lines (NCI-H146 and NCI-H209) and in T-cell leukemia-derived Jurkat cells. These studies
492 demonstrate that ligand-mediated inhibition of Sig-1R results in cell cycle arrest in G_1 phase with no stimulation
493 of apoptosis (197). This effect correlates with the reduction of voltage-operated K^+ currents in all these cell
494 lines, indicating that Sig-1R-mediated cell growth arrest is the result of the negative regulation of K^+ channels
495 (197). Cell cycle arrest also correlates with the negative modulation of Volume-Regulated Chloride Channels
496 (VRCC), both effects leading to the overall inhibition of the Regulated Volume Decrease (RVD) process, (196),
497 which is known to govern cell cycle progression and numerous cellular functions such as proliferation,
498 apoptosis, migration, ... (202). The *Xenopus* oocyte system allowed direct reconstitution (in the absence of Sig-
499 1R ligands) of the interaction that exists between Sig-1R and potassium channels (Kv1.4 and Kv1.5) and
500 established that Sig-1R directly acts as an auxiliary subunit of a few channels to govern several cell functions
501 (12). In order to provide further evidence of this direct interaction, I participated in a study (47) showing that in
502 chronic myeloid leukemia (CML)-derived cells (K562) – which overexpress Sig-1R – the abnormally expressed
503 cardiac K^+ channel hERG co-immunoprecipitated with Sig-1R. In this model, Sig-1R extinction by a specific
504 shRNA or Sig-1R inhibition by Sig-1R ligands led to the reduction of hERG K^+ currents, whereas co-injection
505 of both hERG and Sig-1R in *Xenopus* oocytes increased hERG current when compared to oocytes injected with

506 hERG alone. This modulation proceeds from the regulation of hERG channel maturation and stability, which is
507 in adequation with the role Sig-1R as a chaperone of hERG K⁺ channel in cancer cells (47).

508 Recently, using the technique of atomic force microscopy, Sig-1R was shown to physically associate, in a four-
509 fold symmetry (1 channel molecule vs. 4 Sig-1R molecules), with the voltage-gated Na⁺ channel Na_v1.5 (13),
510 previously proven to promote invasiveness in breast cancer cells (24, 86). With the same technique, the same
511 team demonstrated physical interaction between Sig-1R and acid-sensing ion channels (ASIC1a; (36)) and
512 provided evidence of Sig-1R association with the heteromeric GluN1/GluN2A NMDA receptor, via specific
513 binding of Sig1R to the GluN1 subunit (14).

514 These data raise the following question: how can an ER protein interact with plasma membrane ion channels of
515 so many different families and govern so many cell functions in oncogenesis? A plausible explanation resides in
516 the following observations: 1) Sig-1R is localized in ER, a cell area where Ca²⁺ signaling is crucial; 2) Sig1R is
517 highly mobile in the ER membrane and Sig-1R overexpression, cellular stress or pharmacological treatment lead
518 to dissociation of Sig-1R from its chaperone BiP and subsequent translocation to the plasma membrane (100,
519 149, 221), where it interacts with its client proteins (ion channels, receptors). Indeed, during cancer progression,
520 the tumor cells undergo harsh conditions (hypoxia, low pH, starvation in nutrients) or even treatments that
521 create cell stress and alter ER protein expression levels or generate abnormal secretion of ER-resident
522 chaperones (157). Indeed, inasmuch as ER chaperones such as Sig-1R contribute to the correct folding of newly
523 synthesized or conformationally distorted proteins, or to protect proteins that are prone to de misfolded, their
524 role becomes crucial in case of cell stress. Moreover, it is worth pointing out that the BiP-Sig-1R association is
525 tightly regulated in an ER Ca²⁺/Mn²⁺-dependent manner. Any cell signal that results in lowering ER Ca²⁺
526 concentration below physiological concentration (0.5 mM) leads to disassembly the two chaperones, thus
527 increasing the activity of each (100). Under cell stress conditions, Sig-1R stabilizes the conformation of type3-
528 IP3 receptor in the ER, thus stimulating Ca²⁺ transport from the ER to the mitochondria (100) and to the cytosol
529 (252). Hence, Sig-1R orchestrates Ca²⁺ signaling between ER and mitochondria, by fueling the production of
530 ATP by the mitochondria, which conversely regulates Ca²⁺ concentration in the ER (99). Sig1R also maintains
531 proper folding and proper function of the plasma membrane proteins challenged by cell stress: it thus plays a
532 major role in cell survival (Figure 3). Sig-1R has since then been proposed as a modulator of inter-organelle
533 signaling (220).

534 Subsequently, several lines of evidence demonstrate that Sig-1R interacts with different types of proteins
535 including ER stress sensors, cytokines, ion channels and that this interaction constitutes the mechanism whereby
536 Sig-1R governs cancer cell behavior in response to alterations of the tumor microenvironment. Sig-1R
537 modulates cell survival (160), apoptosis (60, 151) and cell cycle (196, 197). Recently, I participated in a study
538 presenting evidence that Sig-1R is a key regulator of the signaling events that occur between the tumor cell and
539 its microenvironment. Indeed, in chronic myeloid leukemia- and colon carcinoma-derived cells (resp. K562 and
540 HCT-116 cells), we showed that extracellular matrix triggered a Sig-1R-dependent increase in cell adhesion,
541 cell migration, cell invasiveness, as well as VEGF secretion (48) and that a hERG K⁺ channel-β1 integrin
542 interaction mediated these effects in both cell lines. Silencing of Sig-1R in K562 and HCT-116 cells resulted in
543 *in vivo* reduction of angiogenesis, invasion and extravasation (48). Sig-1R thus appears as a major regulator of
544 tumor development. Further studies are necessary to assess whether Sig-1R might be envisioned as a therapeutic
545 target and whether Sig-1R ligands might be designed as antitumoral treatments.

546

547 **V. Antiangiogenic therapy and therapy resistance**

548 Cancer is one of the deadliest pathologies, with approximately 14 million of new cases and 8 million of cancer-
549 related deaths in 2012; negative WHO predictions expect annual cancer cases to rise from 14 million in 2012 to
550 22 million in the next two decades. Carcinomas (lung, liver, stomach, colorectal, breast, esophageal) are among
551 the most common causes of cancer deaths <http://www.who.int/mediacentre/factsheets/fs297/en/>.

552 Conventional global anti-tumoral therapies (chemotherapies) are only partly efficient because of the lack of
553 responsiveness of some cancers and because these therapies sometimes induce systemic toxicity or resistance,
554 ending up in treatment discontinuation. For vascularized cancers, a more precise strategy of angiogenesis
555 inhibition has been developed, which focuses on the neutralization of the effects of angiogenic factors (FGF,
556 VEGF) or on the inhibition of their receptors (70, 122). This strategy leads to vasculature destruction. Given the
557 drawbacks of such therapies (89, 90), which can lead to more severe condition or even to death (194, 195), solid
558 tumor therapies evolved towards normalization rather than destruction of the abnormal vasculature, the
559 objective being to restore a close to normal blood flux and pO₂, to favor normal immune cell function and

560 finally, to allow efficient delivery of anticancer agents (106, 109, 190, 219). Thus, antiangiogenic therapies were
561 used to treat several vascularized cancers in the last 40 years. However, too many carcinomas are still refractory
562 to antiangiogenic-based treatments.

563 Hence, in parallel to existing therapeutic strategies, it is important to identify predictive biomarkers of stage and
564 aggressiveness for the different types of tumors, as well as further targets for the development of
565 complementary treatments to be used in combination with VEGF-targeted therapies. Carcinogenesis is a
566 multistep process in which each step constitutes a potential angle of attack for a new therapeutic strategy
567 development. Ion channels are implicated in the growth, migration, invasion and metastasis of tumors and
568 therefore cancers can be included in the category of “channelopathies” (pathologies characterized by alterations
569 in channel function). Consequently, recent studies present ion channels as potential targets for the development
570 of future treatments. Several factors are in favor of the use of ion channels as therapeutic targets: i) cell
571 expression of ion channels is often modified in cancers; ii) this expression varies with the type of tumor and, in
572 a given type of tumor, with the specific step of tumor progression; iii) ion channels are accessible from the
573 extracellular side of the cell and the potential pharmacological tools (available or yet to be designed) are
574 multiple.

575 We do not intend here to review how ion channels can be targeted in each step of cancer development or in
576 different cancers to design potential therapeutic strategies. Several excellent reviews have already been written
577 from these standpoints (9, 49, 155, 203, 239). Rather, we will take a few examples of global cancer-associated
578 clinical dysfunctions in which ion channels may be targeted in order to restore normal function or diminish
579 cancer-associated disagreements and, finally, to improve patient’s condition.

580

581 *Restoring a less toxic microenvironment*

582 Concomitant with gradual accumulation of mutations in tumor cells, changes in the physical parameters of the
583 microenvironment also contribute to reinforce the cell aggressiveness. Among these physical changes, as
584 discussed in §II.A of this review, the microenvironment undergoes hostile conditions characterized by low pO₂,
585 high interstitial pressure, low extracellular glucose and high extracellular lactate – all of these create high
586 extracellular acidity. Regulating the establishment of such harsh conditions would reduce the metastatic
587 potential of tumor cells. Cancer therapies thus include trials based on the regulation of extracellular pH. They

588 consist of systemic buffering using sodium bicarbonate to increase extracellular pH or take advantage of the
589 proton transport systems involved in tumor cells (150). For the latter, inhibition of carbonic anhydrases
590 (especially CAIX), of the Na^+/H^+ exchanger, using amiloride or EIPA are efficient but have not been developed
591 for clinical use. Conversely, inhibition of the vacuolar H^+ -ATPase, using omeprazole (already used to suppress
592 gastric acidity) was proven efficient as a pH normalizer in cancer treatment. Interestingly, the opposite treatment
593 consisting in hyperacidification of tumor cells or of their microenvironment is also a promising treatment.
594 Intracellular hyperacidification promotes tumor cell apoptosis whereas extracellular hyperacidification may
595 provide an important tool to deliver therapeutic agents directly to the tumor site using acid-accumulating
596 peptides (69).

597

598 *Normalizing the vasculature*

599 Carcinoma progression is associated with the production, by tumoral and stromal cells, of angiogenic factors,
600 which participate in the development of a non-productive angiogenesis, due to blood vessel disorganization and
601 to vessel wall high permeability. Antiangiogenic treatments, which tend to block the development of new blood
602 vessels and harvest the tumor from oxygen and nutrients, are inadequate or even counter-productive to treat
603 cancers. Other targets are now under scrutiny, which could participate in the emerging process known as
604 vascular normalization, and among these targets are ion channels. Physical properties of the ECM have been
605 shown to influence normal and malignant cell behavior. Especially, angiogenic response is influenced by the
606 stiffness of tumor ECM (134). Ca^{2+} influx is one of the main responses of endothelial cells to the mechanical
607 stress resulting from the growth of new vessel tips in such a stiff environment. In a recent paper, it has been
608 shown that in a model of mouse prostate adenocarcinoma, mechanosensitive transient receptor potential
609 vanilloid 4 (TRPV4) calcium channels regulate angiogenesis and tumor vessel maturation by the regulation of
610 tumor endothelial cell mechanosensitivity via the reduction in basal Rho activity (2). This confirms that ion
611 channels must be included in the set of molecules to be targeted in the design of the new tools aiming at
612 normalizing vasculature, hence allowing better delivery of complementary antitumoral therapies. To further
613 reinforce the biological relevance of Ca^{2+} channels as molecular targets for antitumoral treatments, we can
614 mention that all the families of Ca^{2+} channels have recently been the focus point of researches demonstrating the
615 involvement of these channels in pro-angiogenic processes. Many of these channels are now the object of

616 patents that describe the use of modulators of their function in order to treat the symptoms associated with
617 cancer, i.e., proliferation, angiogenesis and metastasis; for review, see (163). The use of K^+ channel or pH
618 modulators to fight cancer development has been patented as well (49, 198).

619

620 *Fighting cancer-induced pain*

621 Cancer-related pain is one of the most devastating side effects of tumor development. Indeed, hypoxia and,
622 consequently, tumor acidosis are known to be responsible for the generation of pain (108). Cancer-related pain
623 can also be the result of tumor metastasis especially in the case of bone dissemination. Indeed, cytokines
624 produced by the tumor cells or by the bone microenvironment activate osteoclasts and thus contribute to the
625 signaling of cancer-associated pain in bones (225). Anti-tumoral treatment administration can, by itself, be a
626 factor of pain, whether the treatment is traditional chemotherapy, radiation, or surgery.

627 Moreover, it now seems that, except for morphine, for which the study results are not conclusive, conventional
628 pain-reducing therapies (NSAIDs, cannabinoids, opioids, GABA-ergic drugs) may have a negative impact on
629 cancer evolution (135). Consequently, the research of other analgesics and their targets is required. Current
630 research on this topic focuses on ion channels associated with pain, such as ASIC (57, 258), especially ASIC3
631 (56), and such as TRP channels of several sub-families, especially TRPV1 (192). Indeed, both types of channels
632 are known to be expressed in sensory neurons and to be activated by an acid load. These channels represent
633 relevant targets for the future development of new analgesics. Treatments based on TRPV1 pharmacological
634 inhibitors are already being clinically tested with some showing promising results (22). Voltage-gated Ca^{2+}
635 channels and voltage-gated Na^+ channels are also under study, especially $Ca_v2.2$, $Ca_v3.2$, $Na_v1.7$, $Na_v1.8$.
636 Specific antagonists derived from conotoxins have been developed and now seem to have good results in
637 clinical tests (242).

638 This field of research is still wide open and progressing very quickly. It is interesting to note that most of the ion
639 channels involved in the perception of pain are also implicated in one step or another of tumor development.
640 This leaves open the possibility that a specific treatment designed against an ion channel in order to treat cancer-
641 associated pain might also be used to prevent cancer progression. In particular, cannabinoids demonstrate
642 antitumoral effect in several cancer types (hepatocellular carcinoma, ccRCC, etc.), which is not mediated by the
643 regular CB1 and CB2 cannabinoid receptors (131, 235). Since cannabinoids can bind to many different ion

644 channel types, such as GPCR, NMDA receptors, GABA_A receptors (182), any of them can mediate the effect of
645 cannabinoids in tumor cells, thus shedding a new light on the yet unexplained anti-oncogenic effect of
646 cannabinoids.

647 In this topic, it is tempting to propose the potential development of Sig-1R ligands as treatments to fight cancer-
648 induced pain. Indeed, as we mentioned earlier, Sig1R promotes the action of ASIC channels, which are involved
649 with nociception (36). Sig-1R-inhibiting agents might be a way to obtain relief in this matter.

650

651 **VI. Conclusion**

652 A new concept in oncogenesis has emerged for the last 20 to 25 years, which states that ion channels and
653 transporters regulate different aspects of the neoplastic cell physiology. A growing body of evidence now
654 demonstrates that they govern the so-called “hallmarks of cancer”. As for neoangiogenesis, ion channels are
655 involved at all stages, in tumor cells, as well as endothelial cells or stromal cells. Ion channels thus represent a
656 new and promising field of research concerning the development of novel therapeutic agents. TRP channels for
657 instance, which are highly expressed in endothelial cells and involved in thermosensation, osmoregulation and
658 mechanoreception, are under current scrutiny for novel therapeutic strategy developments. Similarly, whereas
659 monocarboxylate transporters (MCT1 and MCT4) have now emerged as anticancer targets, the lactate shuttle
660 mechanism governing tumor cell pH regulation is still under study (55, 59). However, it is important to take into
661 account that most ion channels, though overexpressed in many cancer types, still exist in healthy organs. The
662 use of ion channel inhibitor-based therapies might thus induce many side effects, which will need to be carefully
663 prevented or minimized. As an example, blocking hERG1 K⁺ channel prevents the angiogenic switch in
664 colorectal cancer and, in turn, reduces metastasis (46). However, hERG1 is one of the main heart-expressed K⁺
665 channels, which certainly hinders the use of any of the current hERG1-targeted molecule as a cancer therapeutic
666 agent. It might thus be important to develop non-cardiotoxic hERG1 blockers to be used as complementary anti-
667 angiogenic treatments. The ion channel-based antitumoral therapies, though very attractive and potentially
668 powerful, probably need more research time in order to be carefully designed for safe clinical use. It might be
669 more fruitful to develop pharmacological treatments against Sig-1R, taking into account that this protein is

670 present in many tissues (CNS, lung, liver, pancreas, spleen), in membrane compartments. Targeted
671 pharmacological action on Sig-1R at the MAM or at the plasma membrane represents a potential new axis for
672 the development of therapies against numerous diseases including cancer.

673

674

675 **ACKNOWLEDGEMENTS**

676 I am very grateful to Ryan Arant, H el ene Guizouarn, Gilles Pag es and Olivier Soriani for their careful reading
677 of the manuscript and useful comments.

678

679 **REFERENCES**

- 680 1. **Aalkjaer C, Boedtkjer E, Choi I, and Lee S.** Cation-coupled bicarbonate transporters. *Compr Physiol*
681 4: 1605-1637, 2014.
- 682 2. **Adapala RK, Thoppil RJ, Ghosh K, Cappelli HC, Dudley AC, Paruchuri S, Keshamouni V,**
683 **Klagsbrun M, Meszaros JG, Chilian WM, Ingber DE, and Thodeti CK.** Activation of mechanosensitive ion
684 channel TRPV4 normalizes tumor vasculature and improves cancer therapy. *Oncogene* 2015.
- 685 3. **Adinolfi E, Capece M, Amoroso F, De Marchi E, and Franceschini A.** Emerging roles of P2X
686 receptors in cancer. *Curr Med Chem* 22: 878-890, 2015.
- 687 4. **Adinolfi E, Capece M, Franceschini A, Falzoni S, Giuliani AL, Rotondo A, Sarti AC, Bonora M,**
688 **Syberg S, Corigliano D, Pinton P, Jorgensen NR, Abelli L, Emionite L, Raffaghello L, Pistoia V, and Di**
689 **Virgilio F.** Accelerated tumor progression in mice lacking the ATP receptor P2X7. *Cancer Res* 75: 635-644,
690 2015.
- 691 5. **Adinolfi E, Raffaghello L, Giuliani AL, Cavazzini L, Capece M, Chiozzi P, Bianchi G, Kroemer G,**
692 **Pistoia V, and Di Virgilio F.** Expression of P2X7 receptor increases in vivo tumor growth. *Cancer Res* 72:
693 2957-2969, 2012.
- 694 6. **Airley RE, and Mobasher A.** Hypoxic regulation of glucose transport, anaerobic metabolism and
695 angiogenesis in cancer: novel pathways and targets for anticancer therapeutics. *Chemotherapy* 53: 233-256,
696 2007.
- 697 7. **Amoroso F, Capece M, Rotondo A, Cangelosi D, Ferracin M, Franceschini A, Raffaghello L,**
698 **Pistoia V, Varesio L, and Adinolfi E.** The P2X7 receptor is a key modulator of the PI3K/GSK3 /VEGF
699 signaling network: evidence in experimental neuroblastoma. *Oncogene* 2015.
- 700 8. **Arcangeli A.** Ion channels and transporters in cancer. 3. Ion channels in the tumor cell-
701 microenvironment cross talk. *Am J Physiol Cell Physiol* 301: C762-771, 2011.
- 702 9. **Arcangeli A, Crociani O, Lastraioli E, Masi A, Pillozzi S, and Becchetti A.** Targeting ion channels in
703 cancer: a novel frontier in antineoplastic therapy. *Curr Med Chem* 16: 66-93, 2009.
- 704 10. **Aydar E, Onganer P, Perrett R, Djamgoz MB, and Palmer CP.** The expression and functional
705 characterization of sigma (sigma) 1 receptors in breast cancer cell lines. *Cancer Letters* 242: 245-257, 2006.
- 706 11. **Aydar E, Palmer CP, and Djamgoz MBA.** Sigma receptors and cancer: possible involvement of ion
707 channels. *Cancer Research* 64: 5029-5035, 2004.
- 708 12. **Aydar E, Palmer CP, Klyachko VA, and Jackson MB.** The sigma receptor as a ligand-regulated
709 auxiliary potassium channel subunit. *Neuron* 34: 399-410, 2002.

- 710 13. **Balasuriya D, Stewart AP, Crottès D, Borgese F, Soriani O, and Edwardson JM.** The sigma-1
711 receptor binds to the Nav1.5 voltage-gated Na⁺ channel with 4-fold symmetry. *J Biol Chem* 287: 37021-37029,
712 2012.
- 713 14. **Balasuriya D, Stewart AP, and Edwardson JM.** The σ -1 receptor interacts directly with GluN1 but
714 not GluN2A in the GluN1/GluN2A NMDA receptor. *J Neurosci* 33: 18219-18224, 2013.
- 715 15. **Banumathi E, O'Connor A, Gurunathan S, Simpson DA, McGeown JG, and Curtis TM.** VEGF-
716 induced retinal angiogenic signaling is critically dependent on Ca²⁺ signaling by Ca²⁺/calmodulin-dependent
717 protein kinase II. *Invest Ophthalmol Vis Sci* 52: 3103-3111, 2011.
- 718 16. **Bates RC, and Mercurio AM.** The epithelial-mesenchymal transition (EMT) and colorectal cancer
719 progression. *Cancer Biol Ther* 4: 365-370, 2005.
- 720 17. **Benga G.** The first discovered water channel protein, later called aquaporin 1: molecular characteristics,
721 functions and medical implications. *Mol Aspects Med* 33: 518-534, 2012.
- 722 18. **Bergers G, Brekken R, McMahon G, Vu TH, Itoh T, Tamaki K, Tanzawa K, Thorpe P, Itohara S,
723 Werb Z, and Hanahan D.** Matrix metalloproteinase-9 triggers the angiogenic switch during carcinogenesis.
724 *Nat Cell Biol* 2: 737-744, 2000.
- 725 19. **Bieche I, Vacher S, Vallerand D, Richon S, Hatem R, De Plater L, Dahmani A, Némati F, Angevin
726 E, Marangoni E, Roman-Roman S, Decaudin D, and Dangles-Marie V.** Vasculature analysis of patient
727 derived tumor xenografts using species-specific PCR assays: evidence of tumor endothelial cells and atypical
728 VEGFA-VEGFR1/2 signalings. *BMC Cancer* 14: 178, 2014.
- 729 20. **Bortner CD, and Cidlowski JA.** Ion channels and apoptosis in cancer. *Philos Trans R Soc Lond B Biol
730 Sci* 369: 20130104, 2014.
- 731 21. **Brahimi-Horn MC, and Pouyssegur J.** Oxygen, a source of life and stress. *FEBS Lett* 581: 3582-3591,
732 2007.
- 733 22. **Brederson JD, Kym PR, and Szallasi A.** Targeting TRP channels for pain relief. *Eur J Pharmacol*
734 716: 61-76, 2013.
- 735 23. **Brisson L, Driffort V, Benoist L, Poet M, Counillon L, Antelmi E, Rubino R, Besson P, Labbal F,
736 Chevalier S, Reshkin SJ, Gore J, and Roger S.** Nav1.5 Na⁺ channels allosterically regulate the NHE-1
737 exchanger and promote the activity of breast cancer cell invadopodia. *J Cell Sci* 126: 4835-4842, 2013.
- 738 24. **Brisson L, Gillet L, Calaghan S, Besson P, Le Guennec JY, Roger S, and Gore J.** Na(V)1.5
739 enhances breast cancer cell invasiveness by increasing NHE1-dependent H(+) efflux in caveolae. *Oncogene* 30:
740 2070-2076, 2011.
- 741 25. **Brown RS, Goodman TM, Zasadny KR, Greenson JK, and Wahl RL.** Expression of hexokinase II
742 and Glut-1 in untreated human breast cancer. *Nucl Med Biol* 29: 443-453, 2002.
- 743 26. **Burg ED, Remillard CV, and Yuan JX.** K⁺ channels in apoptosis. *J Membr Biol* 209: 3-20, 2006.
- 744 27. **Burrell K, and Zadeh G.** Molecular Mechanisms of Tumor Angiogenesis. In: *Tumor Angiogenesis*,
745 edited by Ran DSInTech, 2012, p. 296.
- 746 28. **Bury M, Girault A, Mégalizzi V, Spiegl-Kreinecker S, Mathieu V, Berger W, Evidente A,
747 Kornienko A, Gailly P, Vandier C, and Kiss R.** Ophiobolin A induces paraptosis-like cell death in human
748 glioblastoma cells by decreasing BKCa channel activity. *Cell Death Dis* 4: e561, 2013.
- 749 29. **Busco G, Cardone RA, Greco MR, Bellizzi A, Colella M, Antelmi E, Mancini MT, Dell'Aquila
750 ME, Casavola V, Paradiso A, and Reshkin SJ.** NHE1 promotes invadopodial ECM proteolysis through
751 acidification of the peri-invadopodial space. *FASEB J* 24: 3903-3915, 2010.
- 752 30. **Buttigieg J, Pan J, Yeger H, and Cutz E.** NOX2 (gp91phox) is a predominant O₂ sensor in a human
753 airway chemoreceptor cell line: biochemical, molecular, and electrophysiological evidence. *Am J Physiol Lung
754 Cell Mol Physiol* 303: L598-607, 2012.
- 755 31. **Camacho J.** Ether a go-go potassium channels and cancer. *Cancer Letters* 233: 1-9, 2006.
- 756 32. **Campbell TM, Main MJ, and Fitzgerald EM.** Functional expression of the voltage-gated Na⁺-channel
757 Nav1.7 is necessary for EGF-mediated invasion in human non-small cell lung cancer cells. *J Cell Sci* 126: 4939-
758 4949, 2013.
- 759 33. **Cardone RA, Casavola V, and Reshkin SJ.** The role of disturbed pH dynamics and the Na⁺/H⁺
760 exchanger in metastasis. *Nat Rev Cancer* 5: 786-795, 2005.
- 761 34. **Carmeliet P, and Jain RK.** Molecular mechanisms and clinical applications of angiogenesis. *Nature*
762 473: 298-307, 2011.

- 763 35. **Carmeliet P, and Jain RK.** Principles and mechanisms of vessel normalization for cancer and other
764 angiogenic diseases. *Nat Rev Drug Discov* 10: 417-427, 2011.
- 765 36. **Carnally SM, Johannessen M, Henderson RM, Jackson MB, and Edwardson JM.** Demonstration of
766 a direct interaction between sigma-1 receptors and acid-sensing ion channels. *Biophys J* 98: 1182-1191, 2010.
- 767 37. **Chantome A, Girault A, Potier M, Collin C, Vaudin P, Pages J-C, Vandier C, and Joulin V.**
768 KCa2.3 channel-dependent hyperpolarization increases melanoma cell motility. *Experimental Cell Research*
769 315: 3620-3630, 2009.
- 770 38. **Chappell JC, Taylor SM, Ferrara N, and Bautch VL.** Local guidance of emerging vessel sprouts
771 requires soluble Flt-1. *Dev Cell* 17: 377-386, 2009.
- 772 39. **Chen CH, Lin H, Chuang SM, Lin SY, and Chen JJ.** Acidic stress facilitates tyrosine
773 phosphorylation of HLJ1 to associate with actin cytoskeleton in lung cancer cells. *Exp Cell Res* 316: 2910-2921,
774 2010.
- 775 40. **Chen KD, Li YS, Kim M, Li S, Yuan S, Chien S, and Shyy JY.** Mechanotransduction in response to
776 shear stress. Roles of receptor tyrosine kinases, integrins, and Shc. *J Biol Chem* 274: 18393-18400, 1999.
- 777 41. **Chen M, Wang WP, Jia WR, Tang L, Wang Y, Zhan WW, and Fei XC.** Three-dimensional
778 contrast-enhanced sonography in the assessment of breast tumor angiogenesis: correlation with microvessel
779 density and vascular endothelial growth factor expression. *J Ultrasound Med* 33: 835-846, 2014.
- 780 42. **Cheng HW, James AF, Foster RR, Hancox JC, and Bates DO.** VEGF activates receptor-operated
781 cation channels in human microvascular endothelial cells. *Arterioscler Thromb Vasc Biol* 26: 1768-1776, 2006.
- 782 43. **Chetham PM, Babál P, Bridges JP, Moore TM, and Stevens T.** Segmental regulation of pulmonary
783 vascular permeability by store-operated Ca²⁺ entry. *Am J Physiol* 276: L41-50, 1999.
- 784 44. **Chouaib S, Messai Y, Couve S, Escudier B, Hasmim M, and Noman MZ.** Hypoxia promotes tumor
785 growth in linking angiogenesis to immune escape. *Front Immunol* 3: 21, 2012.
- 786 45. **Chung AS, Lee J, and Ferrara N.** Targeting the tumour vasculature: insights from physiological
787 angiogenesis. *Nat Rev Cancer* 10: 505-514, 2010.
- 788 46. **Crociani O, Zanieri F, Pillozzi S, Lastraioli E, Stefanini M, Fiore A, Fortunato A, D'Amico M,
789 Masselli M, De Lorenzo E, Gasparoli L, Chiu M, Bussolati O, Becchetti A, and Arcangeli A.** hERG1
790 channels modulate integrin signaling to trigger angiogenesis and tumor progression in colorectal cancer. *Sci Rep*
791 3: 3308, 2013.
- 792 47. **Crottès D, Martial S, Rapetti-Mauss R, Pisani DF, Loriol C, Pellissier B, Martin P, Chevet E,
793 Borgese F, and Soriani O.** Sig1R protein regulates hERG channel expression through a post-translational
794 mechanism in leukemic cells. *J Biol Chem* 286: 27947-27958, 2011.
- 795 48. **Crottès D, Rapetti-Mauss R, Alcaraz-Perez F, Tichet M, Gariano G, Martial S, Guizouarn H,
796 Pellissier B, Loubat A, Popa A, Paquet A, Presta M, Tartare-Deckert S, Cayuela ML, Martin P, Borgese
797 F, and Soriani O.** SigmaR1 regulates membrane electrical activity in response to extracellular matrix
798 stimulation to drive cancer cell invasiveness. *Cancer Res* 2015.
- 799 49. **D'Amico M, Gasparoli L, and Arcangeli A.** Potassium channels: novel emerging biomarkers and
800 targets for therapy in cancer. *Recent Pat Anticancer Drug Discov* 8: 53-65, 2013.
- 801 50. **Damaghi M, Wojtkowiak JW, and Gillies RJ.** pH sensing and regulation in cancer. *Front Physiol* 4:
802 370, 2013.
- 803 51. **Daschil N, Kniewallner KM, Obermair GJ, Hutter-Paier B, Windisch M, Marksteiner J, and
804 Humpel C.** L-type calcium channel blockers and substance P induce angiogenesis of cortical vessels associated
805 with beta-amyloid plaques in an Alzheimer mouse model. *Neurobiol Aging* 36: 1333-1341, 2015.
- 806 52. **Davis FM, Azimi I, Faville RA, Peters AA, Jalink K, Putney JW, Goodhill GJ, Thompson EW,
807 Roberts-Thomson SJ, and Monteith GR.** Induction of epithelial-mesenchymal transition (EMT) in breast
808 cancer cells is calcium signal dependent. *Oncogene* 33: 2307-2316, 2014.
- 809 53. **Davis GE, Stratman AN, Sacharidou A, and Koh W.** Molecular basis for endothelial lumen
810 formation and tubulogenesis during vasculogenesis and angiogenic sprouting. *Int Rev Cell Mol Biol* 288: 101-
811 165, 2011.
- 812 54. **De Bock K, De Smet F, Leite De Oliveira R, Anthonis K, and Carmeliet P.** Endothelial oxygen
813 sensors regulate tumor vessel abnormalization by instructing phalanx endothelial cells. *J Mol Med (Berl)* 87:
814 561-569, 2009.
- 815 55. **De Saedeleer CJ, Porporato PE, Copetti T, Pérez-Escuredo J, Payen VL, Brisson L, Feron O, and
816 Sonveaux P.** Glucose deprivation increases monocarboxylate transporter 1 (MCT1) expression and MCT1-
817 dependent tumor cell migration. *Oncogene* 33: 4060-4068, 2014.

- 818 56. **Delaunay A, Gasull X, Salinas M, Noël J, Friend V, Lingueglia E, and Deval E.** Human ASIC3
819 channel dynamically adapts its activity to sense the extracellular pH in both acidic and alkaline directions. *Proc*
820 *Natl Acad Sci U S A* 109: 13124-13129, 2012.
- 821 57. **Deval E, Gasull X, Noël J, Salinas M, Baron A, Diochot S, and Lingueglia E.** Acid-sensing ion
822 channels (ASICs): pharmacology and implication in pain. *Pharmacol Ther* 128: 549-558, 2010.
- 823 58. **Dewhirst MW, Cao Y, and Moeller B.** Cycling hypoxia and free radicals regulate angiogenesis and
824 radiotherapy response. *Nat Rev Cancer* 8: 425-437, 2008.
- 825 59. **Dhup S, Dadhich RK, Porporato PE, and Sonveaux P.** Multiple biological activities of lactic acid in
826 cancer: influences on tumor growth, angiogenesis and metastasis. *Curr Pharm Des* 18: 1319-1330, 2012.
- 827 60. **Do W, Herrera C, Mighty J, Shumskaya M, Redenti SM, and Sauane M.** Sigma 1 Receptor plays a
828 prominent role in IL-24-induced cancer-specific apoptosis. *Biochem Biophys Res Commun* 439: 215-220, 2013.
- 829 61. **Downie BR, Sánchez A, Knötgen H, Contreras-Jurado C, Gymnopoulos M, Weber C, Stühmer W,
830 and Pardo LA.** Eag1 expression interferes with hypoxia homeostasis and induces angiogenesis in tumors. *J*
831 *Biol Chem* 283: 36234-36240, 2008.
- 832 62. **Doyen J, Trastour C, Ettore F, Peyrottes I, Toussant N, Gal J, Ilc K, Roux D, Parks SK, Ferrero
833 JM, and Pouysségur J.** Expression of the hypoxia-inducible monocarboxylate transporter MCT4 is increased
834 in triple negative breast cancer and correlates independently with clinical outcome. *Biochem Biophys Res*
835 *Commun* 451: 54-61, 2014.
- 836 63. **Du GJ, Li JH, Liu WJ, Liu YH, Zhao B, Li HR, Hou XD, Li H, Qi XX, and Duan YJ.** The
837 combination of TRPM8 and TRPA1 expression causes an invasive phenotype in lung cancer. *Tumour Biol* 35:
838 1251-1261, 2014.
- 839 64. **Dubois L, Lieuwes NG, Maresca A, Thiry A, Supuran CT, Scozzafava A, Wouters BG, and
840 Lambin P.** Imaging of CA IX with fluorescent labelled sulfonamides distinguishes hypoxic and (re)-
841 oxygenated cells in a xenograft tumour model. *Radiother Oncol* 92: 423-428, 2009.
- 842 65. **Duncan RR, Westwood PK, Boyd A, and Ashley RH.** Rat brain p64H1, expression of a new member
843 of the p64 chloride channel protein family in endoplasmic reticulum. *J Biol Chem* 272: 23880-23886, 1997.
- 844 66. **El Hindy N, Bankfalvi A, Herring A, Adamzik M, Lambertz N, Zhu Y, Siffert W, Sure U, and
845 Sandalcioglu IE.** Correlation of aquaporin-1 water channel protein expression with tumor angiogenesis in
846 human astrocytoma. *Anticancer Res* 33: 609-613, 2013.
- 847 67. **Elble RC, and Pauli BU.** Tumor suppression by a proapoptotic calcium-activated chloride channel in
848 mammary epithelium. *J Biol Chem* 276: 40510-40517, 2001.
- 849 68. **Esteva-Font C, Jin BJ, and Verkman AS.** Aquaporin-1 gene deletion reduces breast tumor growth and
850 lung metastasis in tumor-producing MMTV-PyVT mice. *FASEB J* 28: 1446-1453, 2014.
- 851 69. **Fais S, De Milito A, You H, and Qin W.** Targeting vacuolar H⁺-ATPases as a new strategy against
852 cancer. *Cancer Res* 67: 10627-10630, 2007.
- 853 70. **Fan TP, Jaggar R, and Bicknell R.** Controlling the vasculature: angiogenesis, anti-angiogenesis and
854 vascular targeting of gene therapy. *Trends Pharmacol Sci* 16: 57-66, 1995.
- 855 71. **Fantozzi I, Zhang S, Platoshyn O, Remillard CV, Cowling RT, and Yuan JX.** Hypoxia increases
856 AP-1 binding activity by enhancing capacitative Ca²⁺ entry in human pulmonary artery endothelial cells. *Am J*
857 *Physiol Lung Cell Mol Physiol* 285: L1233-1245, 2003.
- 858 72. **Fearnley GW, Bruns AF, Wheatcroft SB, and Ponnambalam S.** VEGF-A isoform-specific
859 regulation of calcium ion flux, transcriptional activation and endothelial cell migration. *Biol Open* 2015.
- 860 73. **Feng X, Jia S, Martin TA, and Jiang WG.** Regulation and involvement in cancer and pathological
861 conditions of MAGI1, a tight junction protein. *Anticancer Res* 34: 3251-3256, 2014.
- 862 74. **Fiore A, Carraresi L, Morabito A, Polvani S, Fortunato A, Lastraioli E, Femia AP, De Lorenzo E,
863 Caderni G, and Arcangeli A.** Characterization of hERG1 channel role in mouse colorectal carcinogenesis.
864 *Cancer Med* 2: 583-594, 2013.
- 865 75. **Fiorio Pla A, Avanzato D, Munaron L, and Ambudkar IS.** Ion channels and transporters in cancer. 6.
866 Vascularizing the tumor: TRP channels as molecular targets. *Am J Physiol Cell Physiol* 302: C9-15, 2012.
- 867 76. **Fiorio Pla A, and Gkika D.** Emerging role of TRP channels in cell migration: from tumor
868 vascularization to metastasis. *Front Physiol* 4: 311, 2013.
- 869 77. **Folkman J.** Tumor angiogenesis: therapeutic implications. *N Engl J Med* 285: 1182-1186, 1971.
- 870 78. **Fraser SP, and Pardo LA.** Ion channels: functional expression and therapeutic potential in cancer.
871 Colloquium on Ion Channels and Cancer. *EMBO Rep* 9: 512-515, 2008.

- 872 79. **Friday E, Oliver R, Welbourne T, and Turturro F.** Role of epidermal growth factor receptor
873 (EGFR)-signaling versus cellular acidosis via Na⁺/H⁺ exchanger1(NHE1)-inhibition in troglitazone-induced
874 growth arrest of breast cancer-derived cells MCF-7. *Cell Physiol Biochem* 20: 751-762, 2007.
- 875 80. **Gao W, Chang G, Wang J, Jin W, Wang L, Lin Y, Li H, Ma L, Li Q, and Pang T.** Inhibition of
876 K562 leukemia angiogenesis and growth by selective Na⁽⁺⁾/H⁽⁺⁾ exchanger inhibitor cariporide through down-
877 regulation of pro-angiogenesis factor VEGF. *Leukemia Research* 2011.
- 878 81. **Gassmann P, and Haier J.** The tumor cell-host organ interface in the early onset of metastatic organ
879 colonisation. *Clin Exp Metastasis* 25: 171-181, 2008.
- 880 82. **Gatenby RA, and Gillies RJ.** A microenvironmental model of carcinogenesis. *Nat Rev Cancer* 8: 56-
881 61, 2008.
- 882 83. **Gatenby RA, and Gillies RJ.** Why do cancers have high aerobic glycolysis? *Nat Rev Cancer* 4: 891-
883 899, 2004.
- 884 84. **Gatenby RA, Smallbone K, Maini PK, Rose F, Averill J, Nagle RB, Worrall L, and Gillies RJ.**
885 Cellular adaptations to hypoxia and acidosis during somatic evolution of breast cancer. *Br J Cancer* 97: 646-
886 653, 2007.
- 887 85. **Gerhardt H, Golding M, Fruttiger M, Ruhrberg C, Lundkvist A, Abramsson A, Jeltsch M,
888 Mitchell C, Alitalo K, Shima D, and Betsholtz C.** VEGF guides angiogenic sprouting utilizing endothelial tip
889 cell filopodia. *J Cell Biol* 161: 1163-1177, 2003.
- 890 86. **Gillet L, Roger S, Besson P, Lecaille F, Gore J, Bognoux P, Lalmanach G, and Le Guennec JY.**
891 Voltage-gated Sodium Channel Activity Promotes Cysteine Cathepsin-dependent Invasiveness and Colony
892 Growth of Human Cancer Cells. *J Biol Chem* 284: 8680-8691, 2009.
- 893 87. **Gout S, and Huot J.** Role of cancer microenvironment in metastasis: focus on colon cancer. *Cancer*
894 *Microenviron* 1: 69-83, 2008.
- 895 88. **Greco MR, Antelmi E, Busco G, Guerra L, Rubino R, Casavola V, Reshkin SJ, and Cardone RA.**
896 Protease activity at invadopodial focal digestive areas is dependent on NHE1-driven acidic pH. *Oncol Rep* 31:
897 940-946, 2014.
- 898 89. **Grepin R, Guyot M, Jacquin M, Durivault J, Chamorey E, Sudaka A, Serdjebi C, Lacarelle B,
899 Scoazec JY, Negrier S, Simonnet H, and Pages G.** Acceleration of clear cell renal cell carcinoma growth in
900 mice following bevacizumab/Avastin treatment: the role of CXCL cytokines. *Oncogene* 31: 1683-1694, 2012.
- 901 90. **Grépin R, and Pagès G.** [The vascular endothelial growth factor (VEGF): a model of gene regulation
902 and a marker of tumour aggressiveness. An obvious therapeutic target?]. *J Soc Biol* 203: 181-192, 2009.
- 903 91. **Gupta GP, and Massagué J.** Cancer metastasis: building a framework. *Cell* 127: 679-695, 2006.
- 904 92. **Gupta SC, Singh R, Asters M, Liu J, Zhang X, Pabbidi MR, Watabe K, and Mo YY.** Regulation of
905 breast tumorigenesis through acid sensors. *Oncogene* 2015.
- 906 93. **Hackett NR, Shaykhiev R, Walters MS, Wang R, Zwick RK, Ferris B, Witover B, Salit J, and
907 Crystal RG.** The human airway epithelial basal cell transcriptome. *PLoS One* 6: e18378, 2011.
- 908 94. **Hanahan D, and Folkman J.** Patterns and emerging mechanisms of the angiogenic switch during
909 tumorigenesis. *Cell* 86: 353-364, 1996.
- 910 95. **Hanahan D, and Weinberg RA.** Hallmarks of cancer: the next generation. *Cell* 144: 646-674, 2011.
- 911 96. **Hanahan D, and Weinberg RA.** The hallmarks of cancer. *Cell* 100: 57-70, 2000.
- 912 97. **Harris AL.** Hypoxia--a key regulatory factor in tumour growth. *Nat Rev Cancer* 2: 38-47, 2002.
- 913 98. **Hashizume H, Baluk P, Morikawa S, McLean JW, Thurston G, Roberge S, Jain RK, and
914 McDonald DM.** Openings between defective endothelial cells explain tumor vessel leakiness. *Am J Pathol* 156:
915 1363-1380, 2000.
- 916 99. **Hayashi T, Rizzuto R, Hajnoczky G, and Su TP.** MAM: more than just a housekeeper. *Trends Cell*
917 *Biol* 19: 81-88, 2009.
- 918 100. **Hayashi T, and Su T-P.** Sigma-1 receptor chaperones at the ER-mitochondrion interface regulate
919 Ca⁽²⁺⁾ signaling and cell survival. *Cell* 131: 596-610, 2007.
- 920 101. **Hayashi T, and Su TP.** Intracellular dynamics of sigma-1 receptors (sigma(1) binding sites) in NG108-
921 15 cells. *J Pharmacol Exp Ther* 306: 726-733, 2003.
- 922 102. **Hiratsuka S, Minowa O, Kuno J, Noda T, and Shibuya M.** Flt-1 lacking the tyrosine kinase domain
923 is sufficient for normal development and angiogenesis in mice. *Proc Natl Acad Sci U S A* 95: 9349-9354, 1998.
- 924 103. **Hofman P, Cherfils-Vicini J, Bazin M, Ilie M, Juhel T, Hébuterne X, Gilson E, Schmid-Alliana A,
925 Boyer O, Adriouch S, and Vouret-Craviari V.** Genetic and pharmacological inactivation of the purinergic

- 926 P2RX7 receptor dampens inflammation but increases tumor incidence in a mouse model of colitis-associated
927 cancer. *Cancer Res* 75: 835-845, 2015.
- 928 104. **Holash J, Wiegand SJ, and Yancopoulos GD.** New model of tumor angiogenesis: dynamic balance
929 between vessel regression and growth mediated by angiopoietins and VEGF. *Oncogene* 18: 5356-5362, 1999.
- 930 105. **Huang WC, Swietach P, Vaughan-Jones RD, Ansorge O, and Glitsch MD.** Extracellular
931 acidification elicits spatially and temporally distinct Ca²⁺ signals. *Curr Biol* 18: 781-785, 2008.
- 932 106. **Huang Y, Goel S, Duda DG, Fukumura D, and Jain RK.** Vascular normalization as an emerging
933 strategy to enhance cancer immunotherapy. *Cancer Res* 73: 2943-2948, 2013.
- 934 107. **Huber MA, Kraut N, and Beug H.** Molecular requirements for epithelial-mesenchymal transition
935 during tumor progression. *Curr Opin Cell Biol* 17: 548-558, 2005.
- 936 108. **Issberner U, Reeh PW, and Steen KH.** Pain due to tissue acidosis: a mechanism for inflammatory and
937 ischemic myalgia? *Neurosci Lett* 208: 191-194, 1996.
- 938 109. **Jain RK.** Normalizing tumor microenvironment to treat cancer: bench to bedside to biomarkers. *J Clin*
939 *Oncol* 31: 2205-2218, 2013.
- 940 110. **Jelassi B, Chantôme A, Alcaraz-Pérez F, Baroja-Mazo A, Cayuela ML, Pelegrin P, Surprenant A,**
941 **and Roger S.** P2X(7) receptor activation enhances SK3 channels- and cysteine cathepsin-dependent cancer cells
942 invasiveness. *Oncogene* 30: 2108-2122, 2011.
- 943 111. **Jho D, Mehta D, Ahmmed G, Gao XP, Tiruppathi C, Broman M, and Malik AB.** Angiopoietin-1
944 opposes VEGF-induced increase in endothelial permeability by inhibiting TRPC1-dependent Ca²⁺ influx. *Circ*
945 *Res* 96: 1282-1290, 2005.
- 946 112. **Jin C, Ye QH, Yuan FL, Gu YL, Li JP, Shi YH, Shen XM, Bo-Liu, and Lin ZH.** Involvement of
947 acid-sensing ion channel 1 α in hepatic carcinoma cell migration and invasion. *Tumour Biol* 2015.
- 948 113. **John CS, Bowen WD, Varma VM, McAfee JG, and Moody TW.** Sigma receptors are expressed in
949 human non-small cell lung carcinoma. *Life Sci* 56: 2385-2392, 1995.
- 950 114. **Jäger H, Dreker T, Buck A, Giehl K, Gress T, and Grissmer S.** Blockage of intermediate-
951 conductance Ca²⁺-activated K⁺ channels inhibit human pancreatic cancer cell growth in vitro. *Mol Pharmacol*
952 65: 630-638, 2004.
- 953 115. **Kaelin WG.** The von Hippel-Lindau tumour suppressor protein: O₂ sensing and cancer. *Nat Rev*
954 *Cancer* 8: 865-873, 2008.
- 955 116. **Kalluri R, and Weinberg RA.** The basics of epithelial-mesenchymal transition. *J Clin Invest* 119:
956 1420-1428, 2009.
- 957 117. **Kamei M, Saunders WB, Bayless KJ, Dye L, Davis GE, and Weinstein BM.** Endothelial tubes
958 assemble from intracellular vacuoles in vivo. *Nature* 442: 453-456, 2006.
- 959 118. **Kamouchi M, Droogmans G, and Nilius B.** Membrane potential as a modulator of the free
960 intracellular Ca²⁺ concentration in agonist-activated endothelial cells. *Gen Physiol Biophys* 18: 199-208, 1999.
- 961 119. **Kaneko Y, and Szallasi A.** Transient receptor potential (TRP) channels: a clinical perspective. *Br J*
962 *Pharmacol* 171: 2474-2507, 2014.
- 963 120. **Kang BW, Kim JG, Lee SJ, Chae YS, Jeong JY, Yoon GS, Park SY, Kim HJ, Park JS, and Choi**
964 **GS.** Expression of aquaporin-1, aquaporin-3, and aquaporin-5 correlates with nodal metastasis in colon cancer.
965 *Oncology* 88: 369-376, 2015.
- 966 121. **Kato Y, Ozawa S, Tsukuda M, Kubota E, Miyazaki K, St-Pierre Y, and Hata R.** Acidic
967 extracellular pH increases calcium influx-triggered phospholipase D activity along with acidic
968 sphingomyelinase activation to induce matrix metalloproteinase-9 expression in mouse metastatic melanoma.
969 *FEBS J* 274: 3171-3183, 2007.
- 970 122. **Kerbrel RS.** Tumor angiogenesis: past, present and the near future. *Carcinogenesis* 21: 505-515, 2000.
- 971 123. **Kim KY, Kim SO, Lim H, Yoo SE, and Hong KW.** KR-31372 inhibits KDR/Flk-1 tyrosine
972 phosphorylation via K⁺(ATP) channel opening in its antiangiogenic effect. *Eur J Pharmacol* 465: 219-228,
973 2003.
- 974 124. **Kim M, Jiang LH, Wilson HL, North RA, and Surprenant A.** Proteomic and functional evidence for
975 a P2X7 receptor signalling complex. *EMBO J* 20: 6347-6358, 2001.
- 976 125. **Korc M, and Friesel RE.** The role of fibroblast growth factors in tumor growth. *Curr Cancer Drug*
977 *Targets* 9: 639-651, 2009.
- 978 126. **Kourrich S, Su TP, Fujimoto M, and Bonci A.** The sigma-1 receptor: roles in neuronal plasticity and
979 disease. *Trends Neurosci* 35: 762-771, 2012.
- 980 127. **Kunzelmann K.** Ion channels and cancer. *Journal of Membrane Biology* 205: 159-173, 2005.

- 981 128. **Kweon HJ, Yu SY, Kim DI, and Suh BC.** Differential regulation of proton-sensitive ion channels by
982 phospholipids: a comparative study between ASICs and TRPV1. *PLoS One* 10: e0122014, 2015.
- 983 129. **Lai W, Liu L, Zeng Y, Wu H, Xu H, Chen S, and Chu Z.** KCNN4 channels participate in the EMT
984 induced by PRL-3 in colorectal cancer. *Med Oncol* 30: 566, 2013.
- 985 130. **Lang F, Foller M, Lang KS, Lang PA, Ritter M, Gulbins E, Vereninov A, and Huber SM.** Ion
986 channels in cell proliferation and apoptotic cell death. *Journal of Membrane Biology* 205: 147-157, 2005.
- 987 131. **Larrinaga G, Sanz B, Pérez I, Blanco L, Cándenas ML, Pinto FM, Gil J, and López JI.**
988 Cannabinoid CB₁ receptor is downregulated in clear cell renal cell carcinoma. *J Histochem Cytochem* 58: 1129-
989 1134, 2010.
- 990 132. **Lastraioli E, Bencini L, Bianchini E, Romoli MR, Crociani O, Giommoni E, Messerini L,**
991 **Gasperoni S, Moretti R, Di Costanzo F, Boni L, and Arcangeli A.** hERG1 Channels and Glut-1 as
992 Independent Prognostic Indicators of Worse Outcome in Stage I and II Colorectal Cancer: A Pilot Study. *Transl*
993 *Oncol* 5: 105-112, 2012.
- 994 133. **Lastraioli E, Guasti L, Crociani O, Polvani S, Hofmann G, Witchel H, Bencini L, Calistri M,**
995 **Messerini L, Scatizzi M, Moretti R, Wanke E, Olivotto M, Mugnai G, and Arcangeli A.** hERG1 gene and
996 HERG1 protein are overexpressed in colorectal cancers and regulate cell invasion of tumor cells. *Cancer*
997 *Research* 64: 606-611, 2004.
- 998 134. **Lee HJ, Li N, Evans SM, Diaz MF, and Wenzel PL.** Biomechanical force in blood development:
999 extrinsic physical cues drive pro-hematopoietic signaling. *Differentiation* 86: 92-103, 2013.
- 1000 135. **Lee SK, Dawson J, Lee JA, Osman G, Levitin MO, Guzel RM, and Djamgoz MB.** Management of
1001 cancer pain: 1. Wider implications of orthodox analgesics. *Int J Gen Med* 7: 49-58, 2014.
- 1002 136. **Lee TH, Seng S, Sekine M, Hinton C, Fu Y, Avraham HK, and Avraham S.** Vascular endothelial
1003 growth factor mediates intracrine survival in human breast carcinoma cells through internally expressed
1004 VEGFR1/FLT1. *PLoS Med* 4: e186, 2007.
- 1005 137. **Lehen'kyi V, Shapovalov G, Skryma R, and Prevarskaya N.** Ion channels and transporters in
1006 cancer. 5. Ion channels in control of cancer and cell apoptosis. *Am J Physiol Cell Physiol* 301: C1281-1289,
1007 2011.
- 1008 138. **Lemonnier L, Lazarenko R, Shuba Y, Thebault S, Roudbaraki M, Lepage G, Prevarskaya N, and**
1009 **Skryma R.** Alterations in the regulatory volume decrease (RVD) and swelling-activated Cl⁻ current associated
1010 with neuroendocrine differentiation of prostate cancer epithelial cells. *Endocr Relat Cancer* 12: 335-349, 2005.
- 1011 139. **Li J, Cubbon RM, Wilson LA, Amer MS, McKeown L, Hou B, Majeed Y, Tumova S, Seymour**
1012 **VA, Taylor H, Stacey M, O'Regan D, Foster R, Porter KE, Kearney MT, and Beech DJ.** Orail and CRAC
1013 channel dependence of VEGF-activated Ca²⁺ entry and endothelial tube formation. *Circ Res* 108: 1190-1198,
1014 2011.
- 1015 140. **Liu JP, Nakakura T, Tomura H, Tobo M, Mogi C, Wang JQ, He XD, Takano M, Damirin A,**
1016 **Komachi M, Sato K, and Okajima F.** Each one of certain histidine residues in G-protein-coupled receptor
1017 GPR4 is critical for extracellular proton-induced stimulation of multiple G-protein-signaling pathways.
1018 *Pharmacol Res* 61: 499-505, 2010.
- 1019 141. **Liu Y, Zhao L, Ma W, Cao X, Chen H, Feng D, Liang J, Yin K, and Jiang X.** The Blockage of
1020 KCa3.1 Channel Inhibited Proliferation, Migration and Promoted Apoptosis of Human Hepatocellular
1021 Carcinoma Cells. *J Cancer* 6: 643-651, 2015.
- 1022 142. **Lkhagvadorj S, Oh SS, Lee MR, Jung JH, Chung HC, Cha SK, and Eom M.** VEGFR-1 Expression
1023 Relates to Fuhrman Nuclear Grade of Clear Cell Renal Cell Carcinoma. *J Lifestyle Med* 4: 64-70, 2014.
- 1024 143. **Lubarsky B, and Krasnow MA.** Tube morphogenesis: making and shaping biological tubes. *Cell* 112:
1025 19-28, 2003.
- 1026 144. **Lynn KD, Roland CL, and Brekken RA.** VEGF and pleiotrophin modulate the immune profile of
1027 breast cancer. *Cancers (Basel)* 2: 970-988, 2010.
- 1028 145. **Maertens C, Wei L, Tytgat J, Droogmans G, and Nilius B.** Chlorotoxin does not inhibit volume-
1029 regulated, calcium-activated and cyclic AMP-activated chloride channels. *Br J Pharmacol* 129: 791-801, 2000.
- 1030 146. **Manolopoulos VG, Liekens S, Koolwijk P, Voets T, Peters E, Droogmans G, Lelkes PI, De Clercq**
1031 **E, and Nilius B.** Inhibition of angiogenesis by blockers of volume-regulated anion channels. *Gen Pharmacol*
1032 34: 107-116, 2000.

- 1033 147. **Martial S, Giorgelli J-L, Renaudo A, Derijard B, and Soriani O.** SP600125 inhibits Kv channels
1034 through a JNK-independent pathway in cancer cells. *Biochemical and Biophysical Research Communications*
1035 366: 944-950, 2008.
- 1036 148. **Maurice T, and Su T-P.** The pharmacology of sigma-1 receptors. *Pharmacology & Therapeutics* 124:
1037 195-206, 2009.
- 1038 149. **Mavlyutov TA, and Ruoho AE.** Ligand-dependent localization and intracellular stability of sigma-1
1039 receptors in CHO-K1 cells. *Journal of Molecular Signaling* 2: 8, 2007.
- 1040 150. **McCarty MF, and Whitaker J.** Manipulating tumor acidification as a cancer treatment strategy. *Altern*
1041 *Med Rev* 15: 264-272, 2010.
- 1042 151. **Meunier J, and Hayashi T.** Sigma-1 receptors regulate Bcl-2 expression by reactive oxygen species-
1043 dependent transcriptional regulation of nuclear factor kappaB. *Journal of Pharmacology and Experimental*
1044 *Therapeutics* 332: 388-397, 2010.
- 1045 152. **Millership JE, Devor DC, Hamilton KL, Balut CM, Bruce JI, and Fearon IM.** Calcium-activated
1046 K⁺ channels increase cell proliferation independent of K⁺ conductance. *Am J Physiol Cell Physiol* 300: C792-
1047 802, 2011.
- 1048 153. **Miternique-Grosse A, Griffon C, Siegel L, Neuville A, Weltin D, and Stephan D.** Antiangiogenic
1049 effects of spironolactone and other potassium-sparing diuretics in human umbilical vein endothelial cells and in
1050 fibrin gel chambers implanted in rats. *J Hypertens* 24: 2207-2213, 2006.
- 1051 154. **Mittal M, Urao N, Hecquet CM, Zhang M, Sudhahar V, Gao XP, Komarova Y, Ushio-Fukai M,**
1052 **and Malik AB.** Novel role of reactive oxygen species-activated Trp melastatin channel-2 in mediating
1053 angiogenesis and postischemic neovascularization. *Arterioscler Thromb Vasc Biol* 35: 877-887, 2015.
- 1054 155. **Moccia F, Dragoni S, Lodola F, Bonetti E, Bottino C, Guerra G, Laforenza U, Rosti V, and Tanzi**
1055 **F.** Store-dependent Ca(2⁺) entry in endothelial progenitor cells as a perspective tool to enhance cell-based
1056 therapy and adverse tumour vascularization. *Curr Med Chem* 19: 5802-5818, 2012.
- 1057 156. **Moccia F, and Poletto V.** May the remodeling of the Ca(2⁺) toolkit in endothelial progenitor cells
1058 derived from cancer patients suggest alternative targets for anti-angiogenic treatment? *Biochim Biophys Acta*
1059 2014.
- 1060 157. **Moenner M, Pluquet O, Bouchecareilh M, and Chevet E.** Integrated endoplasmic reticulum stress
1061 responses in cancer. *Cancer Res* 67: 10631-10634, 2007.
- 1062 158. **Monet M, Lehen'kyi V, Gackiere F, Firlej V, Vandenberghe M, Roudbaraki M, Gkika D, Pournier**
1063 **A, Bidaux G, Slomianny C, Delcourt P, Rassendren F, Bergerat JP, Ceraline J, Cabon F, Humez S, and**
1064 **Prevarskaya N.** Role of cationic channel TRPV2 in promoting prostate cancer migration and progression to
1065 androgen resistance. *Cancer Res* 70: 1225-1235, 2010.
- 1066 159. **Morelli MB, Amantini C, Nabissi M, Liberati S, Cardinali C, Farfariello V, Tomassoni D, Quaglia**
1067 **W, Piergentili A, Bonifazi A, Del Bello F, Santoni M, Mammana G, Servi L, Filosa A, Gismondi A, and**
1068 **Santoni G.** Cross-talk between alpha1D-adrenoceptors and transient receptor potential vanilloid type 1 triggers
1069 prostate cancer cell proliferation. *BMC Cancer* 14: 921, 2014.
- 1070 160. **Mori T, Hayashi T, Hayashi E, and Su TP.** Sigma-1 receptor chaperone at the ER-mitochondrion
1071 interface mediates the mitochondrion-ER-nucleus signaling for cellular survival. *PLoS One* 8: e76941, 2013.
- 1072 161. **Motiani RK, Hyzinski-Garcia MC, Zhang X, Henkel MM, Abdullaev IF, Kuo YH, Matrougui K,**
1073 **Mongin AA, and Trebak M.** STIM1 and Orai1 mediate CRAC channel activity and are essential for human
1074 glioblastoma invasion. *Pflugers Arch* 465: 1249-1260, 2013.
- 1075 162. **Munaron L.** Systems biology of ion channels and transporters in tumor angiogenesis: An omics view.
1076 *Biochim Biophys Acta* 2014.
- 1077 163. **Munaron L, Genova T, Avanzato D, Antoniotti S, and Fiorio Pla A.** Targeting calcium channels to
1078 block tumor vascularization. *Recent Pat Anticancer Drug Discov* 8: 27-37, 2013.
- 1079 164. **Nakajima S, Doi R, Toyoda E, Tsuji S, Wada M, Koizumi M, Tulachan SS, Ito D, Kami K, Mori**
1080 **T, Kawaguchi Y, Fujimoto K, Hosotani R, and Imamura M.** N-cadherin expression and epithelial-
1081 mesenchymal transition in pancreatic carcinoma. *Clin Cancer Res* 10: 4125-4133, 2004.
- 1082 165. **Nilius B, and Droogmans G.** Ion channels and their functional role in vascular endothelium. *Physiol*
1083 *Rev* 81: 1415-1459, 2001.
- 1084 166. **Nilius B, Eggermont J, Voets T, Buyse G, Manolopoulos V, and Droogmans G.** Properties of
1085 volume-regulated anion channels in mammalian cells. *Prog Biophys Mol Biol* 68: 69-119, 1997.
- 1086 167. **Nilius B, Eggermont J, Voets T, and Droogmans G.** Volume-activated Cl⁻ channels. *Gen Pharmacol*
1087 27: 1131-1140, 1996.

- 1088 168. **O'Grady SM, and Lee SY.** Molecular diversity and function of voltage-gated (Kv) potassium channels
1089 in epithelial cells. *International Journal of Biochemistry and Cell Biology* 37: 1578-1594, 2005.
- 1090 169. **Okamoto Y, Ohkubo T, Ikebe T, and Yamazaki J.** Blockade of TRPM8 activity reduces the invasion
1091 potential of oral squamous carcinoma cell lines. *Int J Oncol* 40: 1431-1440, 2012.
- 1092 170. **Olszewski U, Hlozek M, and Hamilton G.** Activation of Na⁺/H⁺ exchanger 1 by neurotensin signaling
1093 in pancreatic cancer cell lines. *Biochem Biophys Res Commun* 393: 414-419, 2010.
- 1094 171. **Oosterwijk E, and Gillies RJ.** Targeting ion transport in cancer. *Philos Trans R Soc Lond B Biol Sci*
1095 369: 20130107, 2014.
- 1096 172. **Pagès G, Milanini J, Richard DE, Berra E, Gothié E, Viñals F, and Pouyssegur J.** Signaling
1097 angiogenesis via p42/p44 MAP kinase cascade. *Ann N Y Acad Sci* 902: 187-200, 2000.
- 1098 173. **Palmer CP, Mahen R, Schnell E, Djamgoz MB, and Aydar E.** Sigma-1 receptors bind cholesterol
1099 and remodel lipid rafts in breast cancer cell lines. *Cancer Res* 67: 11166-11175, 2007.
- 1100 174. **Panyi G, Beeton C, and Felipe A.** Ion channels and anti-cancer immunity. *Philos Trans R Soc Lond B*
1101 *Biol Sci* 369: 20130106, 2014.
- 1102 175. **Papandreou I, Cairns RA, Fontana L, Lim AL, and Denko NC.** HIF-1 mediates adaptation to
1103 hypoxia by actively downregulating mitochondrial oxygen consumption. *Cell Metab* 3: 187-197, 2006.
- 1104 176. **Papetti M, and Herman IM.** Mechanisms of normal and tumor-derived angiogenesis. *Am J Physiol*
1105 *Cell Physiol* 282: C947-970, 2002.
- 1106 177. **Pardo LA, Gómez-Varela D, Major F, Sansuk K, Leurs R, Downie BR, Tietze LF, and Stühmer**
1107 **W.** Approaches targeting K(V)10.1 open a novel window for cancer diagnosis and therapy. *Curr Med Chem* 19:
1108 675-682, 2012.
- 1109 178. **Paria BC, Vogel SM, Ahmmed GU, Alamgir S, Shroff J, Malik AB, and Tiruppathi C.** Tumor
1110 necrosis factor-alpha-induced TRPC1 expression amplifies store-operated Ca²⁺ influx and endothelial
1111 permeability. *Am J Physiol Lung Cell Mol Physiol* 287: L1303-1313, 2004.
- 1112 179. **Parker LH, Schmidt M, Jin SW, Gray AM, Beis D, Pham T, Frantz G, Palmieri S, Hillan K,**
1113 **Stainier DY, De Sauvage FJ, and Ye W.** The endothelial-cell-derived secreted factor Eglf7 regulates vascular
1114 tube formation. *Nature* 428: 754-758, 2004.
- 1115 180. **Parks SK, and Pouyssegur J.** The Na(+) /HCO₃(-) Co-Transporter SLC4A4 Plays a Role in Growth
1116 and Migration of Colon and Breast Cancer Cells. *J Cell Physiol* 230: 1954-1963, 2015.
- 1117 181. **Pellegatti P, Raffaghello L, Bianchi G, Piccardi F, Pistoia V, and Di Virgilio F.** Increased level of
1118 extracellular ATP at tumor sites: in vivo imaging with plasma membrane luciferase. *PLoS One* 3: e2599, 2008.
- 1119 182. **Pertwee RG, Howlett AC, Abood ME, Alexander SP, Di Marzo V, Elphick MR, Greasley PJ,**
1120 **Hansen HS, Kunos G, Mackie K, Mechoulam R, and Ross RA.** International Union of Basic and Clinical
1121 Pharmacology. LXXIX. Cannabinoid receptors and their ligands: beyond CB₁ and CB₂. *Pharmacol Rev* 62:
1122 588-631, 2010.
- 1123 183. **Pillozzi S, Brizzi MF, Bernabei PA, Bartolozzi B, Caporale R, Basile V, Boddi V, Pegoraro L,**
1124 **Becchetti A, and Arcangeli A.** VEGFR-1 (FLT-1), beta1 integrin, and hERG K⁺ channel for a macromolecular
1125 signaling complex in acute myeloid leukemia: role in cell migration and clinical outcome. *Blood* 110: 1238-
1126 1250, 2007.
- 1127 184. **Pinheiro C, Longatto-Filho A, Ferreira L, Pereira SM, Etlinger D, Moreira MA, Jubé LF, Queiroz**
1128 **GS, Schmitt F, and Baltazar F.** Increasing expression of monocarboxylate transporters 1 and 4 along
1129 progression to invasive cervical carcinoma. *Int J Gynecol Pathol* 27: 568-574, 2008.
- 1130 185. **Plate KH, Breier G, Millauer B, Ullrich A, and Risau W.** Up-regulation of vascular endothelial
1131 growth factor and its cognate receptors in a rat glioma model of tumor angiogenesis. *Cancer Res* 53: 5822-5827,
1132 1993.
- 1133 186. **Pocock TM, Foster RR, and Bates DO.** Evidence of a role for TRPC channels in VEGF-mediated
1134 increased vascular permeability in vivo. *Am J Physiol Heart Circ Physiol* 286: H1015-1026, 2004.
- 1135 187. **Poincloux R, Lizárraga F, and Chavrier P.** Matrix invasion by tumour cells: a focus on MT1-MMP
1136 trafficking to invadopodia. *J Cell Sci* 122: 3015-3024, 2009.
- 1137 188. **Polette M, Nawrocki-Raby B, Gilles C, Clavel C, and Birembaut P.** Tumour invasion and matrix
1138 metalloproteinases. *Crit Rev Oncol Hematol* 49: 179-186, 2004.
- 1139 189. **Poole TJ, Finkelstein EB, and Cox CM.** The role of FGF and VEGF in angioblast induction and
1140 migration during vascular development. *Dev Dyn* 220: 1-17, 2001.

1141 190. **Potente M, Gerhardt H, and Carmeliet P.** Basic and therapeutic aspects of angiogenesis. *Cell* 146:
1142 873-887, 2011.

1143 191. **Potier M, Joulin V, Roger S, Besson P, Jourdan M-L, Le Guennec J-Y, Bougnoux P, and Vandier**
1144 **C.** Identification of SK3 channel as a new mediator of breast cancer cell migration. *Molecular Cancer*
1145 *Therapeutics* 5: 2946-2953, 2006.

1146 192. **Premkumar LS, and Abooj M.** TRP channels and analgesia. *Life Sci* 92: 415-424, 2013.

1147 193. **Qiu Y, Li WH, Zhang HQ, Liu Y, Tian XX, and Fang WG.** P2X7 mediates ATP-driven invasiveness
1148 in prostate cancer cells. *PLoS One* 9: e114371, 2014.

1149 194. **Ranpura V, Hapani S, Chuang J, and Wu S.** Risk of cardiac ischemia and arterial thromboembolic
1150 events with the angiogenesis inhibitor bevacizumab in cancer patients: a meta-analysis of randomized controlled
1151 trials. *Acta Oncol* 49: 287-297, 2010.

1152 195. **Ranpura V, Hapani S, and Wu S.** Treatment-related mortality with bevacizumab in cancer patients: a
1153 meta-analysis. *JAMA* 305: 487-494, 2011.

1154 196. **Renaudo A, L'Hoste S, Guizouarn H, Borgese F, and Soriani O.** Cancer cell cycle modulated by a
1155 functional coupling between sigma-1 receptors and Cl⁻ channels. *Journal of Biological Chemistry* 282: 2259-
1156 2267, 2007.

1157 197. **Renaudo A, Watry V, Chassot A-A, Ponzio G, Ehrenfeld J, and Soriani O.** Inhibition of tumor cell
1158 proliferation by sigma Ligands is associated with K⁺ channel inhibition and p27kip1 accumulation. *Journal of*
1159 *Pharmacology and Experimental Therapeutics* 311: 1105-1114, 2004.

1160 198. **Reshkin SJ, Cardone RA, and Harguindey S.** Na⁺-H⁺ exchanger, pH regulation and cancer. *Recent*
1161 *Pat Anticancer Drug Discov* 8: 85-99, 2013.

1162 199. **Ribatti D, and Djonov V.** Intussusceptive microvascular growth in tumors. *Cancer Lett* 316: 126-131,
1163 2012.

1164 200. **Roger S, Potier M, Vandier C, Le Guennec JY, and Besson P.** Description and role in proliferation
1165 of iberiotoxin-sensitive currents in different human mammary epithelial normal and cancerous cells. *Biochim*
1166 *Biophys Acta* 1667: 190-199, 2004.

1167 201. **Rosenthal R, Heimann H, Agostini H, Martin G, Hansen LL, and Strauss O.** Ca²⁺ channels in
1168 retinal pigment epithelial cells regulate vascular endothelial growth factor secretion rates in health and disease.
1169 *Mol Vis* 13: 443-456, 2007.

1170 202. **Rouzaire-Dubois B, Ouanounou G, O'Regan S, and Dubois JM.** Sodium-dependent activity of
1171 aquaporin-1 in rat glioma cells: a new mechanism of cell volume regulation. *Pflugers Arch* 457: 1187-1198,
1172 2009.

1173 203. **Santoni G, and Farfariello V.** TRP channels and cancer: new targets for diagnosis and chemotherapy.
1174 *Endocr Metab Immune Disord Drug Targets* 11: 54-67, 2011.

1175 204. **Santoni G, Santoni M, and Nabissi M.** Functional role of T-type calcium channels in tumour growth
1176 and progression: prospective in cancer therapy. *Br J Pharmacol* 166: 1244-1246, 2012.

1177 205. **Sawano A, Iwai S, Sakurai Y, Ito M, Shitara K, Nakahata T, and Shibuya M.** Flt-1, vascular
1178 endothelial growth factor receptor 1, is a novel cell surface marker for the lineage of monocyte-macrophages in
1179 humans. *Blood* 97: 785-791, 2001.

1180 206. **Schneider L, Klausen TK, Stock C, Mally S, Christensen ST, Pedersen SF, Hoffmann EK, and**
1181 **Schwab A.** H-ras transformation sensitizes volume-activated anion channels and increases migratory activity of
1182 NIH3T3 fibroblasts. *Pflugers Arch* 455: 1055-1062, 2008.

1183 207. **Schonherr R.** Clinical relevance of ion channels for diagnosis and therapy of cancer. *Journal of*
1184 *Membrane Biology* 205: 175-184, 2005.

1185 208. **Schornack PA, and Gillies RJ.** Contributions of cell metabolism and H⁺ diffusion to the acidic pH of
1186 tumors. *Neoplasia* 5: 135-145, 2003.

1187 209. **Schreiber R.** Ca²⁺ signaling, intracellular pH and cell volume in cell proliferation. *Journal of*
1188 *Membrane Biology* 205: 129-137, 2005.

1189 210. **Schwab A, Nechyporuk-Zloy V, Fabian A, and Stock C.** Cells move when ions and water flow.
1190 *Pflugers Arch* 453: 421-432, 2007.

1191 211. **Shapovalov G, Lehen'kyi V, Skryma R, and Prevarskaya N.** TRP channels in cell survival and cell
1192 death in normal and transformed cells. *Cell Calcium* 50: 295-302, 2011.

1193 212. **Sharma SK, Chintala NK, Vadrevu SK, Patel J, Karbowniczek M, and Markiewski MM.**
1194 Pulmonary alveolar macrophages contribute to the premetastatic niche by suppressing antitumor T cell
1195 responses in the lungs. *J Immunol* 194: 5529-5538, 2015.

1196 213. **Sidani M, Wessels D, Mounieime G, Ghosh M, Goswami S, Sarmiento C, Wang W, Kuhl S, El-**
1197 **Sibai M, Backer JM, Eddy R, Soll D, and Condeelis J.** Cofilin determines the migration behavior and turning
1198 frequency of metastatic cancer cells. *J Cell Biol* 179: 777-791, 2007.

1199 214. **Smith GAM, Tsui H-W, Newell EW, Jiang X, Zhu X-P, Tsui FWL, and Schlichter LC.** Functional
1200 up-regulation of HERG K⁺ channels in neoplastic hematopoietic cells. *Journal of Biological Chemistry* 277:
1201 18528-18534, 2002.

1202 215. **Soroceanu L, Manning TJ, and Sontheimer H.** Modulation of glioma cell migration and invasion
1203 using Cl⁻ and K⁺ ion channel blockers. *J Neurosci* 19: 5942-5954, 1999.

1204 216. **Spinelli AM, González-Cobos JC, Zhang X, Motiani RK, Rowan S, Zhang W, Garrett J, Vincent**
1205 **PA, Matrougui K, Singer HA, and Trebak M.** Airway smooth muscle STIM1 and Orail are upregulated in
1206 asthmatic mice and mediate PDGF-activated SOCE, CRAC currents, proliferation, and migration. *Pflugers Arch*
1207 464: 481-492, 2012.

1208 217. **Stock C, and Schwab A.** Ion channels and transporters in metastasis. *Biochim Biophys Acta* 2014.

1209 218. **Stock C, and Schwab A.** Protons make tumor cells move like clockwork. *Pflugers Arch* 458: 981-992,
1210 2009.

1211 219. **Stylianopoulos T, and Jain RK.** Combining two strategies to improve perfusion and drug delivery in
1212 solid tumors. *Proc Natl Acad Sci U S A* 110: 18632-18637, 2013.

1213 220. **Su TP, Hayashi T, Maurice T, Buch S, and Ruoho AE.** The sigma-1 receptor chaperone as an inter-
1214 organelle signaling modulator. *Trends Pharmacol Sci* 31: 557-566, 2010.

1215 221. **Su TP, Hayashi T, and Vaupel DB.** When the endogenous hallucinogenic trace amine N,N-
1216 dimethyltryptamine meets the sigma-1 receptor. *Sci Signal* 2: pe12, 2009.

1217 222. **Svastova E, Witariski W, Csaderova L, Kosik I, Skvarkova L, Hulikova A, Zatovicova M,**
1218 **Barathova M, Kopacek J, Pastorek J, and Pastorekova S.** Carbonic anhydrase IX interacts with bicarbonate
1219 transporters in lamellipodia and increases cell migration via its catalytic domain. *J Biol Chem* 287: 3392-3402,
1220 2012.

1221 223. **Svastová E, Hulíková A, Rafajová M, Zat'ovicová M, Gibadulinová A, Casini A, Cecchi A,**
1222 **Scozzafava A, Supuran CT, Pastorek J, and Pastoreková S.** Hypoxia activates the capacity of tumor-
1223 associated carbonic anhydrase IX to acidify extracellular pH. *FEBS Lett* 577: 439-445, 2004.

1224 224. **Szigligeti P, Neumeier L, Duke E, Chougnat C, Takimoto K, Lee SM, Filipovich AH, and Conforti**
1225 **L.** Signalling during hypoxia in human T lymphocytes--critical role of the src protein tyrosine kinase p56Lck in
1226 the O₂ sensitivity of Kv1.3 channels. *Journal of Physiology* 573: 357-370, 2006.

1227 225. **Theriault RL.** Biology of bone metastases. *Cancer Control* 19: 92-101, 2012.

1228 226. **Thiery JP, and Sleeman JP.** Complex networks orchestrate epithelial-mesenchymal transitions. *Nat*
1229 *Rev Mol Cell Biol* 7: 131-142, 2006.

1230 227. **Toschi A, Lee E, Gadir N, Ohh M, and Foster DA.** Differential dependence of hypoxia-inducible
1231 factors 1 alpha and 2 alpha on mTORC1 and mTORC2. *J Biol Chem* 283: 34495-34499, 2008.

1232 228. **Tsai FC, Seki A, Yang HW, Hayer A, Carrasco S, Malmersjö S, and Meyer T.** A polarized Ca²⁺,
1233 diacylglycerol and STIM1 signalling system regulates directed cell migration. *Nat Cell Biol* 16: 133-144, 2014.

1234 229. **Tung JJ, Hobert O, Berryman M, and Kitajewski J.** Chloride intracellular channel 4 is involved in
1235 endothelial proliferation and morphogenesis in vitro. *Angiogenesis* 12: 209-220, 2009.

1236 230. **Tung JJ, and Kitajewski J.** Chloride intracellular channel 1 functions in endothelial cell growth and
1237 migration. *J Angiogenesis Res* 2: 23, 2010.

1238 231. **Turturro F, Friday E, Fowler R, Surie D, and Welbourne T.** Troglitazone acts on cellular pH and
1239 DNA synthesis through a peroxisome proliferator-activated receptor gamma-independent mechanism in breast
1240 cancer-derived cell lines. *Clin Cancer Res* 10: 7022-7030, 2004.

1241 232. **Ulmasov B, Bruno J, Gordon N, Hartnett ME, and Edwards JC.** Chloride intracellular channel
1242 protein-4 functions in angiogenesis by supporting acidification of vacuoles along the intracellular tubulogenic
1243 pathway. *Am J Pathol* 174: 1084-1096, 2009.

1244 233. **Valenzuela SM, Martin DK, Por SB, Robbins JM, Warton K, Bootcov MR, Schofield PR,**
1245 **Campbell TJ, and Breit SN.** Molecular cloning and expression of a chloride ion channel of cell nuclei. *J Biol*
1246 *Chem* 272: 12575-12582, 1997.

1247 234. **van Waarde A, Rybczynska AA, Ramakrishnan NK, Ishiwata K, Elsinga PH, and Dierckx RA.**
1248 Potential applications for sigma receptor ligands in cancer diagnosis and therapy. *Biochim Biophys Acta* 1848:
1249 2703-2714, 2015.

- 1250 235. **Vara D, Morell C, Rodríguez-Henche N, and Diaz-Laviada I.** Involvement of PPAR γ in the
1251 antitumoral action of cannabinoids on hepatocellular carcinoma. *Cell Death Dis* 4: e618, 2013.
- 1252 236. **Vaupel P.** Tumor microenvironmental physiology and its implications for radiation oncology. *Semin*
1253 *Radiat Oncol* 14: 198-206, 2004.
- 1254 237. **Vaupel P, and Harrison L.** Tumor hypoxia: causative factors, compensatory mechanisms, and cellular
1255 response. *Oncologist* 9 Suppl 5: 4-9, 2004.
- 1256 238. **Veliceasa D, Ivanovic M, Hoepfner FT, Thumbikat P, Volpert OV, and Smith ND.** Transient
1257 potential receptor channel 4 controls thrombospondin-1 secretion and angiogenesis in renal cell carcinoma.
1258 *FEBS J* 274: 6365-6377, 2007.
- 1259 239. **Verkman AS, Anderson MO, and Papadopoulos MC.** Aquaporins: important but elusive drug targets.
1260 *Nat Rev Drug Discov* 13: 259-277, 2014.
- 1261 240. **Villaamil MV, Gallego AG, Rubira VL, Campelo GR, Valladares-Ayerbes M, Pulido GE, Bolos**
1262 **VM, Cainzos SI, and Aparicio AL.** Fructose transporter GLUT5 expression in clear renal cell carcinoma.
1263 *Oncology Reports* 25: 315-323, 2011.
- 1264 241. **Vilner BJ, John CS, and Bowen WD.** Sigma-1 and sigma-2 receptors are expressed in a wide variety
1265 of human and rodent tumor cell lines. *Cancer Res* 55: 408-413, 1995.
- 1266 242. **Vink S, and Alewood PF.** Targeting voltage-gated calcium channels: developments in peptide and
1267 small-molecule inhibitors for the treatment of neuropathic pain. *Br J Pharmacol* 167: 970-989, 2012.
- 1268 243. **Voloshyna I, Besana A, Castillo M, Matos T, Weinstein IB, Mansukhani M, Robinson RB,**
1269 **Cordon-Cardo C, and Feinmark SJ.** TREK-1 is a novel molecular target in prostate cancer. *Cancer Res* 68:
1270 1197-1203, 2008.
- 1271 244. **Walia V, Yu Y, Cao D, Sun M, McLean JR, Hollier BG, Cheng J, Mani SA, Rao K, Premkumar**
1272 **L, and Elble RC.** Loss of breast epithelial marker hCLCA2 promotes epithelial-to-mesenchymal transition and
1273 indicates higher risk of metastasis. *Oncogene* 31: 2237-2246, 2012.
- 1274 245. **Walker JM, Bowen WD, Walker FO, Matsumoto RR, De Costa B, and Rice KC.** Sigma receptors:
1275 biology and function. *Pharmacological Reviews* 42: 355-402, 1990.
- 1276 246. **Walters DK, Arendt BK, and Jelinek DF.** CD147 regulates the expression of MCT1 and lactate
1277 export in multiple myeloma cells. *Cell Cycle* 12: 3175-3183, 2013.
- 1278 247. **Wang X, and van Breemen C.** Depolarization-mediated inhibition of Ca(2+) entry in endothelial cells.
1279 *Am J Physiol* 277: H1498-1504, 1999.
- 1280 248. **Wang Y, Kaiser MS, Larson JD, Nasevicius A, Clark KJ, Wadman SA, Roberg-Perez SE, Ekker**
1281 **SC, Hackett PB, McGrail M, and Essner JJ.** Moesin1 and Ve-cadherin are required in endothelial cells
1282 during in vivo tubulogenesis. *Development* 137: 3119-3128, 2010.
- 1283 249. **Wannous R, Bon E, Gillet L, Chamouton J, Weber G, Brisson L, Goré J, Bognoux P, Besson P,**
1284 **Roger S, and Chevalier S.** Suppression of PPAR β , and DHA treatment, inhibit Nav1.5 and NHE-1 pro-
1285 invasive activities. *Pflugers Arch* 2014.
- 1286 250. **Welch-Reardon KM, Ehsan SM, Wang K, Wu N, Newman AC, Romero-Lopez M, Fong AH,**
1287 **George SC, Edwards RA, and Hughes CC.** Angiogenic sprouting is regulated by endothelial cell expression
1288 of Slug. *J Cell Sci* 127: 2017-2028, 2014.
- 1289 251. **Wrzosek A.** Endothelium as target for large-conductance calcium-activated potassium channel openers.
1290 *Acta Biochim Pol* 56: 393-404, 2009.
- 1291 252. **Wu Z, and Bowen WD.** Role of sigma-1 receptor C-terminal segment in inositol 1,4,5-trisphosphate
1292 receptor activation: constitutive enhancement of calcium signaling in MCF-7 tumor cells. *Journal of Biological*
1293 *Chemistry* 283: 28198-28215, 2008.
- 1294 253. **Xia J, Yu X, Tang L, Li G, and He T.** P2X7 receptor stimulates breast cancer cell invasion and
1295 migration via the AKT pathway. *Oncol Rep* 2015.
- 1296 254. **Xiao N, Jiang LM, Ge B, Zhang TY, Zhao XK, and Zhou X.** Over-expression of TRPM8 is
1297 associated with poor prognosis in urothelial carcinoma of bladder. *Tumour Biol* 35: 11499-11504, 2014.
- 1298 255. **Yang AD, Camp ER, Fan F, Shen L, Gray MJ, Liu W, Somcio R, Bauer TW, Wu Y, Hicklin DJ,**
1299 **and Ellis LM.** Vascular endothelial growth factor receptor-1 activation mediates epithelial to mesenchymal
1300 transition in human pancreatic carcinoma cells. *Cancer Res* 66: 46-51, 2006.
- 1301 256. **Yang S, Zhang JJ, and Huang XY.** Orai1 and STIM1 are critical for breast tumor cell migration and
1302 metastasis. *Cancer Cell* 15: 124-134, 2009.
- 1303 257. **Ye JH, Gao J, Wu YN, Hu YJ, Zhang CP, and Xu TL.** Identification of acid-sensing ion channels in
1304 adenoid cystic carcinomas. *Biochem Biophys Res Commun* 355: 986-992, 2007.

- 1305 258. **Yingjun G, and Xun Q.** Acid-sensing ion channels under hypoxia. *Channels (Austin)* 7: 231-237, 2013.
- 1306 259. **Yoneda T, Hiasa M, Nagata Y, Okui T, and White F.** Contribution of acidic extracellular
- 1307 microenvironment of cancer-colonized bone to bone pain. *Biochim Biophys Acta* 2015.
- 1308 260. **Zamora-León SP, Golde DW, Concha II, Rivas CI, Delgado-López F, Baselga J, Nualart F, and**
- 1309 **Vera JC.** Expression of the fructose transporter GLUT5 in human breast cancer. *Proc Natl Acad Sci U S A* 93:
- 1310 1847-1852, 1996.
- 1311 261. **Zhang DX, and Gutterman DD.** Transient receptor potential channel activation and endothelium-
- 1312 dependent dilation in the systemic circulation. *J Cardiovasc Pharmacol* 57: 133-139, 2011.
- 1313 262. **Zhang J, Wei J, Kanada M, Yan L, Zhang Z, Watanabe H, and Terakawa S.** Inhibition of store-
- 1314 operated Ca²⁺ entry suppresses EGF-induced migration and eliminates extravasation from vasculature in
- 1315 nasopharyngeal carcinoma cell. *Cancer Lett* 336: 390-397, 2013.
- 1316 263. **Zhao CX, Luo CL, and Wu XH.** Hypoxia promotes 786-O cells invasiveness and resistance to
- 1317 sorafenib via HIF-2 α /COX-2. *Med Oncol* 32: 419, 2015.
- 1318
- 1319

1320 **FIGURE LEGENDS**

1321

1322 **Fig. 1:** Schematic representation of the different transport systems activated in tumor cells by hypoxia.

1323 Hypoxia induces activation of the α subunit of HIF1 (hypoxia-induced factor), which dimerizes with HIF1 β to
1324 result in HIF1. HIF1, in turn, stimulates a set of proteins involved in membrane ion or solute transport. NHE:
1325 Na⁺/H⁺ exchanger; ASIC: Acid-Sensing Ion Channel; NBC: Na⁺/HCO₃⁻ cotransporter; CAIX: carbonic
1326 anhydrase type IX; GLUT: Glucose transporter; MCT: Monocarboxylate transporter.

1327

1328 **Fig. 2:** Schematic representation of Ca²⁺ signaling in endothelial cells activated by the tumor environment
1329 physical and biochemical conditions. In a hypoxic environment, acidification of the microenvironment triggers
1330 the synthesis and release of VEGF (blue circles) by both tumor and stromal (CAF) cells. VEGF binds to its
1331 specific endothelial cell receptor (VEGF-R2), which, in turn, activates Ca²⁺ entry by two main processes. Store-
1332 operated Ca²⁺ entry (SOCE) results from 1) activation of Ca²⁺ release from endoplasmic reticulum (ER),
1333 especially via inositol 1,4,5-trisphosphate (IP₃R), and subsequent replenishment of these stores by Ca²⁺ entry via
1334 TRPC1 or TRPC4 channels; 2) activation of the stimulating unit (STIM) of the STIM-ORAI complex and
1335 subsequent Ca²⁺ influx by the ORAI transporting unit. Receptor-operated Ca²⁺ entry (ROCE) results from the
1336 direct activation of TRPC3 or TRPC6 Ca²⁺ channels. The ensuing increase in intracellular [Ca²⁺] is the
1337 mainspring of mobilization, organization and permeabilization of new blood vessels.

1338

1339 **Fig. 3:** Schematic representation of plasma membrane ion channel chaperoning and Ca²⁺ signaling enhancement
1340 by Sig-1R in cancer cells (adapted after Su et al, 2010, Trends Pharmacol. Sci.31: 557-566). During cancer
1341 progression, tumor cells are submitted to stressful conditions caused by low pO₂, low pH, and few nutrients.
1342 Under these conditions, Sig-1R (blue wide arrow) released from its ER co-chaperone BiP, activates the ER
1343 IP₃R3 Ca²⁺ channel, thus enhancing Ca²⁺ flux from the ER to the mitochondria and increasing the production of
1344 energy by the mitochondria. Sig-1R also stimulates the *bcl2* oncogene transcription, which reduces ROS
1345 production by the mitochondria and is in favor of cell survival. Under prolonged conditions of stress, ER
1346 chaperones are up-regulated; Sig-1R is overexpressed and translocates to the plasma membrane area where it
1347 becomes able to chaperone and activate membrane proteins such as ion channels (hERG, Na_v1.5, ASIC1a, ...) or
1348 enzymes (e.g. Src kinase).

Table 1. Main ion channels and solute transporters involved in tumor development and their role in these processes.

TRANSPORT PROTEIN	GENE	OFFICIAL NAME	FUNCTION IN ONCOGENESIS
Water channel			
AQP-1	<i>Aqp1</i>	Aquaporin	Angiogenesis, proliferation, migration (66, 68, 120)
Solute carriers (Na⁺/solute exchangers)			
GLUT-1	<i>Slc2a1</i>	Glucose transporter	Tumor cell glucose metabolism (6, 84, 132)
GLUT-5	<i>Slc2a5</i>	Fructose transporter	Tumor cell glucose metabolism (240, 260)
NHE1	<i>Slc9a1</i>	Na ⁺ /H ⁺ exchanger isoform 1	Cell volume and pH homeostasis (33, 170) Tumor angiogenesis (80) Metastasis (88, 218)
Bicarbonate transporters			
AE1-3	<i>Slc4a1-3</i>	Anion exchangers	pH homeostasis (222) Cell migration (222)
NBC transporters	<i>Slc4a4, 5, 7, 10</i>	Sodium-bicarbonate cotransporters	pH homeostasis (180) Tumor growth (180) Metastasis (180)
MCT family of monocarboxylate transporters			
MCT 1-4	<i>Slc16a1-4</i>	Solute carrier family 16 members 1-4	Cell migration and metastasis (184)
TRP superfamily of cation channels			
TRPA (ankyrin) family	<i>Trpa1</i>	Transient receptor potential cation channel, subfamily A	Cell migration and metastasis (63) Nociception, inflammation (22)
TRPC (canonical) family	<i>Trpc1-7</i>	Transient receptor potential cation channel, subfamily C	Cell migration and metastasis (76) Tumor angiogenesis (261)
TRPM (melastatin) family	<i>Trpm1-8</i>	Transient receptor potential cation channel, subfamily M	Cell migration and metastasis (63) Tumor angiogenesis (75)
TRPV (vanilloid) family	<i>Trpv1-6</i>	Transient receptor potential cation channel, subfamily V	Cell migration and metastasis (119) Tumor angiogenesis (75) Nociception (22)
Voltage-gated family of K⁺ channels			
Kv1.1-1.8	<i>Kcna1-7, 10</i>	Potassium channel, voltage gated, shaker related subfamily A, members 1-7	Proliferation of cancer cells (168) Apoptosis resistance (168) Immunoresponsiveness (224)
EAG1, 2	<i>Kcnh1, 5</i>	Ether-a-go-go K ⁺ channels	Proliferation of cancer cells (31) Migration of breast cancer cells (183)
hERG 1-3 or Kv11.1-11.3	<i>Kcnh2, 6, 7</i>	Ether-a-go-go-related gene K ⁺ channels	Proliferation of cancer cells (214) Crosstalk with cell adhesion molecules (183) Tumor angiogenesis (183)

Table 1. cont'd

Two-Pore K⁺ channel family			
K _{2p} 2.1 or TREK	<i>Kcnk2</i>	Arachidonic acid-sensitive K ⁺ channel	Cell proliferation (243)
K _{2p} 3.1 or TASK	<i>Kcnk3</i>	Acid-sensitive two-pore K ⁺ channel	Cell proliferation (127, 243) Evasion from apoptosis (127)
Calcium-activated K⁺ channels			
K _{Ca} 1.1 or BK _{Ca}	<i>Kcnn1</i>	Large conductance Ca ²⁺ -activated K ⁺ channel	Proliferation (200) Apoptosis (28)
K _{Ca} 2.1-3 or SK1-3	<i>Kcnn1-3</i>	Small conductance Ca ²⁺ -activated K ⁺ channels	Cell migration and metastasis (37, 110, 191)
K _{Ca} 3.1 or IK _{Ca} or SK4	<i>Kcnn4</i>	Intermediate conductance Ca ²⁺ -activated K ⁺ channel	Proliferation of cancer cells (141) Cell migration (141) Tumor angiogenesis (141) Evasion from immune destruction (174)
Voltage-gated Na⁺ channels			
Na _v	<i>Kcnn1</i>	Voltage-gated Na ⁺ channel	Tumor cell migration, invasion, endocytosis (32) Metastasis (32)
Voltage-gated family of Ca²⁺ channels			
Ca _v 1.1-1.4	<i>Cacna1s, c, d, f</i>	L-type Ca ²⁺ current	Tumor angiogenesis (51)
Ca _v 2.1-2.3	<i>Cacna1a, b, e</i>	P/Q-, N-, R-type Ca ²⁺ currents	Tumor angiogenesis (154)
Ca _v 3.1-3.3	<i>Cacna1g, h, i</i>	T-type Ca ²⁺ currents	Tumor angiogenesis (204)
P2X Receptor (P2XR) family of Na⁺, K⁺, Ca²⁺ ion channels			
P2X ₇ receptor	<i>P2rx7</i>	Ligand-gated cation channel	Proliferation of cancer cells (3, 103) Apoptosis resistance (3, 103)
Acid-sensing ion channel family			
ASIC1-3	<i>Asic1-3</i>	Acid-sensing ion channels	pH homeostasis (92) Cell migration and invasion (92)
Chloride intracellular channel family			
CLIC1-6	<i>Clic1-6</i>	Nuclear and intracellular organelle Cl ⁻ channels	Tumor angiogenesis (232)
Calcium release-activated calcium channels (CRAC channels)			
Orai1 (works with STIM)	<i>Orai1</i>	Calcium release-activated calcium channel protein 1	Cell migration and metastasis (161, 216) Tumor angiogenesis (139) Immunosuppression (174)





