

## Reply to 'Waves do not contribute to global sea-level rise'

Angelique Melet, Benoit Meyssignac, Rafael Almar, Gonéri Le Cozannet

### ▶ To cite this version:

Angelique Melet, Benoit Meyssignac, Rafael Almar, Gonéri Le Cozannet. Reply to 'Waves do not contribute to global sea-level rise'. Nature Climate Change, 2019, 9, pp.3. 10.1038/s41558-018-0378-4 . hal-02380691

## HAL Id: hal-02380691 https://hal.science/hal-02380691

Submitted on 6 Dec 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Reply to 'Waves do not contribute to global sea-level rise'

Melet et al. reply — The quantification of adaptation needs to sea-level hazards at global-to-regional scales is a major research challenge. Our global-scale analysis<sup>1</sup> contributes to understanding the drivers of interannual-to-multidecadal coastal water level changes by reporting that at first order, wave contributions are sizeable. Typically these contributions have not been accounted for, and thus we advocate for their inclusion to provide more accurate assessments<sup>2</sup> of the vulnerability of the coasts. We clarify below three main points regarding our analysis and its implications.

First, coastal water-level changes result from different processes acting over a range of timescales. Wind-wave-induced coastal water level changes are largest for short timescales (wave period or wave-group timescales). However, wave set-up and swash can be modulated at interannualto-multidecadal timescales by: changes in offshore wave height, period and direction due to surface wind changes3; changes in geomorphology<sup>4</sup> (such as beach orientation, nearshore bathymetry); and changes in currents and water depth<sup>5-7</sup>. Our analysis focused on the first point, highlighting that wave contributions should be considered not only for extreme events, but also in analyses of interannual-to-multidecadal coastal sealevel changes and rise.

Second, policy-relevant water-level changes are those that result from a combination of processes that cause shoreline changes, including wave set-up and swash, and from both climate change and internal variability. But as wave set-up is always positive at the coast, whereas swash is oscillatory, their contributions were shown separately, providing a more direct comparison between set-up and altimetric sea level<sup>1</sup>. Trends induced by internal variability were highlighted to discuss the future evolution of wave set-up and altimetric sea level. It should be noted that even estimates of coastal sea-level rise due to steric effects and melting glaciers have been substantial due to internal variability over the past two decades<sup>8-10</sup>. Although

wave climate projections remain uncertain, some robust patterns of longer-term waveregime changes emerge in climate change projections<sup>11</sup>. In any case, wave climate changes are expected in response to climate change and will be transmitted to water-level changes at the coast through changes in wave set-up and swash.

Third, we acknowledge that our estimates are associated with large uncertainties. We aimed to be transparent on this point by mentioning the main limitations of our methodology, separating the contributions from wave set-up and swash, and reporting the large sensitivity of our estimates to choices in foreshore beach slope values and empirical parameterizations. Although Aucan et al.<sup>12</sup> criticize the parameterization values used, no worldwide estimate of foreshore beach slopes is available<sup>13–15</sup>. In the SHOM report they cite<sup>16</sup>, beach slopes rely on a parametric beach profile model and on assumptions regarding sediment grain sizes, using only four beaches as a benchmark. As recognized by that report's authors, this method does not deliver a reliable global estimate of beach slopes.

Beach slopes generally range between 0.01 and 0.20 (ref. 17). Although a slope of 0.1 might lie in the upper range for intermediate beaches, and despite limitations in empirical formulations of wave set-up and swash, this value allows an order of magnitude estimate for a global-scale analysis. A sensitivity analysis using slopes ranging from 0.05 to 0.15 in the generic, state-ofthe-art empirical formulation of set-up and swash<sup>1,18</sup> completes our main analysis. For dissipative conditions (generally including beaches with lower slopes, <0.02) we did use the dissipative-specific formulation<sup>1,18</sup> also used by Vitousek and colleagues<sup>5</sup>. Using a slope of 0.05 (or 0.02) instead of 0.10 with the generic formula reduces wave set-up by a factor of 2.0 (5.0), swash by a factor of 1.3 (1.5) and run-up by a factor of 1.55 (2.1). Using the dissipative-specific formulation instead of the generic one with a beach slope of 0.1 reduces set-up by a factor of 2.4 and swash by a factor of 2.0 (not 5 as stated by

Aucan et al.<sup>12</sup>). The main message of our study is therefore robust regardless of the formulation used, but our results call for future research efforts to refine estimates of wave set-up and swash.

Only long-term observations of wave set-up and swash could validate interannualto-multidecadal changes, which are the focus of our study. However, despite numerous coastal wave measurements (from buoys, for example) existing, sustained efforts are still required to collect observations such as video monitoring or LIDAR to monitor run-up over longer periods<sup>19</sup>.

### Angélique Melet<sup>1\*</sup>, Benoît Meyssignac<sup>2</sup>, Rafaël Almar<sup>2</sup> and Gonéri Le Cozannet<sup>3</sup>

<sup>1</sup>Mercator Ocean International, Ramonville St Agne, France. <sup>2</sup>LEGOS, Université de Toulouse, CNES, CNRS, IRD, UPS, Toulouse, France. <sup>3</sup>BRGM / French Geological Survey, Orléans, France. \*e-mail: angelique.melet@mercator-ocean.fr

### Published online: 18 December 2018 https://doi.org/10.1038/s41558-018-0378-4

#### References

- Melet, A., Meyssignac, B., Almar, R. & Le Cozannet, G. Nat. Clim. Change 8, 234–239 (2018).
- 2. Bouwer, L. M. Nat. Clim. Change 8, 765-766 (2018).
- 3. Semedo, A., Suselj, K., Rutgersson, A. & Sterl, A. J. Clim. 24,
- 1461–1479 (2011).
- Cohn, N. & Ruggiero, P. Coast. Eng. 115, 79–92 (2016).
  Vitousek, S. et al. Sci. Rep. 7, 1399 (2017).
- Vitousek, S. et al. Sci. Rep. 7, 1399 (2017)
  Arns, A. et al. Sci. Rep. 7, 40171 (2017).
- Anderson, T. et al. Sci. Rep. 8, 14484 (2018).
- Carson, M., Koehl, A. & Stammer, D. J. Clim. 28, 853–861 (2015).
- Bilbao, R. A. F., Gregory, J. M. & Bouttes, N. Clim. Dynam. 45, 2647–2666 (2015).
- Marzeion, B., Cogley, J. G., Richter, K. & Parkes, D. Science 345, 919–921 (2014).
- Morim, J., Hemer, M. A., Cartwright, N., Strauss, D. & Andutta, F. Glob. Planet. Change 167, 160–171 (2018).
- 12. Aucan, J. et al. Nat. Clim. Change https://doi.org/10.1038/s41558-018-0377-5 (2018).
- 13. Vousdoukas, M. I. et al. Nat. Commun. 9, 2360 (2018).
- 14. Wright, L. D. & Short, A. D. Mar. Geol. 56, (1984).
- 15. Luijendijk, A. et al. Sci. Rep. 8, 6641 (2018).
- Deshoulières, A. Prévision des Pentes de Plage et Estimation Statistique Mondiale (SHOM, 2009).
- 17. Komar, P. *Beach Processes and Sedimentation* 2nd edn (Prentice Hall, Upper Saddle River, 1998).
- Stockdon, H. F., Holman, R. A., Howd, P. A. & Sallenger, A. H. Coast. Eng. 53, 573–588 (2006).
- Blenkinsopp, C., Mole, M., Turner, I. & Peirson, W. Coast. Eng. 57, 1059–1065 (2010).