

Exploring the cognitive workload during a visual search task in Parkinson's Disease

Maud Ranchet, John Morgan, Abiodun Emmanuel Akinwuntan, Hannes Devos

▶ To cite this version:

Maud Ranchet, John Morgan, Abiodun Emmanuel Akinwuntan, Hannes Devos. Exploring the cognitive workload during a visual search task in Parkinson's Disease. International Congress of Parkinson's Disease and Movement Disorders, Sep 2019, NICE, France. International Congress of Parkinson's Disease and Movement Disorders, 1 p, 2019. hal-02384125v1

HAL Id: hal-02384125 https://hal.science/hal-02384125v1

Submitted on 28 Nov 2019 (v1), last revised 23 Mar 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.









Exploring the cognitive workload during a visual search task in Parkinson's Disease

M. Ranchet¹, J.C. Morgan², A. E. Akinwuntan³, H. Devos⁴

¹Univ Lyon, F-69000 Lyon, France-IFSTTAR, TS2, LESCOT, F-69500 Bron

²Parkinson's Foundation Center of Excellence, Movement and Memory Disorder Programs, Department of Neurology, Medical College of Georgia, Augusta University, Augusta, GA, USA

³Dean's office, School of Health Professions, The University of Kansas Medical Center, Kansas City, KS, USA

⁴Department of Physical Therapy and Rehabilitation Science, School of Health Professions, The University of Kansas Medical Center, Kansas City, KS, USA

- Visual search impairments in patients with Parkinson's Disease (PD) may negatively affect daily life activities, such as driving^{1,2}
- The index of cognitive activity (ICA) is a pupillary response that accurately estimates cognitive workload during various tasks in individuals with cognitive decline^{3,4,5}

Objective

To investigate cognitive workload during a visual search task in patients with PD

Methods

✓ 20 patients with PD (69 yrs-old ± 8; 16 Males) and 15 controls (61 yrs-old ± 11; 8 Males)

VISUAL SEARCH TASK

✓ Participants were instructed to search for a target stimulus (road sign) among distractor stimuli (other road signs) (Figure 1)




Figure 1. Example of a target-present trial. Gray circle represents the pre-identified target and the gray square represents the target.

- ✓ The task included 42 trials (half of them had the target present)
- ✓ ICA was recorded by two synchronized, remote eye trackers (FOVIO, SeeingMachines Inc, Canberra, Australia) at 60 Hz. Values ranged between 0 and 1
- ✓ Cognitive workload (ICA) over the course of the visual search task was explored

References

1.Devos, H., Vandenberghe, W., Tant, M., Akinwuntan, A. E., De Weerdt, W., Nieuwboer, A., & Uc, E. Y. (2013). Driving and off-road impairments underlying failure on road testing in Parkinson's disease. Movement Disorders: Official Journal Movement Disorder 28(14), the Society, 1949–1956. https://doi.org/10.1002/mds.25701

2. Uc, E. Y., Rizzo, M., Anderson, S. W., Sparks, J., Rodnitzky, R. L., & Dawson, J. D. (2006). Impaired visual search in drivers with Parkinson's disease. Annals of Neurology, 60(4), 407-413. https://doi.org/10.1002/ana.20958


3. Kahneman, D. (1973). Attention and Effort. Englewood Cliffs, NJ: Prentice-Hall. 4. Marshall, S. P. (2007). Identifying cognitive state from eye metrics. Aviation,

Space, and Environmental Medicine, 78(5 Suppl), B165-175.

5. Ranchet, M., Orlosky, J., Morgan, J., Qadir, S., Akinwuntan, A. E., & Devos, H. (2017). Pupillary response to cognitive workload during saccadic tasks in Parkinson's disease. Behavioural Brain Research, 327, 162–166. https://doi.org/10.1016/j.bbr.2017.03.043

Results

✓ PD patients were slower than controls to detect correctly the target (7s. \pm 2 vs 5 s. \pm 1, p = 0.004) and missed more targets than controls (p = 0.04)



Each asterisk represents significant differences between cognitive workload values at a certain percentage of completion time and the baseline corresponding to 5% before the start of the trial. The black asterisks are for the patients and the gray asterisks are for the controls.

*p<0.05 ** p<0.01, ***p<0.001

Figure 2. Change in ICA from baseline to completion of the visual search of all target-present trials for the left eye


✓ PD patients had increased values in cognitive workload throughout the entire duration of the task when compared to baseline (p < 0.05). By contrast, few significant differences were observed in controls (at 5% and 15%).

Conclusion

- ✓ PD patients required increased and longer effort to correctly perform the visual search task compared with baseline cognitive workload
- ✓ The use of ICA in patients with PD while performing a static visual search task provides new information into the effort (cognitive workload) required by patients on a moment-to-moment basis

Acknowledgement

✓ We thank the American Parkinson's Disease Association for funding this research. We also thank Adam Bruetsch, MS, for his help with data analysis.



www.ifsttar.fr