



HAL
open science

Analysis and numerical solvability of backward-forward conservation laws

Thibault Liard, Enrique Zuazua

► **To cite this version:**

Thibault Liard, Enrique Zuazua. Analysis and numerical solvability of backward-forward conservation laws. 2022. hal-02389808v3

HAL Id: hal-02389808

<https://hal.science/hal-02389808v3>

Preprint submitted on 16 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Analysis and numerical solvability of backward-forward conservation laws

Thibault Liard* Enrique Zuazua †‡§

Abstract

In this paper, we study the problem of initial data identification for weak-entropy solutions of the one-dimensional Burgers equation. This problem consists in identifying the set of initial data evolving to a given target at a final time. Due to the time-irreversibility of the Burgers equation, some target functions are unattainable from solutions of this equation, making the identification problem under consideration ill-posed. To get around this issue, we introduce a non-smooth optimization problem which consists in minimizing the difference between the predictions of the Burgers equation and the observations of the system at a final time in $L^2(\mathbb{R})$ norm. Here, we fully characterize the set of minimizers of the aforementioned non-smooth optimization problem. One of minimizers is the backward entropy solution, constructed using a backward-forward method. Some simulations are given using a wave-front tracking algorithm.

Keywords: Backward-forward method, Identification problems; Conservation Laws; Weak-entropy solutions; Non-smooth optimization problem; Wave-front tracking algorithm.

AMS classification: 35L65, 35F20, 93B30, 35R30.

1 Introduction

1.1 Presentation of the Problem

Initial data identification problem consists in finding the origin of a physical phenomenon, governed for instance by partial differential equations (PDEs), from a set of observations at a given time. This arises naturally in meteorology, oceanography or climatology [32, 45, 23, 44, 30, 5, 19] to improve the forecasts of a model. Finding optimal positions or shapes of sensors [40, 41, 42] also lead to the study of identification problems.

Initial identification problems need to be carefully addressed, depending on each type of PDEs.

*Univ. Limoges, CNRS, XLIM, UMR 7252, F-87000 Limoges, France.

†Chair in Applied Analysis, Alexander von Humboldt-Professorship, Department of Mathematics Friedrich-Alexander-Universität, Erlangen-Nürnberg, 91058 Erlangen, Germany.

‡Departamento de Matemáticas, Universidad Autónoma de Madrid, 28049 Madrid, Spain.

§Chair of Computational Mathematics, Fundación Deusto Av. de las Universidades 24, 48007 Bilbao, Basque Country, Spain.

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement NO. 694126-DyCon). This work was partially supported by the ELKARTEK project KK-2018/00083 ROAD2DC of the Basque Government, by the Grant MTM2017-92996-C2-1-R/2-R COSNET of MINECO (Spain), by the Air Force Office of Scientific Research (AFOSR) under Award NO. FA9550-18-1-0242, by the Alexander von Humboldt-Professorship program, by the European Union's Horizon 2020 research and the innovation programme under the Marie Skłodowska-Curie grant agreement NO. 765579-ConFlex and by the grant ICON-ANR-16-ACHN-0014 of the French ANR.

- In the case of parabolic PDEs, the high and instant regularization effect induces the non-existence of initial data for which the corresponding solution evolves to given not-necessary regular target functions, and causes numerical instabilities when solving the PDE backwards in time. In [34], the authors solve an identification problem for the heat equation with applications in pollution source localization. Note however that, when the target is attainable, the initial datum whose the corresponding trajectory evolves to this target, is unique as seen in [36].
- In the case of nonlinear hyperbolic PDE as (1), the backward uniqueness property fails due to the presence of discontinuities (so-called *shocks*), i.e multiple initial data may evolve to the same attainable target function. Moreover, due to the time-irreversibility of nonlinear hyperbolic PDEs, a target function u^T can be unattainable, that is to say that there is non-existence of initial data leading to the target function u^T .

In this paper, we study the latter case. More precisely, we consider the scalar conservation laws

$$\begin{cases} \partial_t u(t, x) + \partial_x f(u(t, x)) = 0, & (t, x) \in \mathbb{R}^+ \times \mathbb{R}, \\ u(0, x) = u_0(x), \end{cases} \quad (1)$$

where the flux function f is defined by $f : u \mapsto \frac{u^2}{2}$. Kruzkov's theory [31] provides existence and uniqueness of a weak-entropy solution u of (1) with initial datum $u_0 \in L^\infty(\mathbb{R})$. Let $T > 0$ and $u^T \in L^\infty(\mathbb{R})$ a given function, the goal is to find the set of initial data generating weak-entropy solutions of (1) that are as close as possible to u^T at time T in $L^2(\mathbb{R})$ -norm, i.e to solve the following non-smooth optimization problem

$$\inf_{u_0 \in \mathcal{U}_{\text{ad}}^0} J_0(u_0) := \|u^T(\cdot) - u(T, \cdot)\|_{L^2(\mathbb{R})}, \quad (\mathcal{O}_T)$$

where u is the weak-entropy solution of (1) and $\mathcal{U}_{\text{ad}}^0$ is the class of admissible initial data defined in (14). The study of initial data identification for (1) is motivated by the minimization of the sonic boom effects generated by supersonic aircrafts which are modeled by an augmented Burgers equation [15, 3, 2, 35].

1.2 State of the art and main results

Initial data identification for (1) in the case of attainable targets has already been studied in [12, 13, 24, 1, 28, 16, 35]. In [16, Theorem 3.1, Corollary 3.2], [28, Corollary 1] or [24], the authors prove that u^T is truly attainable in an exact manner by a solution of (1) if and only if u^T satisfies the one-sided Lipschitz condition [8, 25, 39, 21], i.e

$$\partial_x u^T \leq \frac{1}{T} \text{ in the sense of distributions.} \quad (2)$$

When u^T is an attainable target, the authors in [28] prove that the set of initial data evolving to u^T is a convex set. Later on, the aforementioned set was fully characterized in [16, 22] using the classical Lax-Hopf formula. In [35], an alternative proof is given using backward generalized characteristics.

The optimization problem (\mathcal{O}_T) with attainable targets has also been studied in [12, 13, 1] from numerical points of view. Since the weak-entropy solution u of (1) may contain shocks even if the initial datum is a smooth function, this generates important added difficulties that have been

the object of intensive study in the past, see [38, 37, 9, 10, 6, 7, 4] and the references therein. In [9, 10, 6, 7], the derivative of the cost function J_0 in (\mathcal{O}_T) is regarded in a weak sense by requiring strong conditions on the set of initial data. This leads to require that weak-entropy solutions of (1) have a finite number of non-interacting jumps. When J_0 is weakly differentiable, gradient descent methods have been implemented in [12, 13, 1] to solve numerically the optimization problem (\mathcal{O}_T) . In the cases where it was applied successfully, only one possible initial datum emerges, namely the backward entropy solution, see Remark 1. This is mainly due to the numerical viscosity that numerical schemes introduce to gain stability. To find some multiple minimizers, the authors in [28] use a filtering step in the backward adjoint solution.

In this article, we give a full characterization of the set of minimizers for the optimization problem (\mathcal{O}_T) . More precisely, we prove that the backward entropy solution, denoted by $S_T^-(u^T)$, is a minimizer of (\mathcal{O}_T) using a backward-forward method described in Section 2.1. Then, we show that u_0^* is a minimizer of (\mathcal{O}_T) if and only if the weak-entropy solution of (1) with initial datum u_0^* coincides, at time T , with the weak-entropy solution of (1) with initial datum $S_T^-(u^T)$ using variational methods. Moreover, we construct numerically random minimizers of (\mathcal{O}_T) based on a wave-front tracking algorithm. Note that, contrary to [12, 13, 1], we do not require strong assumptions on the set of initial data, i.e weak-entropy solutions of (1) may have a countable number of interacting jumps.

The article is organized as follows. In section 2, we describe the backward-forward method and we recall known results on initial data identification for (1). In Section 3, we state the main results where the proofs are given in Section 4. In Section 5, we run some simulations using a wave-front tracking method.

1.3 Some related open problems

Let us address some related open questions and possible extensions of this work.

- It would be interesting to study the optimization problem (\mathcal{O}_T) in L^1 -norm, which is the natural distance in the framework of conservation laws. This problem leads to additional difficulties since $x \mapsto \|x\|_{L^1(\mathbb{R})}$ is not a differentiable function.
- It would be also interesting to consider a convex-concave function as a flux function in (1) which is, for instance, a more realistic choice to describe the flow of pedestrian [17, 14]. The main difficulty comes from the existence of discontinuities (called non-classical shocks) violating standard admissibility entropy conditions such that the Oleinik inequality.
- We could also study a Burgers equation with source terms. In this case, some suitable conditions on source terms have to be determined to use the backward-forward method described in this paper. For instance, the backward operator $S_t^-(u^T)$ defined in Section 2.1 associated to

$$\begin{cases} \partial_t u(t, x) + \partial_x f(u(t, x)) = -u^3(t, x), & (t, x) \in \mathbb{R}^+ \times \mathbb{R}, \\ u(T, \cdot) = u^T(x), & x \in \mathbb{R}. \end{cases}$$

may blow up at time $t < T$.

- We can also investigate systems of conservation laws in one dimension (Euler equations, Saint-Venant equations, Aw-Rascle-Zhang traffic flow model). Note that, as soon as the backward-forward operator $S_T^+(S_T^-)$ is well-defined, $S_T^+(S_T^-)(u^T)$ may give a good candidate to solve the inverse design of systems of conservation laws.

- We may consider a multi-dimensional equation of conservation of laws in a numerical point of view. For instance, a fractional steps method [18, 33, 29] (or splitting method) may be implemented to solve an identification problem of a two-dimensional equation of conservation laws.

2 Notations and comments

In the sequel, we denote by $BV(\mathbb{R})$ the class of functions of bounded variation, see [21, Definition 1.7.1]. If $g \in BV(\mathbb{R})$, we use the notation $g(x-) := \lim_{y \rightarrow x} g(y)$ and $g(x+) := \lim_{y \rightarrow x} g(y)$. Let $f \in BV(\mathbb{R})$, we denote by $X(f)$ the set defined by

$$X(f) := \{x \in \mathbb{R} / f(x-) = f(x+)\}, \quad (3)$$

and $\text{supp}(f)$ stands for the support of the function f . Let Ω be a domain in \mathbb{R} , $\mathcal{D}(\Omega) := C_c^\infty(\Omega)$ denotes the set of infinitely differentiable functions with compact support. Let two distributions $T_1, T_2 \in \mathcal{D}'(\Omega)$, we say that $T_1 \leq T_2$ in the sense of distributions if

$$\forall \varphi \in \mathcal{D}(\Omega), \varphi \geq 0, \quad \langle T_1, \varphi \rangle \leq \langle T_2, \varphi \rangle,$$

where $\langle \cdot, \cdot \rangle$ is a duality bracket between \mathcal{D}' and \mathcal{D} .

2.1 The backward-forward method

For a sake of completeness, we recall the definition of a weak-entropy solution of (1).

Definition 2.1 • We say that $u \in L^\infty(\mathbb{R}^+ \times \mathbb{R}) \cap C^0(\mathbb{R}^+, L_{loc}^1(\mathbb{R}))$ is a weak solution if for all $\varphi \in C_c^1(\mathbb{R}^2, \mathbb{R})$,

$$\int_{\mathbb{R}^+} \int_{\mathbb{R}} (u \partial_t \varphi + f(u) \partial_x \varphi) dx dt + \int_{\mathbb{R}} u_0(x) \varphi(0, x) dx = 0.$$

- We say that $u \in L^\infty(\mathbb{R}^+ \times \mathbb{R}) \cap C^0(\mathbb{R}^+, L_{loc}^1(\mathbb{R}))$ is a weak-entropy solution if u is a weak solution and for every $k \in \mathbb{R}$, for all $\varphi \in C_c^1(\mathbb{R}^2, \mathbb{R}^+)$,

$$\int_{\mathbb{R}^+} \int_{\mathbb{R}} (|u - k| \partial_t \varphi + \text{sgn}(u - k)(f(u) - f(k)) \partial_x \varphi) dx dt + \int_{\mathbb{R}} |u_0 - k| \varphi(0, x) dx \geq 0.$$

Kruzkov's theory [31] provides existence and uniqueness of a weak-entropy solution $(t, x) \rightarrow S_t^+(u_0)(x)$ of (1) with initial datum $u_0 \in L^\infty(\mathbb{R})$. For a given function u^T , we introduce the function $(t, x) \rightarrow S_t^-(u^T)(x)$ as follows: for every $t \in [0, T]$, for a.e $x \in \mathbb{R}$,

$$S_t^-(u^T)(x) = S_t^+(x \rightarrow u^T(-x))(-x). \quad (4)$$

Remark 1 The solutions $S_t^+(u_0)$ and $S_t^-(u^T)$ may be regarded as the zero viscosity limit of the solutions $S_t^{+, \epsilon}(u_0)$ and $S_T^{-, \epsilon}(u^T)$ respectively where $S_t^{+, \epsilon}(u_0)$ and $S_t^{-, \epsilon}(u^T)$ are defined as follows: $S_t^{+, \epsilon}(u_0)$ is the solution of the following viscous Burgers equation

$$\begin{cases} \partial_t u(t, x) + \partial_x f(u(t, x)) = +\epsilon \partial_{xx}^2 u(t, x), & (t, x) \in \mathbb{R}^+ \times \mathbb{R}, \\ u(0, \cdot) = u_0(x), & x \in \mathbb{R}, \end{cases}$$

and $S_t^{-\epsilon}(u^T)$ is the solution of the following backward equation

$$\begin{cases} \partial_t u(t, x) + \partial_x f(u(t, x)) = -\epsilon \partial_{xx}^2 u(t, x), & (t, x) \in \mathbb{R}^+ \times \mathbb{R}, \\ u(T, \cdot) = u^T(x), & x \in \mathbb{R}. \end{cases}$$

Using the change of variable $(t, x) \rightarrow (T - t, -x)$, we notice that the backward equation above is well-defined. Thus, $S_T^-(u^T)$ is called the backward entropy solution. Note that the construction involving S_t^\pm is also related to scattering theory [27, 26]

The backward-forward method consists in solving backward in time the PDE (1) with final target u^T and then solving it forward in time with initial datum $S_T^-(u^T)$, the solution of the backward PDE.

For any attainable target u^T , we have $S_T^+(S_T^-(u^T)) = u^T$ as seen in [16, Theorem 3.1, Corollary 3.2] and [28, Corollary 1]. However, there exist some target functions u^T verifying $S_T^+(S_T^-(u^T)) \neq u^T$ as seen in Example 1.

Example 1 Assuming that u^T is defined by $u^T(\cdot) = -\mathbb{1}_{(-\infty, 0)}(\cdot) + \mathbb{1}_{(0, \infty)}(\cdot)$ then the weak-entropy solution v of (1) with initial datum $v(0, x) = u^T(-x)$ is defined by

$$v(t, x) = \begin{cases} 1 & \text{if } x < 0, \\ -1 & \text{if } x > 0. \end{cases}$$

Thus, $S_{T-t}^-(u^T) : x \rightarrow v(T - t, -x)$ is a weak solution of (1) verifying that $u(T) = u^T$. The weak-entropy solution u_e with initial datum $v(T, -x)$ is defined by

$$u_e(t, x) = \begin{cases} -1 & \text{if } x < -t, \\ \frac{x}{t} & \text{if } -t \leq x \leq t, \\ 1 & \text{if } t < x. \end{cases}$$

In particular, $S_T^+(S_T^-(u^T)) := u_e(T) \neq u^T$. Note that u^T is an unattainable target.

2.2 Identification problem for attainable targets

Fix $u^T \in L^\infty(\mathbb{R})$, we introduce the set

$$\mathcal{I}^+(u^T) = \{u_0 \in L^\infty(\mathbb{R}) : S_T^+(u_0) = u^T\}. \quad (5)$$

From [16, Corollary 3.2], $\mathcal{I}^+(u^T) \neq \emptyset$ if and only if a suitable representative of u^T satisfies the Oleinik condition [8, 25, 39, 21], i.e for every $x \in \mathbb{R}$ and $y \in \mathbb{R}^+ \setminus \{0\}$,

$$f'(u^T(x+y)) - f'(u^T(x)) \leq \frac{y}{T}. \quad (6)$$

The following theorem stated in [35, Theorem 1] (see also [16, 22]) gives a full characterization of the set of initial data $u_0 \in L^\infty(\mathbb{R})$ such that $S_T^+(u_0) = u^T$.

Theorem 2.1 ([35]) *Let $T > 0$ and a suitable representation of $u^T \in L^\infty(\mathbb{R})$ satisfies the Oleinik condition (2). Then the initial data $u_0 \in L^\infty(\mathbb{R})$ verifies $S_T^+(u_0) = u^T$ if and only if the following statements holds. For any $(x, y) \in X(u^T) \times \mathbb{R}$*

$$\int_{x-Tf'(u^T(x))}^y S_T^-(u^T)(s) ds \leq \int_{x-Tf'(u^T(x))}^y u_0(s) ds, \quad (7)$$

For any $(x, y) \in X(u^T)^2$,

$$\int_{x-Tf'(u^T(x))}^{y-Tf'(u^T(y))} S_T^-(u^T)(s) ds = \int_{x-Tf'(u^T(x))}^{y-Tf'(u^T(y))} u_0(s) ds, \quad (8)$$

where $X(u^T)$ is defined in (3) and $S_T^-(u^T)$ is defined in (4).

Remark 2 When $u^T \in L^\infty(\mathbb{R})$ satisfies the Oleinik condition (2), then $u^T \in BV_{loc}(\mathbb{R})$. Thus, $X(u^T)$ is well-defined.

Theorem 2.1 points out the richness and the diversity of initial data evolving to the same target at time T (see Figure 1).

- There exists $u_0 \in L^\infty(\mathbb{R})$ such that $S_T^+(u_0) = u^T$ with $\min_{x \in \mathbb{R}} u_0(x) < \min_{x \in \mathbb{R}} u^T(x)$ and/or $\max_{x \in \mathbb{R}} u^T(x) < \max_{x \in \mathbb{R}} u_0(x)$, see Figure 9.
- The set $\mathcal{I}^+(u^T)$ defined in (5) is a convex cone having as unique extremal point at its vertex the map $S_T^-(u^T)$, see [16, Proposition 5.2]; for any $u_0 \in \mathcal{I}^+(u^T)$, for every $\eta > 0$, $u_0^\eta = S_T^-(u^T) + \eta(u_0 - S_T^-(u^T)) \in \mathcal{I}^+(u^T)$.




Figure 1: Three initial data $u_0^*(-)$, $u_1(- -)$ and $u_2(\cdot \cdot \cdot)$ leading to a attainable target $u^T(\cdot) := \mathbb{1}_{(-\infty, 0)}(\cdot) - \mathbb{1}_{(0, +\infty)}(\cdot)$ at time $T = 1$ along forward entropic evolution.

The following theorem will be a essential tool to prove Theorem 3.1. Fix $u_0 \in L^\infty(\mathbb{R})$, we introduce the set

$$\mathcal{I}^-(u_0) = \{u^T \in L^\infty(\mathbb{R}) : S_T^-(u^T) = u_0\}. \quad (9)$$

From (4) and [16, Corollary 3.2], $\mathcal{I}^-(u_0) \neq \emptyset$ if and only if for every $x \in \mathbb{R}$ and $y \in \mathbb{R}^+ \setminus \{0\}$,

$$f'(u_0(x+y) - f'(u_0(x))) \geq -\frac{y}{T}. \quad (10)$$

Theorem 2.2 gives a full characterization of the set of function $u_T \in L^\infty(\mathbb{R})$ such that $S_T^-(u_T) = u_0$.

Theorem 2.2 *Let $T > 0$ and a suitable representation of $u_0 \in L^\infty(\mathbb{R})$ satisfies (10). Then a map $u_T \in L^\infty(\mathbb{R})$ satisfies $S_T^-(u_T) = u_0$ if and only if the following statements holds. For any $(x, y) \in X(u_0) \times \mathbb{R}$*

$$\int_{x+Tf'(u_0(x))}^y S_T^+(u_0)(s) ds \geq \int_{x+Tf'(u_0(x))}^y u_T(s) ds, \quad (11)$$

For any $(x, y) \in X(u_0)^2$,

$$\int_{x+Tf'(u_0(x))}^{y+Tf'(u_0(y))} S_T^+(u_0)(s) ds = \int_{x+Tf'(u_0(x))}^{y+Tf'(u_0(y))} u_T(s) ds, \quad (12)$$

where $X(u_0) = \{x \in \mathbb{R}, u_0(x-) = u_0(x+)\}$.

Theorem 2.2 is a consequence of Theorem 2.1 noticing that $S_T^-(u_T) : x \rightarrow S_T^+(x \rightarrow u_T(-x))(-x)$.

3 Main results

Let $T > 0$ and $u^T \in L^\infty(\mathbb{R})$ with compact support, that is to say there exists a compact set $K^T := [a^T, b^T] \subset \mathbb{R}$ and a constant $C > 0$ verifying

$$\text{supp}(u^T) \subset K^T \text{ and } \|u\|_{L^\infty(\mathbb{R})} \leq C \quad (13)$$

Let $K_0 := [a_0, b_0]$ such that $[a^T - Tf'(C), b^T + Tf'(C)] \subset K_0$ (see an illustration in Figure 3), we consider the non-smooth optimization problem

$$\inf_{u_0 \in \mathcal{U}_{\text{ad}}^0} J_0(u_0) := \|u^T(\cdot) - S_T^+(u_0)(\cdot)\|_{L^2(\mathbb{R})}, \quad (\mathcal{O}_T)$$

where S_T^+ is defined in Section 2.1 and $\mathcal{U}_{\text{ad}}^0$ is the class of admissible initial data defined by

$$\mathcal{U}_{\text{ad}}^0 = \{u_0 \in L^\infty(\mathbb{R}) / \|u_0\|_{L^\infty(\mathbb{R})} \leq C \text{ and } \text{supp}(u_0) \subset K_0\}, \quad (14)$$

Theorem 3.1 characterizes the set of minimizers of (\mathcal{O}_T) (see an illustration in Figure 2).

Theorem 3.1 *Let $T > 0$ and $u^T \in L^\infty(\mathbb{R})$ verifying (13). The initial datum $u_0^* \in L^\infty(\mathbb{R})$ is a minimizer of (\mathcal{O}_T) if and only if $u_0^* \in \mathcal{U}_{\text{ad}}^0$ satisfies $S_T^+(u_0^*) = S_T^+(S_T^-(u^T))$.*

Remark 3 *The constraints $\|u_0\|_{L^\infty(\mathbb{R})} \leq C$ and $\text{Supp}(u_0) \subset K_0$ in (14) are used to guarantee the existence of minimizers of (\mathcal{O}_T) . Moreover, the assumption $[a^T - Tf'(C), b^T + Tf'(C)] \subset K_0$ is required to have $S_T^-(u^T) \in \mathcal{U}_{\text{ad}}^0$.*

Corollary is a direct consequence of Theorem 3.1 and the full characterization of the set $\{u_0 \in L^\infty(\mathbb{R}) / S_T^+(u_0) = S_T^+(S_T^-(u^T))\}$ given in Theorem 2.1.

Corollary 3.1 *Let $T > 0$ and $u^T \in L^\infty(\mathbb{R})$ verifying (13). The map $u_0^* \in L^\infty(\mathbb{R})$ is a minimizer of (\mathcal{O}_T) if and only if the following statements holds. For any $(x, y) \in X(S_T^+(S_T^-(u^T))) \times \mathbb{R}$*

$$\int_{x-Tf'(S_T^+(S_T^-(u^T)))(x)}^y S_T^-(u^T)(s) ds \leq \int_{x-Tf'(S_T^+(S_T^-(u^T)))(x)}^y u_0^*(s) ds, \quad (15)$$

For any $(x, y) \in X(S_T^+(S_T^-(u^T)))^2$,

$$\int_{x-Tf'(S_T^+(S_T^-(u^T)))(x)}^{y-Tf'(S_T^+(S_T^-(u^T)))(y)} S_T^-(u^T)(s) ds = \int_{x-Tf'(S_T^+(S_T^-(u^T)))(x)}^{y-Tf'(S_T^+(S_T^-(u^T)))(y)} u_0^*(s) ds, \quad (16)$$

where $X(S_T^+(S_T^-(u^T)))$ is defined in (3) and $S_T^-(u^T)$ is defined in (4).




Figure 2: The backward-forward solution $S_T^+(S_T^-(u^T))$ is the projection of u^T onto the set of attainable target functions. The shaded area in red at time $t = 0$ represents the set of minimizers of (\mathcal{O}_T) .

Remark 4 From (26), $S_T^+(S_T^-(u^T)) \in BV(\mathbb{R})$. Thus, for any $x \in \mathbb{R}$, $S_T^+(S_T^-(u^T))(x-)$ and $S_T^+(S_T^-(u^T))(x+)$ exist and then $X(S_T^+(S_T^-(u^T)))$ is well-defined.

The proof of Theorem 3.1 is structured as follows. From [16, Theorem 3.1, Corollary 3.2], [28, Corollary 1] or [24], there exists $u_0 \in L^\infty(\mathbb{R})$ such that $S_T^+(u_0) = q$ if and only if $q \in L^\infty(\mathbb{R})$ satisfies the one-sided Lipschitz condition (2). Let $K_1 \subset \mathbb{R}$ such that $[a_0 - Tf'(C), b_0 + Tf'(C)] \subset K_1$, the optimal problem (\mathcal{O}_T) is equivalent to

$$\min_{q \in \mathcal{U}_{\text{ad}}^T} J_1(q) := \|u^T - q\|_{L^2(\mathbb{R})}, \quad (17)$$

where the admissible set $\mathcal{U}_{\text{ad}}^T$ is defined by

$$\mathcal{U}_{\text{ad}}^T = \{q \in L^\infty(\mathbb{R}) / \partial_x q \leq \frac{1}{T}, \|q\|_{L^\infty(\mathbb{R})} \leq C \text{ and } \text{Supp}(q) \subset K_1\}. \quad (18)$$

The optimization problem (17) admits a unique minimizer using Hilbert projection Theorem. We prove that $q = S_T^+(S_T^-(u^T))$ is the minimizer of (17) using the first-order optimality conditions applied to (17) and the full characterization of the set $\{u_0 \in BV(\mathbb{R}) / S_T^-(u_0) = S_T^-(u^T)\}$ given in Theorem 2.2.

Remark 5 The optimal problem (17) is not related to the PDE model (1). As a consequence, unlike J_0 in (\mathcal{O}_T) , the cost function J_1 in (17) is a differentiable function.

Assuming that the given target u^T is attainable. Since $S_T^+(S_T^-(u^T)) = u^T$, Theorem 3.1 and Corollary 3.1 give a fully characterization of initial data leading to u^T along forward entropic evolution, as in [22, 35]. Note that there exists initial data yielding weak solutions u that coincide with u^T such that the inequalities (15) and (16) do not hold, see Example 2.

Example 2 Let $T = 1$ and assuming that u^T is defined by $u^T(\cdot) = \mathbb{1}_{(-\infty, 0)}(\cdot) - \mathbb{1}_{(0, +\infty)}(\cdot)$ then the weak solution u defined by

$$u(t, x) = \begin{cases} \mathbb{1}_{(-\infty, 4t-2)}(x) + 7\mathbb{1}_{(4t-2, 3t-\frac{3}{2})}(x) - \mathbb{1}_{(3t-\frac{3}{2}, +\infty)}(x) & \text{if } t < \frac{1}{2}, \\ \mathbb{1}_{(-\infty, 0)}(x) - \mathbb{1}_{(0, +\infty)}(x) & \text{if } \frac{1}{2} \leq t, \end{cases}$$

satisfies $u(T, \cdot) = u^T$ and $S_T^+(u(0, \cdot)) \neq u^T$.

4 Proof of Theorem 3.1

The proof of Theorem 3.1 is based on the following Lemma.

Lemma 4.1 *The optimal problem (17) admits a unique minimizer $S_T^+(S_T^-(u^T))$.*

PROOF. The proof is divided in two steps.

Step 1: Existence of minimizers of (17).

By definition of J_1 in (17), it is enough to prove that $\mathcal{U}_{\text{ad}}^T$ defined in (18) is a closed convex set of $L^2(\mathbb{R})$ using Hilbert projection Theorem.

- Assuming that $q_1, q_2 \in \mathcal{U}_{\text{ad}}^T$, we immediately have, for every $\alpha \in [0, 1]$, $\alpha q_1 + (1 - \alpha)q_2 \in \mathcal{U}_{\text{ad}}^T$. Thus, $\mathcal{U}_{\text{ad}}^T$ is a convex set.
- Assuming that $q_n \in \mathcal{U}_{\text{ad}}^T$ converges to q in $L^2(\mathbb{R})$ then q_n converges to q in the sense of distributions and by passing to the limit in $\partial_x q_n \leq \frac{1}{T}$, we have $\partial_x q \leq \frac{1}{T}$. Since $\|q_n\|_{L^\infty(\mathbb{R})} \leq C$ and using that the closed ball $B_{L^\infty(\mathbb{R})}$ is compact in the weak* topology $\sigma(L^\infty, L^1)$ [11, Theorem 3.16], q_n converges, (up to a subsequence, still denoted by q_n) to $q \in L^\infty(\mathbb{R})$ in the weak* topology of $L^\infty(\mathbb{R})$. Moreover, from [11, Proposition 3.13], $\|q\|_{L^\infty(\mathbb{R})} \leq \liminf_n \|q_n\|_{L^\infty(\mathbb{R})} \leq C$. Using that q_n converges to q in $L^2(\mathbb{R})$, q_n converges a.e to q . Moreover, since $\text{supp}(q_n) \subset K_1$, we have $q_n(x) = 0$ for a.e $x \in \mathbb{R} \setminus K_1$. Therefore, we have $\text{Supp}(q) \subset K_1$ and we conclude that $q \in \mathcal{U}_{\text{ad}}^T$. Thus, $\mathcal{U}_{\text{ad}}^T$ is a closed set.

Step 2: First-order optimality conditions.

Our aim is to prove that,

$$S_T^+(S_T^-(u^T)) \in \mathcal{U}_{\text{ad}}^T, \quad (19)$$

and for any admissible perturbation $h \in \mathcal{T}_{\mathcal{U}_{\text{ad}}^T}(S_T^+(S_T^-(u^T)))$,

$$- \int_{\mathbb{R}} (u^T(x) - S_T^+(S_T^-(u^T))(x)) h(x) dx \geq 0. \quad (20)$$

Above, $\mathcal{T}_{\mathcal{U}_{\text{ad}}^T}(S_T^+(S_T^-(u^T)))$ is a set of functions $h \in L^\infty(\mathbb{R})$ such that, for any sequence of positive real numbers ϵ_n decreasing to 0, there exists a sequence of functions $h_n \in L^\infty(\mathbb{R})$ converging to h as $n \rightarrow \infty$ and $S_T^+(S_T^-(u^T)) + \epsilon_n h_n \in \mathcal{U}_{\text{ad}}^T$ for every $n \in \mathbb{N}$. If (19) and (20) hold then $S_T^+(S_T^-(u^T))$ is a critical point of (17). Since J_1 is strictly convex, $S_T^+(S_T^-(u^T))$ is the unique minimizer of (17).

We now prove (19): since u^T satisfies (13), we have $\|u^T\|_{L^\infty(\mathbb{R})} \leq C$. By using the definition of S_T^+ and S_T^- and the maximum principle fulfilled by weak-entropy solutions [43, Theorem 2.3.5], we have

$$\|S_T^-(u^T)\|_{L^\infty(\mathbb{R})} \leq C \text{ and } \|S_T^+(S_T^-(u^T))\|_{L^\infty(\mathbb{R})} \leq C. \quad (21)$$




Figure 3: Illustration of K^T (---) and K_0 (—) defined in Section 3.

From (13), by definition of S_T^- and using the finite velocity of propagation, we have $\text{supp}(S_T^-(u^T)) \subset [a^T - Tf'(C), b^T + Tf'(C)] \subset K_0$ (see Figure 3). Therefore, together with (21),

$$S_T^-(u^T) \in \mathcal{U}_{\text{ad}}^0. \quad (22)$$

Moreover, by definition of S_T^+ and using $[a_0 - Tf'(C), b_0 + Tf'(C)] \subset K_1$ and [21, Theorem 6.2.3], we have

$$S_T^+(u_0) \in \mathcal{U}_{\text{ad}}^T \text{ for any } u_0 \in \mathcal{U}_{\text{ad}}^0, \quad (23)$$

where $\mathcal{U}_{\text{ad}}^T$ is defined in (18). We replace u_0 in (23) by $S_T^-(u^T)$ and we deduce that (19) holds.

We now prove (20) : let $x \in X(S_T^-(u^T))$ with X defined in (3) and we introduce the function $F : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$F : y \mapsto \int_{x+Tf'(S_T^-(u^T)(x))}^y (u^T(s) - S_T^+(S_T^-(u^T))(s)) ds. \quad (24)$$

Since $u^T \in L^\infty(\mathbb{R})$ satisfies (13), we have

$$u^T \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R}). \quad (25)$$

By definition of S_T^- and S_T^+ (see Section 2.1), from [21, Theorem 11.2.2] and $u^T \in L^\infty(\mathbb{R})$, we have $S_T^+ S_T^-(u^T) \in BV_{\text{loc}}(\mathbb{R})$. Therefore, together with (19), we deduce that for any $T > 0$,

$$S_T^+ S_T^-(u^T) \in BV(\mathbb{R}) \text{ and } S_T^+ S_T^-(u^T) \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R}). \quad (26)$$

From (24), (25) and (26), we have that

$$F \in W^{1,1}(\mathbb{R}) \cap W^{1,\infty}(\mathbb{R}). \quad (27)$$

and for a.e $y \in X(S_T^+ S_T^-(u^T))$,

$$F'(y) = u^T(y) - S_T^+(S_T^-(u^T))(y). \quad (28)$$

We now introduce the function $p : X(S_T^+ S_T^-(u^T)) \rightarrow \mathbb{R}$ defined by

$$p(y) = y - Tf'(S_T^+(S_T^-(u^T))(y)). \quad (29)$$

From [21, Theorem 11.1.3], $p(y) = \xi_+(0) = \xi_-(0)$ where ξ_- and ξ_+ denote respectively the minimal and the maximal backward generalized characteristics associated with the solution $S_t^+(S_T^-(u^T))$




Figure 4: Plotting of the forward generalized characteristic $\xi(\cdot)$ emanating from $(x, 0)$ and the extremal backward generalized characteristics $\xi_-(\cdot)$ and $\xi_+(\cdot)$ emanating from (y, T) associated with the solution $S_t^+(S_T^-(u^T))$. We have $p(y) := y - Tf'(S_T^+(S_T^-(u^T)))(y) = \xi_+(0) = \xi_-(0)$.

emanating from (y, T) (see Figure 4). From (24), (26) and [21, Theorem 1.7.4, Theorem 11.3.4], $X(S_T^+S_T^-(u^T))$ has full Lebesgue measure and

$$\begin{aligned}
-\int_{\mathbb{R}}(u^T(y) - S_T^+(S_T^-(u^T))(y))h(y) dy &= -\int_{X(S_T^+S_T^-(u^T))}(u^T(y) - S_T^+(S_T^-(u^T))(y))h(y) dy, \\
&= -\int_{p^{-1}(X(S_T^-(u^T)))} F'(y)h(y) dy \\
&= -\int_{X(S_T^+S_T^-(u^T)) \setminus p^{-1}(X(S_T^-(u^T)))} F'(y)h(y) dy.
\end{aligned} \tag{30}$$

where $p^{-1}(X(S_T^-(u^T))) := \{y \in X(S_T^+S_T^-(u^T)) / p(y) \in X(S_T^-(u^T))\}$ (see an illustration in Figure 5). By definition of S_T^- , using that $u^T \in L^\infty(\mathbb{R})$ satisfies (13) and [21, Theorem 6.2.6], we have




Figure 5: Illustration of $(t, x) \rightarrow S_t^+S_T^-(u^T)(x)$, $p^{-1}(X(S_T^-(u^T))) \subset X(S_T^+S_T^-(u^T))$ (—), $X(S_T^+S_T^-(u^T)) \setminus p^{-1}(X(S_T^-(u^T)))$ (-) and discontinuous points of $S_T^+S_T^-(u^T)$ (\times). Here, $y \in X(S_T^+S_T^-(u^T)) \setminus p^{-1}(X(S_T^-(u^T)))$, $p(y)$ is defined in (29) and at any discontinuous points $(z_k)_{k \in \mathbb{N}}$ of $S_T^-(u^T)$ verifying $S_T^-(u^T)(z_k^-) < S_T^-(u^T)(z_k^+)$, a rarefaction wave is created at time $t = 0$.

$S_T^-(u^T) \in BV(\mathbb{R})$. As a consequence, $S_T^-(u^T)$ has a countable number of discontinuous points $(z_k)_{k \in \mathbb{N}}$ verifying $S_T^-(u^T)(z_k^-) < S_T^-(u^T)(z_k^+)$. Moreover, if $y \in X(S_T^+S_T^-(u^T)) \setminus p^{-1}(X(S_T^-(u^T)))$, from [21, Theorem 11.1.3] associated with the solution $S_t^+(S_T^-(u^T))$, we have $S_T^-(u^T)(p(y)-) < S_T^-(u^T)(p(y)+)$. Thus, a rarefaction wave is created at time $t = 0$ and at the position $p(y)$, i.e

$y \in [p(y) + Tf'(S_T^-(u^T)(p(y)-)), p(y) + Tf'(S_T^-(u^T)(p(y)+))]$. We conclude that

$$X(S_T^+ S_T^-(u^T)) \setminus p^{-1}(X(S_T^-(u^T))) = \cup_{k \in \mathbb{N}} [z_k + Tf'(S_T^-(u^T)(z_k-)), z_k + Tf'(S_T^-(u^T)(z_k+))]. \quad (31)$$

Thus, (30) can be written as

$$\begin{aligned} - \int_{\mathbb{R}} (u^T(y) - S_T^+(S_T^-(u^T))(y)) h(y) dy &= - \int_{p^{-1}(X(S_T^-(u^T)))} F'(y) h(y) dy \\ &= - \sum_{k \in \mathbb{N}} \int_{\mathcal{I}_k} F'(y) h(y) dy. \end{aligned} \quad (32)$$

with

$$\mathcal{I}_k := (z_k + Tf'(S_T^-(u^T)(z_k-)), z_k + Tf'(S_T^-(u^T)(z_k+))). \quad (33)$$

We now study each term on the right side of the equality (32).

- Let $x \in X(S_T^-(u^T))$ and $y \in p^{-1}(X(S_T^-(u^T)))$. Applying Theorem 2.2 with $u_0 = S_T^-(u^T)$ and $u_T = u^T$, the equality (12) holds, i.e for any $(x, p(y)) \in X(S_T^-(u^T))^2$,

$$\int_{x+Tf'(S_T^-(u^T)(x))}^{p(y)+Tf'(S_T^-(u^T)(p(y)))} S_T^+(S_T^-(u^T))(s) ds = \int_{x+Tf'(S_T^-(u^T)(x))}^{p(y)+Tf'(S_T^-(u^T)(p(y)))} u^T(s) ds, \quad (34)$$

Using $y \in p^{-1}(X(S_T^-(u^T)))$ and [21, Theorem 11.1.3, Theorem 11.3.2] associated with the solution $S_t^+(S_T^-(u^T))$, there exists a unique forward generalized characteristic $\xi(\cdot)$ emanating from $(p(y), 0)$ and $\xi(T) = p(y) + Tf'(S_T^-(u^T)(p(y))) = y$. From (24) and (34), we conclude that for any $y \in p^{-1}(X(S_T^-(u^T)))$, $F(y) = 0$. From (27), F is a continuous function on \mathbb{R} and from (26) the set of discontinuous points of $S_T^+ S_T^-(u^T)$ is countable. Then, together with (31), we have for any $y \in \mathbb{R} \setminus (\cup_{k \in \mathbb{N}} \mathcal{I}_k)$,

$$F(y) = 0. \quad (35)$$

Therefore, for ϵ small enough, for any $y \in \mathbb{R} \setminus (\cup_{k \in \mathbb{N}} \overline{\mathcal{I}_k})$, we deduce that

$$0 = \frac{F(y + \epsilon) - F(y)}{\epsilon} = \frac{1}{\epsilon} \int_y^{y+\epsilon} F'(s) ds. \quad (36)$$

Combining (36) with Lebesgue differentiation Theorem, we have for a.e $y \in p^{-1}(X(S_T^-(u^T)))$

$$F'(y) = 0. \quad (37)$$

Thus, from (37), for every $h \in \mathcal{T}_{\text{ad}}^r(S_T^+(S_T^-(u^T)))$,

$$- \int_{p^{-1}(X(S_T^-(u^T)))} F'(y) h(y) dy = 0. \quad (38)$$

- Let $x \in X(S_T^-(u^T))$ and $y \in \cup_{k \in \mathbb{N}} \mathcal{I}_k$ with \mathcal{I}_k defined in (33). Since a rarefaction is created at $(p(y), 0)$ (see Figure 5), we have

$$\partial_y S_T^+(S_T^-(u^T))(y) = \frac{1}{T}. \quad (39)$$

Since $h \in \mathcal{T}_{\text{ad}}^r(S_T^+(S_T^-(u^T)))$ is an admissible perturbation, for every $\epsilon_n > 0$ such that $\epsilon_n \rightarrow 0$ when $n \rightarrow \infty$ there exists $h_n \in L^\infty(\mathbb{R})$ such that $\lim_{n \rightarrow \infty} h_n = h$ in $L^\infty(\mathbb{R})$ and $S_T^+(S_T^-(u^T)) + \epsilon_n h_n \in \mathcal{U}_{\text{ad}}^T$. Thus,

$$\partial_y S_T^+(S_T^-(u^T))(y) + \epsilon_n \partial_y h_n(y) \leq \frac{1}{T} \quad \text{in the sense of distributions.} \quad (40)$$

Using (39) and (40), we have $\partial_y h_n(y) \leq 0$ in the sense of distributions. Since $\lim_{n \rightarrow \infty} h_n = h$ in $L^\infty(\mathbb{R})$, h_n tends to h in the sense of distributions and we conclude that for any admissible perturbation $h \in \mathcal{T}_{\mathcal{U}_{\text{ad}}^T}(S_T^+(S_T^-(u^T)))$,

$$\partial_y h(y) \leq 0 \quad \text{in the sense of distributions.} \quad (41)$$

Applying Theorem 2.2 with $u_0 = S_T^-(u^T)$, $u_T = u^T$, the inequality (15) holds, i.e

$$\int_{x+Tf'(S_T^-(u^T)(x))}^y u^T(s) ds \leq \int_{x+Tf'(S_T^-(u^T)(x))}^y S_T^+(S_T^-(u^T))(s) ds,$$

From (24), we conclude that for any $y \in \cup_k \mathcal{I}_k$

$$F(y) \leq 0 \quad (42)$$

Let $k \in \mathbb{N}$. Using (27) and (35), we have $F \in W_0^{1,1}(\mathcal{I}_k)$. Thus, there exists $F_n \in C_c^\infty(\mathcal{I}_k)$ such that F_n converges to F in $W^{1,1}(\mathcal{I}_k)$. Moreover, $F_n := \rho_n * F$ where ρ_n is a sequence of positive mollifiers (see details in [11, Section 8]). Therefore, together with (42), we have $F_n(y) \leq 0$ for any $y \in \mathcal{I}_k$. Besides, for every $n \in \mathbb{N}$,

$$- \int_{\mathcal{I}_k} F_n'(y) h(y) dy = \langle \partial_y h, F_n \rangle, \quad (43)$$

where $\langle \cdot, \cdot \rangle$ is a duality bracket between the distribution $\partial_y h$ and the test function $F_n \in C_c^\infty(\mathcal{I}_k)$. Using (41) and (42), we have $\langle \partial_y h, F_n \rangle \geq 0$. From (43),

$$- \int_{\mathcal{I}_k} F_n'(y) h(y) dy \geq 0. \quad (44)$$

Since F_n converges to F in $W^{1,1}(\mathcal{I}_k)$, by passing to the limit in (44), we conclude that, for any admissible perturbation $h \in \mathcal{T}_{\mathcal{U}_{\text{ad}}^T}(S_T^+(S_T^-(u^T)))$,

$$- \int_{\mathcal{I}_k} F'(y) h(y) dy \geq 0. \quad (45)$$

Using (32), (38) and (45), the inequality (20) holds. □

Proof of Theorem 3.1: From Lemma 4.1, for every $q \in \mathcal{U}_{\text{ad}}^T$, we have

$$\|u^T - S_T^+(S_T^-(u^T))\|_{L^2(\mathbb{R})} \leq \|u^T - q\|_{L^2(\mathbb{R})}. \quad (46)$$

From (23) and (46), we deduce that, for any $u_0 \in \mathcal{U}_{\text{ad}}^0$,

$$\|u^T - S_T^+(S_T^-(u^T))\|_{L^2(\mathbb{R})} \leq \|u^T - S_T^+(u_0)\|_{L^2(\mathbb{R})}, \quad (47)$$

with $S_T^-(u^T) \in \mathcal{U}_{\text{ad}}^0$ using (22). Thus, $S_T^-(u^T)$ is a minimizer of (\mathcal{O}_T) .

- Let u_0^* a minimizer of (\mathcal{O}_T) . Then $u_0^* \in \mathcal{U}_{\text{ad}}^0$ and for any $u_0 \in \mathcal{U}_{\text{ad}}^0$ we have

$$\|u^T - S_T^+(u_0^*)\|_{L^2(\mathbb{R})} \leq \|u^T - S_T^+(u_0)\|_{L^2(\mathbb{R})}. \quad (48)$$

From (47) and (48), we immediately have $S_T^+(u_0^*) = S_T^+(S_T^-(u^T))$.

- Let $u_0^* \in \mathcal{U}_{\text{ad}}^0$ satisfying $S_T^+(u_0^*) = S_T^+(S_T^-(u^T))$. From (47), for any $u_0 \in \mathcal{U}_{\text{ad}}^0$,

$$\|u^T - S_T^+(u_0^*)\|_{L^2(\mathbb{R})} \leq \|u^T - S_T^+(u_0)\|_{L^2(\mathbb{R})}.$$

Thus, u_0^* is a minimizer of (\mathcal{O}_T) .

5 Numerical investigations

5.1 A wave-front tracking algorithm

In this section, we present an algorithm that allows us to construct randomly a minimizer $u_0^* \in BV(\mathbb{R})$ of (\mathcal{O}_T) . To simplify the presentation, we assume that

$$u^T = u_L \mathbb{1}_{(-\infty, \bar{x})} + u_R \mathbb{1}_{(\bar{x}, \infty)}, \quad (49)$$

with $u_L > u_R$, $\bar{x} \in \mathbb{R}$, $T > 0$. We introduce the set

$$\Gamma(u_L, u_R, \bar{x}, T) := \left\{ \gamma \in W^{1,1}([\bar{x} - Tf'(u_L), \bar{x} - Tf'(u_R)], \mathbb{R}) / \gamma \text{ satisfies (A1), (A2), (A3) and (A4)} \right\} \quad (50)$$

with

$$(A1) \quad \dot{\gamma} \in BV(\mathbb{R})$$

$$(A2) \quad \gamma(\bar{x} - Tf'(u_L)) = 0,$$

$$(A3) \quad \gamma(\bar{x} - Tf'(u_R)) = T(u_L f'(u_L) - f(u_L) - u_R f'(u_R) + f(u_R)),$$

$$(A4) \quad \text{For every } x \in [\bar{x} - Tf'(u_L), \bar{x} - Tf'(u_R)],$$

$$\gamma(x) \geq \gamma_*(x) := -T \int_{u_L}^{(f')^{-1}(\frac{\bar{x}-x}{T})} s f''(s) ds.$$

An illustration of the set $\Gamma(u_L, u_R, \bar{x}, T)$ is given in Figure 6.




Figure 6: Let u^T defined in (49). The set $\Gamma(u_L, u_R, \bar{x}, T)$ is illustrated by the shaded area. The function γ_* is defined by $\gamma_*(x) = -T \int_{u_L}^{(f')^{-1}(\frac{\bar{x}-x}{T})} s f''(s) ds = S_T^+(S_T^-(u^T))(x)$ for a.e $x \in [\bar{x} - Tf'(u_L), \bar{x} - Tf'(u_R)]$. We have $\gamma_1 \in \Gamma(u_L, u_R, \bar{x}, T)$ and $\gamma_2 \notin \Gamma(u_L, u_R, \bar{x}, T)$. From Theorem 2.2, $u_0^{\gamma_1}$ defined in (51) is a minimizer of (\mathcal{O}_T) while $u_0^{\gamma_2}$ is not.

Algorithm. First, we construct the backward-forward solution $S_T^+(S_T^-(u^T))$ using a wave-front tracking algorithm (see [20]). Second, we pick a random path $\gamma \in \Gamma(u_L, u_R, \bar{x}, T)$ and from Theorem 2.2, the initial data $u_0^\gamma \in BV(\mathbb{R})$ defined by

$$u_0^\gamma = \begin{cases} u_L & \text{if } x < \bar{x} - Tf'(u_L) \\ \dot{\gamma}(x) & \text{if } \bar{x} - Tf'(u_L) < x < \bar{x} - Tf'(u_R) \\ u_R & \text{if } \bar{x} - Tf'(u_R) < x \end{cases} \quad (51)$$

is a minimizer of (\mathcal{O}_T) (see an illustration in Figure 6).

Example 3 Let $T = 1$ and $u^T(\cdot) := \mathbb{1}_{(-\infty, 0)}(\cdot) - \mathbb{1}_{(0, +\infty)}(\cdot)$. In Figure 1, two different initial data defined by $u_1(x) = \mathbb{1}_{(-\infty, 0)}(x) - \mathbb{1}_{(0, +\infty)}(x)$ and $u_2(x) = \mathbb{1}_{(-\infty, -\frac{1}{4})}(x) + 2\mathbb{1}_{(-\frac{1}{4}, -\frac{1}{12})}(x) - \mathbb{1}_{(-\frac{1}{12}, +\infty)}(x)$ are constructed. The two $\gamma_1 : [-1, 1] \rightarrow \mathbb{R}$ and $\gamma_2 : [-1, 1] \rightarrow \mathbb{R}$ defined almost everywhere by $\hat{\gamma}_1(\cdot) = u_1(\cdot)$ and $\hat{\gamma}_2(\cdot) = u_2(\cdot)$ belongs to $\Gamma(1, -1, 0, 1)$, see Figure 7. From Theorem 2.2, $S_T^+(u_1) = S_T^+(u_2) = u^T$.

5.2 Construction of the set of minimizers of (\mathcal{O}_T)

In this section, (\mathcal{O}_T) is solved numerically with attainable and unattainable targets using the algorithm described in Section 5.1.

5.2.1 With attainable target u^T

Let $T = 1$. We consider the target u^T defined by $u^T(\cdot) = 0.6875\mathbb{1}_{(-\infty, 4.6)}(\cdot) - \mathbb{1}_{(4.6, \infty)}(\cdot)$ (Figure 8). Since $u^T(4.6+) < u^T(4.6-)$, the inequality (2) holds and so $u^T(\cdot)$ is an attainable function. As a consequence, we have $u^T = S_T^+(S_T^-(u^T))$. In Figure 8 and Figure 9, six minimizers u_0^* of (\mathcal{O}_T) are constructed. From Theorem 3.1, for any minimizer u_0^* , we have $S_T^+(u_0^*) = u^T$. Note that in top left corner of Figure 9, $S_T^-(u^T)$ is plotted with respect to x .




Figure 7: Plotting of γ_1 and γ_2 belonging to $\Gamma(1, -1, 0, 1)$. For a.e $x \in [-1, 1]$, $\hat{\gamma}_*(x) = S_T^+(S_T^-(u^T))(x)$, $\hat{\gamma}_1(x) = u_1(x)$ and $\hat{\gamma}_2(x) = u_2(x)$ where $u^T(\cdot)$, $u_1(\cdot)$ and $u_2(\cdot)$ are defined in Example 3.

5.2.2 With unattainable target u^T

In Example 4 and Example 5, The optimization problem (\mathcal{O}_T) is solved numerically with two unattainable targets.

Example 4 Let $T = 2$. We consider the target u^T defined as

$$u^T(x) = \begin{cases} 2 & \text{if } x \in (-0.2, 1.1) \cup (2, 3.1) \cup (4.1, 5.3) \cup (6.1, 7.2), \\ -1 & \text{otherwise.} \end{cases}$$

Since for every $x \in \{-0.2, 2, 4.1, 6.1\}$, we have $u^T(x-) < u^T(x+)$, and so the inequality (2) does not hold. Thus, u^T is an unattainable target.

- In Figure 10a), the approximate function $(x, t) \rightarrow S_t^-(u^T)(-x)$ is plotted.




Figure 8: The attainable target u^T defined by $u^T(\cdot) = 0.6875\mathbb{1}_{(-\infty, 4.6)}(\cdot) - \mathbb{1}_{(4.6, \infty)}(\cdot)$

- In Figure 10b), the approximate minimizer $S_T^-(u^T)$ of (\mathcal{O}_T) is plotted.
- In Figure 10c), the approximate function $(x, t) \rightarrow S_t^+(S_T^-(u^T))(x)$ is plotted.
- In Figure 10d), the function u^T and the approximate function $x \rightarrow S_T^+(S_T^-(u^T))(x)$ are plotted.

Four minimizers u_0^* of (\mathcal{O}_T) are constructed in Figure 11. From Theorem 3.1, for any u_0^* , $S_T^+(u_0^*) = S_T^+(S_T^-(u^T)) \neq u^T$.


Example 5 Let $T = 1$. We consider the target u^T defined as

$$u^T = -\mathbb{1}_{(-\infty, -0.2)} + 2\mathbb{1}_{(-0.2, 1.1)} + 0.16\mathbb{1}_{(1.1, 2)} + 1.33\mathbb{1}_{(2, 3.1)} - 0.77\mathbb{1}_{(3.1, 4.1)} \\ - 0.42\mathbb{1}_{(4.1, 5.3)} - \mathbb{1}_{(5.3, 6.1)} + 1.91\mathbb{1}_{(6.1, 7.2)} - \mathbb{1}_{(7.2, \infty)}.$$


The function u^T is an unattainable target. In Figure 12, the function u^T and $x \mapsto S_T^+(S_T^-(u^T))(x)$ are plotted.

References


- [1] Navid Allahverdi, Alejandro Pozo, and Enrique Zuazua. Numerical aspects of large-time optimal control of burgers equation. *ESAIM: Mathematical Modelling and Numerical Analysis*, 50(5):1371–1401, 2016.
- [2] Navid Allahverdi, Alejandro Pozo, and Enrique Zuazua. Numerical aspects of sonic-boom minimization. *A Panorama of Mathematics: Pure and Applied*, 658:267, 2016.
- [3] Juan J Alonso and Michael R Colonno. Multidisciplinary optimization with applications to sonic-boom minimization. *Annual Review of Fluid Mechanics*, 44:505–526, 2012.
- [4] Claude Bardos, Olivier Pironneau, et al. Data assimilation for conservation laws. *Methods and Applications of Analysis*, 12(2):103–134, 2005.
- [5] Andrew F Bennett. *Inverse modeling of the ocean and atmosphere*. Cambridge University Press, 2005.
- [6] François Bouchut and François James. One-dimensional transport equations with discontinuous coefficients. *Nonlinear Analysis*, 32(7):891, 1998.
- [7] François Bouchut and François James. Differentiability with respect to initial data for a scalar conservation law. In *Hyperbolic problems: theory, numerics, applications*, pages 113–118. Springer, 1999.
- [8] Alberto Bressan and Rinaldo M Colombo. Decay of positive waves in nonlinear systems of conservation laws. *Annali della Scuola Normale Superiore di Pisa-Classe di Scienze*, 26(1):133–160, 1998.
- [9] Alberto Bressan and Andrea Marson. A maximum principle for optimally controlled systems of conservation laws. *Rendiconti del Seminario Matematico della Università di Padova*, 94:79–94, 1995.
- [10] Alberto Bressan and Andrea Marson. A variational calculus for discontinuous solutions of systems of conservation laws. *Communications in partial differential equations*, 20(9):1491–1552, 1995.




$M = 1$ discontinuous point




$M = 2$ discontinuous points




$M = 3$ discontinuous points



$M = 4$ discontinuous points



$M = 5$ discontinuous points



$M = 6$ discontinuous points

Figure 9: Construction of six minimizers u_0^* of (\mathcal{O}_T) having $M \in \{1, \dots, 6\}$ discontinuous points. $T = 1$, $u^T = u_L \mathbb{1}_{(-\infty, \bar{x})} + u_R \mathbb{1}_{(\bar{x}, \infty)}$ with $u_L = 0.6875$, $u_R = -1$, $\bar{x} = 4.6$.

- [11] Haim Brezis and Haim Brézis. *Functional analysis, Sobolev spaces and partial differential equations*, volume 2. Springer, 2011.
- [12] Carlos Castro, Francisco Palacios, and Enrique Zuazua. An alternating descent method for the optimal control of the inviscid burgers equation in the presence of shocks. *Mathematical Models and Methods in Applied Sciences*, 18(03):369–416, 2008.
- [13] Carlos Castro, Francisco Palacios, and Enrique Zuazua. Optimal control and vanishing viscosity for the burgers equation. In *Integral Methods in Science and Engineering, Volume 2*, pages 65–90. Springer, 2010.




Figure 10: $T = 2$. Construction of the minimizer $S_t^+(S_T^-(u^T))$ of (17) with u^T an unattainable target defined in Example 4 using a wave-front tracking algorithm.

- [14] Christophe Chalons. Transport-equilibrium schemes for pedestrian flows with nonclassical shocks. In *Traffic and Granular Flow'05*, pages 347–356. Springer, 2007.
- [15] Robin Olav Cleveland. *Propagation of sonic booms through a real, stratified atmosphere*. PhD thesis, Citeseer, 1995.
- [16] Rinaldo M Colombo and Vincent Perrollaz. Initial data identification in conservation laws and hamilton–jacobi equations. *Journal de Mathématiques Pures et Appliquées*, 138:1–27, 2020.
- [17] Rinaldo M Colombo and Massimiliano D Rosini. Pedestrian flows and non-classical shocks. *Mathematical methods in the applied sciences*, 28(13):1553–1567, 2005.
- [18] Michael G Crandall and Andrew Majda. Monotone difference approximations for scalar conservation laws. *Mathematics of Computation*, 34(149):1–21, 1980.
- [19] Dacian N Daescu and Ionel Michael Navon. Adaptive observations in the context of 4d-var data assimilation. *Meteorology and Atmospheric Physics*, 85(4):205–226, 2004.
- [20] Constantine M Dafermos. Polygonal approximations of solutions of the initial value problem for a conservation law. *Journal of Mathematical Analysis and Applications*, 38(1):33–41, 1972.
- [21] Constantine M Dafermos. *Hyperbolic conservation laws in continuum physics*, volume 325 of *grundlehren der mathematischen wissenschaften [fundamental principles of mathematical sciences]*, 2010.
- [22] Carlos Esteve and Enrique Zuazua. The inverse problem for hamilton–jacobi equations and semiconcave envelopes. *SIAM Journal on Mathematical Analysis*, 52(6):5627–5657, 2020.
- [23] Michael Ghil and Paola Malanotte-Rizzoli. Data assimilation in meteorology and oceanography. In *Advances in geophysics*, volume 33, pages 141–266. Elsevier, 1991.




Figure 11: $T = 2$. Four minimizers u_0^* of (\mathcal{O}_T) with u^T defined in Example 4.




Figure 12: u^T and $x \rightarrow S_T^+(S_T^-(u^T))(x)$ with u^T defined in Example 5

- [24] Shyam Sundar Ghoshal, GD Veerappa Gowda, et al. Exact controllability of scalar conservation laws with strict convex flux. *Mathematical Control & Related Fields*, 4(4):401, 2014.
- [25] Olivier Glass. An extension of oleinik's inequality for general 1d scalar conservation laws. *Journal of Hyperbolic Differential Equations*, 5(1):113, 2008.
- [26] James Glimm. The interaction of nonlinear hyperbolic waves. *Communications on Pure and Applied Mathematics*, 41(5):569–590, 1988.
- [27] James Glimm and DH Sharp. An s matrix theory for classical nonlinear physics. *Foundations of physics*, 16(2):125–141, 1986.

- [28] Laurent Gosse and Enrique Zuazua. Filtered gradient algorithms for inverse design problems of one-dimensional burgers equation. In *Innovative algorithms and analysis*, pages 197–227. Springer, 2017.
- [29] Helge Holden and Nils Henrik Risebro. A method of fractional steps for scalar conservation laws without the cfl condition. *Mathematics of computation*, 60(201):221–232, 1993.
- [30] Eugenia Kalnay. *Atmospheric modeling, data assimilation and predictability*. Cambridge university press, 2003.
- [31] Stanislav N Kružkov. First order quasilinear equations in several independent variables. *Mathematics of the USSR-Sbornik*, 10(2):217, 1970.
- [32] François-Xavier Le Dimet and Olivier Talagrand. Variational algorithms for analysis and assimilation of meteorological observations: theoretical aspects. *Tellus A: Dynamic Meteorology and Oceanography*, 38(2):97–110, 1986.
- [33] Randall J LeVeque. *Numerical methods for conservation laws*, volume 132. Springer, 1992.
- [34] Yingying Li, Stanley Osher, and Richard Tsai. Heat source identification based unconstrained minimization. *Inverse Problems and Imaging*, 8(1):199–221, 2014.
- [35] Thibault Liard and Enrique Zuazua. Initial data identification for the one-dimensional burgers equation. *IEEE Transactions on Automatic Control*, pages 1–1, 2021.
- [36] J-L Lions and Bernard Malgrange. Sur l’unicité rétrograde dans les problèmes mixtes paraboliques. *Mathematica Scandinavica*, 8(2):277–286, 1961.
- [37] Andrew Majda. *The existence of multi-dimensional shock fronts*, volume 281. American Mathematical Soc., 1983.
- [38] Andrew Majda. *The stability of multi-dimensional shock fronts*, volume 275. American Mathematical Soc., 1983.
- [39] Olga Arsen’evna Oleinik. Discontinuous solutions of non-linear differential equations. *Uspekhi Matematicheskikh Nauk*, 12(3):3–73, 1957.
- [40] Yannick Privat, Emmanuel Trélat, and Enrique Zuazua. Optimal location of controllers for the one-dimensional wave equation. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 30(6):1097–1126, 2013.
- [41] Yannick Privat, Emmanuel Trélat, and Enrique Zuazua. Optimal observation of the one-dimensional wave equation. *J. Fourier Anal. Appl.*, 19(3):514–544, 2013.
- [42] Yannick Privat, Emmanuel Trélat, and Enrique Zuazua. Optimal shape and location of sensors for parabolic equations with random initial data. *Archive for Rational Mechanics and Analysis*, 216(3):921–981, 2014.
- [43] Denis Serre. *Systems of Conservation Laws 1: Hyperbolicity, entropies, shock waves*. Cambridge University Press, 1999.
- [44] Kyle Swanson, Robert Vautard, and Carlos Pires. Four-dimensional variational assimilation and predictability in a quasi-geostrophic model. *Tellus A: Dynamic Meteorology and Oceanography*, 50(4):369–390, 1998.
- [45] Olivier Talagrand and Philippe Courtier. Variational assimilation of meteorological observations with the adjoint vorticity equation. i: Theory. *Quarterly Journal of the Royal Meteorological Society*, 113(478):1311–1328, 1987.