



HAL
open science

Physiological and behavioral responses of phytoplankton communities to nutrient availability in a disturbed Mediterranean coastal lagoon

A Leruste, V. Pasqualini, M. Garrido, N. Malet, Rutger de Wit, B. Bec

► **To cite this version:**

A Leruste, V. Pasqualini, M. Garrido, N. Malet, Rutger de Wit, et al.. Physiological and behavioral responses of phytoplankton communities to nutrient availability in a disturbed Mediterranean coastal lagoon. *Estuarine, Coastal and Shelf Science*, 2019, 219, pp.176-188. 10.1016/j.ecss.2019.02.014 . hal-02390967

HAL Id: hal-02390967

<https://hal.science/hal-02390967>

Submitted on 16 Dec 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Physiological and behavioral responses of phytoplankton communities to nutrient availability
2 in a disturbed Mediterranean coastal lagoon

3 A. Leruste^{a, b*}, V. Pasqualini^a, M. Garrido^{a, c}, N. Malet^d, R. De Wit^b, B. Bec^b

4 ^a UMR SPE CNRS/UMS Stella Mare CNRS, Université de Corse, 20250 Corte, France

5 ^b MARine Biodiversity Exploitation and Conservation MARBEC, Université de Montpellier,
6 IRD, Ifremer, CNRS. Bât. 24, Place Eugène Bataillon, 34095 Montpellier cedex 5, France,

7 ^c Environmental agency of Corsica, 7 Avenue Jean Nicoli, 20250 Corte, France

8 ^d Ifremer, Laboratoire Environnement Ressources Provence-Azur-Corse (LER/PAC), Station
9 de Bastia, Z.I. Furiani, Immeuble Agostini, 20600 Bastia, France

10


11 *Corresponding author: leruste-calpena_a@univ-corse.fr

12 Running page: Phytoplankton functional response to nutrient availability

13 Keywords: growth rate, dilution experiment, eutrophication, functional traits

14 Abstract

15 Short-term bioassays were conducted in Biguglia lagoon (Corsica) to study the physiological
16 and behavioral responses of phytoplankton to N- and P-availability. Natural communities
17 were collected in two stations representative of the two sub-basins, at three periods with
18 contrasting environmental characteristics to address the impact of seasonal variability. These
19 samples were separately enriched with a full N and P enrichment, and with enrichments
20 minus N or minus P. Phytoplankton size structuration, diversity, and growth of the total
21 phytoplankton, the micro-, nano- and ultraphytoplankton were evaluated using
22 spectrofluorimetry, and optical microscopy. Results showed that the communities were fueled
23 by NO_3^- in the wet periods (autumn and spring) and NH_4^+ in summer. The phytoplankton
24 communities displayed highest cell size in autumn, with high abundances of nanoflagellates,
25 and smallest cell size in summer with a large dominance of phycocyanin-rich
26 picocyanobacteria. Blooms of dinoflagellates also occurred during the wet periods, coinciding
27 with high N:P ratios. The full enrichment has not stimulated phytoplankton growth in autumn,
28 suggesting the importance of other controlling factors such as light, a possible NH_4^+ inhibition
29 or the use of mixotrophic abilities. In spring, communities have displayed single P-limitation
30 in the northern basin and different N and P co-limitations in the southern basin. In summer,
31 the full enrichment consistently stimulated the growth of all cell sizes. The communities
32 showed high N and P co-limitations, which is consistent with growing observations in aquatic
33 ecosystems, and reflects the different functional responses of phytoplankton communities to
34 the nutrient availability.



35

36

37 Graphical abstract. Chlorophyll *a* (Chl *a*) concentration, DIN:DIP ratio, main
 38 taxa/morphotypes, response to the full nutrient enrichment, and N- and/or P-limitation during
 39 the three bioassay periods in the two stations representatives of the Northern and the Southern
 40 basins of Biguglia lagoon.

41

42 Highlights

43 • Freshwater inputs from the watershed strongly influenced the phytoplankton of
 44 Biguglia lagoon

45 • Nanoflagellates dominated phytoplankton communities in autumn and spring with
 46 high N:P ratios

47 • Phycocyanin-rich picocyanobacteria bloomed in the sampling in September with low
 48 N:P ratios

49 • Phytoplankton showed differential responses to nutrient enrichments and induced
 50 limitations

- 51 • Phytoplankton was generally co-limited by N and P, with exceptions depending on
52 seasons
53

54 1. Introduction

55 Coastal lagoons are very productive ecosystems at the sea-land interface hosting a high
56 degree of biodiversity and providing numerous ecosystem services for human wellbeing
57 (Barbier et al., 2011). These services confer a high economical value for the coastal lagoons,
58 *e.g.* as water and food supply (fishing, shellfish farming), tourism and recreation (Franco et
59 al., 2010; Rochette et al., 2010). However, because of their location in densely populated
60 littoral areas and the services they provide, coastal lagoons and their surroundings have been
61 increasingly exploited for human uses. Overexploitation combined with the hydromorphology
62 of these semi-enclosed systems and climate change have increased their vulnerability to
63 anthropogenic impacts. Hence, coastal lagoons are among the most threatened ecosystems
64 (Nixon, 2009). This threat is mainly associated with changes in their hydrology and with an
65 increase of nutrients and pollutants inputs from sewage effluents and watershed run-off
66 (Mouillot et al., 2000; Cloern, 2001). Due to shallowness and confinement, these inputs result
67 in high concentration of nutrients and pollutants in coastal lagoons, thereby exacerbating their
68 degradation (Donald et al., 2013).

69

70 An excessive or unbalanced input of nutrients disturbs the community of primary producers,
71 leading to an increase of the eutrophication processes and to drastic changes in the
72 biodiversity of the autotrophic compartment. Thus, it may promote fast-growing opportunistic
73 algae in the first place and finally phytoplankton at the expense of benthic organisms such as
74 macrophytes, and may result in an increase of the frequency of harmful algal blooms
75 (Schramm, 1999; Livingston, 2000; Le Fur et al., 2018). These changes in the composition of
76 the autotrophic communities impact all the trophic levels, altering the ecosystem functioning
77 (Cloern, 2001). High nutrient inputs can also, through an accumulation of macroalgal or
78 phytoplankton biomass, lead to the development of dystrophic crisis and hypoxia and anoxia

79 events, having dramatic impacts on all the living organisms (Cloern, 2001). Therefore, the
80 management of a eutrophicated coastal lagoon needs to target the reduction of nutrient
81 loadings. In addition, knowledge is required about the availability of nutrients within lagoons
82 and the autotrophic communities responses (Duarte et al., 2000; Glibert et al., 2015). This
83 allows to assess the resilience of the aquatic communities, and to estimate the critical nutrient
84 loading for the ecosystem (Pasqualini et al., 2017; Le Fur et al., 2018).

85
86 Phytoplankton represents the basis of food webs and biochemical cycles, and is generally the
87 first autotrophic compartment responding to a change of nutrient availability (Leruste et al.,
88 2016). Understanding how the ecosystem functioning influences the composition of
89 phytoplankton communities and their growth mechanisms is indispensable (Livingston, 2000;
90 Paerl et al., 2003). Hence, it is particularly important to understand and predict how N and P
91 inputs affect the growth and community assembly of phytoplankton. These responses are
92 indicators of water quality, and are consequently informative for water management (Duarte
93 et al., 2000; Reed et al., 2016). Morphological and physiological traits such as cell size and
94 maximal growth rates shape the phytoplankton functional responses to nutrient availability.
95 Small-size fast-growing algae are competitive at low nutrient concentrations and low light,
96 while larger cells can generally thrive under a pulsed nutrient supply and higher light
97 intensities (Litchman et al., 2007; Bec et al., 2011; Andersen et al., 2015). The community
98 composition also reflects which nutrients are available. Diatoms are particularly competitive
99 for nitrate, while green algae and picocyanobacteria are more competitive for ammonium. The
100 latter often originates from regenerated internal sources. Dinoflagellates can develop even
101 under dissolved inorganic nutrient limitation, because of their relatively low growth rates and
102 potential mixotrophic abilities (Litchman et al., 2007; Glibert et al., 2015).

103

104 The general aim of this study was to explore the responses of the phytoplankton communities
105 of a disturbed Mediterranean coastal lagoon to the nutrient availability. Therefore, we choose
106 Biguglia lagoon, which is the largest lagoon on the island of Corsica, as the study site. This
107 Mediterranean lagoon has been increasingly impacted since 1980 by a strong increase of the
108 human population density, industrial and agricultural uses in its catchment and its surrounding
109 areas. This lagoon receives particularly high amounts of nitrate from runoffs, tributaries, and
110 from groundwater flows from a coastal aquifer connected with the lagoon (Erostate et al.,
111 2018). Biguglia lagoon also displays a high confinement and restricted exchanges with the
112 Tyrrhenian Sea that have induced higher levels of nutrient concentration than in most other
113 French Mediterranean lagoons these last decades (Souchu et al., 2010; Pasqualini et al.,
114 2017). Hence, the lagoon has experienced strong changes in its ecological state, *i.e.* increasing
115 eutrophication and pollutant levels linked with changes in the composition of its autotrophic
116 community (Mouillot et al., 2000; Pasqualini et al., 2017). Moreover, a dystrophic crisis
117 associated with a bloom of toxic cyanobacteria in 2007 has raised the awareness of the water
118 quality in the lagoon among the responsible management authorities. Therefore, they have
119 taken remediation measures to reduce the confinement in order to decrease the concentration
120 of nutrients. This included the modification of the lagoon hydrology since 2009 to increase
121 dilution processes (Cecchi et al., 2016; Garrido et al., 2016). However, so far, no efficient
122 direct management measures have been implemented to reduce the external nutrient loadings
123 that remain important and uncontrolled. Moreover, although the measures taken in 2009 may
124 have helped the development of aquatic angiosperms (Pasqualini et al., 2017), these may have
125 also promoted the occurrence of blooms of potentially toxic and or harmful dinoflagellate
126 species (Cecchi et al., 2016).

127

128

129 The specific objectives of this study were to describe in different seasons with contrasting
130 environmental conditions including nutrient form and origin, (1) the structure of
131 phytoplankton communities in Biguglia lagoon submitted to a strong anthropogenic impact in
132 term of size classes and species diversity, and (2) their growth rate in response to various
133 forms of nutrients inputs. These two observations illustrate phytoplankton functional
134 responses to contrasting nutrient availability, and the latter specially reflects the vulnerability
135 of the whole ecosystem to a short-term nutrient pulse. We experimentally incubated
136 phytoplankton communities under *in situ* light and temperature conditions with enrichments
137 containing N (NO_3^- , NH_4^+) and P (PO_4^{3-}), either in combination or separately to selectively
138 induce either N- or P-limitation. Exploring the responses of phytoplankton communities to a
139 change in nutrient availability helped understanding the impact of these changes on the
140 ecosystem functioning.

141

142 2. Material and Methods

143 2.1. *Characterization of the study site*


144 Biguglia lagoon (42°36'N; 9°28'E) is a shallow brackish coastal lagoon (average depth 1.2
145 m), located on the East coast of Corsica, and separated from the Tyrrhenian Sea by a sandy
146 beach barrier (Fig. 1). This choked lagoon *sensu* Kjerfve (1994) covers 14.5 km² with a single
147 inlet in the north to the sea that consists of a 1.5 km long, narrow and shallow natural channel.
148 Biguglia lagoon is divided into two sub-basins by a peninsula in its middle. Each of these sub-
149 basins displays a specific hydrological functioning resulting in differences in salinity, nutrient
150 concentrations and phytoplankton biomasses (Garrido et al., 2016).

151

152 The microtidal conditions, and the morphology of this inlet and its natural inclination to silt
153 up restrict the exchanges with the sea and thus limit marine water input in the lagoon

154 (Mouillot et al., 2000). This lagoon is highly influenced by freshwater inputs from its
155 watershed (182 km²), and its salinity remains particularly low (Pasqualini et al., 2017). For
156 example, this ranged from 2.0 to 18.5 between November 2013 and December 2014.
157 Freshwater comes from rainfall, sewage plants, several rivers draining the watershed and
158 flowing in the north-west of the lagoon, the Golu River connected to the lagoon by the
159 Fossone channel in the south-western part, and from pumping stations draining the
160 agricultural plain in the southern part of the watershed (Erostate et al., 2018) (Fig. 1).
161 Therefore, Biguglia lagoon displays a salinity gradient from the north to the south (Garrido et
162 al., 2016).

163



164

165 Fig. 1. Location of the two stations representative of the Northern Basin (NB) and the
166 Southern Basin (SB) of Biguglia lagoon.

167

168 Biguglia lagoon has been increasingly eutrophicated since the 1980s due to an intensification
169 of human activities and urbanization on its watershed southward of city of Bastia, including
170 four towns hosting approximately 35,000 inhabitants in winter (Lafabrie et al., 2013). During
171 summer touristic period, the population doubles, which results in increasing anthropogenic
172 pressure. Hence, this lagoon displays very high nutrients concentrations (NH_4^+ , NO_2^- , NO_3^- ,
173 DIN, Si, TP, TN) in the water column, enhanced by the reduced exchanges with the sea
174 (Orsoni et al., 2001). The sediment compartment shows an important silting with high nutrient
175 concentrations and organic matter content compared to other French Mediterranean coastal
176 lagoons (Souchu et al., 2010). An eutrophication gradient is also observed from the south to
177 the north because of substantial anthropogenic inputs in the southern part of the lagoon.

178

179 The lagoon has a Mediterranean climate with a succession of a dry and warm summer and wet
180 autumn, winter and spring seasons with sudden and plentiful rainfalls causing flash floods
181 (Pasqualini et al., 2017). The salinity and nutrients inputs are strongly influenced by
182 theseasonal variability of the climate (Collos et al., 2003). This generally determines three
183 hydrological seasonal periods characterized by differences in their nutrient origin and
184 availability, in phytoplankton biomass and size class structure, and in photosynthetic
185 performance (Cecchi et al., 2016; Garrido et al., 2016).

186

187 *2.2. Sampling procedures*

188 Water samples were collected in two stations representatives of the northern (NB) and the
189 southern (SB) basins (Fig. 1, $42^\circ38'12''\text{N}$, $9^\circ27'15''\text{E}$ and $42^\circ35'00''\text{N}$, $9^\circ29'18''\text{E}$,
190 respectively).

191

192 Experiments were carried out in autumn 2013 (26 November to 5 December), spring 2014 (2
193 to 8 April), and summer 2014 (9 to 12 September). In each of the two NB and SB stations, 70
194 l of pre-filtered water through 1000 μm mesh to remove larger debris without removing
195 zooplankton or larger phytoplankton cells (Collos et al., 2005) were sampled in sub-surface
196 (20 cm depth) with a pump from a small boat between 7.00 am and 8.00 am, and kept in the
197 dark. Sub-surface salinity, temperature, percentage of dissolved oxygen (DO) and turbidity
198 were measured *in situ* with a multi parameter Water Quality Probe (YSI® 6600 V2-2). Upon
199 return to the laboratory, water samples were homogenized by gentle shaking, aliquoted and
200 immediately stored at -20°C for the characterization of phytoplankton communities. Measures
201 of NH_4^+ , PO_4^{3-} , NO_3^- , NO_2^- , TN, and TP concentrations (μM) were performed on duplicates
202 of 80 ml, filtered on 20 μm meshes for the dissolved inorganic nutrients (Aminot and
203 Chaussepied, 1983).

204

205 2.3. Characterization of phytoplankton biomass

206 Chlorophyll *a* (Chl *a*) concentrations ($\mu\text{g l}^{-1}$) were used as a proxy for phytoplankton
207 biomass. Chl *a* measurements were performed on the total and the fractioned water samples.
208 This allowed assessing the contribution of the ultraphytoplankton $<5 \mu\text{m}$, the
209 nanophytoplankton between 5 and 20 μm and the microphytoplankton $>20 \mu\text{m}$ to the total
210 biomass. Upon return to the laboratory, triplicates of water samples (maximum 100 ml,
211 depending on phytoplankton abundance) and of samples pre-filtered on 20 μm and on 5 μm
212 meshes were analyzed by spectrofluorimetry (Neveux and Lantoiné, 1993). The biomasses of
213 the three size classes were estimated by subtracting the concentrations obtained for the
214 fraction $<20 \mu\text{m}$ to the total water (microphytoplankton), and the fraction $<5 \mu\text{m}$ to the
215 fraction $<20 \mu\text{m}$ (nanophytoplankton). The Chl *a* concentration of the fraction $<5 \mu\text{m}$
216 estimated the ultraphytoplankton biomass.

217

218 *2.4. Assessment of phytoplankton community composition*

219 Composition and abundance of the ultraphytoplankton <5 µm were studied using flow
220 cytometry. Triplicates of 1 ml of sample fixed with 2 % formaldehyde (filtered on 0.2 µm,
221 final concentration) were submitted to a flash freeze in liquid nitrogen and then stored at -
222 80°C until analysis. Cells counts were performed with a FACSCalibur flow cytometer
223 (Becton Dickinson), using beads for size calibration. Populations of coccoid phycoerythrin-
224 rich picocyanobacteria (PE-cyanos) were identified by light diffraction (FSC) and their
225 orange fluorescence emissions, while phycocyanin-rich picocyanobacteria (PC-cyanos) were
226 distinguished by their low red fluorescence and their absence of orange fluorescence.
227 Photosynthetic picoeukaryotes <3 µm were identified by their red fluorescence emissions
228 (Chl *a*, wavelength >650 nm).

229

230 Optical microscopy was used to describe nano- and microphytoplankton (>5 µm) composition
231 and abundance. Triplicates of 1 l of sample fixed with formaldehyde (5 % final concentration)
232 were stored at obscurity prior to analysis. A modified method of the Utermöhl protocol was
233 used (Leruste et al., 2018). Samples were examined with an optic microscope Olympus
234 AX10. At least 400 cells per sample were counted to obtain a relevant assessment of the
235 assemblage. Taxonomic resolution was realized at species level whenever possible, and
236 identification was verified according to several books (Bellinger and Sigee, 2015; Bérard-
237 Therriault et al., 1999; Loir, 2004; Tomas, 1997) and database such as the World Register of
238 Marine Species (<http://www.marinespecies.org/>, databases available online). Taxonomic
239 diversity was explored using species richness and Shannon index (Leruste et al., 2018).

240

241 2.5. *Experimental procedure of bioassay experiments*

242 To measure maximal growth rate, five dilutions of water sample (9, 17, 43, 74 and 100 %) in
243 0.2 μm filtered sample in duplicate were incubated with an enrichment containing vitamins,
244 silica, metals trace, nitrogen and phosphorus (NaH_2PO_4 , final concentration 0.8 μM) (Landry
245 and Hassett, 1982). Nitrogen was supplied as nitrate and/or ammonium (f/2 and h/2 media,
246 respectively, Guillard & Ryther, 1962) depending on season. In April 2014, N was supplied as
247 nitrate (20 μM final concentration), assuming that nitrate inputs from the watershed were the
248 main nitrogen source. In September 2014, N was supplied as ammonium (20 μM final
249 concentration), assuming that this may be the predominant form provided from
250 remineralization processes in the sediment (Collos et al., 2003). In November-December
251 2013, N was supplied as a mix of nitrate and ammonium (10 μM final concentration each),
252 assuming that sufficiently high temperatures allow the regeneration of ammonium from the
253 sediments, associated with flash floods bringing nitrate from the watershed. This nitrogen
254 concentration has been chosen to avoid phytoplankton growth limitation during the
255 incubation, which has been occasionally observed at 10 μM in another Mediterranean lagoon
256 (Bec et al., 2005). To highlight the potential growth limitation of phytoplankton by N and/or
257 P, two other dilution series were enriched either without nitrogen or phosphorus (Andersen et
258 al., 1991). Two bottles of sample without dilution and enrichment were incubated as control,
259 and the all bottles were simultaneously incubated for 24 h in Biguglia lagoon under *in situ*
260 temperature and light conditions at 30 cm depth.

261

262 2.6. *Estimation of maximal growth rate and N-/P-limitation*

263 After 24 h incubation, the temporal changes of the total and size-fractionated Chlorophyll *a* in
264 each bottle were used to calculate their apparent specific growth rate $k(x)$ for each dilution x .

265 The relationship between the apparent growth rates $k(x)$ in the fully-enriched bottles and
266 dilution factor x were fitted to the linear equation (1):

$$267 \quad k(x) = \mu_{\max} - gx \quad (1)$$

268 Where $k(x)$ is the phytoplankton apparent growth rate in the bottles at dilution x , μ_{\max} is the
269 maximal growth rate under nutrient replete conditions, and g is the grazing rate. All rates
270 were expressed on a per day basis (d^{-1}). The maximal growth rate μ_{\max} was obtained from the
271 Y axis intercept, and the phytoplankton mortality rates g as the slope of the linear regressions
272 assuming that mortality was due to grazing (Landry and Hassett, 1982). Equations were fitted
273 with the 'lmer' function in the 'lme4' library (version 1.1-10, (Bates et al., 2015)) to evaluate
274 if the linear model best fitted the trend of the relationship between the apparent growth rate
275 and the dilution factor. To check whether the linear equation (1) was acceptable, we also
276 considered a quadratic function by adding the term ax^2 . The quadratic function has a parabolic
277 pattern and could reflect the contribution of regenerated resources or a saturation of the
278 grazing rate in the less diluted bottles (Andersen et al., 1991). All combinations of the
279 parameters of the quadratic function were considered, and model selection was based on
280 parsimony using the small-sample corrected Akaike's information criterion (AIC_c) (Burnham
281 and Anderson, 2004). Therefore, we calculated the difference between the AIC_c of the linear
282 model and the model having the lowest AIC_c , *i.e.* the most parsimonious model, to obtain a
283 ΔAIC_c value. The models with a $\Delta\text{AIC}_c < 2$ were considered as having equivalent levels of
284 support (Burnham and Anderson, 2004), using the 'dredge' function of the MuMIn package
285 (Bartón, 2013). Hence for $\Delta\text{AIC}_c < 2$ we selected the linear model as it allows calculating μ_{\max}
286 and g according to equation (1) (Landry and Hassett, 1982), while the quadratic relationship
287 did not allow calculating μ_{\max} and g .

288

289 Phytoplankton net growth rate in controls (μ_0) was calculated by adding g to the apparent
290 growth rate (k_0) *i.e.* $\mu_0 = g + k_0$. Hence, it was assumed that grazing rate was not impacted by
291 nutrient enrichments. The $\mu_0:\mu_{\max}$ ratio assessed the impact of inorganic nutrient enrichment
292 on growth, and estimated the nutrient sufficiency for phytoplankton growth (Landry and
293 Hassett, 1982; Landry et al., 1998).

294

295 In the selectively enriched bioassays, we took account that apparent growth rate is not
296 necessarily a linear function of the dilution factor, because the instantaneous growth rate may
297 decline as internal and external nutrient pools become depleted. μ_{-N} and μ_{-P} were calculated as
298 followed: $\mu_{-N} = g + k_{-N}$ and $\mu_{-P} = g + k_{-P}$. To estimate how much phytoplankton growth was N-
299 and/or P-limited, the ratio between growth rate with the enrichments minus N or minus P (μ_{-N}
300 or μ_{-P}) and maximal growth rate in theoretically non-limiting nutrient condition (μ_{\max}) was
301 calculated. Differences between growth rates with full enrichment, enrichment minus N and
302 enrichment minus P helped distinguish different types of co-limitation (Harpole et al., 2011;
303 Burson et al., 2016).

304

305 2.7. Statistical analyses

306 Data analysis was performed on R (R Core Team, 2013).

307 Two-way ANOVAs assessed if the total phytoplankton biomass significantly changed among
308 the three samplings and between NB and SB. Posteriori pairwise mean comparisons between
309 stations and samplings were made using Tukey-Kramer's post-hoc analysis (alpha level =
310 0.05). Permanovas assessed if the size class structure varied among stations and samplings,
311 according to differences of biomass, temperature, salinity, dissolved inorganic nutrients, TN
312 and TP concentrations, turbidity and DO (Anderson, 2001). Taxonomic diversity indices *i.e.*
313 the Shannon index on the relative abundance of taxa and morphotypes (H), and the species

314 richness (S) were estimated using Vegan package (Oksanen et al., 2018). ANOVAs assessed
315 if S and H significantly changed among stations and samplings. Spearman's rank correlations
316 between S, H, total Chl *a* biomass and environmental variables (temperature, salinity,
317 inorganic nutrient concentrations, TN and TP concentrations, turbidity and DO) were assessed
318 for the two stations and the three samplings.

319

320 Regression analyses determined which mixed models (*i.e.* combinations of explanatory
321 factors) best fitted the biomass of total phytoplankton and the three size classes. Among all
322 the possible fixed effects (stations, samplings, salinity, temperature, nutrients concentrations,
323 dissolved oxygen, and turbidity), the most relevant were chosen according to a correlation
324 matrix. Values of fixed effects were scaled prior to analysis. Because of pseudo-replication
325 (each triplicate was taken from the same bottle), a random effect was added as a bottle effect.
326 Models were fitted with the 'lmer' function and model selection was assessed using the AIC_c
327 (see below). To estimate the relative importance of each fixed effect, the AIC_c-w of every
328 model was summed (Patiño et al., 2013), and to estimate the variable force effect, we used the
329 mean of the relative coefficient weighted by the AIC_c-w.

330

331 3. Results

332 3.1. *Environmental variables and phytoplankton biomass*

333 Salinities were low during the three dilution experiments, ranging from 2.0 in NB in April, to
334 10.9 in September 2014 (Table 1). The percentage of DO was the lowest in September 2014
335 for both stations, and was negatively correlated with the temperature for the three sampling
336 periods (Table 2, Spearman's rank correlations, *p*-value <0.05). Turbidity was highest in the
337 autumn sampling (Table 1).

338

339 Concentrations of dissolved inorganic nutrients (NH_4^+ , NO_3^- , NO_2^- , DIN, PO_4^{3-}) and of TN
340 and TP were extremely variable. In the sampling in November-December 2013, both stations
341 presented the highest PO_4^{3-} concentrations, and NB displayed the highest concentration of
342 dissolved inorganic nutrients, DIN being mainly composed of NO_3^- (Table 1). In April 2014,
343 N was mainly represented by NO_3^- in the two stations, and N levels were especially high in
344 SB, with 106.5 μM of TN and 95.4 μM of DIN. Inversely, in September 2014, both stations
345 displayed low NO_3^- concentrations ($<0.15 \mu\text{M}$) and NH_4^+ was the main nitrogen form. TP was
346 higher in SB than in NB in all three samplings (Table 1). All dissolved inorganic nutrient
347 concentrations and TN and TP concentrations were negatively correlated with the
348 temperature. Moreover, NH_4^+ concentrations were also negatively correlated with the salinity.
349 DIN forms were positively correlated between them, and NO_3^- and NO_2^- concentrations were
350 also positively correlated with TN and TP concentrations. PO_4^{3-} concentrations were
351 positively correlated with the percentage of DO (Table 2). The TN:TP and DIN:DIP ratios
352 displayed highest values in April 2014, respectively 71.0 in SB and 124.5 in NB.

353

354 The biomass of the total phytoplankton significantly differed among stations and samplings,
355 and the differences among samplings also changed between the two stations (Fig. 2, ANOVA,
356 p -value <0.05). Chlorophyll *a* (Chl *a*) concentration generally ranged between 3.5 and 6.0 μg
357 l^{-1} , with the exception in SB in December 2013 when it peaked at 20.6 $\mu\text{g l}^{-1}$ (Fig. 2A). Taking
358 all samplings collectively, we found that Chl *a* concentration showed a negative relationship
359 with the salinity and NH_4^+ concentrations, the latter effect changing among stations
360 (Regression analyses, force of the salinity effect: -4.5, and of the NH_4^+ concentration: -8.1).
361 Total biomass was significantly higher (i) in SB than in NB, (ii) in the autumn sampling than
362 in April and in September 2014, as well as (iii) in April than in September samplings (Fig. 2,
363 pairwise Tukey multiple comparisons, $p <0.05$). The total biomass was negatively correlated

364 with temperature, and positively correlated with NO_2^- , NO_3^- , PO_4^{3-} concentrations and with
365 the percentage of DO, reflecting the peak of biomass observed in December in SB (Table 2,
366 Spearman's rank correlations, p -value <0.05).


367 Table 1. Mean values of environmental parameters in the two NB and SB stations of Biguglia lagoon in November-December 2013, April and
 368 September 2014. DO: dissolved oxygen. Dissolved inorganic nitrogen is expressed as $DIN = NH_4 + NO_3 + NO_2$. Dissolved inorganic phosphorus
 369 corresponds to PO_4^{3-} concentration.

Sampling periods	Stations	Date	T°(C)	Salinity	DO (%)	Turbidity (NTU)	TN	NH ₄ ⁺	NO ₃ ⁻	NO ₂ ⁻	DIN	TP	PO ₄ ³⁻	TN:TP	DIN:DIP
							(μM)								
Nov -Dec 2013	NB	26/11/13	8.3	4.7	100.8	2.87	84.37	7.52	70.01	0.49	78.03	1.37	0.73	61.4	107.3
	SB	04/12/13	8.2	6.1	101.5	16.83	61.87	0.69	40.72	0.29	41.70	1.77	0.64	34.9	65.5
April 2014	NB	07/04/14	15.0	2.0	113.0	0.80	34.55	2.18	17.47	0.16	19.80	0.82	0.16	41.9	124.5
	SB	02/04/14	15.5	6.0	93.2	4.63	106.47	1.85	93.22	0.30	95.38	1.50	0.00	71.0	-
September 2014	NB	11/09/14	24.8	10.9	97.0	1.83	38.51	0.37	0.03	0.00	0.40	0.89	0.03	43.3	13.3
	SB	09/09/14	25.9	6.0	86.2	10.50	29.95	1.24	0.11	0.09	1.44	1.17	0.11	25.6	13.1

370

371 Table 2. Spearman's rank correlation between species richness (S), Shannon index (H), Chl *a* concentrations and environmental variables at the
 372 95% confidence interval for the three sampling periods in both NB and SB stations of Biguglia lagoon. Significant correlations are in bold. DO:
 373 dissolved oxygen. ****p*-value <0.001, ***p*-value <0.01, **p*-value <0.05.

	S	H	Chl <i>a</i>	Temperature	Salinity	NH ₄ ⁺	NO ₂ ⁻	NO ₃ ⁻	PO ₄ ³⁻	TN	TP	Turbidity
H	-0.314											
Chl <i>a</i>	-0.312	0.626**										
Temperature	0.469*	-0.661**	-0.900***									
Salinity	-0.051	-0.693**	-0.242	0.203								
NH ₄ ⁺	-0.066	0.680**	0.260	-0.314	-0.928***							
NO ₂ ⁻	-0.513*	0.536*	0.530*	-0.657**	-0.464	0.714***						
NO ₃ ⁻	-0.494*	0.335	0.492*	-0.543*	-0.377	0.600**	0.943***					
PO ₄ ³⁻	-0.405	0.836***	0.709***	-0.880***	-0.400	0.516*	0.637**	0.395				
TN	-0.280	-0.009	0.360	-0.486*	0.058	0.257	0.771***	0.829***	0.273			
TP	-0.809***	0.047	0.398	-0.486*	0.319	-0.086	0.600**	0.657*	0.273	0.657**		
Turbidity	-0.828***	-0.041	0.122	0.143	0.406	-0.314	0.200	0.257	0.030	0.143	0.829***	
DO	0.041	0.624**	0.774***	-0.771***	-0.348	0.257	0.200	0.086	0.698**	0.029	-0.143	-0.371



374

375 Fig. 2. Mean chlorophyll *a* (Chl *a*) concentration ($\mu\text{g l}^{-1}$) (A) in the three samplings at the NB
 376 and SB stations in Biguglia lagoon. Error bar indicates standard deviations. Proportions of the
 377 different size classes of phytoplankton contributing to phytoplankton biomass (B), *i.e.*
 378 microphytoplankton $>20 \mu\text{m}$ (dark grey), nanophytoplankton between 5 and $20 \mu\text{m}$ (grey) and
 379 ultraphytoplankton $<5 \mu\text{m}$ (light grey).

380

381 3.1. Phytoplankton community composition

382 The size class structuration of phytoplankton communities was significantly different between
 383 the NB and SB stations among the three samplings (Fig. 2B), and the differences between
 384 samplings changed between stations (Permanova, p -value <0.05). Taking all samplings
 385 collectively, the ultra- and the nanophytoplankton biomasses showed a positive relationship
 386 with the salinity and the phosphorus concentrations. In contrast, the microphytoplankton
 387 biomasses showed a positive relationship with NO_3^- concentrations and a negative

388 relationship with NH_4^+ concentrations (Regression analyses, $\Delta\text{AICc} < 2$).
389 Ultraphytoplankton was the major group in the phytoplankton communities (up to 87 % of
390 Chl *a* in September 2014) (Fig. 2B), except for NB in November 2013, when a higher
391 proportion of nanophytoplankton (40 %) than ultraphytoplankton (34 %) was observed. For
392 both stations, the nanophytoplankton biomass was highest in November 2013.

393

394 The taxonomic diversity indices showed different patterns at the two stations and among the
395 three samplings. Both the species richness (S) and the Shannon index (H) significantly
396 changed among stations and samplings (ANOVAs, both p -values < 0.05). S was the lowest in
397 the autumn sampling and highest in the sampling in September for the two stations, while H
398 showed the inverse trend. Hence, S was positively correlated with temperature, and negatively
399 correlated with NO_2^- , NO_3^- , TP concentrations and turbidity (Table 2). H was positively
400 correlated with total biomass, NH_4^+ , NO_2^- and PO_4^{3-} concentrations and percentage of DO,
401 while it was negatively correlated with temperature and salinity (Table 2).

402

403 Fig. 3 gives the contributions of four size classes ($< 3 \mu\text{m}$, between 3 and 5 μm , between 5 and
404 20 μm and $> 20 \mu\text{m}$) to the total abundances of cell counts in the two stations at the three
405 sampling periods, and Fig. 4 specifies the main taxonomic classes represented in each of the
406 size classes. For the three samplings, small cells dominated the phytoplankton communities
407 (Fig. 3). The NB station showed higher Bacillariophyceae abundances than SB, while the SB
408 station showed higher Chlorophyta and Cryptophyceae abundances (Fig. 4). NB has also
409 displayed higher dinoflagellates abundances than SB, except in November-December 2013
410 (Fig. 4).

411


412 In the sampling in November 2013, the phytoplankton community in NB showed a bloom of
413 the dictyochophycean *Apedinella radians* (Lohmann) P.H. Campbell, 1973 (1.3×10^6 cells.l⁻¹,
414 *i.e.* 41.6 % of total abundances, Fig. 3-4). This community also comprised high abundances of
415 eukaryotes with flagella between 3 and 5 μ m (6.9×10^5 cells.l⁻¹, Fig. 3-4), and of
416 *Prorocentrum cordatum* (Ostenfeld) J.D.Dodge, 1975 *alias Prorocentrum minimum*
417 (Pavillard, 1916) J.Schiller, 1933 (2.9×10^5 cells.l⁻¹, Fig. 4). Dinoflagellates dominated the
418 community in SB in December, mainly represented by *Heterocapsa minima* Pomroy, 1989
419 (5.1×10^6 , *i.e.* 34.2 % of total abundances Fig. 3), associated with *A. radians* (4.8×10^6 cells.l⁻¹,
420 Fig. 4). Microphytoplanktonic cells were scarce (*e.g.* *Kryptoperidinium foliaceum*
421 Lindemann, 1924: 3.7×10^3 cells.l⁻¹; *Cocconeis* sp.: 3.7×10^3 cells.l⁻¹; *Prorocentrum micans*
422 Ehrenberg 1834: 1.5×10^3 cells.l⁻¹). Conversely, we observed a bloom of *Mesodinium rubrum*
423 Lohmann, 1908 (2.3×10^5 cells.l⁻¹, *i.e.* 78 % of the microplanktonic cells, Fig. 4).

424

425 In the sampling in April 2014, dinoflagellates dominated the NB community mainly due to a
426 bloom of *H. minima* (3.7×10^6 cells.l⁻¹, *i.e.* 35.8 % of total abundances, Fig. 3-4).
427 Bacillariophyceae were also abundant and distributed among the three size classes $>3 \mu$ m (3.1
428 $\times 10^6$ cells.l⁻¹, Fig. 4) with *e.g.* *Thalassiosira* sp., *Diatoma* sp., and *Chaetoceros* spp.. The SB
429 community was dominated by phycocyanin-rich picocyanobacteria (PC-cyanos, 6.4×10^7
430 cells.l⁻¹, *i.e.* 49.6 % of total abundances, Fig. 3-4), Picoeukaryotes (Peuk) without flagella (5.6
431 $\times 10^7$ cells.l⁻¹) and eukaryotes with flagella between 3 and 5 μ m (5.9×10^6 cells.l⁻¹, Fig. 4).
432 The two latest morphotypes probably belonged to the green algae according to the pigment
433 composition (data not shown). *M. rubrum* also bloomed as in December 2013 (1.6×10^5
434 cells.l⁻¹, Fig. 4).

435


436 In the sampling in September 2014 the communities of the two stations were dominated by
437 PC-cyanos (9.4×10^8 cells.l⁻¹, *i.e.* 92.7 % of total abundances in NB, 3.0×10^8 cells.l⁻¹ *i.e.*
438 91.9 % of total abundances in SB, Fig. 3-4) and Peuk without flagella (6.5×10^7 cells.l⁻¹ in
439 NB, 1.7×10^7 cells.l⁻¹ in SB, Fig. 3-4), that most likely belonged to the green algae according
440 to the pigment composition (data not shown). The nanophytoplankton in NB displayed high
441 abundances of *Tetraselmis* sp. (2.0×10^5 cells.l⁻¹), *P. cordatum* (8.2×10^4 cells.l⁻¹) and *H.*
442 *minima* (3.2×10^4 cells.l⁻¹). Dinoflagellates dominated the microphytoplankton with
443 *Prorocentrum* sp. *cf. mexicanum* Osorio-Tafall, 1942, and *Gonyaulax* sp. (10^4 cells.l⁻¹). We
444 also observed high abundances of *Chaetoceros* spp., and *Ceratoneis closterium* Ehrenberg,
445 1839 (10^4 cells.l⁻¹). The ultra- and the nanophytoplankton in SB also contained high
446 abundances of *A. radians* (1.6×10^6 cells.l⁻¹), *Pyramimonas* sp. (8.9×10^5 cells.l⁻¹),
447 *Plagioselmis* sp., *H. minima*, *P. minimum* and nanoplanktonic centric diatoms (10^4 cells.l⁻¹).
448 *C. closterium*, and *Gonyaulax* sp. *cf. spinifera* (10^4 cells.l⁻¹) were the most abundant
449 microplanktonic cells in SB (Fig. 4).



450

451 Fig. 3. Contribution of the size classes $<3 \mu\text{m}$, $[3-5 \mu\text{m}]$, $[5-20 \mu\text{m}]$ and $>20 \mu\text{m}$ to the total
 452 abundances of cell counts in the two NB and SB of Biguglia lagoon at the three sampling
 453 periods. Total abundances are specified on the top of the bars.

454



456 Fig. 4. Contribution of the main taxonomic classes to the abundances of the four size classes
457 <3 μm , [3-5 μm], [5-20 μm] and >20 μm in the two stations of Biguglia lagoon at the three
458 sampling periods.

459

460 3.2. *Phytoplankton net growth and mortality rates without enrichment*

461 Phytoplankton net growth rates (μ_0) without enrichment ranged from negative values for the
462 microphytoplankton of NB in November 2013 and April 2014, to 1.91 d^{-1} for the
463 nanophytoplankton of SB in April 2014 (Table 3). In November-December 2013, the total
464 phytoplankton displayed low net growth rates (0.45 d^{-1} in NB and 0.48 d^{-1} in SB), and similar
465 mortality rates (0.62 d^{-1} in NB to 0.69 d^{-1} in SB). In NB the ultraphytoplankton dominated by
466 picoeukaryotes (Peuk) displayed a two-fold higher growth rate, while its mortality rate was
467 three times higher than those of the total phytoplankton (Table 3). The nanophytoplankton,
468 mainly comprising *A. radians* and *P. cordatum*, and particularly the microphytoplankton
469 comprising diatoms and dinoflagellates species, showed a loss of biomass after 24 h
470 incubation (Table 3). In SB, only the net growth rate (μ_0) of ultraphytoplankton could have
471 been estimated, but this size class dominated by Peuk has shown a loss of biomass after the 24
472 h incubation (Table 3, Fig. S2A).

473

474 In April, we were able to calculate the μ_0 only for the nanophytoplankton (Table 3, Fig. S2B).
475 In NB, this size class dominated by *H. minima* displayed a high net growth rate ($\mu_0 = 1.13 \text{ d}^{-1}$)
476 and a relatively low mortality rate (0.28 d^{-1}), while in SB, it was dominated by Chlorophyta
477 and Cryptophyta cells, which displayed the highest estimated μ_0 (1.91 d^{-1}) and a high
478 mortality rate (0.83 d^{-1}).

479

480 In September, the net growth rates in NB ranged between 0.50 d⁻¹ for the ultraphytoplankton
481 dominated by PC-cyanos and 1.84 d⁻¹ for the microphytoplankton dominated by
482 dinoflagellates and Bacillariophyceae. The nano-, dominated by *Plagioselmis* sp., *Tetraselmis*
483 sp., and *P. cordatum*, and the microphytoplankton showed the highest growth rates ($\mu_0 = 1.6$
484 and $\mu_0 = 1.8$ d⁻¹, respectively) with similar mortality rates ($g = 0.57$ d⁻¹). In SB, the total
485 phytoplankton and the ultraphytoplankton also dominated by PC-cyanos experienced a loss of
486 biomass after 24 h incubation (Table 3). The nanophytoplankton dominated by *A. radians* and
487 *Plagioselmis* sp. displayed high growth and mortality rates ($\mu_0 = 1.4$ d⁻¹; $g = 1.1$ d⁻¹). The
488 biomass of microphytoplankton at T0 was too low to estimate their apparent growth rate, we
489 were not able to truly calculate its specific growth and mortality rates.

490

491 Table 3. Net growth rate without enrichment (μ_0) and maximal growth rates, with (i) full
492 enrichment (μ_{\max}), (ii) enrichment minus N (μ_{-N}), (iii) enrichment minus P (μ_{-P}) for the NB
493 and SB stations of Biguglia lagoon in November-December 2013, April and September 2014.
494 g represents the mortality rate. Bold values specify when the relationship between apparent
495 growth rate as a function of the dilution factor can be explained as a linear model, although
496 for the minus N and minus P enrichments other models can also fit this relationship; the
497 $\mu_0:\mu_{\max}$ ratio illustrates the impact of the full enrichment on phytoplankton growth; the ratios
498 between growth rates with selectively N- or P-enrichment (μ_{-N} or μ_{-P}) and μ_{\max} highlight N- or
499 P-limitations ($\mu_{-X}:\mu_{\max} = 1$ no limitation, $\mu_{-X}:\mu_{\max} < 1$ limitation of the lacking nutrient X, μ_{-X}
500 $:\mu_{\max} > 1$ limitation of the added nutrient).

Stations	Samplings	Fraction	μ_0	g	μ_{\max}	$\mu_0:\mu_{\max}$	μ_{-N}	$\mu_{-N}:\mu_{\max}$	μ_{-P}	$\mu_{-P}:\mu_{\max}$
			(d ⁻¹)				(d ⁻¹)		(d ⁻¹)	
NB	Nov - Dec 2013	Total	0.45	0.62	<0	-	<0	-	0.30	<0
		Micro-	<0	1.45	<0	-	<0	-	<0	-
		Nano-	0.37	0.26	0.46	0.79	0.62	1.34	1.13	2.44
		Ultra-	0.96	1.82	<0	-	<0	-	<0	-
	April 2014	Total	<0	<0	0.25	-	0.51	1.99	0.07	0.29
		Micro-	<0	1.01	0.24	-	1.59	6.70	1.01	4.24
		Nano-	1.13	0.28	1.83	0.62	1.95	1.06	1.54	0.84
		Ultra-	<0	0.43	<0	-	<0	-	<0	-
	September 2014	Total	0.64	0.30	1.78	0.36	0.86	0.48	1.03	0.58

	Micro-	1.84	0.57	3.40	0.54	1.78	0.52	2.24	0.66
	Nano-	1.60	0.57	2.70	0.59	1.77	0.65	2.13	0.79
	Ultra-	0.51	0.45	1.12	0.45	0.51	0.46	0.52	0.46
SB Nov - Dec 2013	Total	0.48	0.69	0.41	1.19	0.26	0.64	0.53	1.31
	Micro-	1.12	1.12	0.45	2.48	1.09	2.41	1.05	2.32
	Nano-	1.17	0.50	0.90	1.30	0.78	0.87	1.07	1.19
	Ultra-	<0	1.10	0.14	<0	1.85	13.68	2.02	14.94
April 2014	Total	0.17	<0	0.19	0.92	0.06	0.31	0.26	1.35
	Micro-	-	-	-	-	-	-	-	-
	Nano-	1.91	0.83	2.00	0.95	1.64	0.82	1.64	0.82
	Ultra-	<0	<0	0.01	<0	0.10	8.73	0.34	27.96
September 2014	Total	<0	0.16	1.79	<0	0.28	0.16	1.34	0.75
	Micro-	-	-	-	-	-	-	-	-
	Nano-	1.42	1.14	3.23	0.44	1.58	0.48	<0	<0
	Ultra-	<0	<0	0.87	<0	<0	<0	<0	<0

501

502 3.3. Bioassay experiments

503 In November and December 2013, the full enrichment including NO_3^- , NH_4^+ and PO_4^{3-} did
504 not enhance phytoplankton growth rates in the two stations. The total phytoplankton, the
505 micro- and ultraphytoplankton in NB experienced a loss of biomass after 24 h incubation
506 compared with the controls. Only the growth rate of the ultraphytoplankton of SB was higher
507 with the enrichment, and showed a linear relationship between the apparent growth rate and
508 the dilution factor. This indicates a nutrient limitation for this class size (Table 3). In NB, the
509 enrichment minus N has only resulted in a higher growth rate of the nanophytoplankton. The
510 non-linear relationship between the apparent growth rate and the dilution factor for the
511 ultraphytoplankton (Fig. S2A) suggests the use of other resources than the externally supplied
512 nutrients, or a saturation of predation in the less diluted samples. The enrichment minus P
513 partly released the total phytoplankton growth from N-limitation, allowing a positive growth
514 rate. Ratios between μ_{-N} or μ_{-P} and μ_{max} for the nanophytoplankton indicate an independent
515 co-limitation by N and P. The negative growth rates of the micro- and ultraphytoplankton, and
516 the non-linear relationship between their apparent growth rate and the dilution factor with the
517 enrichments minus N and minus P more likely suggest a co-limitation by N and P (Table 3,

518 Fig. S2A). In SB, ratios between μ_{-N} or μ_{-P} and μ_{max} of the total phytoplankton and of the
519 nanophytoplankton, as well as their non-linear relationship between apparent growth rates and
520 dilution factors suggest a single N-limitation (Table 3, Fig. S2B). The higher growth rate of
521 microphytoplankton with the enrichments minus N and minus P compared with μ_{max} (Table 3)
522 suggests a co-limitation by N and P. Ratios between μ_{-N} or μ_{-P} and μ_{max} for the
523 ultraphytoplankton suggest an independent co-limitation by N and P (Table 3).

524

525 In April 2014, the full enrichment with NO_3^- and PO_4^{3-} resulted in an increase of the growth
526 rate of the nanophytoplankton by 62 % and 5 % in NB and SB respectively. The non-linear
527 relationship between the apparent growth rate and the dilution factor indicates that other
528 fractions in the two stations may have been limited despite the full nutrient enrichment (Table
529 3, Fig. S3). In NB, ratios between μ_{-N} or μ_{-P} and μ_{max} of the total phytoplankton and of the
530 nanophytoplankton suggest a single P-limitation. The microphytoplankton displayed an
531 independent co-limitation of N and P, which was more pronounced for P. The negative μ_{-N}
532 and μ_{-P} of the ultraphytoplankton, and the relationship between their apparent growth rate and
533 the dilution factor in the enrichments minus N and minus P also suggest a co-limitation by N
534 and P (Table 3, Fig. S3A). In SB, ratios between μ_{-N} or μ_{-P} and μ_{max} of the total phytoplankton
535 indicate a single N-limitation, while those of the nano- and of the ultraphytoplankton indicate
536 a simultaneous and an independent co-limitation by N and P respectively (Table 3, Fig. S3).
537 This is also supported by the non-linear relationship between their apparent growth rate and
538 the dilution factor that may highlight the use of other N and P resources to cope with the
539 nutrient limitation (Table 3, Fig. S3B).

540

541 In September 2014, the enrichment with NH_4^+ and PO_4^{3-} led to more than the doubling of the
542 growth rate of most of phytoplankton size classes in NB (Table 3), indicating a strong nutrient

543 limitation. The microphytoplankton of NB displayed the highest μ_{max} of the two communities
544 for the three bioassays. In SB, the full enrichment has allowed a release from nutrient
545 limitation of the entire community, especially for the ultraphytoplankton and the
546 nanophytoplankton (Table 3). In NB, ratios between μ_{-N} or μ_{-P} and μ_{max} of the total
547 phytoplankton and the three size classes suggest a N and P co-limitation, that was serial for
548 the micro- and for the ultraphytoplankton, and independent for the nanophytoplankton (Table
549 3). In SB, ratios between μ_{-N} or μ_{-P} and μ_{max} of the three size classes indicate an independent
550 co-limitation by N and P for the total phytoplankton, a serial co-limitation by N and P for the
551 nanophytoplankton and a simultaneous co-limitation by N and P for the ultraphytoplankton
552 (Table 3, Fig. S4B).

553

554 4. Discussion

555 In this study, we aimed to describe functional traits of phytoplankton communities in Biguglia
556 lagoon submitted to a strong anthropogenic impact in different seasons with contrasting
557 environmental conditions, and the response of these communities to various forms of
558 nutrients inputs simulating nutrient pulses to predict their impact on ecosystem functioning.

559

560 4.1. *Functional traits of phytoplankton communities in response to environmental* 561 *conditions*

562 Biguglia lagoon was globally strongly influenced by freshwater inputs from the watershed
563 and from the connection with one of the main river in Corsica in the south of the lagoon.
564 These freshwater inputs have contributed to maintain a low salinity that prevailed in the two
565 sub-basins during the year 2014, despite their distinct hydrological dynamics. This freshwater
566 influence has led to the dominance of species associated with brackish or fresh waters in
567 phytoplankton assemblages. Phycocyanin-rich picocyanobacteria (PC-cyanos) and flagellates

568 from the Dictyochophyceae, dinoflagellates, Chlorophytes, and Cryptophytes were the main
569 representative groups in the two stations at the three samplings. Their size structure,
570 community composition and productivity illustrate their functional responses to lagoon
571 variability.

572

573 Biguglia lagoon was principally enriched by high nitrogen inputs, especially during the wet
574 periods. Indeed, high nitrate inputs from a huge historical accumulation are carried to the
575 lagoon by groundwater flows during strong episodic rainfall events (Erostate et al., 2018).
576 The negative correlations, both between temperature and nutrient concentrations, and between
577 salinity and ammonium concentration also support that freshwater discharge was the main
578 source for nutrient enrichment in the lagoon at the three sampling periods (Garrido et al.,
579 2016; Pasqualini et al., 2017). The important precipitations in November 2013 and Marsh
580 2014 (150.4 mm and 92.6 mm respectively, <https://www.infoclimat.fr/>, Station of Bastia
581 Poretta) explain the high NO_3^- concentrations and high TN:TP and DIN:DIP values in the
582 autumn and spring samplings. In contrast, low dissolved inorganic nitrogen and phosphorus
583 concentrations were observed in September, although the summer period corresponds to the
584 most important period for the recycling of ammonium and phosphate in the lagoon (Collos et
585 al., 2003).

586

587 Phytoplankton cell size decreased with the decrease of DIN availability, which may be
588 explained by higher uptake affinities of small cells under nutrient-limiting conditions
589 (Litchman et al., 2007). High nanophytoplankton biomasses in Biguglia lagoon have already
590 been related to the availability of oxidized forms of nitrogen (Cecchi et al., 2016). In this
591 study, ultra- and nanophytoplankton biomasses were rather positively related to PO_4^{3-}
592 availability. Dominance of fast-growing pico- and nanoflagellates can be frequent in shallow

593 lagoons exposed to high nutrient loads and submitted to continuous inputs of freshwater or
594 discontinuous floods. These nutrient loads allow the release of the limitation of cells $>3 \mu\text{m}$
595 that can coexist with fast growing and highly competitive picoeukaryotes (Bec et al., 2011;
596 Coelho et al., 2015).

597

598 Nanoplanktonic dinoflagellates were particularly abundant during the samplings in the wet
599 seasons (autumn and spring). We particularly observed blooms ($>10^5 \text{ cells.l}^{-1}$) of *P. cordatum*
600 in NB in November 2013, and of *H. minima* in SB in December 2013 and in NB in April
601 2014. Blooms of *P. cordatum* are frequently related to high N:P ratios in coastal transitional
602 ecosystems characterized by brackish waters (Johnson, 2015), and to freshwater inflows that
603 cause low salinities and high nitrate concentrations (Heil et al., 2005; Coelho et al., 2007;
604 Cecchi et al., 2016). Moreover, *P. cordatum* and some *Heterocaspa* species have
605 demonstrated potential mixotrophic abilities, especially under N- or P-starvation (Legrand et
606 al., 1998; Jeong et al., 2005; Johnson, 2015). The presence of potential preys such as
607 picocyanobacteria, Cryptophytes and small diatoms may thus have stimulated their
608 development despite the potential nutrient limitation.

609

610 Biomass of the microphytoplankton, mainly composed of Bacillariophyceae and
611 dinoflagellates in the three samplings, has been positively related to the high NO_3^-
612 concentrations. Indeed, pulses of NO_3^- especially favor fast-growing diatoms during spring
613 blooms in river-dominated coastal ecosystems (Paerl et al., 2010; Donald et al., 2011; Glibert
614 et al., 2015). They were generally most abundant in the northern basin, probably due to their
615 resistance to turbulent environments, their high affinity for nitrate, and the high freshwater
616 inputs (Garrido et al., 2016).

617

618 Picophytoplankton (picocyanobacteria and picoeukaryotes) often represents an important
619 component of phytoplankton communities in brackish lagoons. Particularly, phycoerythrin-rich
620 picocyanobacteria have shown blooms after floods when salinities achieved their minimal
621 values (Bec et al., 2011; Lafabrie et al., 2013). Small cells have physiological advantages that
622 confer them a high competitiveness to acquire and use nutrients under limiting conditions
623 (Raven, 1998). They are especially competitive in warmer waters and under low N-
624 availability as observed during the summer period (Bec et al., 2005; Leruste et al., 2016;
625 Domingues et al., 2017). Moreover, they are able to use nitrate, ammonium as well as organic
626 N-sources in a highly efficient way (Domingues et al., 2011), and are hardly inhibited by
627 occasionally occurring high ammonium concentrations (Collos and Harrison, 2014). Bloom of
628 picophytoplankton in summer coincides with high regeneration process, and a dominance of
629 ammonium that picophytoplankton can preferentially use, while diatoms and dinoflagellates
630 preferentially use NO_3^- , and are able to show high growth rate in nitrate-rich waters
631 (Domingues et al., 2011; Glibert et al., 2015; Reed et al., 2016).

632

633 These competitive trade-offs may thus explain the competitive exclusion of the larger cells by
634 the PC-cyanos in September. Shift between diatoms and dinoflagellates to picophytoplankton
635 from spring to late summer frequently occur in coastal aquatic systems with the shift of
636 available N forms, from NO_3^- (>88% of the DIN in the two stations in November-December
637 2013 and April 2014) to NH_4^+ (>85 % of the DIN in the two stations in September 2014)
638 (Glibert et al., 2015; Domingues et al., 2017). However, ambient nutrient concentrations are
639 measures of the residual nutrients after biogeochemical activity, and may not reflect the real
640 available nutrient concentrations for phytoplankton growth. The bioassay experiments gave
641 further insights about the relationship between their functional responses to the nutrient

642 availability. This especially specified which nutrient was potentially limiting for
643 phytoplankton growth at the three sampling periods.

644

645 4.2. *Phytoplankton behavior under induced nutrient limitation*

646 Temperate coastal ecosystems generally display seasonal changes of nutrient availability and
647 limitation, characterized by shifts from a P-limitation during the wet seasons after high
648 nitrate-rich freshwater inputs, to N-limitation during the dry season (Lomas and Glibert, 2000;
649 Kemp et al., 2005). In contrast, we did not observe this pattern in Biguglia lagoon, despite the
650 high DIN inputs from the watershed during the wet seasons, reflected by high TN:TP and
651 DIN:DIP⁻ values. The results of the bioassay experiments more often show a co-limitation by
652 N and P for the phytoplankton communities. Nevertheless, some size fractions were limited
653 by one of these elements alone, *e.g.* the nanophytoplankton of the NB station limited by P in
654 April 2014, and the microphytoplankton of the SB station limited by N in December 2013. A
655 review showed that co-limitation by N and P in freshwater and marine ecosystems is more
656 common than previously thought (Elser et al., 2007). Interestingly, the single limitation for
657 some size fractions within a co-limited community, such as observed in SB in September
658 2014 may indicate niche differentiation. We also observed that the phytoplankton in SB was
659 more often N-limited than in NB despite high DIN:DIP ratios. Conclusions from the dilution
660 experiment should be drawn with caution, because incubations have been conducted on small
661 spatial and temporal scales. The responses of phytoplankton communities at these scales can
662 differ from the overall ecology of the entire ecosystem (Piehler et al., 2004; Reed et al.,
663 2016). However, these results support that ambient nutrient ratios do not necessarily reflect
664 the availability of nutrients for autotrophic communities, particularly because of the known
665 variability of nutrient stoichiometry of phytoplankton. The communities of the two stations
666 showed the strongest simultaneous or serial co-limitation by N and P in summer while in the


667 wet seasons they showed single N- or P-limitations or independent co-limitation depending on
668 fractions. The figure 5 conceptualizes our observations for the three samplings, and links the
669 environmental specificities, the differences in community structure and composition, with the
670 calculated growth rates and the observed N- and P-limitation.

671

672 Picocyanobacteria, dinoflagellates and other phytoplankton groups have demonstrated
673 numerous ways to acclimate to temporally and varying N- and P-limitation (Lin et al., 2015).
674 For example, luxury PO_4^{3-} uptake prior to sampling for the bioassays could have provided the
675 necessary intracellular P for phytoplankton growth during the 24 h incubation.
676 Picocyanobacteria may have used high P-storage abilities that could have allowed them to
677 maintain their bloom despite low P-availability (Sorokin and Dallochio, 2008). Moreover,
678 they have shown high inorganic P-uptake efficiency at low levels of available soluble reactive
679 phosphorus (Collos et al., 2009). Alternatively, the high dissolved and particulate organic N
680 and P stocks in the sediment of Biguglia lagoon may have been used by the phytoplankton to
681 thrive despite low dissolved inorganic N- and P-availability. Indeed, the very low dissolved
682 inorganic phosphorus concentrations at the three samplings may have stimulated the
683 production of alkaline phosphatase, allowing the use of dissolved organic phosphorus (Piehler
684 et al., 2004; Lin et al., 2015). Moreover, high abundances of potentially mixotrophic species
685 among *Prorocentrum*, *Heterocapsa*, *Gonyaulax* and other genera, and of *Mesodinium rubrum*
686 (Collos et al., 2014; Fischer et al., 2016; Johnson, 2015) have been observed during these
687 experiments. Mixotrophy is an advantageous behavioral response under nutrient limiting
688 conditions, allowing cells to reduce direct competition for inorganic nutrients (Jeong et al.,
689 2005; Mitra et al., 2014; Johnson, 2015). Hence, the high abundances of picocyanobacteria,
690 physiologically able to cope with P-limitation, and of potentially mixotrophic dinoflagellates
691 and nanoflagellates, could explain the N and P co-limitation of the communities despite high

692 DIN:DIP ratios (Burson et al., 2016). Indeed, these physiological and behavioral capacities
 693 may have offset an inorganic P-limitation, either by the ingestion of starving preys that may
 694 benefit from the added inorganic nutrients, or of dissolved organic substances. This may have
 695 led to a N and P co-limitation rather than a single P-limitation (Burson et al., 2016; Fischer et
 696 al., 2016). Mixotrophy is also stimulated by nutrient limitation, and would have been
 697 particularly competitive in Biguglia lagoon during the three sampling periods. This may
 698 explain the bloom of the mixotrophic *Mesodinium rubrum* (Tong et al., 2015; Seong et al.,
 699 2017) in the southern basin in the autumn and spring samplings (Fig. 5).

700



701

702 Fig. 5. Schematic representation of the environmental characteristics, phytoplankton
 703 community composition and their relationship with nutrient availability during the three
 704 bioassays periods in the two stations of Biguglia lagoon. For each sampling, the schematic
 705 representation specifies the water temperature, the salinity, the freshwater influence, and the

706 nutrients inputs whether they come from the watershed or from the sediments. Main
707 phytoplankters are specified as pictograms. Phytoplankton co-limitation by N and P is
708 inferred according to ratios between growth rate under experimentally induced N- or P-
709 limitation and maximal growth rate (see Results).

710

711 *4.3. Assumptions about responses of phytoplankton communities to a nutrient pulse*

712 Bioassay experiments can provide valuable insights about relationship between nutrient
713 dynamics and phytoplankton physiological responses. This can simulate a nutrient pulse,
714 helping to predict algal-based expression of eutrophication (Xu et al., 2012).

715

716 The composition of nutrient enrichments used at the three sampling periods was based on
717 previous studies and assumptions about ecosystem functioning, and may have not always
718 reflected the real environmental conditions or the potential nutrient pulses threatening
719 Biguglia lagoon due to its high variability. These enrichments may thus have not always been
720 the most suitable to involve an optimum response of the phytoplankton communities. Hence,
721 the addition of the full enrichment has not always allowed the estimation of maximal growth
722 and mortality rates probably because the underlying assumptions of this method were not
723 always fulfilled. For almost the half of the fractions (11 out of 24) the relationship between
724 the apparent growth rates and the dilution factor was non-linear. Non-linear trends may
725 indicate microzooplankton feeding threshold or saturation depending on the curve pattern
726 (Chen et al., 2014). Indeed, the shape of some of these plots (Fig. S1, red lines) may indicate a
727 strong nutrient limitation and suggests the use of other nutrient resources such as those
728 regenerated through cell lyses, grazing and mineralization that increase with the proportion of
729 undiluted sample (Andersen et al., 1991).

730

731 The responses of the phytoplankton communities to the nutrient enrichment in the three
732 bioassays have reflected contrasting environmental vulnerability of Biguglia lagoon to short-
733 term nutrient enrichment. Nutrient limitation can have many indirect effects on phytoplankton
734 productivity, *e.g.* a modification of the internal cell stoichiometry that changes its nutritional
735 quality and modify predation rates (Harpole et al., 2011). In November-December 2013, the
736 addition of the full enrichment comprising a mixture of nitrate and ammonium (10 μ M each)
737 seems to have had a negative impact. In certain cases, ammonium can negatively affect nitrate
738 uptake and assimilation, especially in nitrate-dominated systems (Domingues et al., 2011;
739 2017), and particularly for dinoflagellates, the least ammonium-tolerant group (Collos and
740 Harrison, 2014). Hence, the addition of ammonium when nitrogen was mainly available as
741 nitrate, and dinoflagellates were the most abundant, may have inhibited its uptake and led to a
742 decrease of growth rate (Dodds and Whiles, 2010; Glibert et al., 2015). As nutrient addition is
743 expected to increase limitation by other resources, a negative response may also reflect
744 strongly constrained N:P stoichiometry. This is suggested by high TN:TP and DIN:DIP
745 values. Rather than re-equilibrating these ratios, nutrient addition may have led to an
746 exacerbated internal stoichiometric imbalance of N and P nutrients with negative impact on
747 phytoplankton growth (Harpole et al., 2011).

748

749 In general, the phytoplankton maximal growth rates followed a predictable seasonal trend,
750 with minima in autumn and maxima in summer. These changes are correlated with water
751 temperatures and photoperiods length, providing more favorable growth conditions in
752 summer. Lower temperatures and light availability in autumn may have altered the uptake rate
753 of phytoplankton (Dodds and Whiles, 2010; Domingues et al., 2011; Reed et al., 2016). This
754 is also consistent with the community composition, the picophytoplankton dominating the
755 community in September potentially having higher growth rate than the nanophytoplankton

756 dominating the autumn and spring samplings (Bec et al., 2011). Phytoplankton was thus more
757 reactive to a nutrient addition during the summer season when the nutrient demand is the
758 highest. This suggests that Biguglia lagoon is the most vulnerable to nutrient inputs during
759 summer.

760

761 Conclusion

762 Experimental enrichments bioassays are particularly appropriate to explore cause-effect
763 relationships between nutrient availability and phytoplankton dynamics, which can reflect the
764 vulnerability of the whole ecosystem. This is an important issue for managers to draw up
765 appropriate nutrient loading budgets, and to help evaluating the efficiency of subsequent
766 nutrient reduction strategies. In our study, N inputs, particularly NO_3^- during the wet periods
767 and NH_4^+ in summer, stimulated phytoplankton growth in Biguglia lagoon emphasizing the
768 need to mitigate these inputs. The phytoplankton community was strongly influenced by
769 freshwater and nutrients inputs that have led to different physiological and behavioral
770 responses over seasons. During autumn and spring, high DIN:DIP ratios have been associated
771 with blooms of potentially harmful dinoflagellates, which are increasingly observed in
772 Biguglia lagoon.

773

774 The absence of a significant positive effect of the enrichment could be explained through
775 several hypotheses. The first is the presence of sufficient amounts of dissolved inorganic
776 nutrients in the water column to sustain the phytoplankton growth during the incubation time.
777 The second could be a more intensive use of other resources than the external pool, such as
778 internal or regenerated inorganic nutrients. The third could be a strong competition for
779 available nutrient with the bacterial communities. Of increasing concern is thus the potential
780 for organic nutrients and high abundances of small preys to promote mixotrophic species,

781 some of which are potentially harmful. It would be important to quantify the availability of
782 dissolved organic nutrients and its effect on phytoplankton growth and trophic modes. This
783 should be linked with a study about the interaction between phytoplankton and bacteria, such
784 as competition for organic and inorganic available nutrients. The impact of bacterial activity
785 on the available nutrients stocks has also to be explored. Indeed, bacterial communities can
786 play an essential role in nutrient cycling, *e.g.* through the remobilization of organic matter,
787 leading to a tight phytoplankton-bacterioplankton coupling (Caroppo, 2002). The possible use
788 of external, internal and regenerated N and P pools, as well as the mixotrophic abilities of the
789 phytoplankton communities also need to be studied.

790

791 5. Acknowledgements

792 Amandine Leruste was supported in 2018 by a Postdoctoral fellowship of the University of
793 Corsica Pasquale Paoli (SPE CNRS 6134 Laboratory and UMS Stella Mare) and the
794 Collectivity of Corsica, and from 2013-2016 by a PhD fellowship of the Doctoral School at
795 the University of Montpellier. This work has been financed by Biguglia lagoon nature
796 reserve, the Collectivity of Corsica and the University of Corsica Pasquale Paoli, that authors
797 are grateful for their cooperation. We thank the Ifremer station of Bastia and the UMS Stella
798 Mare for technical support and help during the experiments. This work was (co)funded by the
799 Labex DRIIHM, French programme "Investissements d'Avenir" (ANR-11-LABX-0010)
800 which is managed by the ANR. Thanks are extended to everyone who participated in the
801 experiments. The authors thank the reviewers for their helpful and constructive comments.
802 This research did not receive any specific grant from funding agencies in the public,
803 commercial, or not-for-profit sectors.

804

805 All authors have approved the final article.

806

807 6. References

- 808 Aminot, A., Chaussepied, M.C., 1983. Manuel des analyses chimiques en milieu marin.
809 CNEXO, Brest.
- 810 Andersen, K.H., Aksnes, D.L., Berge, T., Fiksen, O., Visser, A., 2015. Modelling emergent
811 trophic strategies in plankton. *Journal of Plankton Research* 37, 862–868.
812 <https://doi.org/10.1093/plankt/fbv054>
- 813 Andersen, T., Schartau, A., Paasche, E., 1991. Quantifying External and Internal Nitrogen and
814 Phosphorus Pools, as Well. *Mar. Ecol.-Prog. Ser.* 69, 67–80.
815 <https://doi.org/10.3354/meps069067>
- 816 Anderson, M.J., 2001. A new method for non-parametric multivariate analysis of variance.
817 *Austral Ecology* 26, 32–46. <https://doi.org/10.1111/j.1442-9993.2001.01070.pp.x>
- 818 Barbier, E.B., Hacker, S.D., Kennedy, C., Koch, E.W., Stier, A.C., Silliman, B.R., 2011. The
819 value of estuarine and coastal ecosystem services. *Ecol. Monogr.* 81, 169–193.
820 <https://doi.org/10.1890/10-1510.1>
- 821 Bartón, K., 2013. MuMIn: Multi-modal inference. Model selection and model averaging
822 based on information criteria (AICc and alike). [http://cran.r-](http://cran.r-project.org/web/packages/MuMIn/index.html)
823 [project.org/web/packages/MuMIn/index.html](http://cran.r-project.org/web/packages/MuMIn/index.html).
- 824 Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models
825 Using lme4. *Journal of Statistical Software* 67. <https://doi.org/10.18637/jss.v067.i01>

- 826 Bec, B., Collos, Y., Souchu, P., Vaquer, A., Lautier, J., Fiandrino, A., Benau, L., Orsoni, V.,
827 Laugier, T., 2011. Distribution of picophytoplankton and nanophytoplankton along an
828 anthropogenic eutrophication gradient in French Mediterranean coastal lagoons.
829 *Aquat. Microb. Ecol.* 63, 29–45. <https://doi.org/10.3354/ame01480>
- 830 Bec, B., Hussein-Ratrema, J., Collos, Y., Souchu, P., Vaquer, A., 2005. Phytoplankton
831 seasonal dynamics in a Mediterranean coastal lagoon: emphasis on the picoeukaryote
832 community. *J. Plankton Res.* 27, 881–894. <https://doi.org/10.1093/plankt/fbi061>
- 833 Bellinger, E.G., Sigeo, D.C. (Eds.), 2015. *Freshwater Algae: Identification, Enumeration and*
834 *Use as Bioindicators.* John Wiley & Sons, Inc., Hoboken, NJ, USA.
835 <https://doi.org/10.1002/9781118917152>
- 836 Bérard-Therriault, L., Poulin, M., Bossé, L., 1999. *Guide d'identification du phytoplancton*
837 *marin de l'estuaire et du golfe du Saint-Laurent incluant également certains*
838 *protozoaires* Canadian Special Publication of Fisheries and Aquatic Sciences No. 128.
839 NRC Research Press.
- 840 Burnham, K.P., Anderson, D.R. (Eds.), 2004. *Model Selection and Multimodel Inference.*
841 Springer New York, New York, NY.
- 842 Burson, A., Stomp, M., Akil, L., Brussaard, C.P.D., Huisman, J., 2016. Unbalanced reduction
843 of nutrient loads has created an offshore gradient from phosphorus to nitrogen
844 limitation in the North Sea. *Limnology and Oceanography* 61, 869–888.
845 <https://doi.org/10.1002/lno.10257>
- 846 Caroppo, C., 2002. Variability and interactions of phytoplankton and bacterioplankton in
847 Varano lagoon (Adriatic Sea). *J Plankton Res* 24, 267–273.
848 <https://doi.org/10.1093/plankt/24.3.267>
- 849 Cecchi, P., Garrido, M., Collos, Y., Pasqualini, V., 2016. Water flux management and
850 phytoplankton communities in a Mediterranean coastal lagoon. Part II: Mixotrophy of
851 dinoflagellates as an adaptive strategy? *Mar. Pollut. Bull.*
852 <https://doi.org/10.1016/j.marpolbul.2016.04.041>
- 853 Chen, B., Laws, E.A., Liu, H., Huang, B., 2014. Estimating microzooplankton grazing
854 half-saturation constants from dilution experiments with nonlinear feeding kinetics.
855 <https://doi.org/10.4319/lo.2014.59.3.0639> 59, 639–644.
- 856 Cloern, J.E., 2001. Our evolving conceptual model of the coastal eutrophication problem.
857 *Mar. Ecol.-Prog. Ser.* 210, 223–253. <https://doi.org/10.3354/meps210223>
- 858 Coelho, S., Gamito, S., Perez-Ruzafa, A., 2007. Trophic state of Foz de Almagem coastal
859 lagoon (Algarve, South Portugal) based on the water quality and the phytoplankton
860 community. *Estuar. Coast. Shelf Sci.* 71, 218–231.
861 <https://doi.org/10.1016/j.ecss.2006.07.017>
- 862 Coelho, S., Perez-Ruzafa, A., Gamito, S., 2015. Phytoplankton community dynamics in an
863 intermittently open hypereutrophic coastal lagoon in Southern Portugal. *Estuarine*
864 *Coastal and Shelf Science* 167, 102–112. <https://doi.org/10.1016/j.ecss.2015.07.022>

- 865 Collos, Y., Vaquer, A., Bibent, B., Souchu, P., Slawyk, G., Garcia, N., 2003. Response of
866 coastal phytoplankton to ammonium and nitrate pulses: seasonal variations of nitrogen
867 uptake and regeneration. *Aquat. Ecol.* 37, 227–236.
868 <https://doi.org/10.1023/A:1025881323812>
- 869 Collos, Y., Hussein-Ratrema, J., Bec, B., Vaquer, A., Hoai, T.L., Rougier, C., Pons, V.,
870 Souchu, P., 2005. Pheopigment dynamics, zooplankton grazing rates and the autumnal
871 ammonium peak in a Mediterranean lagoon. *Hydrobiologia* 550, 83–93.
872 <https://doi.org/10.1007/s10750-005-4365-1>
- 873 Collos, Y., Bec, B., Jauzein, C., Abadie, E., Laugier, T., Lautier, J., Pastoureaud, A., Souchu,
874 P., Vaquer, A., 2009. Oligotrophication and emergence of picocyanobacteria and a
875 toxic dinoflagellate in Thau lagoon, southern France. *J. Sea Res.* 61, 68–75.
876 <https://doi.org/10.1016/j.seares.2008.05.008>
- 877 Collos, Y., Harrison, P.J., 2014. Acclimation and toxicity of high ammonium concentrations
878 to unicellular algae. *Mar. Pollut. Bull.* 80, 8–23.
879 <https://doi.org/10.1016/j.marpolbul.2014.01.006>
- 880 Collos, Y., Jauzein, C., Ratmaya, W., Souchu, P., Abadie, E., Vaguer, A., 2014. Comparing
881 diatom and *Alexandrium catenella/tamarensis* blooms in Thau lagoon: Importance of
882 dissolved organic nitrogen in seasonally N-limited systems. *Harmful Algae* 37, 84–91.
883 <https://doi.org/10.1016/j.hal.2014.05.008>
- 884 Dodds, W., Whiles, M., 2010. Chapter 16: Nutrient Use and Remineralization, in: *Freshwater
885 Ecology - Concepts and Environmental Applications*. pp. 437–467.
- 886 Domingues, R.B., Barbosa, A.B., Sommer, U., Galvao, H.M., 2011. Ammonium, nitrate and
887 phytoplankton interactions in a freshwater tidal estuarine zone: potential effects of
888 cultural eutrophication. *Aquat. Sci.* 73, 331–343. <https://doi.org/10.1007/s00027-011-0180-0>
889
- 890 Domingues, R.B., Guerra, C.C., Barbosa, A.B., Galvão, H.M., 2017. Will nutrient and light
891 limitation prevent eutrophication in an anthropogenically-impacted coastal lagoon?
892 *Continental Shelf Research* 141, 11–25.
- 893 Donald, D.B., Bogard, M.J., Finlay, K., Bunting, L., Leavitt, P.R., 2013. Phytoplankton-
894 Specific Response to Enrichment of Phosphorus-Rich Surface Waters with
895 Ammonium, Nitrate, and Urea. *PLoS One* 8.
896 <https://doi.org/10.1371/journal.pone.0053277>
- 897 Donald, D.B., Bogard, M.J., Finlay, K., Leavitt, P.R., 2011. Comparative effects of urea,
898 ammonium, and nitrate on phytoplankton abundance, community composition, and
899 toxicity in hypereutrophic freshwaters. *Limnol. Oceanogr.* 56, 2161–2175.
900 <https://doi.org/10.4319/lo.2011.56.6.2161>
- 901 Duarte, C.M., Agust, S., Gasol, J.M., Vaqu, D., VazquezDominguez, E., 2000. Effect of
902 nutrient supply on the biomass structure of planktonic communities: an experimental
903 test on a Mediterranean coastal community. *Mar Ecol Prog Ser* 206, 87–95.
904 <https://doi.org/10.3354/meps206087>

- 905 Elser, J.J., Bracken, M.E.S., Cleland, E.E., Gruner, D.S., Harpole, W.S., Hillebrand, H., Ngai,
906 J.T., Seabloom, E.W., Shurin, J.B., Smith, J.E., 2007. Global analysis of nitrogen and
907 phosphorus limitation of primary producers in freshwater, marine and terrestrial
908 ecosystems. *Ecology Letters* 10, 1135–1142. [https://doi.org/10.1111/j.1461-](https://doi.org/10.1111/j.1461-0248.2007.01113.x)
909 [0248.2007.01113.x](https://doi.org/10.1111/j.1461-0248.2007.01113.x)
- 910 Erostate, M., Huneau, F., Garel, E., Lehmann, M.F., Kuhn, T., Aquilina, L., Vergnaud-
911 Ayraud, V., Labasque, T., Santoni, S., Robert, S., Provitolo, D., Pasqualini, V., 2018.
912 Delayed nitrate dispersion within a coastal aquifer provides constraints on land-use
913 evolution and nitrate contamination in the past. *Science of The Total Environment*
914 644, 928–940. <https://doi.org/10.1016/j.scitotenv.2018.06.375>
- 915 Fischer, R., Giebel, H., Ptacnik, R., 2016. The identity of the limiting nutrient (N vs. P)
916 affects the competitive success of mixotrophs. *Marine Ecology Progress Series*.
917 <https://doi.org/10.3354/meps11968>
- 918 Franco, A., Fiorin, R., Zucchetta, M., Torricelli, P., Franzoi, P., 2010. Flounder growth and
919 production as indicators of the nursery value of marsh habitats in a Mediterranean
920 lagoon. *J. Sea Res.* 64, 457–464. <https://doi.org/10.1016/j.seares.2010.01.006>
- 921 Garrido, M., Cecchi, P., Collos, Y., Agostini, S., Pasqualini, V., 2016. Water flux
922 management and phytoplankton communities in a Mediterranean coastal lagoon. Part
923 I: How to promote dinoflagellate dominance? *Mar. Pollut. Bull.* 104, 139–152.
924 <https://doi.org/10.1016/j.marpolbul.2016.01.049>
- 925 Glibert, P.M., Wilkerson, F.P., Dugdale, R.C., Raven, J.A., Dupont, C.L., Leavitt, P.R.,
926 Parker, A.E., Burkholder, J.M., Kana, T.M., 2015. Pluses and minuses of ammonium
927 and nitrate uptake and assimilation by phytoplankton and implications for productivity
928 and community composition, with emphasis on nitrogen-enriched conditions. *Limnol.*
929 *Oceanogr.* n/a-n/a. <https://doi.org/10.1002/lno.10203>
- 930 Guillard, R.R.L., Ryther, J.H., 1962. Studies of Marine Planktonic Diatoms: I. *Cyclotella*
931 *Nana* Hustedt, and *Detonula Confervacea* (cleve) Gran. *Can. J. Microbiol.* 8, 229–239.
932 <https://doi.org/10.1139/m62-029>
- 933 Harpole, W.S., Ngai, J.T., Cleland, E.E., Seabloom, E.W., Borer, E.T., Bracken, M.E.S.,
934 Elser, J.J., Gruner, D.S., Hillebrand, H., Shurin, J.B., Smith, J.E., 2011. Nutrient co-
935 limitation of primary producer communities. *Ecology Letters* 14, 852–862.
936 <https://doi.org/10.1111/j.1461-0248.2011.01651.x>
- 937 Heil, C.A., Glibert, P.M., Fan, C.L., 2005. *Prorocentrum minimum* (Pavillard) Schiller - A
938 review of a harmful algal bloom species of growing worldwide importance. *Harmful*
939 *Algae* 4, 449–470. <https://doi.org/10.1016/j.hal.2004.08.003>
- 940 Jeong, H.J., Park, J.Y., Nho, J.H., Park, M.O., Ha, J.H., Seong, K.A., Jeng, C., Seong, C.N.,
941 Lee, K.Y., Yih, W.H., 2005. Feeding by red-tide dinoflagellates on the
942 cyanobacterium *Synechococcus*. *Aquat. Microb. Ecol.* 41, 131–143.
943 <https://doi.org/10.3354/ame041131>
- 944 Johnson, M.D., 2015. Inducible Mixotrophy in the Dinoflagellate *Prorocentrum minimum*.
945 *Journal of Eukaryotic Microbiology* 62, 431–443.

- 946 Kemp, W.M., Boynton, W.R., Adolf, J.E., Boesch, D.F., Boicourt, W.C., Brush, G.,
 947 Cornwell, J.C., Fisher, T.R., Glibert, P.M., Hagy, J.D., Harding, L.W., Houde, E.D.,
 948 Kimmel, D.G., Miller, W.D., Newell, R.I.E., Roman, M.R., Smith, E.M., Stevenson,
 949 J.C., 2005. Eutrophication of Chesapeake Bay: historical trends and ecological
 950 interactions. *Marine ecology. Progress series* 303, 1–29.
- 951 Kjerfve, B., 1994. *Coastal Lagoon Processes*. Elsevier.
- 952 Lafabrie, C., Garrido, M., Leboulanger, C., Cecchi, P., Grégori, G., Pasqualini, V., Pringault,
 953 O., 2013. Impact of contaminated-sediment resuspension on phytoplankton in the
 954 Biguglia lagoon (Corsica, Mediterranean Sea). *Estuarine, Coastal and Shelf Science*,
 955 Pressures, Stresses, Shocks and Trends in Estuarine Ecosystems 130, 70–80.
 956 <https://doi.org/10.1016/j.ecss.2013.06.025>
- 957 Landry, M., Hassett, R., 1982. Estimating the Grazing Impact of Marine Micro-Zooplankton.
 958 *Mar. Biol.* 67, 283–288. <https://doi.org/10.1007/BF00397668>
- 959 Landry, M.R., Brown, S.L., Campbell, L., Constantinou, J., Liu, H., 1998. Spatial patterns in
 960 phytoplankton growth and microzooplankton grazing in the Arabian Sea during
 961 monsoon forcing. *Deep-Sea Research Part II-Topical Studies in Oceanography* 45,
 962 2353–2368. [https://doi.org/10.1016/s0967-0645\(98\)00074-5](https://doi.org/10.1016/s0967-0645(98)00074-5)
- 963 Le Fur, I., De Wit, R., Plus, M., Oheix, J., Simier, M., Ouisse, V., 2018. Submerged benthic
 964 macrophytes in Mediterranean lagoons: distribution patterns in relation to water
 965 chemistry and depth. *Hydrobiologia* 175–200. <https://doi.org/10.1007/s10750-017-3421-y>
- 967 Legrand, C., Graneli, E., Carlsson, P., 1998. Induced phagotrophy in the photosynthetic
 968 dinoflagellate *Heterocapsa triquetra*. *Aquat. Microb. Ecol.* 15, 65–75.
 969 <https://doi.org/10.3354/ame015065>
- 970 Leruste, A., Malet, N., Munaron, D., Derolez, V., Hatey, E., Collos, Y., De Wit, R., Bec, B.,
 971 2016. First steps of ecological restoration in Mediterranean lagoons: Shifts in
 972 phytoplankton communities. *Estuarine, Coastal and Shelf Science* 180, 190–203.
 973 <https://doi.org/10.1016/j.ecss.2016.06.029>
- 974 Leruste, A., Villéger, S., Malet, N., Wit, R.D., Bec, B., 2018. Complementarity of the
 975 multidimensional functional and the taxonomic approaches to study phytoplankton
 976 communities in three Mediterranean coastal lagoons of different trophic status.
 977 *Hydrobiologia* 815. <https://doi.org/10.1007/s10750-018-3565-4>
- 978 Lin, S., Litaker, R.W., Sunda, W.G., 2015. Phosphorus physiological ecology and molecular
 979 mechanisms in marine phytoplankton. *Journal of Phycology* 52, 10–36.
- 980 Litchman, E., Klausmeier, C.A., Schofield, O.M., Falkowski, P.G., 2007. The role of
 981 functional traits and trade-offs in structuring phytoplankton communities: scaling from
 982 cellular to ecosystem level. *Ecol. Lett.* 10, 1170–1181. <https://doi.org/10.1111/j.1461-0248.2007.01117.x>
- 984 Livingston, R.J., 2000. *Eutrophication Processes in Coastal Systems: Origin and Succession
 985 of Plankton Blooms and Effects on Secondary Production in Gulf Coast Estuaries*.
 986 CRC Press.

- 987 Loir, M., 2004. Guide des diatomées: Plus de 200 micro-algues silicieuses photographiées.
988 Delachaux & Niestlé, Paris.
- 989 Lomas, M.W., Glibert, P.M., 2000. Comparisons of nitrate uptake, storage, and reduction in
990 marine diatoms and flagellates. *J. Phycol.* 36, 903–913. <https://doi.org/10.1046/j.1529-8817.2000.99029.x>
991
- 992 Maurice, C.F., Bouvier, C., de Wit, R., Bouvier, T., 2013. Linking the lytic and lysogenic
993 bacteriophage cycles to environmental conditions, host physiology and their
994 variability in coastal lagoons. *Environ. Microbiol.* 15, 2463–2475.
995 <https://doi.org/10.1111/1462-2920.12120>
- 996 Mitra, A., Flynn, K.J., Berge, T., Calbet, A., Raven, J.A., Graneli, E., Glibert, P.M., Hansen,
997 P., Stoecker, D.K., Thingstad, T.F., Tillmann, U., Våge, S., Wilken, S., Zubkov, M.V.,
998 2014. The role of mixotrophic protists in the biological carbon pump. *Biogeosciences*
999 11, 995–1005.
- 1000 Mouillot, D., Titeux, A., Migon, C., Sandroni, V., Frodello, J.-P., Viale, D., 2000.
1001 Anthropogenic influences on a mediterranean Nature Reserve: modelling and
1002 forecasting. *Environmental Modeling & Assessment* 5, 185–192.
1003 <https://doi.org/10.1023/A:1011533811237>
- 1004 Neveux, J., Lantoiné, F., 1993. Spectrofluorometric assay of chlorophylls and phaeopigments
1005 using the least squares approximation technique. *Deep Sea Research Part I:*
1006 *Oceanographic Research Papers* 40, 1747–1765. [https://doi.org/10.1016/0967-0637\(93\)90030-7](https://doi.org/10.1016/0967-0637(93)90030-7)
1007
- 1008 Nixon, S.W., 2009. Eutrophication and the macroscope. *Hydrobiologia* 629, 5–19.
1009 <https://doi.org/10.1007/s10750-009-9759-z>
- 1010 Oksanen, J., Blanchet, F., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.,
1011 O’Hara, R., Simpson, G., Stevens, M., Wagner, H., 2018. *vegan: Community Ecology*
1012 *Package*.
- 1013 Orsoni, V., Souchu, P., Sauzade, D., 2001. Caractérisation de l’état d’eutrophisation des trois
1014 principaux étangs corses (Biguglia, Diana et Urbino), et proposition de renforcement
1015 de leur surveillance. Rapport final.
- 1016 Paerl, H.W., Dyble, J., Moisaner, P.H., Noble, R.T., Piehler, M.F., Pinckney, J.L., Steppe,
1017 T.F., Twomey, L., Valdes, L.M., 2003. Microbial indicators of aquatic ecosystem
1018 change: current applications to eutrophication studies. *FEMS Microbiol. Ecol.* 46,
1019 233–246. [https://doi.org/10.1016/S0168-6496\(03\)00200-9](https://doi.org/10.1016/S0168-6496(03)00200-9)
- 1020 Paerl, H.W., Rossignol, K.L., Hall, N.S., Peierls, B.L., Wetz, M.S., 2010. Phytoplankton
1021 Community Indicators of Short- and Long-term Ecological Change in the
1022 Anthropogenically and Climatically Impacted Neuse River Estuary, North Carolina,
1023 USA. *Estuaries and Coasts* 33, 485–497. <https://doi.org/10.1007/s12237-009-9137-0>
- 1024 Pasqualini, V., Derolez, V., Garrido, M., Orsoni, V., Baldi, Y., Etourneau, S., Leoni, V.,
1025 Rébillout, P., Laugier, T., Souchu, P., Malet, N., 2017. Spatiotemporal dynamics of
1026 submerged macrophyte status and watershed exploitation in a Mediterranean coastal

- 1027 lagoon: Understanding critical factors in ecosystem degradation and restoration.
1028 Ecological Engineering 102, 1–14. <https://doi.org/10.1016/j.ecoleng.2017.01.027>
- 1029 Patiño, J., Guilhaumon, F., Whittaker, R.J., Triantis, K.A., Gradstein, S.R., Hedenäs, L.,
1030 González-Mancebo, J.M., Vanderpoorten, A., 2013. Accounting for data heterogeneity
1031 in patterns of biodiversity: an application of linear mixed effect models to the oceanic
1032 island biogeography of spore-producing plants. *Ecography* 36, 904–913.
1033 <https://doi.org/10.1111/j.1600-0587.2012.00020.x>
- 1034 Piehler, M.F., Twomey, L.J., Hall, N.S., Paerl, H.W., 2004. Impacts of inorganic nutrient
1035 enrichment on phytoplankton community structure and function in Pamlico Sound,
1036 NC, USA. *Estuarine, Coastal and Shelf Science* 61, 197–209.
1037 <https://doi.org/10.1016/j.ecss.2004.05.001>
- 1038 R Core Team, 2013. R: A language and environment for statistical computing. R Foundation
1039 for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- 1040 Raven, J.A., 1998. The twelfth Tansley Lecture. Small is beautiful: the picophytoplankton.
1041 *Funct. Ecol.* 12, 503–513. <https://doi.org/10.1046/j.1365-2435.1998.00233.x>
- 1042 Reed, M., Pinckney, J.L., Keppler, C., Brock, L., Hogan, S., Greenfield, D., 2016. The
1043 influence of nitrogen and phosphorus on phytoplankton growth and assemblage
1044 composition in four coastal, southeastern USA systems. *Estuarine Coastal and Shelf
1045 Science* 177, 71–82.
- 1046 Rochette, S., Rivot, E., Morin, J., Mackinson, S., Riou, P., Le Pape, O., 2010. Effect of
1047 nursery habitat degradation on flatfish population: Application to *Solea solea* in the
1048 Eastern Channel (Western Europe). *Journal of Sea Research* 64, 34–44.
1049 <https://doi.org/10.1016/j.seares.2009.08.003>
- 1050 Schramm, W., 1999. Factors influencing seaweed responses to eutrophication: some results
1051 from EU-project EUMAC. *Journal of Applied Phycology* 11, 69–78.
1052 <https://doi.org/10.1023/A:1008076026792>
- 1053 Seong, K.A., Myung, G., Jeong, H.J., Yih, W., Kim, H.S., Park, J.Y., Yoo, Y.D., 2017.
1054 Ingestion rate and grazing impact by the mixotrophic ciliate *Mesodinium rubrum* on
1055 natural populations of marine heterotrophic bacteria in the coastal waters of Korea.
1056 *Algae* 32, 47–55.
- 1057 Sorokin, Y., Dalocchio, F., 2008. Dynamics of phosphorus in the Venice lagoon during a
1058 picocyanobacteria bloom. *Journal of Plankton Research* 30, 1019–1026.
- 1059 Souchu, P., Bec, B., Smith, V.H., Laugier, T., Fiandrino, A., Benau, L., Orsoni, V., Collos,
1060 Y., Vaquer, A., 2010. Patterns in nutrient limitation and chlorophyll a along an
1061 anthropogenic eutrophication gradient in French Mediterranean coastal lagoons. *Can.
1062 J. Fish. Aquat. Sci.* 67, 743–753. <https://doi.org/10.1139/F10-018>
- 1063 Tomas, C.R., 1997. *Identifying Marine Phytoplankton*. Academic Press.
- 1064 Tong, M., Smith, J., Anderson, D.M., 2015. Role of dissolved nitrate and phosphate in
1065 isolates of *Mesodinium rubrum* and toxin-producing *Dinophysis acuminata*. *Aquatic
1066 Microbial Ecology* 75, 169–185.


1067 Xu, J., Yin, K., Lee, J.H.W., Anderson, D.M., Jiang, Y., Yuan, X., Ho, A.Y.T., Harrison, P.J.,
1068 2012. Resistance of Hong Kong waters to nutrient enrichment: assessment of the role
1069 of physical processes in reducing eutrophication. *J. Oceanogr.* 68, 545–560.
1070 <https://doi.org/10.1007/s10872-012-0118-8>.

1071

1072 7. Supplementary material

1073 Table S1. Net growth rates (μ_0) measured in incubations without enrichment and maximal
 1074 growth (μ_{\max}) and mortality rates (g) subsequently calculated using full N and P enrichment in
 1075 the two NB and SB stations of Biguglia lagoon. Values were estimated according the linear
 1076 Landry and Hassett (1982) equation. This model has been compared to alternative models by
 1077 mixed-effect multiple linear regressions of apparent growth rates, based on Chl *a*
 1078 concentration from dilution experiments. The parsimony of the different models was checked
 1079 by the AIC criterion. The linear model was only accepted when it presented a delta-AIC_c, <2
 1080 with respect to the most parsimonious model (see Methods). AIC_{c-w} describe the strength of
 1081 explanation by the model of the model. R²: coefficient of determination. The $\mu_0:\mu_{\max}$ ratio
 1082 reflects the impact of the nutrient enrichment on the phytoplankton growth (1: no effect; 0:
 1083 high enhancement of growth), highlighting the nutrient limitation under natural conditions.
 1084 ****p*-value <0.001, ***p*-value <0.01, **p*-value <0.05.

Stations	Samplings	Fraction	μ_0 μ_{\max} g			n	AIC _c	Δ AIC _c	AIC _{c-w}	R ²	$\mu_0:\mu_{\max}$	
			(d ⁻¹)									
NB	Nov - Dec 2013	Tot	0.45	<0	0.62	8	11.5	1.45	0.33	0.31	<0	
		Micro	<0	<0	1.45	7	37.1	6.05	0.04	-0.05	-	
		Nano	0.37	0.46	0.26	8	6.4	3.54	0.13	0.10	0.79	
		Ultra	0.96	<0	1.82	8	24.7	0.00	0.55	0.45*	<0	
	April 2014	Tot	<0	0.25	<0	7	1.4	4.25	0.10	0.19	<0	
		Micro	<0	0.24	1.01	8	24.0	2.39	0.19	0.13	<0	
		Nano	1.13	1.83	0.28	8	-5.2	0.00	0.50	0.53*	0.62	
		Ultra	<0	<0	0.43	8	9.9	2.26	0.21	0.23	-	
	September 2014	Tot	0.64	1.78	0.30	9	-10.7	0.00	0.68	0.64**	0.36	
		Micro	1.84	3.40	0.57	9	3.2	0.00	1.00	0.58*	0.54	
		Nano	1.60	2.70	0.57	9	-4.1	0.00	1.00	0.76**	0.59	
		Ultra	0.51	1.12	0.45	8	2.20	0.86	0.39	0.48*	0.45	
	SB	Nov - Dec 2013	Tot	0.48	0.41	0.69	10	1.7	0.00	0.54	0.65**	1.19
			Micro	1.12	0.45	1.12	9	45.3	4.40	0.09	-0.09	2.48
			Nano	1.17	0.90	0.50	10	20.3	2.62	0.18	0.05	1.30
			Ultra	<0	0.14	1.10	8	9.3	0.00	0.93	0.70**	<0
April 2014		Tot	0.17	0.19	<0	7	15.1	5.18	0.07	0.08	0.92	
		Micro	-	-	-	-	-	-	-	-	-	
		Nano	1.91	2.00	0.83	7	12.8	0.37	0.33	0.59*	0.95	
		Ultra	<0	0.01	<0	8	14.3	4.46	0.09	-0.01	<0	
September 2014		Tot	<0	1.79	0.16	7	-21.1	0.45	0.44	0.88**	<0	
		Micro	0.39	10.08	0.39	9	27.9	4.80	0.05	-0.09	0.04	
		Nano	1.42	3.23	1.14	10	11.6	0.69	0.40	0.65**	0.44	
		Ultra	<0	0.87	<0	10	9.6	0.67	0.32	0.38*	<0	



1085

1086

1087

1088

1089

1090

1091

1092

1093

Fig. S1. Apparent growth rate as a function of the dilution factor in bioassays with full N and P enrichment in NB and SB station of Biguglia lagoon in (A) November-December 2013, (B) April 2014, and (C) September 2014. From the left to the right, total phytoplankton, microphytoplankton $>20 \mu\text{m}$, nanophytoplankton between 5 and $20 \mu\text{m}$, and ultraphytoplankton $<5 \mu\text{m}$. The line corresponds to the linear equation $k(x) = \mu_{\text{max}} - gx$, with k the apparent growth rate, x the fraction of undiluted plankton, μ_{max} (as Y axis intercept) the maximum growth rate, and g the grazing rate. Blue lines highlight when the linear model best-fitted the relationship between the apparent growth rates and the dilution factors.


1094 Table S2. Net growth (μ_N) and mortality rates (g) subsequently calculated using enrichment
1095 minus N in the two NB and SB stations of Biguglia lagoon. Values were estimated according
1096 the equations from Andersen et al. (1991). The linear model has been compared to alternative
1097 models by mixed-effect multiple linear regressions of apparent growth rates, based on Chl *a*
1098 concentration from dilution experiments. The parsimony of the different models was checked
1099 by the AIC criterion. The linear model was only accepted when it presented a delta-AIC_c, <2
1100 with respect to the most parsimonious model (see Methods). AIC_{c-w} describe the strength of
1101 explanation by the model of the model. R²: coefficient of determination. ****p*-value <0.001,
1102 ***p*-value <0.01, **p*-value <0.05.

Stations	Samplings	Fraction	μ_N	g	n	AIC _c	Δ AIC _c	AIC _{c-w}	R ₂	
			(d ⁻¹)							
NB	Nov - Dec 2013	Total	<0	0.19	8	-5.3	1.75	0.30	0.28	
		Micro-	<0	1.70	6	-	-	-	-0.13	
		Nano-	0.62	0.29	8	-9	0,00	0.93	0.70**	
		Ultra-	<0	1.82	8	24.7	0,00	0.55	0.45*	
	April 2014	Total	0.51	0.71	9	-5	0,00	1,00	0.84***	
		Micro-	1.59	2.57	8	19.4	0,00	0.93	0.70**	
		Nano-	1.95	0.83	10	-6.1	0,00	1,00	0.86***	
		Ultra-	-1.59	0.13	8	-	-	-	-0.03	
	September 2014	Total	0.86	0.71	10	-1.3	0,00	0.99	0.73**	
		Micro-	1.78	0.55	9	10	0.02	0.50	0.33 .	
		Nano-	1.77	0.99	10	12.6	0,00	0.93	0.56**	
		Ultra-	0.51	0.73	9	1.3	0,00	0.98	0.73**	
	SB	Nov - Dec 2013	Total	0.26	0.39	10	-9.7	0,00	0.98	0.65**
			Micro-	1.09	2.14	9	35.8	0.38	0.45	0.30 .
			Nano-	0.78	0.15	9	-	-	-	-0.08
			Ultra-	1.85	0.82	10	3.1	0,00	0.99	0.69**
April 2014		Total	0.06	0.03	9	-	-	-	-0.14	
		Micro-	-	-	-	-	-	-	-	
		Nano-	1.64	0.42	10	-	-	-	0.08	
		Ultra-	0.10	0.50	9	4.1	0,00	0.71	0.45*	
September 2014		Total	0.28	1.06	10	1.2	0,00	1.00	0.83***	
		Micro-	9.80	1.11	10	8.7	0,00	0.99	0.70**	
		Nano-	1.58	1.01	10	-3.6	0,00	1,00	0.87***	
		Ultra-	<0	1.25	10	7.1	0,00	1.00	0.78***	

1103 Table S3. Net growth (μ_p) and mortality rates (g) subsequently calculated using enrichment
 1104 minus P in the two NB and SB stations of Biguglia lagoon. Values were estimated according
 1105 the equations from Andersen et al. (1991). The linear model has been compared to alternative
 1106 models by mixed-effect multiple linear regressions of apparent growth rates, based on Chl *a*
 1107 concentration from dilution experiments. The parsimony of the different models was checked
 1108 by the AIC criterion. The linear model was only accepted when it presented a delta-AIC_c, <2
 1109 with respect to the most parsimonious model (see Methods). AIC_{c-w} describe the strength of
 1110 explanation by the model of the model. R²: coefficient of determination. ****p*-value <0.001,
 1111 ***p*-value <0.01, **p*-value <0.05.

Stations	Samplings	Fraction	μ_p	g	n	AIC _c	Δ AIC _c	AIC _{c-w}	R ²
			(d ⁻¹)						
NB	Nov - Dec 2013	Total	0.30	0.24	9	-1.9	1.16	0.36	0.24
		Micro-	<0	<0	10	-	-	-	-0.12
		Nano-	1.13	0.44	10	1.9	0,00	0.74	0.40*
		Ultra-	<0	<0	10	-	-	-	-0.12
	April 2014	Total	0.07	0.41	8	1.9	0,00	0.65	0.50*
		Micro-	1.01	1.46	4	-	-	-	0.27
		Nano-	1.54	0.44	8	8.2	1.35	0.34	0.31
		Ultra-	<0	0.53	6	7.4	1.14	0.36	0.71*
	September 2014	Total	1.03	0.24	10	-4.8	0.15	0.48	0.26 .
		Micro-	2.24	0.58	10	12.3	0.01	0.50	0.27
		Nano-	2.13	0.29	10	-5.2	0,00	0.67	0.37*
		Ultra-	0.52	0.31	9	-0.1	0.12	0.49	0.32 .
SB	Nov - Dec 2013	Total	0.53	0.78	10	-0.6	0,00	1.00	0.75***
		Micro-	1.05	2.65	9	25.7	0,00	0.95	0.65**
		Nano-	1.07	0.48	10	3.5	0,00	0.74	0.40*
		Ultra-	2.02	0.83	10	5.2	0,00	0.98	0.65**
	April 2014	Total	0.26	0.57	9	-9.5	0,00	1.00	0.86***
		Micro-	-	-	-	-	-	-	-
		Nano-	1.64	0.52	9	3	0,00	0.83	0.53*
		Ultra-	0.34	0.69	9	-7	0,00	1.00	0.87***
	September 2014	Total	1.34	0.03	9	-	-	-	-0.14
		Micro-	11.72	-0.17	10	-	-	-	-0.06
		Nano-	<0	2.45	8	-	-	-	0.15
		Ultra-	<0	0.19	9	-4.6	1.65	0.30	0.19


1112



1113

1114 Fig. S2. Apparent growth rate as a function of the dilution factor in bioassays with full FH
 1115 enrichment minus N and minus P, in the NB (A) and SB (B) stations of Biguglia lagoon in
 1116 November-December 2013. From the left to the right, total phytoplankton,
 1117 microphytoplankton $>20 \mu\text{m}$, nanophytoplankton between 5 and $20 \mu\text{m}$, and
 1118 ultraphytoplankton $<5 \mu\text{m}$. The line corresponds to the linear equation $k(x) = \mu - gx$, with k
 1119 the apparent gross growth rate, x the fraction of undiluted plankton, μ (as Y axis intercept) the
 1120 gross growth rate, and g the grazing rate. However, we assume that this model can not best-fit
 1121 the relationship between apparent growth rate and fraction of undiluted plankton in nutrient
 1122 limiting conditions, due to the potential use of alternative resources (Andersen et al., 1991).


1123



1124

1125 Fig. S3. Apparent growth rate as a function of the dilution factor in bioassays with full F
 1126 enrichment minus N and minus P, in the NB (A) and SB (B) stations of Biguglia lagoon in
 1127 April 2014. From the left to the right, total phytoplankton, microphytoplankton >20 μm
 1128 (absent in SB), nanophytoplankton between 5 and 20 μm , and ultraphytoplankton <5 μm . The
 1129 line corresponds to the linear equation $k(x) = \mu - gx$, with k the apparent gross growth rate, x
 1130 the fraction of undiluted plankton, μ (as Y axis intercept) the gross growth rate, and g the
 1131 grazing rate. However, we assume that this model can not best-fit the relationship between
 1132 apparent growth rate and fraction of undiluted plankton in nutrient limiting conditions, due to
 1133 the potential use of alternative resources (Andersen et al., 1991).

1134



1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

Fig. S4. Apparent growth rate as a function of the dilution factor in bioassays with full H enrichment minus N and minus P, in the NB (A) and SB (B) stations of Biguglia lagoon in September 2014. From the left to the right, total phytoplankton, microphytoplankton >20 μm , nanophytoplankton between 5 and 20 μm , and ultraphytoplankton <5 μm . The line corresponds to the linear equation $k(x) = \mu - gx$, with k the apparent growth rate, x the fraction of undiluted plankton, μ (as Y axis intercept) the growth rate, and g the grazing rate. However, we assume that this model can not best-fit the relationship between apparent growth rate and fraction of undiluted plankton in nutrient limiting conditions, due to the potential use of alternative resources (Andersen et al., 1991).