



HAL
open science

Conformality for a robust class of non-conformal attractors

Beatrice Pozzetti, Andres Sambarino, Anna Wienhard

► **To cite this version:**

Beatrice Pozzetti, Andres Sambarino, Anna Wienhard. Conformality for a robust class of non-conformal attractors. *J. für die reine und ange. Math.*, 2020, 10.1515/crelle-2020-0029. hal-02391752v2

HAL Id: hal-02391752

<https://hal.science/hal-02391752v2>

Submitted on 27 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CONFORMALITY FOR A ROBUST CLASS OF NON-CONFORMAL ATTRACTORS

BEATRICE POZZETTI, ANDRÉS SAMBARINO, AND ANNA WIENHARD

ABSTRACT. In this paper we investigate the Hausdorff dimension of limit sets of Anosov representations. In this context we revisit and extend the framework of hyperconvex representations and establish a convergence property for them, analogue to a differentiability property. As an application of this convergence, we prove that the Hausdorff dimension of the limit set of a hyperconvex representation is equal to a suitably chosen critical exponent.

CONTENTS

1.	Introduction	1
2.	Preliminaries	6
3.	Anosov representations	16
4.	An upper bound on the Hausdorff dimension of the limit set	20
5.	Local conformality and Hausdorff dimension	21
6.	(p, q, r) -hyperconvexity	29
7.	Differentiability properties	34
8.	Examples of locally conformal representations	37
9.	Fundamental groups of surfaces	42
	References	46

1. INTRODUCTION

In his seminal paper, Sullivan [42] describes the Hausdorff dimension of the limit set L_Γ , of a discrete group Γ acting on the real hyperbolic n -space, in terms of the Dirichlet series

$$s \mapsto \sum_{\gamma \in \Gamma} e^{-sd(o, \gamma o)}.$$

More precisely, the *critical exponent* of such a series is

$$h_\Gamma = \inf \left\{ s : \sum_{\gamma \in \Gamma} e^{-sd(o, \gamma o)} < \infty \right\} = \sup \left\{ s : \sum_{\gamma \in \Gamma} e^{-sd(o, \gamma o)} = \infty \right\}$$

and Sullivan shows:

Theorem (Sullivan). *If Γ is a convex co-compact subgroup of $\text{PSO}(1, n)$ then the Hausdorff dimension of L_Γ is h_Γ .*

A.S. was partially financed by ANR DynGeo ANR-16-CE40-0025. B.P. and A.W. acknowledge funding by the Deutsche Forschungsgemeinschaft within the Priority Program SPP 2026 “Geometry at Infinity”. A.W. acknowledges funding by the European Research Council under ERC-Consolidator grant 614733, and by the Klaus-Tschira- Foundation.

This is related to understanding the Hausdorff dimension of a hyperbolic set in dynamical terms. Indeed, the non-wandering set of the geodesic flow of $\Gamma \backslash \mathbb{H}^n$ is, by definition, a maximal isolated compact hyperbolic set, h_Γ is its topological entropy and Sullivan's result can be interpreted in terms of the Ledrappier-Young formula [32].

Describing the Hausdorff dimension of a hyperbolic repeller as a dynamical quantity is today well understood in the *conformal* setting, i.e. when the derivative of the dynamics, restricted to the unstable distribution, acts as a conformal map (see Chen-Pesin's survey [10] and references therein). Analogously, Sullivan's result has been generalized to convex-cocompact groups of a CAT(-1)-space X (see for example Bourdon [5] and Yue [44]). The metric on the visual boundary ∂X used to compute the Hausdorff dimension is the *visual metric*, for which the action of $\text{Isom } X$ is conformal (i.e. sends balls to balls).

However, other natural metrics on ∂X appear in very common situations: if X is a rank 1 symmetric space of non-compact type, then its visual boundary carries the structure of a differentiable manifold and thus one would also like to understand the Hausdorff dimension of limit sets for a (any) Riemannian metric on ∂X . Unless X is the real hyperbolic n -dimensional space, the Riemannian structure behaves differently from the visual structure: the action of $\text{Isom } X$ is no longer conformal.

The dynamical characterization of Hausdorff dimension in a non-conformal setting is still not completely understood. We refer the reader again to Chen-Pesin's survey [10]. Let us also note that only very recently Bárány-Hochman-Rapaport [1] provided a complete answer for Iterated-Function-Systems on the plane. On the discrete groups side, Dufloux [16] has studied a class of Schottky subgroups of isometries of the complex hyperbolic n -space, that he calls *well positioned*, and proves the analogue of Sullivan's result for the Hausdorff dimension of the limit set with respect to any Riemannian metric.

1.1. This paper. In this paper we are interested in describing the Hausdorff dimension of the limit set of discrete subgroups of a semi-simple Lie group G , for a Riemannian structure on the flag spaces (or boundaries) of G . The groups we will consider, called *Anosov representations*, are in many ways similar to convex cocompact subgroups of $\text{SO}(1, n)$, but do not act conformally on the boundaries of G .

Anosov representations were introduced by Labourie [31] for fundamental groups of negatively curved closed manifolds and the definition was extended by Guichard-W. [23] to any hyperbolic group. Such representations provide the appropriate generalization of the class of convex co-compact subgroups in the context of Lie groups of higher rank [23, 27, 28].

We will not use the original definition but follow a more recent approach, developed by Kapovich-Leeb-Porti [28], G3ritaud-Guichard-Kassel-W. [20] and in particular Bochi-Potrie-S. [4], that provides a simplified definition and gives better quantitative control of Anosov representations.

Let $\mathbb{K} = \mathbb{R}$ or \mathbb{C} , consider an inner (or Hermitian if $\mathbb{K} = \mathbb{C}$) product in \mathbb{K}^d and, for $g \in \text{GL}_d(\mathbb{K})$, denote by $g \mapsto g^*$ the corresponding adjoint operator. The singular values of g , i.e. the square root of the modulus of the eigenvalues of gg^* , are denoted by

$$\sigma_1(g) \geq \dots \geq \sigma_d(g).$$

Let Γ be a finitely generated discrete group, consider a finite symmetric generating set S and denote by $||$ the associated word metric on Γ . Given $p \in \llbracket 1, d-1 \rrbracket$ denote by $\mathcal{G}_p(\mathbb{K}^d)$ the Grassmannian of p -dimensional subspaces of \mathbb{K}^d . For a homomorphism $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$, the following are equivalent:

i) There exist positive constants c, μ such that for all $\gamma \in \Gamma$ one has

$$\frac{\sigma_{p+1}}{\sigma_p}(\rho(\gamma)) \leq ce^{-\mu|\gamma|},$$

ii) The group Γ is word-hyperbolic and there exist ρ -equivariant maps $(\xi^p, \xi^{d-p}) : \partial\Gamma \rightarrow \mathcal{G}_p(\mathbb{K}^d) \times \mathcal{G}_{d-p}(\mathbb{K}^d)$ such that for every $x \neq y \in \partial\Gamma$ one has

$$\xi^p(x) \oplus \xi^{d-p}(y) = \mathbb{K}^d,$$

and a suitable associated flow is contracting.

If either condition is satisfied we will say that ρ is an $\{\mathfrak{a}_p\}$ -Anosov representation¹. For such a representation, the critical exponent $h_\rho^{\mathfrak{a}_p}$ of the Dirichlet series

$$\Phi_\rho^{\mathfrak{a}_p}(s) = \sum_{\gamma \in \Gamma} \left(\frac{\sigma_{p+1}}{\sigma_p}(\rho(\gamma)) \right)^s \quad (1)$$

is well defined. By definition, the series is convergent for every $s > h_\rho^{\mathfrak{a}_p}$ and divergent for every $0 < s < h_\rho^{\mathfrak{a}_p}$.

If ρ is furthermore $\{\mathfrak{a}_{p+1}\}$ -Anosov then $h_\rho^{\mathfrak{a}_p}$ is analytic with respect to ρ , and agrees with the entropy of a suitably defined flow (see for example Bridgeman-Canary-Labourie-S. [6] and Potrie-S. [35, Corollary 4.9]). But in general little is known about $h_\rho^{\mathfrak{a}_p}$ without this extra assumption.

We will mainly focus on $\{\mathfrak{a}_1\}$ -Anosov representations. The chosen inner product on \mathbb{K}^d induces a metric on $\mathbb{P}(\mathbb{K}^d)$, we will denote by $\mathrm{Hff}(A)$ the Hausdorff dimension of a subset $A \subset \mathbb{P}(\mathbb{K}^d)$ for this metric. As a first result we obtain the following, independently obtained by Glorieux-Monclair-Tholozan [19].

Proposition (Proposition 4.1). *Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be $\{\mathfrak{a}_1\}$ -Anosov. Then*

$$\mathrm{Hff}(\xi^1(\partial\Gamma)) \leq h_\rho^{\mathfrak{a}_1}.$$

In order to discuss situations in which equality holds, we introduce the notion of *locally conformal points* of ρ (Definition 5.5), these are points of $\partial\Gamma$ designed to detect some asymptotic conformality of the non-conformal action of $\rho(\Gamma)$ when restricted to the limit set $\xi^1(\partial\Gamma)$. Using Patterson's construction we then obtain a (not necessarily quasi-invariant) measure $\mu_\rho^{\mathfrak{a}_1}$ on $\partial\Gamma$. Following Sullivan, we then prove the following.

Theorem (Theorem 5.14). *If the set of locally conformal points of ρ has positive $\mu_\rho^{\mathfrak{a}_1}$ -measure, then*

$$\mathrm{Hff}(\xi^1(\partial\Gamma)) = h_\rho^{\mathfrak{a}_1}.$$

Interestingly, for a rich class of Anosov representations, a 3-point transversality condition, inspired by Labourie [31], forces asymptotic conformality:

¹The implication ii) \Rightarrow i) comes from Labourie [31] and Guichard-W. [23]. The implication i) \Rightarrow ii) is more recent and due to Kapovich-Leeb-Porti [28], see also Guéritaud-Guichard-Kassel-W. [20] and Bochi-Potrie-S. [4] for different approaches. In the language of Bochi-Potrie-S. [4, Section 3.1] a representation verifying condition i) is called *p-dominated*.

Definition. Consider $p, q, r \in \llbracket 1, d-1 \rrbracket$ such that $p + q \leq r$. A $\{\mathfrak{a}_p, \mathfrak{a}_q, \mathfrak{a}_r\}$ -Anosov representation $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ is called (p, q, r) -hyperconvex if for every triple of pairwise distinct points $x, y, z \in \partial\Gamma$ one has

$$(\xi^p(x) \oplus \xi^q(y)) \cap \xi^{d-r}(z) = \{0\}.$$

(Note that p and q are not required to be distinct.)

The main result of this paper is the following.

Theorem A (Corollary 6.9 and Corollary 7.3). *Let ρ be $(1, 1, 2)$ -hyperconvex. Then*

$$h_\rho^{\mathfrak{a}_1} = \mathrm{Hff}(\xi^1(\partial\Gamma)) \leq \mathrm{Hff}(\mathbb{P}(\mathbb{K}^2)).$$

The aforementioned analyticity result for $h_\rho^{\mathfrak{a}_p}$, together with Theorem A, has the following consequence:

Corollary 1.1. *Let $\{\rho_u : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})\}_{u \in D}$ be an analytic family of $(1, 1, 2)$ -hyperconvex representations, then $u \mapsto \mathrm{Hff}(\xi_u^1(\partial\Gamma))$ is analytic.*

In fact Theorem A holds in greater generality. We can replace 2's by any $p \in \llbracket 2, d-1 \rrbracket$ if we additionally require that for every $\gamma \in \Gamma$ one has

$$\sigma_2(\rho(\gamma)) = \sigma_p(\rho(\gamma)),$$

see Corollary 6.10 and Corollary 7.3. This extra condition on the singular values should be interpreted as a restriction on the Zariski closure of the representation (see subsection 8.1 for situations such as $\mathrm{PSp}(1, n)$ and $\mathrm{PU}(1, n)$, and subsection 8.2 for the group $\mathrm{PSO}(p, q)$).

A key ingredient for the proof of Theorem A is the following convergence property for hyperconvex representations, from the inequality readily follows.

Theorem B (Theorem 7.1). *Let ρ be (p, q, r) -hyperconvex, then for every $(w, y) \in \partial^{(2)}\Gamma$ one has*

$$\lim_{(w, y) \rightarrow (x, x)} d(\xi^p(w) \oplus \xi^q(y), \xi^r(x)) = 0.$$

We further investigate how vast the class of hyperconvex representations is. On the one hand one has the following remarks that provide many examples by the *represent and deform* method (see subsection 7.2):

- if $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{R})$ is hyperconvex then, by complexifying, one obtains a hyperconvex representation over \mathbb{C} : this is direct from the definition;
- the space of (p, q, r) -hyperconvex representations is open in $\mathrm{hom}(\Gamma, \mathrm{PGL}_d(\mathbb{K}))$ (Proposition 6.2).

On the other hand there are some ‘verifiable’ restrictions imposed by the hyperconvexity condition. For example, a $(1, 1, p)$ -hyperconvex representation of Γ induces a continuous injective map

$$\partial\Gamma - \{\text{point}\} \rightarrow \mathbb{P}(\mathbb{K}^p),$$

(see Corollary 6.6), and there might be topological obstructions for the existence of such a map. More interesting restrictions arise when $\mathbb{K} = \mathbb{R}$ and $\partial\Gamma$ is a manifold:

Corollary (Proposition 7.4). *Let Γ be such that $\partial\Gamma$ is homeomorphic to a $(p-1)$ -dimensional sphere. If $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{R})$ is $(1, 1, p)$ -hyperconvex then $\xi^1(\partial\Gamma)$ is a C^1 sphere.*

Using openness of hyperconvexity we find new explicit examples of Zariski dense groups with C^1 limit set.

Corollary (Corollary 7.7). *There exist Zariski dense subgroups $\Gamma < \mathrm{PGL}_{d(d+1)}(\mathbb{R})$ whose limit set is a C^1 sphere of dimension $d - 1$.*

Sharper results of similar nature were obtained by Zhang-Zimmer [45].

We now turn to the special situation when $\partial\Gamma$ is a circle. Then Theorem A gives the following computation of $h_\rho^{\mathfrak{a}_1}$:

Corollary. *Assume $\partial\Gamma$ is homeomorphic to a circle, if $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{R})$ is $(1, 1, 2)$ -hyperconvex then $h_\rho^{\mathfrak{a}_1} = 1$.*

This implies Potrie-S. [35, Theorem B] and further generalizes it to the Hitchin component of $\mathrm{PSO}(p, p)$. The proof of Potrie-S. [35, Theorem A] applies then verbatim also to the Hitchin component of $\mathrm{PSO}(p, p)$ and we thus obtain a rigid inequality for the critical exponent in the symmetric space of $\mathrm{PSO}(p, p)$. We refer the reader to Sections 9.2 and 9.3 for more details on Hitchin representations.

While the property of having constant $h_\rho^{\mathfrak{a}_1}$ was expected to be a rare phenomenon, peculiar to Hitchin components, or possibly higher rank Teichmüller theories, we provide, in Section 6.3, many more examples of representations of fundamental groups of surfaces for which Theorem A applies. Interestingly enough, when $\partial\Gamma$ is a circle (and $\mathbb{K} = \mathbb{R}$), $(1, 1, 2)$ -hyperconvexity is not only a local condition, but it can be pushed far away. We say that an $\{\mathfrak{a}_1\}$ -Anosov representation is *weakly irreducible* if $\xi^1(\partial\Gamma)$ is not contained in a proper subspace of $\mathbb{P}(\mathbb{R}^d)$.

Proposition (Proposition 9.3). *Assume that $\partial\Gamma$ is homeomorphic to a circle. Then the space of real weakly irreducible $(1, 1, 2)$ -hyperconvex representations of Γ is closed among real weakly irreducible $\{\mathfrak{a}_1, \mathfrak{a}_2\}$ -Anosov representations.*

Throughout the paper we allow \mathbb{K} to be a local field (not necessarily Archimedean, as we required in this introduction). Originally Anosov representations were only defined over Archimedean fields as it is possible to show that if Γ admits a Anosov representation $\rho : \Gamma \rightarrow \mathrm{PSL}_d(\mathbb{K})$ for non-Archimedean \mathbb{K} , then Γ is virtually free. The main result of our paper, however, associates to such an action an interesting geometric quantity, the Hausdorff dimension of the limit set, which we are able to relate to a dynamical data, the orbit growth rate. We find this very interesting, and this justifies the extra work needed to develop the theory in this more general setting.

The main results go through in this generality, except the analyticity of Hausdorff dimension: the key step is to show that for a $\{\mathfrak{a}_1\}$ -Anosov representation its *entropy*, defined by

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log \#\{\gamma \in \Gamma : \log \sigma_1(\rho(\gamma)) \leq t\}$$

is analytic with ρ . We don't know if this is true, but one can use the thermodynamical formalism to prove that the Hausdorff dimension depends continuously on the representation (and is actually as regular as the map $\rho \mapsto \xi_\rho$ is).

Outline of the paper. The preliminaries of the paper, collected in Section 2 come from three different areas: quantitative linear algebra, dynamics and geometric group theory. In § 2.1 we recall relations between the singular values of an element in $\mathrm{PGL}_d(\mathbb{K})$ and metric properties of its action on Grassmannian manifolds, in the general context of a local field \mathbb{K} . In § 2.2 we discuss the dynamical backgrounds and indicate how to extend Bochi-Gourmelon’s theorem as well as the theory of dominated splittings to general local fields. § 2.3 collects the facts about hyperbolic groups and cone types that we will need in the paper.

Section 3 concerns Anosov representations: we extend to the non-Archimedean setup the definition and the results we will need, particularly concerning the definition and properties of the equivariant boundary maps. Our discussion follows the lines of Bochi-Potrie-S. [4].

In Section 4 we prove that for any Anosov representation the Hausdorff dimension of the limit curve provides a lower bound for the critical exponent for the first root. In Section 5 we give a condition guaranteeing that such bound is optimal, namely the abundance of *locally conformal points* with respect to a suitable measure.

Section 6 concerns the notion of (p, q, r) -hyperconvexity, an open condition (Proposition 6.2) that guarantees abundance of locally conformal points: this is the content of Proposition 6.7, the main technical result of the paper. Using the theory of SL_2 representations we provide in § 6.3 many examples of hyperconvex representations of fundamental groups of surfaces and hyperbolic three manifolds.

In Section 7 we discuss another interesting consequence of hyperconvexity: such property guarantees a weak differentiability property for the limit set (Theorem 7.1) which allows us, on the one hand, to obtain good bounds on the Hausdorff dimension (Proposition 7.3), and on the other to provide examples of Zariski dense subgroups whose limit set in the projective space is a C^1 manifold: we obtain these through the *represent and deform* method explained, in a concrete example, in Proposition 7.5.

In Section 8 we discuss in detail two families of representations for which all our results apply: on the one hand we detail the geometric meaning of our notions in the case of convex cocompact subgroups of rank one groups, rediscovering and generalizing results of Dufloux (§ 8.1), on the other we give a concrete criterion that guarantees hyperconvexity for subgroups of $\mathrm{SO}(p, q)$ and provide examples of groups that satisfy it (§ 8.2).

The last section of the paper (Section 9) concerns representations of fundamental groups of hyperbolic surfaces (or more generally compact hyperbolic orbifolds). For these we show that hyperconvexity is also a closed condition (Proposition 9.3), and discuss a new proof and generalization of a result of Potrie-S. [35].

2. PRELIMINARIES

In the paper we will need preliminaries from three different sources: quantitative linear algebra, dynamics and particularly the work of Bochi-Gourmelon [3] and Bochi-Potrie-S. [4] on dominated sequences, and algebraic and metric properties of hyperbolic groups. We recall the results we need here.

2.1. Quantitative linear algebra. As anticipated at the end of the introduction, in the paper we will be dealing with representations of finitely generated groups on finite dimensional vector spaces over local fields. We recall here some quantitative

results we will need. More details on algebraic groups over local fields can be found in Quint [38].

2.1.1. *Angles and distances on Grassmannians.* We denote by \mathbb{K} a local field, and by $|\cdot| : \mathbb{K} \rightarrow \mathbb{R}^+$ its absolute value. Recall that if \mathbb{K} is \mathbb{R} or \mathbb{C} then $|\cdot|$ is the usual modulus, if, instead, \mathbb{K} is non-Archimedean, we require that $|\omega| = \frac{1}{q}$ where ω denotes the *uniformizing element*, namely a generator of the maximal ideal of the valuation ring \mathcal{O} , and q is the cardinality of the residue field $\mathcal{O}/\omega\mathcal{O}$ (this is finite because \mathbb{K} is, by assumption, local).

Given a finite dimensional vector space V over \mathbb{K} , we denote by $\|\cdot\| : V \rightarrow \mathbb{R}^+$ a *good norm*: for an Archimedean field \mathbb{K} this means that $\|\cdot\|$ is induced from an Hermitian product, if \mathbb{K} is non-Archimedean this means that there exists a basis $\{e_1, \dots, e_n\}$ such that $\|\sum a_i e_i\| = \max\{|a_i|\}$. In this second case we say that a decomposition $V = V_1 \oplus V_2$ is *orthogonal* if $\|v_1 + v_2\| = \max\{\|v_1\|, \|v_2\|\}$ for all $v_1 \in V_1$ and $v_2 \in V_2$. In general, since \mathbb{K} is locally compact, any two norms on V are equivalent.

The choice of a good norm $\|\cdot\|$ on V induces a good norm on every exterior power of V (this is discussed in Quint [38]). This allows to generalize the notion of angle to the non-Archimedean setting: for $v, w \in V$, we define $\sphericalangle(v, w)$ to be the unique number in $[0, \pi]$ such that

$$\sin \sphericalangle(v, w) = \frac{\|v \wedge w\|}{\|v\| \|w\|}.$$

Observe that the angle crucially depends on the choice of the norm. Following Bochi-Potrie-S. [4] we define the angle of two subspaces $P, Q < \mathbb{K}^d$ as

$$\sphericalangle(P, Q) = \min_{v \in P^\times} \min_{w \in Q^\times} \sphericalangle(v, w),$$

where $P^\times = P \setminus \{0\}$, $Q^\times = Q \setminus \{0\}$.

The sine of the angle gives a distance, that we sometimes denote by d , on the projective space $\mathbb{P}(V)$, and more generally on every Grassmannian $\mathcal{G}_k(V)$: we set for $P, Q \in \mathcal{G}_k(V)$

$$d(P, Q) := \max_{v \in P^\times} \min_{w \in Q^\times} \sin \sphericalangle(v, w) = \min_{v \in P^\times} \max_{w \in Q^\times} \sin \sphericalangle(v, w),$$

this corresponds to the Hausdorff distance of $\mathbb{P}(P), \mathbb{P}(Q)$ regarded as subsets of $\mathbb{P}(V)$ with the aforementioned distance. Observe that

$$d(P, Q) \geq \sin \sphericalangle(P, Q)$$

and the latter inequality is, apart from very special cases, strict.

More generally we extend the distance to subspaces of possibly different dimension: for $P \in \mathcal{G}_k(V)$, $Q \in \mathcal{G}_l(V)$, $k \leq l$ we set

$$d(P, Q) := \max_{v \in P^\times} \min_{w \in Q^\times} \sin \sphericalangle(v, w) = \min_{W \in \mathcal{G}_k(Q)} d(P, W).$$

Such distance vanishes if and only if $P \subset Q$.

2.1.2. *Singular values.* Assume now that \mathbb{K} is commutative. Given a \mathbb{K} -norm on V we say that $g \in \mathrm{GL}(V, \mathbb{K})$ is a *semi-homothecy* if there exists a g -invariant \mathbb{K} -orthogonal decomposition $V = V_1 \oplus \dots \oplus V_k$ and $\sigma_1, \dots, \sigma_k \in \mathbb{R}_+$ such that for every $i \in \llbracket 1, k \rrbracket$ and every $v_i \in V_i$ one has

$$\|gv_i\| = \sigma_i \|v_i\|.$$

The numbers σ_i are called the ratios of the semi-homothecy g .

Consider a maximal abelian subgroup of diagonalizable matrices $A \subset \mathrm{GL}(V, \mathbb{K})$, let $K \subset \mathrm{GL}(V, \mathbb{K})$ be a compact subgroup such that if $N_{\mathrm{GL}}(A)$ is the normalizer of A in $\mathrm{GL}(V, \mathbb{K})$ then $N_{\mathrm{GL}}(A) = (N_{\mathrm{GL}}(A) \cap K)A$. Following Quint [37, Théorème 6.1] there exists a \mathbb{K} -norm $\|\cdot\|$ on V such that

- $\|\cdot\|$ is preserved by K ,
- A acts on $(V, \|\cdot\|)$ by semi-homothecies with respect to a common \mathbb{K} -orthogonal decomposition of V in one dimensional subspaces.

Whenever such a norm is fixed, for every $g \in \mathrm{GL}(V)$ we denote the *norm* and its *co-norm* by

$$\|g\| := \max_{v \in V^\times} \frac{\|gv\|}{\|v\|} \quad \mathbf{m}(g) = \inf_{v \in V^\times} \frac{\|gv\|}{\|v\|}.$$

Let $d = \dim V$. Keeping notation from Quint [37], we denote by $\mathbf{E} := \mathbb{R}^d$ a real vector space with a restricted root system of $\mathrm{GL}(V)$, and by

$$\mathbf{E}^+ = \{x = (x_1, \dots, x_d) \in \mathbb{R}^d \mid x_1 \geq \dots \geq x_d\}$$

a Weyl chamber of \mathbf{E} . We will denote by $\mathbf{a}_i \in \mathbf{E}^*$ the simple roots of \mathbf{E} , so that

$$\mathbf{a}_i(x) = x_i - x_{i+1} \in \mathbb{R}.$$

The choice of an ordering (e_1, \dots, e_d) of the joint eigenlines of A (the eigenlines are uniquely determined Quint [38, Lemma II.1.3]) induces a map $\nu : A \rightarrow \mathbf{E}$ given by

$$\nu(a) := (\log \sigma_1(a), \dots, \log \sigma_d(a)),$$

where $\sigma_1(a), \dots, \sigma_d(a)$ are the semi-homothecy ratios in the basis $\{e_1, \dots, e_d\}$. We set $A^+ := \nu^{-1}(\mathbf{E}^+)$, so that A^+ consists of those elements $a \in A$ whose corresponding semi-homothecy ratios satisfy $\sigma_1(a) \geq \dots \geq \sigma_d(a)$.

With respect to the basis $\{e_1, \dots, e_d\}$, when \mathbb{K} is non-Archimedean, it holds that $K = \mathrm{GL}(d, \mathcal{O})$, and the map ν extends to the Cartan projection, still denoted ν from the whole $\mathrm{GL}(V, \mathbb{K})$: indeed $\mathrm{GL}(V, \mathbb{K}) = KA^+K$, and, given $a_1, a_2 \in A^+$, the element a_1 belongs to Ka_2K if and only if $\nu(a_1) = \nu(a_2)$. In particular we can set $\nu(g) = \nu(a_g)$ for any element $a_g \in A$ such that there exist $k_g, l_g \in K$ with $g = k_g a_g l_g$ (Bruhat-Tits [8, Section 3.3]).

For every $g \in \mathrm{GL}(V, \mathbb{K})$, we choose a Cartan decomposition $g = k_g a_g l_g$ as above and define, for $p \in \llbracket 1, d-1 \rrbracket$,

$$u_p(g) = k_g \cdot e_p \in V.$$

If \mathbb{K} is Archimedean, the set $\{u_p(g) : p \in \llbracket 1, d-1 \rrbracket\}$ is an *arbitrary* orthogonal choice of axes (ordered in decreasing length) of the ellipsoid $\{Av : \|v\| = 1\}$. Note that for every v that lies in the span of $g^{-1}u_p(g)$ one has $\|gv\| = \sigma_p(g)\|v\|$. With a slight abuse of notation we will often also denote by $u_p(g)$ the corresponding point in $\mathbb{P}V$.

We furthermore denote by $U_p(g)$ the *Cartan attractor* of g :

$$U_p(g) = u_1(g) \oplus \dots \oplus u_p(g) = k_g \cdot (e_1 \oplus \dots \oplus e_p).$$

Definition 2.1. An element $g \in \mathrm{GL}(V, \mathbb{K})$ is said to have a *gap of index* p if $\sigma_p(g) > \sigma_{p+1}(g)$. In that case, if \mathbb{K} is Archimedean, the p -dimensional space $U_p(g)$ is independent of the Cartan decomposition of g .

Note that if g has a gap of index p , then the decomposition

$$U_{d-p}(g^{-1}) \oplus g^{-1}(U_p(g))$$

is orthogonal: this is clear when \mathbb{K} is Archimedean (see Remark 2.4 for the general case)

Remark 2.2. If \mathbb{K} is not Archimedean, the components k_g, l_g in the Cartan decomposition are not uniquely determined even if g has gaps of every index; in particular the spaces $U_p(g)$ always depend on the choice of the Cartan decomposition. For example take $d = 2$; if $|a| > |b|$ we have

$$\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ b/a & 1 \end{pmatrix} \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}$$

and both $\begin{pmatrix} 1 & 0 \\ b/a & 1 \end{pmatrix}$ and $\begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}$ belong to $K = \mathrm{GL}(2, \mathcal{O})$. In this example it is easy to verify that the set of possible Cartan attractors $U_1(g)$ coincides with the ball of center e_1 and radius $|b/a|$. Note that, since \mathbb{K} is non-Archimedean, any point in this ball is a center.

2.1.3. *Quantitative results.* Many of the auxiliary technical results in [4] rely on the min-max characterization of singular values of linear maps from \mathbb{R}^d to \mathbb{R}^d . This characterization in fact generalizes to any local field if one replaces the singular values with the semi-homothety ratios:

$$\sigma_p(A) = \max_{P \in \mathfrak{S}_p(V)} \mathbf{m}(A|_P) \quad \sigma_{p+1}(A) = \min_{Q \in \mathfrak{S}_{d-p}(V)} \|A|_Q\|.$$

Therefore the quantitative linear algebraic facts collected in [4, Appendix 3] carry through. We now state the ones that we will use in the following.

Lemma 2.3 ([4, Lemma A.4]). *Let $g, h \in \mathrm{GL}(V, \mathbb{K})$ have a gap of index p . Then for any possible choice of Cartan attractor $U_p(g)$ (resp. $U_p(gh)$):*

$$d(U_p(gh), U_p(g)) \leq \|h\| \|h^{-1}\| \frac{\sigma_{p+1}(g)}{\sigma_p(g)} \quad (2)$$

$$d(U_p(gh), gU_p(h)) \leq \|g\| \|g^{-1}\| \frac{\sigma_{p+1}(h)}{\sigma_p(h)}. \quad (3)$$

Remark 2.4. If \mathbb{K} is non-Archimedean, the Cartan attractors $U_p(g)$ are not uniquely defined (cfr. Remark 2.2). However it follows from Lemma 2.3 that, given two different Cartan decompositions for g , $g = k_g a_g l_g = k'_g a'_g l'_g$, and denoting $V_p = \langle e_1, \dots, e_p \rangle$, we have

$$d(k_g V_p, k'_g V_p) \leq \frac{\sigma_{p+1}(g)}{\sigma_p(g)},$$

namely all possible different choices for $U_p(g)$ are contained in a ball of radius $\frac{\sigma_{p+1}(g)}{\sigma_p(g)}$. As the distance d is, in this case, non-Archimedean, we deduce, also in this case, that any choice of $U_p(g)$ is orthogonal to $gU_{d-p}(g^{-1})$ for any other choice of $U_{d-p}(g^{-1})$.

Lemma 2.5 ([4, Lemma A.6]). *Let $g \in \mathrm{GL}(V, \mathbb{K})$ have a gap of index p . Then, for all $P \in \mathfrak{S}_p(V)$ transverse to $U_{d-p}(g^{-1})$ we have:*

$$d(g(P), U_p(g)) \leq \frac{\sigma_{p+1}(g)}{\sigma_p(g)} \frac{1}{\sin \angle(P, U_{d-p}(g^{-1}))}.$$

Lemma 2.6 ([4, Lemma A.7]). *Let $g, h \in \mathbf{GL}(V, \mathbb{K})$. Suppose that g and gh have gaps of index p . Let $\alpha := \angle(U_p(h), U_{d-p}(g^{-1}))$. Then:*

$$\begin{aligned}\sigma_p(gh) &\geq (\sin \alpha) \sigma_p(g) \sigma_p(h), \\ \sigma_{p+1}(gh) &\leq (\sin \alpha)^{-1} \sigma_{p+1}(g) \sigma_{p+1}(h).\end{aligned}$$

Given a subspace $P \in \mathcal{G}_p(V)$ we denote by P^\perp a chosen orthogonal complement of P ; this always exists, but is not unique if \mathbb{K} is non-Archimedean. Suppose that $P, W \in \mathcal{G}_p(V)$ satisfy $d(P, W) < 1$. Then $W \cap P^\perp = \{0\}$, and so there exists a unique linear map

$$L_{W,P}: P \rightarrow P^\perp \quad \text{such that} \quad W = \{v + L_{W,P}(v) : v \in P\}. \quad (4)$$

The association $L_{W,P} \mapsto W$ provides an affine chart for $\mathcal{G}_p(V)$. The next lemma states that this chart is 1-Lipschitz, and it is 4-biLipschitz on a sufficiently small neighbourhood of P :

Lemma 2.7 ([4, Lemma A.11]). *Let $P, P_1, P_2 \in \mathcal{G}_p(V)$, with $d(P_i, P) < 1$, then*

$$d(P_1, P_2) \leq \|L_{P_1,P} - L_{P_2,P}\|$$

for all choices of P^\perp . If moreover $d(P_i, P) < 1/\sqrt{2}$ then $\|L_{P_1,P} - L_{P_2,P}\| \leq 4d(P_1, P_2)$.

Proof. The proof of [4, Lemma A.11] smartly combines the triangular inequality for the distance d and the characterization

$$d(P_1, P_2) = \max_{w \in P_1^*} \min_{v \in P_2^*} \frac{\|v - w\|}{\|w\|}.$$

Since both hold when V is a vector space over a local field \mathbb{K} , the proof generalizes without modifications. In case \mathbb{K} is non-Archimedean, one could also deduce the better estimate $\|L_{P_1,P} - L_{P_2,P}\| \leq 2d(P_1, P_2)$. \square

The next lemma is a variation of [4, Lemma A.10]. In [4] there is an assumption on $d(P_i, P)$ depending on g that we replace here with the contraction assumption $d(gP_i, gP) < 1/\sqrt{2}$. Despite the proof is very similar to [4, Lemma A.10], we include it for completeness:

Lemma 2.8. *Let V be a d -dimensional \mathbb{K} -vector space, and $g \in \mathbf{GL}(V)$. Choose $P \in \mathcal{G}_p(V)$ and $Q \in \mathcal{G}_{d-p}(V)$ such that the pairs (P, Q) and (gP, gQ) are orthogonal. Then for every $P_i \in \mathcal{G}_p(V)$, ($i = 1, 2$) with $P_i \cap Q = \{0\}$ and $d(gP_i, gP) < 1/\sqrt{2}$, it holds*

$$d(gP_1, gP_2) \geq \frac{\mathbf{m}(g|_Q)}{4\|g|_P\|} d(P_1, P_2). \quad (5)$$

Proof of Lemma 2.8. Using the same notation as in (4), for each $i = 1, 2$, we consider the linear map $L_i = L_{P_i,P} : P \rightarrow P^\perp$ and $M_i = L_{gP_i, gP} : gP \rightarrow gP^\perp$; these are well defined since $P_i \cap Q = \{0\}$. Clearly the two maps are related by $L_i = (g^{-1}|_{gQ}) \circ M_i \circ (g|_P)$. As a consequence,

$$\|L_1 - L_2\| = \|(g^{-1}|_{gQ}) \circ (M_1 - M_2) \circ (g|_P)\| \leq \frac{\|g|_P\|}{\mathbf{m}(g|_Q)} \|M_1 - M_2\|.$$

Lemma 2.7 gives:

$$\|L_1 - L_2\| \geq d(P_1, P_2).$$

Since by assumption $d(gP_i, gP) < 1/\sqrt{2}$, Lemma 2.7 implies:

$$\|M_1 - M_2\| \leq 4d(gP_1, gP_2).$$

Putting these three estimates together, we get

$$d(gP_1, gP_2) \geq \frac{1}{4}\|M_1 - M_2\| \geq \frac{1}{4} \frac{\mathbf{m}(g|_Q)}{\|g|_P\|} \|L_1 - L_2\| \geq \frac{1}{4} \frac{\mathbf{m}(g|_Q)}{\|g|_P\|} d(P_1, P_2).$$

□

The following corollary of Lemma 2.8 will be useful in Section 5.2:

Corollary 2.9. *Let V be a \mathbb{K} -vector space, $W < V$ a subspace of dimension 2, and $g \in \mathrm{GL}(V)$. Denote by $\sigma_i(g|_W)$ the semi-homothecy ratios of $g : W \rightarrow gW$ where the norm on W (resp. gW) is induced by the norm on V . For every $P_i \in \mathbb{P}W$ with $P_i \cap U_{d-1}(g^{-1}) = \{0\}$, and $d(gP_i, u_1(g|_W)) < 1/\sqrt{2}$ it holds*

$$d(gP_1, gP_2) \geq \frac{\sigma_2(g|_W)}{4\|g|_W\|} d(P_1, P_2).$$

Proof. This follows directly from Lemma 2.8 once we choose $P = u_2(g^{-1}|_{gL})$, $Q = u_1(g^{-1}|_{gL})$. □

Another useful corollary of Lemma 2.8 is the following.

Corollary 2.10. *Given $\alpha > 0$, there exist positive δ and b with the following properties. Let V be a d -dimensional \mathbb{K} -vector space, and $g \in \mathrm{GL}(V)$. Suppose that $P \in \mathcal{G}_p(V)$ and $Q \in \mathcal{G}_{d-p}(V)$ satisfy*

$$\min\{\angle(P, Q), \angle(gP, gQ)\} \geq \alpha. \quad (6)$$

Then for every $P_i \in \mathcal{G}_p(V)$, ($i = 1, 2$) with $P_i \cap Q = \{0\}$ such that $d(gP_i, gP) < \delta$ one has

$$d(gP_1, gP_2) \geq b \frac{\mathbf{m}(g|_Q)}{\|g|_P\|} d(P_1, P_2). \quad (7)$$

Proof. Since all good norms are equivalent, the general case follows from Lemma 2.8 by considering two norms, one for which P and Q orthogonal and one that makes gP and gQ orthogonal, the operator norm and \mathbf{m} are to be computed using both these norms. □

Along the same lines we get a bound on how elements $g \in \mathrm{GL}(V)$ contract on open sets in Grassmannians:

Corollary 2.11. *Let $g \in \mathrm{GL}(V, \mathbb{K})$ have a gap of index p . Then, for every $\alpha > 0$ there is b such that for all $P_1, P_2 \in \mathcal{G}_p(V)$ with $\angle(P_i, g^{-1}U_{d-p}(g^{-1})) > \alpha$ we have:*

$$d(g(P_1), g(P_2)) \leq b \frac{\sigma_{p+1}}{\sigma_p}(g) d(P_1, P_2).$$

Proof. If we assume that $d(P_i, U_p(g)) \geq 1/\sqrt{2}$ the result follows readily from Lemma 2.7 by considering the linear maps $L_i := L_{P_i, U_p(g)}$ and $M_i := L_{gP_i, gU_p(g)}$. As above $L_i = (g^{-1}|_{U_{d-p}(g^{-1})}) \circ M_i \circ (g|_{U_p(g)})$. In this case the result follows as $\mathbf{m}(g|_{U_p(g)}) = \sigma_p(g)$, and $\mathbf{m}(g^{-1}|_{U_{d-p}(g^{-1})}) = 1/\sigma_{p+1}(g)$. The general statement follows by comparison of different norms. □

2.2. Dynamical background. We now turn to the dynamical preliminaries. The goal of this section is to extend the results of Bochi-Gurmelon [3] and Bochi-Potrie-S. [4] to the non-Archimedean setting.

2.2.1. Dominated splittings and Bochi-Gurmelon's Theorem. In this section we recall the definition of dominated splittings and review its connection with cone fields.

Let X be a compact metric space equipped with a continuous homeomorphism $\vartheta : X \rightarrow X$. Let V be a finite dimensional \mathbb{K} -vector space and let $\psi_0 : X \rightarrow \text{GL}(V, \mathbb{K})$ be continuous. We will denote by $\psi : X \times V \rightarrow X \times V$ the induced cocycle defined by

$$\psi_x(v) = \psi(x, v) = (\vartheta(x), \psi_0(x)v).$$

Definition 2.12. Consider a good norm $\|\cdot\|$ on V . Let $\Lambda \subset X$ be a ϑ -invariant subset, then we say that $\psi|_\Lambda$ has a *dominated splitting* if the trivial bundle $\Lambda \times V$ splits as a Whitney sum of two ψ -invariant sub-bundles $V = E \oplus F$ with the following extra condition: there exist positive μ and c such that for every n positive, $x \in \Lambda$, $u \in E_x$ and $w \in F_x$ one has

$$\frac{\|\psi_x^n u\|}{\|\psi_x^n w\|} \leq ce^{-\mu n} \frac{\|u\|}{\|w\|}.$$

In this situation we say moreover that F (resp. E) is the unstable (resp. stable) bundle and that F *dominates* E .

Note that this condition is independent of the chosen norm. The dominated splitting of $\psi|_\Lambda$ is unique provided *its index*, i.e. $\dim_{\mathbb{K}} F$, is fixed and it extends to the closure $\bar{\Lambda}$ of Λ (see Crovisier-Potrie [12, Proposition 2.2 and 2.5] whose proof works verbatim in our setting). Furthermore:

Proposition 2.13 ([12]). *Suppose a linear flow ψ has dominated splittings $E^1 \oplus F^1$ and $E^2 \oplus F^2$ of index $p_1 \leq p_2$. Then $E^2 \subseteq E^1$ and $F^1 \subseteq F^2$.*

In the case when $\mathbb{K} = \mathbb{R}$ Bochi-Gourmelon [3, Theorem A] gave the following criterion for dominated splittings to exist; their the proof generalizes to every local field \mathbb{K} , as Oseledets theorem holds in this generality:

Theorem 2.14 (Bochi-Gourmelon [3]). *Let X be a compact metric space, V a \mathbb{K} -vector space and $\psi : X \times V \rightarrow X \times V$ a linear cocycle. Then the linear flow ψ has a dominated splitting $E \oplus F$ with $\dim F = p$ if and only if there exist $c > 0$, $\mu > 0$ such that for every $x \in X$ and $n \geq 0$ we have*

$$\frac{\sigma_{p+1}}{\sigma_p}(\psi_x^n) < ce^{-\mu n}.$$

Moreover, the bundles² are given by:

$$F_x = \lim_{n \rightarrow +\infty} U_p(\psi_{\vartheta^{-n}(x)}^n) \text{ and } E_x = \lim_{n \rightarrow +\infty} U_{d-p}(\psi_{\vartheta^n(x)}^{-n}),$$

and these limits are uniform.

²For completeness, let us note that the space U_p associated to an operator from a vector space equipped with a good norm to itself, can be defined for an operator between two vector spaces both equipped with good norms.

Proof. Bochi-Gourmelon's proof is based on the one hand on some angle estimates building upon the min-max characterization of singular values of matrices in $\mathrm{GL}_d(\mathbb{R})$, and on the other hand on the multiplicative ergodic theorem (Oseledets theorem). The former hold verbatim in the general local field setting once the singular values are replaced by the semi-homothecy ratios as defined in Section 2.1.2, the required multiplicative ergodic theorem was established (following Oseledets original proof) by Margulis [34, Theorem V.2.1], the integrability of ψ follows from its continuity and the compactness of the base X . With these ingredients at hand, the sketch of the proof explained in [4, Section A.4] applies verbatim. \square

The existence of a dominated splitting can be furthermore characterized in terms of cone fields; this will be crucial to prove openness of Anosov representations in Section 3.1 (note that the non-Archimedean case has not yet been established). Given a decomposition $V = V_1 \oplus V_2$ and a positive a , then the subset defined by

$$\{v \in V : a\|v_1\| \geq \|v_2\|\}$$

is called a a -cone (of dimension $\dim V_1$) on V .

A cone field on $\Lambda \subset X$ is a continuous choice $x \mapsto \mathcal{C}_{a(x),x}$ of a $a(x)$ -cone on V (of fixed dimension) for each $x \in \Lambda$. Cone fields can be used to characterize dominated splittings.

Proposition 2.15 (See Sambarino [40, Proposition 2.2]). *Let $\Lambda \subset X$ be ϑ -invariant. The cocycle $\psi|_\Lambda$ has a dominated splitting of index i if and only if there exists a map $a : \Lambda \rightarrow \mathbb{R}^+$ bounded away from 0 and ∞ , a cone field $\mathcal{C}_{a(x),x}$ on Λ of dimension i , a number $0 < \lambda < 1$ and a positive integer n_0 such that for every $x \in \Lambda$ the closure of $\psi_x^{n_0}(\mathcal{C}_{a(x),x})$ is contained in $\mathcal{C}_{\lambda a(\vartheta^{n_0}(x)),\vartheta^{n_0}(x)}$.*

2.2.2. *Dominated sequences.* Bochi-Potrie-S. [4, Section 2] applied Bochi-Gourmelon's Theorem 2.14 to the compact space of *dominated sequences of matrices*, and got useful implications on the relative position of the axes of the ellipsoid associated to the products of such sequences: we recall now the relevant definitions and results from [4] where these were first established.

Given $C > 1$, define the following compact set:

$$\mathcal{D}(C) := \{g \in \mathrm{GL}(V, \mathbb{K}), \|g\| \leq C, \|g^{-1}\| \leq C\}.$$

If I is a (possibly infinite) interval in \mathbb{Z} , the set $\mathcal{D}(C)^I$ is endowed with the product topology, turning it into a compact metric space.

Let $p \in \llbracket 1, d-1 \rrbracket$, $\mu > 0$, $c > 0$. For each interval $I \subset \mathbb{Z}$, we denote by $\mathcal{D}(C, p, c, \mu, I)$ the set of sequences of matrices $(g_n) \in \mathcal{D}(C)^I$ such that for all $m, n \in I$ with $m \geq n$ we have

$$\frac{\sigma_{p+1}}{\sigma_p}(g_m \cdots g_{n+1} g_n) \leq ce^{-\mu(m-n+1)}.$$

Definition 2.16. An element of $\mathrm{GL}(V, \mathbb{K})^I$ is a *dominated sequence* if it belongs to $\mathcal{D}(C, p, c, \mu, I)$ for some C, p, c , and μ .

Consider the map $\mathrm{shift} : \mathcal{D}(C)^{\mathbb{Z}} \rightarrow \mathcal{D}(C)^{\mathbb{Z}}$ defined by

$$\mathrm{shift}((g_n)_{-\infty}^{\infty}) = (g_{n+1})_{-\infty}^{\infty}$$

and let $\psi_0 : \mathcal{D}(C)^{\mathbb{Z}} \rightarrow \mathrm{GL}(V, \mathbb{K})$ be $\psi_0((g_n)) = g_0$. The subsets $\mathcal{D}(C, p, c, \mu, \mathbb{Z})$ are shift-invariant and automatically verify the hypothesis of Theorem 2.14.

Proposition 2.17 (Bochi-Potrie-S. [4, Proposition 2.4]). *For each sequence $x = (g_n) \in \mathcal{D}(C, p, \mu, c, \mathbb{Z})$, the limits:*

$$F_x := \lim_{n \rightarrow +\infty} U_p(g_{-1}g_{-2} \cdots g_{-n}),$$

$$E_x := \lim_{n \rightarrow +\infty} U_{d-p}(g_0^{-1} \cdots g_{n-2}^{-1}g_{n-1}^{-1}),$$

exist and are uniform over $\mathcal{D}(C, p, \mu, c, \mathbb{Z})$. Moreover, F dominates E and $E \oplus F$ is a dominated splitting for the linear cocycle over the shift defined above.

By a compactness argument, the proposition above ensures transversality for Cartan attractors and repellers computed in finite, but sufficiently long, sequences of matrices:

Lemma 2.18 (Bochi-Potrie-S. [4, Lemma 2.5]). *Given $C > 1$, $\mu > 0$, and $c > 0$, there exist $L \in \mathbb{N}$ and $\delta > 0$ with the following properties. Suppose that $I \subset \mathbb{Z}$ is an interval and $\{g_i\}_{i \in I}$ is an element of $\mathcal{D}(C, p, c, \mu, I)$. If $n < k < m$ all belong to I and $\min\{k - n, m - k\} > L$ then:*

$$\angle(U_p(g_{k-1} \cdots g_{n+1}g_n), U_{d-p}(g_k^{-1}g_{k+1}^{-1} \cdots g_{m-1}^{-1})) > \delta.$$

2.3. Hyperbolic groups. The last source of preliminaries comes from geometric group theory. Here we recall basic facts about hyperbolic groups and cone types.

Let Γ be a finitely generated group. We fix a finite symmetric generating set S and denote by $||$ the associated word length: for $\gamma \in \Gamma - \{e\}$ we denote by $|\gamma|$ the least number of elements of S needed to write γ as a word on S , and define the induced distance $d_\Gamma(\gamma, \eta) = |\gamma^{-1}\eta|$. A geodesic segment on Γ is a sequence $\{\alpha_i\}_0^k$ of elements in Γ such that $d_\Gamma(\alpha_i, \alpha_j) = |i - j|$.

In the paper we will be only interested in word-hyperbolic groups, namely such that the metric space $(\Gamma, ||)$ is Gromov hyperbolic. Following the footprints of [4], our analysis will be based on the study of cone types, and natural objects associated to them.

2.3.1. Cone types. In the paper we follow Cannon's original definition of cone types, which is more convenient for our geometric purposes, but the reader should be warned that the definition used in [4] is slightly different

Definition 2.19. The *cone type* of $\gamma \in \Gamma$ is defined by


$$\mathcal{C}(\gamma) = \{\eta \in \Gamma : |\gamma\eta| = |\eta| + |\gamma|\}.$$

Notice that if $\eta \in \mathcal{C}(\gamma)$ then

$$d_\Gamma(\gamma^{-1}, \eta) = |\gamma\eta| = |\eta| + |\gamma| = |\eta| + |\gamma^{-1}| = d_\Gamma(e, \eta) + d_\Gamma(e, \gamma^{-1}),$$

i.e. there exists a geodesic segment through e with endpoints γ^{-1} and η . Reciprocally, the endpoint of a geodesic segment starting at γ^{-1} and passing through e necessarily belongs to $\mathcal{C}(\gamma)$.

A fundamental result of Cannon is that, provided Γ is hyperbolic, there are only finitely many cone types (see for example Bridson-Heafliger [7, p. 455] or Coornaert-Delzant-Papadopoulos [11, p. 145]).


 FIGURE 1. The cone type of $\gamma \in \Gamma$

2.3.2. *The geodesic automaton.* See Bridson-Haefliger [7, p. 456].

Given a cone type \mathcal{C} and $a \in S \cap \mathcal{C}$, one easily checks that for every $\gamma \in \Gamma$ with $\mathcal{C}(\gamma) = \mathcal{C}$ one gets

$$a\mathcal{C}(\gamma a) \subset \mathcal{C}(\gamma).$$

Furthermore it is easy to verify that in such case the cone type $\mathcal{C}(\gamma a)$ doesn't depend on γ (see for example [11, Lemma 4.3]), and, with a slight abuse of notation we will denote such cone type $a \cdot \mathcal{C}$.

The *geodesic automaton* of Γ (this also depends on S) is the labelled graph \mathcal{G} defined as follows:

- the vertices are the cone types of Γ ;
- there is an edge $\mathcal{C}_1 \xrightarrow{a} \mathcal{C}_2$ from vertex \mathcal{C}_1 to vertex \mathcal{C}_2 , labelled by a generator $a \in S$, iff $a \in \mathcal{C}_1$ and $\mathcal{C}_2 = a \cdot \mathcal{C}_1$.

Since Γ is hyperbolic there are only finitely many cone types and thus the geodesic automaton has a finite number of vertices.

Let us explain the relation with geodesics. Consider a *geodesic segment* $(\gamma_0, \gamma_1, \dots, \gamma_\ell)$ that is, a sequence of elements of Γ such that $d(\gamma_n, \gamma_m) = |n - m|$, and assume that $\gamma_0 = \text{id}$. Then, there are $a_0, \dots, a_{\ell-1}$ in a generating set S such that $\gamma_n = a_0 a_1 \cdots a_{n-1}$. Note that for each n , the following is an edge of the geodesic automaton graph \mathcal{G} :

$$\mathcal{C}(\gamma_n) \xrightarrow{a_n} \mathcal{C}(\gamma_{n+1}).$$

Thus we obtain a (finite) walk on \mathcal{G} starting from the vertex $\mathcal{C}(\text{id})$. Conversely, for each such walk we may associate a geodesic segment starting at the identity.

Let us define also the *recurrent geodesic automaton* as the maximal recurrent subgraph \mathcal{G}^* of \mathcal{G} ; its vertices are called *recurrent cone types*.

Let Λ_Γ be the subset of all bi-infinite labelled sequences of \mathcal{G}^* . It is a closed shift-invariant subset of $(\mathcal{G}^*)^{\mathbb{Z}}$ and the induced dynamical system $\text{shift} : \Lambda_\Gamma \rightarrow \Lambda_\Gamma$ is a sofic shift (as in Lind-Marcus [33]).

The following concept will be useful in Section 5.

Definition 2.20. Given an integer k we say that two cone types $\mathcal{C}_1, \mathcal{C}_2$ are *k-nested* if there is a path of length k in the geodesic automaton from \mathcal{C}_1 to \mathcal{C}_2 . In this case there is an element $\beta \in \Gamma$ with $|\beta| = k$ and such that $\beta\mathcal{C}_2 \subset \mathcal{C}_1$.

Since Γ is hyperbolic, there are only finitely many cone types, therefore, for every k , there are only finitely many k -nested pairs of elements (however, as soon as Γ is non-elementary, the number of k -nested pairs grows exponentially with k). The following is clear from the definitions:

Lemma 2.21. *If $\{\alpha_i\} \subset \Gamma$ is a geodesic, then the pair $(\mathcal{C}(\alpha_i), \mathcal{C}(\alpha_{i+k}))$ is k -nested.*

2.3.3. *Coverings of the Gromov boundary.* Recall that, as Γ is Gromov hyperbolic, its boundary $\partial\Gamma$, consisting of equivalence classes of geodesic rays, is well defined up to homeomorphism. We associate to every cone type \mathcal{C} which is not the cone type of the identity a subset of $\partial\Gamma$, the *cone type at infinity*, by considering limit points of geodesic rays starting on e and totally contained in \mathcal{C} :

$$\mathcal{C}_\infty = \{[(\alpha_i)] | (\alpha_i) \text{ geodesic ray}, \alpha_0 = e, \alpha_i \in \mathcal{C}\}.$$

It follows from the discussion in the previous paragraph that every point in $\partial\Gamma$ is contained in at least one of the sets \mathcal{C}_∞ . As there are only finitely many cone types, we obtain a finite covering of $\partial\Gamma$ by considering $\mathcal{U} = \{\mathcal{C}_\infty(\gamma)\}$. Starting from this covering we will construct new coverings that will serve as our Sullivan shadows:




FIGURE 2. The set $\gamma \cdot \mathcal{C}_\infty(\gamma)$

Lemma 2.22. *Given $T > 0$, the family of open sets*

$$\mathcal{U}_T := \{\gamma \mathcal{C}_\infty(\gamma) : |\gamma| \geq T\}$$

defines an open covering of $\partial\Gamma$.

Proof. We have to check that every point $x \in \partial\Gamma$ is covered, but this is evident since considering a geodesic ray $(\alpha_i)_0^\infty$ in Γ starting from e converging to x , one has that for all i , $x \in \alpha_i \mathcal{C}_\infty(\alpha_i)$, see Figure 2. \square

3. ANOSOV REPRESENTATIONS

Anosov representations from fundamental groups of negatively curved closed manifolds to $\mathrm{PGL}(d, \mathbb{R})$ were introduced by Labourie [31] and generalized by Guichard-W. [23] to any hyperbolic group. In this section we will generalize to non-Archimedean local fields the work of [4], which provides a simplified definition³.

3.1. Anosov representations and dominated splittings. Let Γ be a discrete group of finite type, fix a finite symmetric generating set S_Γ and denote by $||$ the associated word length.

³Morse actions on Euclidean buildings (and thus in particular Anosov subgroups of $\mathrm{PGL}_d(\mathbb{K})$ when \mathbb{K} is non-Archimedean) were already defined by Kapovich-Leeb-Porti [27, Definition 5.35], the interest of such concept was also suggested in G eritaud-Guichard-Kassel-W. [20, Remark 1.6 (a)].

Definition 3.1. Consider $p \in \llbracket 1, d-1 \rrbracket$. A representation $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ is $\{\mathbf{a}_p\}$ -Anosov⁴ if there exist positive constants c, μ such that for all $\gamma \in \Gamma$ one has

$$\frac{\sigma_{p+1}(\rho(\gamma))}{\sigma_p(\rho(\gamma))} \leq ce^{-\mu|\gamma|}. \quad (8)$$

A $\{\mathbf{a}_1\}$ -Anosov representations will be called *projective Anosov*.

One has the following direct remark.

Remark 3.2 (Bochi-Potrie-S.). Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be $\{\mathbf{a}_p\}$ -Anosov. Given a geodesic $\{\alpha_i\}_{i \in \mathbb{Z}}$ let us denote by $\bar{\alpha}_i = \alpha_{i+1}^{-1}\alpha_i \in S_\Gamma$, then we have

$$(\rho(\bar{\alpha}_i))_{i \in \mathbb{Z}} \in \mathcal{D}(C, p, c, \mu, \mathbb{Z})$$

where c, μ come from equation (8) and $C = \max\{\|\rho(a)\| : a \in S_\Gamma\}$. Note also that

$$(\rho(\bar{\alpha}_i))_{i \in \mathbb{Z}} \in \mathcal{D}(C, d-p, c, \mu, \mathbb{Z}),$$

and thus Theorem 2.14 provides the following splittings of \mathbb{K}^d

$$E_{(\rho(\bar{\alpha}_i))}^p \oplus F_{(\rho(\bar{\alpha}_i))}^{d-p} \quad \text{and} \quad E_{(\rho(\bar{\alpha}_i))}^{d-p} \oplus F_{(\rho(\bar{\alpha}_i))}^p,$$

with the obvious inclusions according to dimension. By domination, these four bundles vary continuously⁵ in $\mathcal{D}(C, p, c, \mu, \mathbb{Z}) \cap \mathcal{D}(C, d-p, c, \mu, \mathbb{Z})$. Finally, Proposition 2.17 yields, for $k \in \{p, d-p\}$ and $m \geq 0$

$$U_k(\rho(\alpha_m)) = U_k(\bar{\alpha}_0^{-1} \cdots \bar{\alpha}_m^{-1}) \rightarrow E_{(\rho(\bar{\alpha}_i))}^k$$

and

$$U_k(\rho(\alpha_{-m})) = U_k(\bar{\alpha}_{-1} \cdots \bar{\alpha}_{-m}) \rightarrow F_{(\rho(\bar{\alpha}_i))}^k$$

as $m \rightarrow \infty$.

Using dominated splittings it is possible to deduce strong angle estimates between Cartan attractors along geodesic rays through the origin; for example the next result is a direct consequence of Lemma 2.18.

Proposition 3.3 (Bochi-Potrie-S.). *Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be a $\{\mathbf{a}_p\}$ -Anosov representation. Then there exists $\delta > 0$ and $L \in \mathbb{N}$ such that for every geodesic segment $(\alpha_i)_0^k$ in Γ through e with $|\alpha_0|, |\alpha_k| \geq L$ one has*

$$\angle \left(U_p(\rho(\alpha_k)), U_{d-p}(\rho(\alpha_0)) \right) > \delta.$$

Bochi-Potrie-S. [4] applied the theory of dominated splittings to the sofic shift $\Lambda_\Gamma \rightarrow \Lambda_\Gamma$ induced by the recurrent geodesic automaton (see Section 2.3.2), to get an easy proof of openness of Anosov representations. Their proof easily extends to every local field:

Proposition 3.4. *The set of $\{\mathbf{a}_p\}$ -Anosov representations is open in $\mathrm{hom}(\Gamma, \mathrm{PGL}_d(\mathbb{K}))$.*

Proof. A representation $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ induces a linear cocycle A_ρ over Λ_Γ , which admits a dominated splitting if and only if the representation ρ is $\{\mathbf{a}_p\}$ -Anosov. Observe that the cocycle A_ρ varies continuously with the representation, since it only depends on the value of ρ on a generating set of Γ . Since, by Proposition 2.15, having a dominated splitting is an open condition on the space of cocycles, the result follows. \square

⁴In the language of Bochi-Potrie-S. [4, Section 3.1] a $\{\mathbf{a}_p\}$ -Anosov representation is called *p-dominated*.

⁵This follows from Proposition 2.15, see also, for example, [4, Theorem A.15].

3.2. Boundary maps. From now on we will assume that Γ is a word hyperbolic group. This is not a restriction: Kapovich-Leeb-Porti proved that the only groups admitting Anosov representations are hyperbolic [27, Theorem 6.15] (cfr. also Bochi-Potrie-S. [4] for a different proof in the Archimedean case). We can thus talk freely about the Gromov boundary $\partial\Gamma$.

An important property of $\{\mathfrak{a}_p\}$ -Anosov representations is that they admit equivariant boundary maps:

Proposition 3.5 (Bochi-Potrie-S. [4, Proposition 4.9]). *If $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ is $\{\mathfrak{a}_p\}$ -Anosov, then for any geodesic ray $\{\gamma_n\}$ with endpoint x , the limits*

$$\xi_\rho^p(x) := \lim_{n \rightarrow \infty} U_p(\rho(\gamma_n)) \quad \xi_\rho^{d-p}(x) := \lim_{n \rightarrow \infty} U_{d-p}(\rho(\gamma_n))$$

exist and do not depend on the ray; they define continuous ρ -equivariant transverse maps $\xi^p : \partial\Gamma \rightarrow \mathcal{G}_p(\mathbb{K}^d)$, $\xi^{d-p} : \partial\Gamma \rightarrow \mathcal{G}_{d-p}(\mathbb{K}^d)$.

Proof. The proof in [4, Proposition 4.9] works without modification in our context: despite $U_p(\rho(\gamma_n))$ is not uniquely defined, Lemma 2.3 (2) guarantees that, for every choice of $U_p(\rho(\gamma_n))$, the sequence $\{U_p(\rho(\gamma_n))\}$ is Cauchy, and therefore has a limit; furthermore, since any pair of geodesic rays defining x is at bounded distance, Lemma 2.3 (2) shows that the limit doesn't depend on the chosen sequence, and the maps are continuous. The equivariance follows from Lemma 2.3 (3). \square

The uniformity of the limits in Proposition 3.5 can be quantified explicitly (cfr. [4, Lemma 4.7]). This will be useful in the proof of Theorem 7.1:

Lemma 3.6. *Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be $\{\mathfrak{a}_p\}$ -Anosov. Then there exist constants C, μ such that, for every $\alpha \in \Gamma$ and every $x \in \alpha\mathcal{C}_\infty(\alpha)$,*

$$d(\xi^p(x), U_p(\rho(\alpha))) \leq C e^{-\mu|\alpha|}.$$

In particular, given $\varepsilon > 0$ there exists $L \in \mathbb{N}$ such that

$$\bigcup_{\gamma: |\gamma| \geq L} \overline{U_p(\rho(\gamma))} \subset \mathcal{N}_\varepsilon(\xi_\rho^p(\partial\Gamma)).$$

Proof. If $x \in \alpha\mathcal{C}_\infty(\alpha)$ there exists a geodesic ray $(\alpha_i)_{i \in \mathbb{N}}$ through α with endpoint x . In particular we get

$$d(U_p(\rho(\alpha)), \xi_\rho^p(x)) \leq \sum_{i \geq |\alpha|} d(U_p(\rho(\alpha_i)), U_p(\rho(\alpha_{i+1}))) \leq c \sum_{i \geq |\alpha|} e^{-\mu i}.$$

Here the first inequality is a consequence of the triangular inequality, the second follows from Lemma 2.3 (2).

The second statement follows since, as Γ is word hyperbolic, there is a constant D such that, for every $\gamma \in \Gamma$ we can choose a geodesic ray $(\alpha_i)_{i \in \mathbb{N}}$, and $k \in \mathbb{N}$ such that $d(\gamma, \alpha_k) \leq D$. This implies that $\gamma = \alpha_k h$ with $|h| \leq D$. Let x be the endpoint of the geodesic $(\alpha_i)_{i \in \mathbb{N}}$. Then

$$d(U_p(\rho(\gamma)), \xi_\rho^p(x)) \leq d(U_p(\rho(\alpha_k h)), U_p(\rho(\alpha_k))) + d(U_p(\rho(\alpha_k)), \xi_\rho^p(x)).$$

\square

Bochi-Potrie-S. [4] observed that the boundary map has an explicit characterization in term of the linear cocycle A_ρ over the sofic shift Λ_Γ (described in the proof of Proposition 3.4). Recall from Definition 2.12 that whenever a cocycle

$A_\rho : \Lambda_\Gamma \times \mathbb{K}^d \rightarrow \Lambda_\Gamma \times \mathbb{K}^d$ has a dominated splitting we denote by E (resp. F) the stable (resp. unstable) bundle.

Proposition 3.7 ([4, Proposition 5.2]). *Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be $\{\mathbf{a}_p\}$ -Anosov. Let $x, y \in \partial\Gamma$ and $(\bar{\alpha}_i)_{-\infty}^\infty \in \Lambda_\Gamma$ a geodesic from y to x . Then one has*

$$\xi^p(x) = E_{(\rho(\bar{\alpha}_i))}^p \quad \text{and} \quad \xi^{d-p}(y) = F_{(\rho(\bar{\alpha}_i))}^{d-p}.$$

As a corollary we can follow [4] and generalize to the non-Archimedean case the following important fact originally proved by Labourie [31] and Guichard-W. [23].

Corollary 3.8. *Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be $\{\mathbf{a}_p\}$ -Anosov. The boundary maps $\xi^p : \partial\Gamma \rightarrow \mathcal{G}_p(\mathbb{K}^d)$, $\xi^{d-p} : \partial\Gamma \rightarrow \mathcal{G}_{d-p}(\mathbb{K}^d)$ vary continuously with the representation.*

Proof. This follows at once from the arguments in the proof of Propositions 3.4 and 3.7, as splittings vary continuously with the cocycles by Proposition 2.15 (this is a standard argument, see [4, Theorem A.15] for a proof). \square

The boundary map, which is unique, gives a realization of the boundary $\partial\Gamma$ in $\mathcal{G}_p(\mathbb{K}^d)$, a space where the dynamics is governed by ratios of semi-homothecy ratios of elements in the projective linear group. To stress this fact and the dependence on ρ we introduce the following notation, which will be heavily used in the rest of the paper: if $\rho : \Gamma \rightarrow \mathrm{GL}_d(\mathbb{K})$ is a $\{\mathbf{a}_p\}$ -Anosov representation with equivariant boundary map ξ^p , and $x \in \partial\Gamma$, we will write

$$x_\rho^p := \xi^p(x).$$

We noticed that this notation improves readability of many formulas and conveniently stresses the dependence of ξ on ρ .

3.3. Geometric estimates. We conclude the section on Anosov representations by collecting a number of geometric lemmas that will be useful later on. The first result provides the quantification we will need of the following geometric principle: endpoints of a geodesics through the origin are uniformly far in the visual boundary, the same holds for their image under the boundary map associated to an Anosov representation.

Lemma 3.9. *Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be $\{\mathbf{a}_p\}$ -Anosov representation. Then there exists $\nu > 0$, depending only on ρ such that: if $\{\alpha_i\}_{i \in \mathbb{Z}} \subset \Gamma$ is a geodesic through id with endpoints $x, z \in \partial\Gamma$, then for all $i \in \mathbb{Z}$ one has*

$$\angle(\rho(\alpha_i^{-1})x_\rho^p, \rho(\alpha_i^{-1})z_\rho^{d-p}) > \nu.$$

Proof. Recall that we denote by $\mathcal{D}(C, d-p, c, \mu, \mathbb{Z})$ the compact, shift invariant space of dominated sequences (cfr. Definition 2.16). The bundle $\mathcal{D}(C, d-p, c, \mu, \mathbb{Z}) \times \mathbb{K}^d$ admits a dominated splitting $E^p \oplus F^{d-p}$ and, by compactness, we get

$$\nu = \inf_{\{g_i\} \in \mathcal{D}(C, d-p, c, \mu, \mathbb{Z})} \angle(E_{(g_i)}^p, F_{(g_i)}^{d-p}) > 0.$$

Remark 3.2 implies that, since ρ is $\{\mathbf{a}_p\}$ -Anosov, $(\rho(\alpha_{i+1}^{-1}\alpha_i))_{i \in \mathbb{Z}} \in \mathcal{D}(C, d-p, c, \mu, \mathbb{Z})$, furthermore one directly computes that

$$\psi^n((\rho(\alpha_{i+1}^{-1}\alpha_i))_{i \in \mathbb{Z}}, v) = ((\rho(\alpha_{i+1}^{-1}\alpha_i))_{i-n \in \mathbb{Z}}, \rho(\alpha_n^{-1})v).$$

As we know from Proposition 2.17 that

$$x_\rho^p = \lim_{i \rightarrow \infty} U_p(\rho(\alpha_i)) = E_{(\rho(\bar{\alpha}_i))}^p$$

and

$$z_\rho^{d-p} = \lim_{i \rightarrow \infty} U_{d-p}(\rho(\alpha_{-i})) = F_{(\rho(\bar{\alpha})_i)}^{d-p},$$

we deduce that for all $i \in \mathbb{Z}$ one has

$$\angle(\rho(\alpha_i^{-1})x_\rho^p, \rho(\alpha_i^{-1})z_\rho^{d-p}) > \nu.$$

□

The next lemma will be crucial in Section 6. It quantifies how the inverse of elements in a geodesic expand the distances exponentially in neighbourhoods of their Cartan attractors; this should be compared with [4, Corollary A.14]:

Lemma 3.10. *Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be $\{\mathbf{a}_p\}$ -Anosov. There exist positive constants $\bar{c}, \bar{\mu}, \delta$ depending only on ρ , and $L \in \mathbb{N}$ such that, for every geodesic ray $\{\alpha_i\}_{i \in \mathbb{N}} \subset \Gamma$, with $\alpha_0 = \mathrm{id}$ and endpoint x , every $i \geq L$, and every $z, w \in \partial\Gamma$ satisfying $z_\rho^p, w_\rho^p \in B_\delta(x_\rho^p)$, and $\rho(\alpha_i^{-1})\{z_\rho^p, w_\rho^p\} \subset B_\delta(\rho(\alpha_i^{-1})x_\rho^p)$, we have*

$$d(\rho(\alpha_i^{-1})w_\rho^p, \rho(\alpha_i^{-1})z_\rho^p) \geq \bar{c}e^{\bar{\mu}i}d(w_\rho^p, z_\rho^p).$$

Proof. We complete the ray $\{\alpha_i\}_{i \in \mathbb{N}}$ to a biinfinite geodesic $\{\alpha_i\}_{i \in \mathbb{Z}}$ with second endpoint y . The sequence $s = \{\rho(\alpha_{i+1}^{-1}\alpha_i)\}_{i \in \mathbb{Z}}$ belongs to $\mathcal{D}(C, p, c, \mu, \mathbb{Z})$. It follows from Propositions 2.17 and 3.5 that the sequence s has the dominated splitting $E \oplus F$ where $F_s = x_\rho^p$ and $E_s = y_\rho^{d-p}$. So there exist constants $\bar{\mu}, c_1$ such that

$$\frac{\mathbf{m}(\rho(\alpha_i^{-1})|_{y_\rho^{d-p}})}{\|\rho(\alpha_i^{-1})|_{x_\rho^p}\|} \geq c_1 e^{\bar{\mu}i}.$$

Since, by Lemma 3.9, the angles $\angle(x_\rho^p, y_\rho^{d-p})$ and $\angle(\rho(\alpha_i^{-1})x_\rho^p, \rho(\alpha_i^{-1})y_\rho^{d-p})$ are bounded below by a uniform constant ν , we can apply Corollary 2.10 with $P = x_\rho^p$, $Q = y_\rho^{d-p}$ and $g = \rho(\alpha_i)^{-1}$ and get

$$d(\rho(\alpha_i^{-1})w_\rho^p, \rho(\alpha_i^{-1})z_\rho^p) \geq b \frac{\mathbf{m}(\rho(\alpha_i^{-1})|_{y_\rho^{d-p}})}{\|\rho(\alpha_i^{-1})|_{x_\rho^p}\|} d(w_\rho^p, z_\rho^p).$$

□

4. AN UPPER BOUND ON THE HAUSDORFF DIMENSION OF THE LIMIT SET

In this section we will prove the following upper bound, this result is independently obtained by Glorieux-Monclair-Tholozan [19] for Archimedean \mathbb{K} . Recall from the introduction that if $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ is projective Anosov then $h_\rho^{\mathbf{a}_1}$ is the critical exponent of the Dirchlet series

$$s \mapsto \sum_{\gamma \in \Gamma} \left(\frac{\sigma_2}{\sigma_1}(\rho(\gamma)) \right)^s.$$

We denote by $\mathrm{Hff}(A)$ the Hausdorff dimension of a subset $A \subset \mathbb{P}(\mathbb{K}^d)$ for the metric induced by a good norm on \mathbb{K}^d .

Proposition 4.1. *Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be $\{\mathbf{a}_1\}$ -Anosov, then $\mathrm{Hff}(\xi^1(\partial\Gamma)) \leq h_\rho^{\mathbf{a}_1}$.*

Recall that for a metric space (Λ, d) and for $s > 0$ its s -capacity is defined as

$$\mathcal{H}^s(\Lambda) = \inf_{\varepsilon} \left\{ \sum_{U \in \mathcal{U}} \mathrm{diam} U^s : \mathcal{U} \text{ is a covering of } \Lambda \text{ with } \sup_{U \in \mathcal{U}} \mathrm{diam} U < \varepsilon \right\} \quad (9)$$

and that

$$\text{Hff}(\Lambda) = \inf\{s : \mathcal{H}^s(\Lambda) = 0\} = \sup\{s : \mathcal{H}^s(\Lambda) = \infty\}. \quad (10)$$

In order to prove Proposition 4.1 we will analyze the image, under the boundary map ξ^1 of the covering \mathcal{U}_T described in subsection 2.3.3, whose elements consist of images of cone types at infinity under sufficiently big group elements. The following crucial lemma will allow us to show that the images of the boundaries of cone types transform as expected under group elements:

Lemma 4.2. *Let $\rho : \Gamma \rightarrow \text{PGL}_d(\mathbb{K})$ be a projective Anosov representation. Then there exist $\delta > 0$, $L \in \mathbb{N}$ such that for all $\gamma \in \Gamma$ with $|\gamma| > L$ and every $x \in \mathcal{C}_\infty(\gamma)$ one has*

$$d(x_\rho^1, U_{d-1}(\rho(\gamma^{-1}))) > \delta.$$

Proof. By definition of $\mathcal{C}(\gamma)$, for all $x \in \mathcal{C}_\infty(\gamma)$ there exists a geodesic ray $\{\alpha_i\}_0^\infty$ in Γ with $\alpha_0 = \gamma^{-1}$ and $\alpha_i \rightarrow x$ as $i \rightarrow \infty$. The lemma then follows combining Proposition 3.3 and Proposition 3.5. \square

4.1. Proof of Proposition 4.1. For each $T > 0$ consider the covering \mathcal{U}_T of $\partial\Gamma$ given by Lemma 2.22. By definition, $U = U_\gamma \in \mathcal{U}_T$ is of the form $\gamma\mathcal{C}_\infty(\gamma)$ for some $\gamma \in \Gamma$ with $|\gamma| \geq T$.

Lemma 4.2 implies that there exists δ such that for every $x \in \mathcal{C}_\infty(\gamma)$ one has

$$d(x_\rho^1, U_{d-1}(\rho(\gamma^{-1}))) \geq \delta$$

and thus Lemma 2.5 applied to $\rho(\gamma)$ implies that

$$d(\rho(\gamma)x_\rho^1, U_1(\rho(\gamma))) \leq \frac{1}{\delta} \frac{\sigma_2}{\sigma_1}(\rho(\gamma))$$

which implies that

$$\text{diam} \xi(U_\gamma) \leq \frac{2}{\delta} \frac{\sigma_2}{\sigma_1}(\rho(\gamma)) \leq Ce^{-\mu T}.$$

In particular $\sup_{U \in \mathcal{U}_T} \text{diam} U$ is arbitrarily small as $T \rightarrow \infty$. Hence,

$$\mathcal{H}^s(\xi^1(\partial\Gamma)) \leq \inf_T \sum_{U \in \mathcal{U}_T} (\text{diam} \xi^1(U))^s \leq \left(\frac{2}{\delta}\right)^s \inf_T \sum_{\gamma: |\gamma| \geq T} \left(\frac{\sigma_2}{\sigma_1}(\rho(\gamma))\right)^s.$$

By definition, if $s > h_\rho^{\text{a}_1}$ then the Dirichlet series $\Phi_\rho^{\text{a}_1}(s)$ is convergent (recall eq. (1)). Hence, for every $s > h_\rho^{\text{a}_1}$ one has

$$\lim_{T \rightarrow \infty} \sum_{\gamma: |\gamma| \geq T} \left(\frac{\sigma_2}{\sigma_1}(\rho(\gamma))\right)^s = 0,$$

which implies that the s -capacity $\mathcal{H}^s(\xi^1(\partial\Gamma))$ vanishes and thus $\text{Hff}(\xi^1(\partial\Gamma)) \leq h_\rho^{\text{a}_1}$.

5. LOCAL CONFORMALITY AND HAUSDORFF DIMENSION

The goal of this section is to find a class of representations for which the equality in Proposition 4.1 holds. This happens in three steps.

- In Section 5.1 we study the *thickened cone types* $X_\infty(\alpha)$, these are a thickening, in $\mathbb{P}(\mathbb{K}^d)$, of the image by ξ^1 of $\mathcal{C}_\infty(\alpha)$ for a given α , and define the *locally conformal points*.

- In Section 5.2 we prove that if x is a locally conformal point, then there is a geodesic ray $\alpha_i \rightarrow x$ such that the sets $\rho(\alpha_i)X_\infty(\alpha_i)$ behave coarsely like balls around x ; the harder inequality is the lower containment, which is achieved in Corollary 5.10.
- In Section 5.3 we define a measure that behaves like an Ahlfors regular measure for the sets $\rho X_\infty(\alpha)$. Putting this together with the previous section, arguments coming from Sullivan's original paper allow us to conclude the desired equality, provided we can guarantee existence of many locally conformal points, this is the purpose of Section 5.4.

5.1. Thickened cone types at infinity and locally conformal points. Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be projective Anosov, it follows from Proposition 3.3 that there is a positive lower bound on the distance of Cartan attractors and repellers of geodesic rays through the origin. Such number will play an important role in our study:

Definition 5.1. Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be projective Anosov, and let L be fixed and big enough. The *least angle* δ_ρ is

$$\delta_\rho = \inf \sin \left(\angle \left(U_1(\rho(\alpha_k)), U_{d-1}(\rho(\alpha_{-m})) \right) \right)$$

where $(\alpha_i)_{i \in \mathbb{Z}}$ ranges among biinfinite geodesics through the origin, and $k, m > L$.

We consider coverings of $\xi(\partial\Gamma)$ obtained by translating *thickened cone types at infinity*:

$$X_\infty(\alpha) := \mathcal{N}_{\delta_\rho/2}(\xi^1(\mathcal{C}_\infty(\alpha))) \cap \xi^1(\partial\Gamma).$$

By construction the sets $X_\infty(\alpha)$ are coarsely balls of $\xi^1(\partial\Gamma)$ centered at points in $\xi^1(\mathcal{C}_\infty(\alpha))$:

Remark 5.2. For every α in Γ , and every $x \in \mathcal{C}_\infty(\alpha)$, the thickened cone type at infinity $X_\infty(\alpha)$ contains a ball centered at the point x_ρ^1 of uniform radius:

$$B(x_\rho^1, \delta_\rho/2) \cap \xi^1(\partial\Gamma) \subset X_\infty(\alpha)$$

Thanks to Proposition 3.3 we can control how thickened cone types shrink under the action of group elements:

Lemma 5.3. *Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be projective Anosov. Then there exist K, L such that, for every geodesic ray $(\alpha_i)_{i=0}^\infty$, for every $i > L$, and every $z_\rho^1, w_\rho^1 \in X_\infty(\alpha_i)$,*

$$d(\alpha_i z_\rho^1, \alpha_i w_\rho^1) \leq K \frac{\sigma_2}{\sigma_1}(\rho(\alpha_i)) d(z_\rho^1, w_\rho^1).$$

Proof. As ρ is projective Anosov we have $d(z_\rho^1, U_{d-1}(\rho(\alpha_i)^{-1})) > \delta_\rho/2$ (Lemma 4.2). The result is then a direct consequence of Corollary 2.11. \square

Corollary 5.4. *If $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ is projective Anosov, and $x \in \alpha_i \mathcal{C}_\infty(\alpha_i)$, then*

$$\rho(\alpha_i)X_\infty(\alpha_i) \subset B \left(x_\rho^1, K \frac{\sigma_2}{\sigma_1}(\rho(\alpha_i)) \right) \cap \xi(\partial\Gamma).$$

In particular, if $\{\alpha_i\}_1^\infty$ is a geodesic ray with endpoint x , the sets $\rho(\alpha)X_\infty(\alpha)$ form a fundamental system of open neighbourhoods of x in $\xi(\partial\Gamma)$ (cfr. Figure 3).

Definition 5.5 gives conditions guaranteeing that the sets $\rho(\alpha)X_\infty(\alpha)$ are coarsely balls whose sizes we can precisely estimate. Given $g \in \mathrm{GL}_d(\mathbb{K})$ we denote by

$$1 \leq p_1(g) < \dots < p_{k(g)}(g) < d$$




FIGURE 3. The sets of the form $\rho(\alpha_i)X_\infty(\alpha_i)$ for a geodesic ray $\{\alpha_i\}$ with endpoint x are the intersections of thinner and thinner ellipses with the limit curve.

the indices of the gaps of g (as in Definition 2.1).

Definition 5.5. Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be projective Anosov. We say that $x \in \partial\Gamma$ is a (ε, L) -locally conformal point for ρ if there exists a geodesic ray $\{\alpha_i\}_0^\infty$ in Γ based at the identity and with endpoint x such that the following conditions hold:

- (i) for all big enough i one has $p_2(\alpha_i) = p_2$ does not depend on i ,
- (ii) for every $i > L$, and for every $z \in (\xi_\rho^1)^{-1}(X_\infty(\alpha_i))$ one has

$$\sin \left(\angle \left(z_\rho^1 \oplus \rho(\alpha_i^{-1})x_\rho^1, U_{d-p_2}(\rho(\alpha_i^{-1})) \right) \right) > \varepsilon.$$




FIGURE 4. The second condition in Definition 5.5.

Note that, in general, the index p_2 might depend on the point x and we do not require that the representation ρ is $\{\mathbf{a}_{p_2}\}$ -Anosov. In the special case when ρ is $\{\mathbf{a}_1, \mathbf{a}_2\}$ -Anosov, the condition (i) is automatically satisfied with $p_2 = 2$, but (ii) can only hold if the dimension of $\partial\Gamma$ is very small (cfr. Corollary 6.6).

Remark 5.6. A generic element $g \in \mathrm{PGL}(V)$ has $p_2(g) = 2$. Nevertheless there are many interesting geometric situations in which condition (i) holds for $p_2 > 2$. For example if g is a generic element in $\mathrm{SO}(m, n)$, we have that $\Lambda^m g \in \mathrm{SL}(V)$ has $p_2(\Lambda^m g) = n - m + 1$, so one can enforce $p_2 > 2$ by considering representations in smaller subgroups. In Section 8.1 we will describe another interesting class of examples.

5.2. Neighborhoods of locally conformal points that are coarsely balls.

We will now show that if x is a locally conformal point for ρ , and $\alpha_i \rightarrow x$ is a geodesic ray, then the sets $\rho(\alpha_i)X_\infty(\alpha_i)$ are coarsely balls centered at x_ρ^1 of radius $\sigma_2/\sigma_1(\rho(\alpha_i))$ for the distance on $\xi_\rho^1(\partial\Gamma)$ induced by d , this will be achieved in Corollary 5.10, and motivated the terminology *locally conformal*.

Proposition 5.7. *Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be projective Anosov. There exist \bar{L} such that, for every (ε, L) -locally conformal point x , there exists a geodesic ray $\{\alpha_i\}_0^\infty$ from the identity with endpoint x , such that for every $i > \bar{L}$ and every $z \in X_\infty(\alpha_i)$ it holds*

$$\frac{\varepsilon}{4} \frac{\sigma_2}{\sigma_1}(\rho(\alpha_i)) \cdot d(z_\rho^1, (\alpha_i^{-1}x)_\rho^1) \leq d((\alpha_i z)_\rho^1, x_\rho^1).$$

Proof. Let $W_i := z_\rho^1 \oplus (\alpha_i^{-1}x)_\rho^1$. As x is (ε, L) -locally conformal, for every $i > L$, we have $d(W_i, U_{d-p_2}(\rho(\alpha_i^{-1}))) > \varepsilon$. From Lemma 2.5 one concludes that

$$\not\prec(\rho(\alpha_i)W_i, U_{p_2}(\rho(\alpha_i))) \rightarrow 0$$

as $i \rightarrow \infty$ at a speed only depending on ε and the Anosov constants of ρ , and thus, possibly increasing L , one concludes that for every $i > L$ it holds

$$\frac{\sigma_2}{\sigma_1}(\rho(\alpha_i)|_{W_i}) \geq \varepsilon \frac{\sigma_{p_2}}{\sigma_1}(\rho(\alpha_i)) = \varepsilon \frac{\sigma_2}{\sigma_1}(\rho(\alpha_i)).$$

Here the last equality is due to the fact that p_2 is the first gap for $\rho(\alpha_i)$ and thus $\sigma_{p_2}(\rho(\alpha_i)) = \sigma_2(\rho(\alpha_i))$.

Furthermore $\rho(\alpha_i^{-1})u_2(\rho(\alpha_i)|_{W_i}) \in W_i \cap U_{d-1}(\rho(\alpha_i^{-1}))$, and then, since ρ is projective Anosov and $z_\rho^1 \in X_\infty(\alpha)$, we have $d(z_\rho^1, \rho(\alpha_i^{-1})u_2(\rho(\alpha_i)|_{W_i})) > \delta_\rho/2$, where δ_ρ is the constant from Definition 5.1. This implies that we can find \bar{L} depending on ρ and ε only such that for every $i > \bar{L}$

$$d(\rho(\alpha_i)z_\rho^1, U_1(\rho(\alpha_i)|_{W_i})) < d(\rho(\alpha_i)z_\rho^1, U_1(\rho(\alpha_i))) + d(U_1(\rho(\alpha_i)), U_1(\rho(\alpha_i)|_{W_i})) < 1/\sqrt{2},$$

since both quantities converge to 0 as $i \rightarrow \infty$ at a speed only depending on the Anosov constants of ρ . The proposition then follows from Corollary 2.9. \square

Recall from Definition 2.20 that we say that a pair of cone types $(\mathcal{C}(\alpha_1), \mathcal{C}(\alpha_2))$ of Γ are k -nested if there exists a path in the geodesic automaton of length k between $\mathcal{C}(\alpha_1)$ and $\mathcal{C}(\alpha_2)$. In this case we say that $\beta \in \Gamma$ is a *nesting word* if β labels one such path.

Lemma 5.8. *For every \bar{L} big enough (depending only on ρ) there exists a constant c (depending on ρ and \bar{L}) such that for every \bar{L} -nested pair $(\mathcal{C}(\alpha_1), \mathcal{C}(\alpha_2))$ and any nesting word β it holds*

- (i) $\rho(\beta)X_\infty(\alpha_2) \subset X_\infty(\alpha_1)$
- (ii) for every $z_\rho^1 \in \xi(\beta\mathcal{C}_\infty(\alpha_2))$ and every $w_\rho^1 \in X_\infty(\alpha_1) \setminus \rho(\beta)X_\infty(\alpha_2)$, it holds

$$d(z_\rho^1, w_\rho^1) > c.$$

Proof.

- (i) By definition of δ_ρ and $X_\infty(\alpha_2)$, whenever $|\beta| \geq L$ and β is a nesting word, then $d(x, U_{d-1}(\rho(\beta^{-1}))) \geq \delta_\rho/2$ for every point x in $X_\infty(\alpha_2)$. Here L is as in Definition 5.1. Up to possibly enlarging L we can assume, by Corollary 2.11, that $\rho(\beta)$ contracts distances on $X_\infty(\alpha_2)$ so that

$$\rho(\beta)X_\infty(\alpha_2) \subseteq \mathcal{N}_{\delta_\rho/2}(\rho(\beta)\xi_\rho^1(\mathcal{C}_\infty(\alpha_2))) \cap \xi(\partial\Gamma) \subseteq X_\infty(\alpha_1).$$

- (ii) Since, by construction, $X_\infty(\alpha_2)$ contains the intersection of $\xi_\rho^1(\partial\Gamma)$ with a ball around any point $z_\rho^1 \in \xi_\rho^1(\mathcal{C}_\infty(\alpha_2))$ of radius $\delta_\rho/2$, the set $\rho(\beta)X_\infty(\alpha_2)$ contains the intersection of $\xi_\rho^1(\partial\Gamma)$ with the ball around any point $z_\rho^1 \in \rho(\beta)\xi_\rho^1(\mathcal{C}_\infty(\alpha_2))$ of radius $\frac{\delta_\rho}{2} \frac{\sigma_4}{\sigma_1}(\rho(\beta)) : \frac{\sigma_4}{\sigma_1}(g)$ is the smallest contraction for

the action of $g \in \mathrm{SL}(d, \mathbb{K})$ on $\mathbb{P}(\mathbb{K}^d)$. Recall that only finitely many β can occur, as, by construction, $|\beta| = \bar{L}$. The result follows taking

$$c = \min_{|\beta|=\bar{L}} \frac{\delta_\rho}{2} \frac{\sigma_d}{\sigma_1}(\rho(\beta))$$

□

Combining Proposition 5.7 and Lemma 5.8 we obtain:

Proposition 5.9. *There exists c_1 depending only on ρ such that, if L is as in Lemma 5.8 and $\{\alpha_i\} \subset \Gamma$ is a geodesic ray with endpoint x , for every y with $y_\rho^1 \in \rho(\alpha_n)X_\infty(\alpha_n) \setminus \rho(\alpha_{n+L})X_\infty(\alpha_{n+L})$, it holds*

$$d(y_\rho^1, x_\rho^1) \geq c_1 \frac{\sigma_2}{\sigma_1}(\rho(\alpha_{n+L})).$$

Proof. It follows from Lemma 2.21 that for every n, L the pair $(\mathcal{C}_\infty(\alpha_n), \mathcal{C}_\infty(\alpha_{n+L}))$ is L -nested. Furthermore, up to choosing L large enough, we can apply Lemma 5.8 to the pair $(\mathcal{C}_\infty(\alpha_n), \mathcal{C}_\infty(\alpha_{n+L}))$. If we denote by $z := \alpha_n^{-1}x$ and $w := \alpha_n^{-1}y$ we deduce that $d(z_\rho^1, w_\rho^1) > c$. Proposition 5.7 implies then that

$$d(y_\rho^1, x_\rho^1) \geq \frac{c\varepsilon}{4} \frac{\sigma_2}{\sigma_1}(\alpha_n) \geq c_1 \frac{\sigma_2}{\sigma_1}(\alpha_{n+L})$$

Where in the last inequality we used that, as L is fixed the homothety ratio gap of α_n is uniformly comparable to the one of α_{n+L} . □

As a corollary of Proposition 5.7 we can finally get the main result of the section (cfr. Figure 3):

Corollary 5.10. *Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be a projective Anosov; then for every locally conformal point $x \in \partial\Gamma$ there exists a geodesic ray $\alpha_i \rightarrow x$ with*

$$B\left(x_\rho^1, c_1 \frac{\sigma_2}{\sigma_1}(\rho(\alpha_i))\right) \cap \xi(\partial\Gamma) \subset \rho(\alpha_i)X_\infty(\alpha_i).$$

Proof. This follows from the above proposition by observing that the sets $\rho(\alpha_i)X_\infty(\alpha_i)$ form a fundamental system of neighborhoods of x_ρ^1 in $\xi(\partial\Gamma)$. □

5.3. A regular measure for conformal points. The goal of this section is to construct, following Patterson's original idea, a measure, supported on $\xi_\rho^1(\partial\Gamma)$, for which we can get good estimates on the measure of the cone types. This will be used in Section 5.4 to obtain the desired lower bound on the Hausdorff dimension of the limit set⁶.

Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be a projective Anosov representation. Recall from the introduction that we have defined

$$\Phi_\rho^{\mathfrak{a}_1}(s) = \sum_{\gamma \in \Gamma} \left(\frac{\sigma_2}{\sigma_1}(\rho(\gamma)) \right)^s.$$

We can assume that $\Phi_\rho^{\mathfrak{a}_1}(h_\rho^{\mathfrak{a}_1}) = \infty$: otherwise, as it is standard in Patterson-Sullivan theory, we would carry out the same construction with the aid of the modified Poincaré series

$$\Phi_\rho^{\mathfrak{a}_1}(s) = \sum_{\gamma \in \Gamma} f\left(\mathfrak{a}_1(\rho(\gamma))\right) \left(\frac{\sigma_2}{\sigma_1}(\rho(\gamma)) \right)^{h_\rho^{\mathfrak{a}_1}},$$

⁶See Remark 5.15 for a comparison with the work of Quint [39]

where $f(s)$ is the function constructed (for example) in Quint [39, Lemma 8.5].

We will therefore assume from now on that the Poincaré series diverges at its critical exponent; for every $s > h_\rho^{\text{a}_1}$, we define

$$\mu_\rho^s = \frac{1}{\Phi_\rho^{\text{a}_1}(s)} \sum_{\gamma \in \Gamma} \left(\frac{\sigma_2}{\sigma_1}(\rho(\gamma)) \right)^s \delta_{U_1(\rho(\gamma))}.$$

Recall from Section 2.1 that, for every element $\gamma \in \Gamma$ we chose a Cartan decomposition of $\rho(\gamma)$ and therefore a 1-dimensional subspace $U_1(\rho(\gamma))$.

One easily checks that for every $s > h_\rho^{\text{a}_1}$ the functional $f \mapsto \int f d\mu_\rho^s$ is continuous on $C(\mathbb{P}(\mathbb{K}^d), \mathbb{R})$ with the uniform topology and hence one can take a weak* accumulation point of μ_ρ^s , as $s \rightarrow h_\rho^{\text{a}_1}$, in the space of Radon probability measures on $\mathbb{P}(\mathbb{K}^d)$. We will denote such Radon measure by $\mu_\rho^{\text{a}_1}$, (note that we do not show, nor require, that $\mu_\rho^{\text{a}_1}$ is the only accumulation point of μ_ρ^s).

Lemma 5.11. *For any $\eta \in \Gamma$ the (signed) measure*

$$\varepsilon(\eta, s) := \eta_* \mu_\rho^s - \frac{1}{\Phi_\rho^{\text{a}_1}(s)} \sum_{\gamma \in \Gamma} \left(\frac{\sigma_2}{\sigma_1}(\rho(\gamma)) \right)^s \delta_{U_1(\rho(\eta\gamma))}$$

weakly converges to zero as $s \rightarrow h_\rho^{\text{a}_1}$.*

Proof. Indeed by definition

$$\eta_* \mu_\rho^s = \frac{1}{\Phi_\rho^{\text{a}_1}(s)} \sum_{\gamma \in \Gamma} \left(\frac{\sigma_2}{\sigma_1}(\rho(\gamma)) \right)^s \delta_{\rho(\eta)U_1(\rho(\gamma))}.$$

Furthermore, Lemma 2.3 (3) implies that

$$d(\rho(\eta)U_1(\rho(\gamma)), U_1(\rho(\eta\gamma))) \leq \|\eta\| \|\eta^{-1}\| \frac{\sigma_2}{\sigma_1}(\rho(\gamma)).$$

In order to show that $\varepsilon(\eta, s)$ converges to zero, it is enough to show that for every continuous function $f : \mathbb{P}(V) \rightarrow \mathbb{R}$ the integral of f on $\varepsilon(\eta, s)$ tends to zero as s converges to $h_\rho^{\text{a}_1}$. However every such function f is uniformly continuous, and therefore for every ε we can find δ such that $|f(x) - f(y)| < \varepsilon/2$ if $d(x, y) < \delta$. It is then enough to choose s close enough to $h_\rho^{\text{a}_1}$ so that the mass of μ_ρ^s of the elements γ such that $\sigma_1/\sigma_2(\gamma) < \|\eta\| \|\eta^{-1}\|/\delta$ is smaller than $\frac{\varepsilon}{2\|f\|}$. \square

One has the following proposition (compare with Sullivan's shadow Lemma [42]).

Proposition 5.12. *Let $\rho : \Gamma \rightarrow \text{PGL}_d(\mathbb{K})$ be a projective Anosov representation, then for all $\eta \in \Gamma$ one has*

$$\left(\frac{\sigma_d}{\sigma_1}(\rho(\eta)) \right)^{h_\rho^{\text{a}_1}} \leq \frac{\mu_\rho^{\text{a}_1}(\rho(\eta)X_\infty(\eta))}{\mu_\rho^{\text{a}_1}(X_\infty(\eta))} \leq \frac{4}{\delta_\rho^2} \left(\frac{\sigma_2}{\sigma_1}(\rho(\eta)) \right)^{h_\rho^{\text{a}_1}}.$$

Recall that there are finitely many cone types, so the number $\mu_\rho^{\text{a}_1}(X_\infty(\eta))$ is an irrelevant constant.

Proof. Consider $s > h_\rho^\sigma$, $\eta \in \Gamma$ and a continuous function $f : \mathbb{P}(\mathbb{K}^d) \rightarrow \mathbb{R}$. One has

$$\begin{aligned} \mu_\rho^s(f \circ \rho(\eta)^{-1}) &= \frac{1}{\Phi_\rho^{\mathfrak{a}_1}(s)} \sum_{\gamma \in \Gamma} \left(\frac{\sigma_2}{\sigma_1}(\rho(\gamma)) \right)^s f(\rho(\eta^{-1})U_1(\rho(\gamma))) \\ &= \varepsilon(\eta^{-1}, s)(f) + \frac{1}{\Phi_\rho^{\mathfrak{a}_1}(s)} \sum_{\gamma \in \Gamma} \left(\frac{\sigma_2}{\sigma_1}(\rho(\eta\gamma)) \right)^s f(U_1(\rho(\gamma))) \quad (11) \\ &= \varepsilon(\eta^{-1}, s)(f) + \frac{1}{\Phi_\rho^{\mathfrak{a}_1}(s)} \sum_{\gamma \in \Gamma} \left(\frac{\sigma_2}{\sigma_1}(\rho(\eta\gamma)) \frac{\sigma_1}{\sigma_2}(\rho(\gamma)) \right)^s \left(\frac{\sigma_2}{\sigma_1}(\rho(\gamma)) \right)^s f(U_1(\rho(\gamma))), \end{aligned}$$

where $\varepsilon(\eta^{-1}, s)$ is the term estimated in Lemma 5.11, so that $\varepsilon(\eta^{-1}, s)(f)$ converges to zero when $s \rightarrow h_\rho^{\mathfrak{a}_1}$.

Assume that the support of f contains $X_\infty(\eta)$ in its interior and s is close enough to $h_\rho^{\mathfrak{a}_1}$ so that $\Phi_\rho^{\mathfrak{a}_1}(s)$ is arbitrary large. Then only the tail of the sum involved in $\mu_\rho^s(f \circ \rho(\eta)^{-1})$ is relevant, this is to say:

- only γ 's for which $|\gamma|$ is large matter,
- since we are integrating f , $U_1(\rho(\gamma))$ has to be near to $X_\infty(\eta)$, so that there is a geodesic segment from η^{-1} to γ passing through the identity.

This, together with Proposition 3.3, implies that such γ 's one has

$$\sin(\angle(U_1(\rho(\gamma)), U_{d-1}(\rho(\eta)))) > \varepsilon_f,$$

for some ε_f depending on the support of f . Note that ε_f approaches $\delta_\rho/2$ (recall Definition 5.1) as $\text{supp } f \rightarrow X_\infty(\eta)$. Choosing a sequence $s_k \rightarrow h_\rho^{\mathfrak{a}_1}$ such that $\mu_\rho^{s_k} \rightarrow \mu_\rho^{\mathfrak{a}_1}$ one has, using Lemma 2.6 and equation (11), that, for any such f

$$\mu_\rho^{\mathfrak{a}_1}(f \circ \rho(\eta)^{-1}) = \lim_{s_k \rightarrow h_\rho^{\mathfrak{a}_1}} \mu_\rho^{s_k}(f \circ \rho(\eta)^{-1}) \leq \frac{4}{\delta_\rho^2} \left(\frac{\sigma_2}{\sigma_1}(\rho(\eta)) \right)^{h_\rho^{\mathfrak{a}_1}} \mu_\rho^{\mathfrak{a}_1}(f).$$

By continuity of $f \mapsto \mu_\rho^{\mathfrak{a}_1}(f)$, one concludes the desired upper bound. The lower bound follows similarly. \square

Since cone types shrink to any given point of $\partial\Gamma$ one has the following consequences of Proposition 5.12.

Corollary 5.13. *The measure $\mu_\rho^{\mathfrak{a}_1}$ has total support and no atoms.*

Proof. If α_i is a geodesic ray converging to x then $\rho(\alpha_i)X_\infty(\alpha_i)$ is a family of open neighborhoods decreasing to x , and since ρ is projective Anosov one has $(\sigma_2/\sigma_1)(\rho(\alpha_i)) \rightarrow 0$ as $i \rightarrow \infty$. As $\mu_\rho^{\mathfrak{a}_1}$ is a Radon measure we have on the one hand

$$\mu_\rho^{\mathfrak{a}_1}(\{x\}) = \inf\{\mu_\rho^{\mathfrak{a}_1}(\rho(\alpha_i)X_\infty(\alpha_i))\} \leq \frac{4}{\delta_\rho^2} \left(\frac{\sigma_2}{\sigma_1}(\rho(\eta)) \right)^{h_\rho^{\mathfrak{a}_1}},$$

on the other hand for every open set A intersecting $\xi(\partial\Gamma)$ we can find α such that $\rho(\alpha)X_\infty(\alpha)$ is contained in A , and thus

$$\mu_\rho^{\mathfrak{a}_1}(A) \geq \mu_\rho^{\mathfrak{a}_1}(\rho(\alpha)X_\infty(\alpha)) \geq \left(\frac{\sigma_d}{\sigma_1}(\rho(\eta)) \right)^{h_\rho^{\mathfrak{a}_1}}.$$

\square

5.4. **When conformal points are abundant.** Denote by

$$\text{LC}(\rho) = \{x \in \partial\Gamma : x \text{ is locally conformal for } \rho\}.$$

We can now prove the following.

Theorem 5.14. *Let $\rho : \Gamma \rightarrow \text{PGL}_d(\mathbb{K})$ be a projective Anosov representation. If $\mu_\rho^{\text{a1}}(\text{LC}(\rho)) > 0$, then*

$$\text{Hff}(\xi_\rho^1(\partial\Gamma)) = h_\rho^{\text{a1}}.$$

Proof. As we already established in Proposition 4.1, $\text{Hff}(\xi_\rho^1(\partial\Gamma)) \leq h_\rho^{\text{a1}}$, so we only need to show the reverse inequality. The proof will follow the main ideas in Sullivan's original work [42], using Corollary 5.10 and Proposition 5.12 as key replacement for the conformality of a Kleinian group action on its boundary, and Sullivan's shadow lemma.

Given $x \in \text{LC}(\rho)$ and a geodesic ray $\{\alpha_i\}$ on Γ converging to x , Corollary 5.10 implies that for all $i \geq N_0(x)$ the set $\rho(\alpha_i)X_\infty(\alpha_i)$ is coarsely (with constants independent of x) a ball of radius

$$r_i(x) = \frac{\sigma_2}{\sigma_1}(\rho(\alpha_i))$$

about x_ρ^1 (for the induced metric on $\xi_\rho^1(\partial\Gamma)$).

Proposition 5.12 then states that for all $i \geq N_0(x)$

$$\mu_\rho^{\text{a1}}(B(x, r_i(x))) \leq cr_i(x)h_\rho^{\text{a1}}. \quad (12)$$

Observe that we can extend equation (12) for any $0 < r \leq r_{N_0}(x)$, up to possibly worsening the constant c : Since ρ is projective Anosov, the word length of $\gamma \in \Gamma$ is coarsely $\log \sigma_2/\sigma_1(\rho(\gamma))$, thus

$$r_i(x)/r_{i+1}(x) \leq K_\rho$$

for some constant K_ρ only depending on ρ ; given r it suffices to consider $r_{i+1}(x) \leq r \leq r_i(x)$ and thus

$$\mu_\rho^{\text{a1}}(B(x, r)) \leq c \left(\frac{r_i(x)}{r_{i+1}(x)} \right)^{h_\rho^{\text{a1}}} r_{i+1}(x)h_\rho^{\text{a1}} \leq L_\rho r^{h_\rho^{\text{a1}}}.$$

Furthermore, there exists ε such that the set $X_\varepsilon = \{x \in \text{LC}(\rho) : r_{N_0}(x) \geq \varepsilon\}$ has positive μ_ρ^{a1} -mass: this follows from the general fact that countable union of sets with measure 0 has measure 0, since we assumed $\mu_\rho^{\text{a1}}(\text{LC}(\rho)) > 0$,

The remainder arguments are verbatim as in Haïssinsky [24, Théorème F.4]. We include them for completeness: as X_ε is a subset of $\xi(\partial\Gamma)$, it is enough to verify that $\text{Hff}(\xi(X_\varepsilon)) \geq h_\rho^{\text{a1}}$; we will show that, denoting by $\sigma := h_\rho^{\text{a1}}$, we have $\mathcal{H}^\sigma(X_\varepsilon) > 0$. Indeed let us denote by $d := \frac{\mu_\rho^{\text{a1}}(X_\varepsilon)}{2L_\rho}$. By definition of σ -capacity we can find an open covering $\mathcal{B} = \{B(x_i, r_i)\}$ of X_ε consisting of balls of radius $r_i < \varepsilon$ and such that

$$\sum r_i^\sigma \leq \mathcal{H}^\sigma(X_\varepsilon) + d.$$

Recall from (9) at the beginning of Section 4 that we denote by $\mathcal{H}^\sigma(X_\varepsilon)$ the σ -capacity of the set X_ε . On the other hand we have

$$\mu_\rho^{\text{a1}}(X_\varepsilon) \leq \mu_\rho^{\text{a1}}\left(\bigcup_i B(x_i, r_i)\right) \leq \sum_i \mu_\rho^{\text{a1}}(B(x_i, r_i)) \leq \sum_i L_\rho r_i^\sigma.$$

This shows that $\mathcal{H}^\sigma(X_\varepsilon)$ is positive, and concludes the proof. \square

Remark 5.15. Patterson-Sullivan measures in a setup close to ours were extensively studied by Quint [39]. For our geometric applications it is crucial to have an Ahlfors regular measure of exponent $h_\rho^{\mathfrak{a}_1}$. Let us denote by G the Zariski closure of $\rho(\Gamma)$, assume that G is reductive (despite this is not always the case in the examples we have in mind), and let \mathcal{F}_G denote the full flag space associated to G . Quint [39, Theorem 8.4] provides a quasi-invariant measure μ on \mathcal{F}_G called a $(\rho(\Gamma), h_\rho^{\mathfrak{a}_1} \mathfrak{a}_1)$ -Patterson-Sullivan, with the desired transformation rule, as long as a technical condition is satisfied, namely that the form $h_\rho^{\mathfrak{a}_1} \mathfrak{a}_1$ is tangent to the growth indicator function $\psi_{\rho(\Gamma)}$. In order to guarantee that this is the case we would have to further assume that the representation ρ is $\{\mathfrak{a}_p\}$ -Anosov, and that $p_2(\rho(\gamma)) = p$ for every $\gamma \in \Gamma$. The measure μ could then be pushed forward via the projection $\mathcal{F}_G \rightarrow \mathbb{P}(\mathbb{K}^d)$ and the fact that ρ is $\{\mathfrak{a}_1, \mathfrak{a}_p\}$ -Anosov would imply that the new measure on $\mathbb{P}(\mathbb{K}^d)$ would still be quasi-invariant. However deducing the analogue of Proposition 5.12 in that setting would require some work as our representations are, in most interesting cases, not Zariski dense.

6. (p, q, r) -HYPERCONVEXITY

In this section we introduce (p, q, r) -hyperconvex representations, establish geometric properties and provide the link with local conformality.

6.1. Hyperconvex representations. The following definition is inspired from Labourie [31] for surface groups. Let Γ be a word-hyperbolic group and denote by

$$\partial^{(3)}\Gamma = \{(x, y, z) \in (\partial\Gamma)^3 : \text{pairwise distinct}\}.$$

Definition 6.1. Consider $p, q, r \in \llbracket 1, d-1 \rrbracket$ such that $p + q \leq d$. We say that a representation $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ is (p, q, r) -hyperconvex if it is $\{\mathfrak{a}_p, \mathfrak{a}_q, \mathfrak{a}_r\}$ -Anosov and for every triple $(x, y, z) \in \partial^{(3)}\Gamma$ one has

$$(x_\rho^p \oplus y_\rho^q) \cap z_\rho^{d-r} = \{0\}.$$

Note that, since $p + q < d$ and the representation is $\{\mathfrak{a}_p, \mathfrak{a}_q\}$ -Anosov, the sum $x_\rho^p + y_\rho^q$ is necessarily direct. Hence, hyperconvexity implies that $p + q \leq r$. We will observe in Corollary 6.6 that ρ can only be (p, q, r) -hyperconvex if $r - p - q \geq \dim(\partial\Gamma) - 1$. Note that we do not require p and q to be different.

Proposition 6.2. *The space of (p, q, r) -hyperconvex representations is open in*

$$\mathrm{hom}(\Gamma, \mathrm{PGL}_d(\mathbb{K})).$$

Proof. The proof follows the same lines as Labourie [31, Proposition 8.2]. Since the action of Γ on $\partial^{(3)}\Gamma$ is properly discontinuous and co-compact, given a triple $(x, y, z) \in \partial^{(3)}\Gamma$ there exists $\gamma \in \Gamma$ such that the the points $\gamma x, \gamma y$ and γz are pairwise far apart. Considering a (p, q, r) -hyperconvex representation ρ , one concludes that the angles between any pair of the spaces $(\gamma x)_\rho^p, (\gamma y)_\rho^q$ and $(\gamma z)_\rho^{d-r}$ are bounded away from zero. Corollary 3.8 states that the Anosov condition is open and that equivariant maps vary continuously with the representation, hence, since the map $\partial^{(2)}\Gamma \rightarrow \mathcal{G}_{p+q}(\mathbb{K}^d)$

$$(a, b) \mapsto a_\rho^p \oplus b_\rho^q$$

is continuous away from the diagonal the result follows. \square

Since hyperconvexity is an open property, one can provide interesting examples of hyperconvex representations by looking at representations of the form $\Gamma \rightarrow G \rightarrow \mathrm{GL}_d(\mathbb{K})$, where the first arrow is convex co-compact (see Section 7.2) furthermore hyperconvexity behaves well with field extensions:

Lemma 6.3. *Let $\mathbb{K} \subset \mathbb{F}$ be a field extension, if $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ is (p, q, r) -hyperconvex then so is $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{F})$.*

We conclude the subsection providing obstructions to the existence of $(1, 1, r)$ -hyperconvex representations. A useful tool for this is the stereographic projection:

Definition 6.4. Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be $\{\mathbf{a}_1, \mathbf{a}_r\}$ -Anosov. Given $z \in \partial\Gamma$, the *stereographic projection* defined by z (and ρ) is the continuous map

$$\pi_{z,\rho} : \partial\Gamma - \{z\} \rightarrow \mathbb{P}(\mathbb{K}^d/z_\rho^{d-r})$$

defined as follows: since ρ is $\{\mathbf{a}_1\}$ -Anosov, for every point $x \in \partial\Gamma$ different from z , the vector space $x_\rho^1 \oplus z_\rho^{d-r}$ has dimension $d - r + 1$ and projects to a line in the quotient space $\mathbb{K}^d/z_\rho^{d-r}$; we define $\pi_{z,\rho}(x) \in \mathbb{P}(\mathbb{K}^d/z_\rho^{d-r})$ to be the projectivisation of this line.

The following is immediate from the definitions:

Lemma 6.5. *If the representation ρ is $(1, 1, r)$ -hyperconvex then for every $z \in \partial\Gamma$ the map $\pi_{z,\rho}$ is continuous and injective.*

Proof. The stereographic projection $\pi_{z,\rho}$ is the composition of the boundary map $\xi^1 : \partial\Gamma \setminus \{z\} \rightarrow \mathbb{P}(\mathbb{K}^d)$ with the projection $\mathbb{P}(\mathbb{K}^d \setminus z_\rho^{d-r}) \rightarrow \mathbb{P}(\mathbb{K}^d/z_\rho^{d-r})$, which is algebraic outside the $(d-r)$ -dimensional subspace z_ρ^{d-r} ; it is well defined as ρ is \mathbf{a}_1 -Anosov, and is therefore continuous. Injectivity follows directly from the definition of hyperconvexity. \square

Corollary 6.6. *If there is no continuous injective map $\partial\Gamma - \{z\} \rightarrow \mathbb{P}(\mathbb{K}^r)$, then there is no $(1, 1, r)$ -hyperconvex representation $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$.*

6.2. From hyperconvexity to local conformality. We now find a link between hyperconvexity and local conformality. The following statement is the main technical result of Section 6, and will be crucial in the proof of Theorem 7.1.

Recall from Section 5.1 that we defined, for every projective Anosov representation $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$, the thickened cone type at infinity $X_\infty(\alpha)$ as the intersection of the $\delta_\rho/2$ -neighbourhood of $\xi_\rho^1(\mathcal{C}_\infty(\alpha))$ with the image of the boundary map. In a similar way, if ρ is $\{\mathbf{a}_p\}$ -Anosov, we set

$$X_\infty^p(\alpha) := \mathcal{N}_{\delta_{p,\rho}/2} \xi_\rho^p(\mathcal{C}_\infty(\alpha)) \cap \xi_\rho^p(\partial\Gamma),$$

where $\delta_{p,\rho}$ is the number δ from Proposition 3.3.

Proposition 6.7. *Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be (p, q, r) -hyperconvex. Then there exist constants L, ε such that for every $\alpha \in \Gamma$ with $|\alpha| > L$, for every $x \in \mathcal{C}_\infty(\alpha)$ and every $y \in (\xi_\rho^q)^{-1} X_\infty^q(\alpha)$, it holds*

$$\sin \angle \left(x_\rho^p \oplus y_\rho^q, U_{d-r}(\rho(\alpha^{-1})) \right) > \varepsilon. \quad (13)$$

Observe that the conclusion of the proposition is the second condition required for a locally conformal point (Definition 5.5).

Before proving the Proposition let us fix a distance d on $\partial\Gamma$ inducing its topology and for $\nu > 0$ define a triple of points $x, y, z \in \partial\Gamma$ is ν -separated, if all distances $d(x, y)$, $d(y, z)$ and $d(x, z)$ are bounded below by ν . The following lemma follows from the convergence property of hyperbolic groups, see for example Tukia [43].

Lemma 6.8. *Let $(\alpha_i)_{i \in \mathbb{Z}}$ be a bi-infinite geodesic through $e \in \Gamma$ with $\alpha_i \rightarrow x$ and $\alpha_{-i} \rightarrow z$ say, as $i \rightarrow +\infty$. Then the function $y \mapsto d(\alpha_i^{-1}y, \alpha_i^{-1}z)$ converges to 0 uniformly on compact sets of $\partial\Gamma - \{x\}$ as $i \rightarrow +\infty$. Consequently, for fixed ν , the positive integers n such that the triple $\alpha_n^{-1}x, \alpha_n^{-1}y, \alpha_n^{-1}z$ is ν -separated is bounded above uniformly on compact sets of $\partial\Gamma - \{x\}$. Finally, there exists $\nu_0 > 0$ such that for every $0 < \varepsilon < \nu_0$ and $y \in \partial\Gamma - \{x\}$ with $d(x, y) < \varepsilon$ there exists $n \in \mathbb{N}$ such that $\alpha_n^{-1}x, \alpha_n^{-1}y, \alpha_n^{-1}z$ are ν_0 -separated.*

Proof. Let us give an idea of the proof in our situation, i.e. assuming that Γ admits a projective Anosov representation ρ . We focus on finding ν_0 and $n \in \mathbb{N}$ so that the last sentence of the statement holds.

Consider the distance d induced by our chosen distance on $\mathbb{P}(\mathbb{K}^d)$ through the boundary map ξ_ρ^1 . The fact that there is a lower bound on the values $d(\alpha_n^{-1}x, \alpha_n^{-1}z)$ for all n follows from Lemma 3.9, and the fact that we can find a suitable n , such that both $d(\alpha_n^{-1}z, \alpha_n^{-1}y) > \nu_0$ and $d(\alpha_n^{-1}x, \alpha_n^{-1}y) > \nu_0$ is a consequence of Lemma 3.10 combined with the fact that the action of the images of the generators on $\mathbb{P}(\mathbb{K}^d)$ is uniformly Lipschitz. \square

Proof of Proposition 6.7. Since the representation $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ is (p, q, r) -hyperconvex we can find ε_0 such that if $s, w, t \in \partial\Gamma$ are ν_0 -separated one has

$$\sin \sphericalangle (s_\rho^p \oplus t_\rho^q, w_\rho^{d-r}) > \varepsilon_0 : \quad (14)$$

this is guaranteed since the set of ν_0 -separated triples is precompact as the group is hyperbolic.

Let us first show that if y is close enough to x (depending on ε_0 , as well as the representation ρ), we can find ε_1, L_1 for which Equation (13) holds.

In order to do so, observe that, since the group is hyperbolic, and thus the cone-type graph is finite, there exists K smaller than the diameter of the cone-type graph such that, if $x \in \mathcal{C}_\infty(\alpha)$, there exists a bi-infinite geodesic $(\alpha_i)_{i \in \mathbb{Z}}$ passing through the identity, and an integer M such that $d(\alpha_{-M}, \alpha^{-1}) < K$; of course in this case $|\alpha| - M < K$. We denote by z be the second endpoint of such geodesic.




FIGURE 5. The first step in the proof of Proposition 6.7

By Lemma 6.8 we can choose $N \in \mathbb{N}$ such that $\alpha_N^{-1}x, \alpha_N^{-1}y, \alpha_N^{-1}z$ are ν_0 -separated. The size of N measures how close y is to x . Using the triangular inequality we get

$$\begin{aligned} & \sin \angle (x_\rho^p \oplus y_\rho^q, U_{d-r}(\rho(\alpha^{-1}))) \geq \sin \angle (U_r(\rho(\alpha_N)), U_{d-r}(\rho(\alpha_{-M}))) \\ & - d \left(U_{d-r}(\rho(\alpha_{-M})), U_{d-r}(\rho(\alpha^{-1})) \right) - d \left(\rho(\alpha_N) \left((\alpha_N^{-1}x)_\rho^p \oplus (\alpha_N^{-1}y)_\rho^q \right), U_r(\rho(\alpha_N)) \right). \end{aligned}$$

The first term of the expression is bigger than $\delta_{r,\rho}$ provided $|\alpha|$ is big enough, by Lemma 3.3. The second term is smaller than $\delta_{r,\rho}/3$ if $|\alpha|$ is big enough by Lemma 2.3 (2): indeed $\alpha^{-1} = \alpha_{-M}a$ for some $a \in \Gamma$ with $|a| < K$. We chose L_1 so that these two conditions are satisfied. In order to prove our claim it is enough to verify that we can find N_0 big enough, depending on the representation only, such that for every $N \geq N_0$, it holds

$$d(\rho(\alpha_N) \left((\alpha_N^{-1}x)_\rho^p \oplus (\alpha_N^{-1}y)_\rho^q \right), U_r(\rho(\alpha_N))) < \delta_\rho/3.$$

Since $z \neq x$ are fixed, the subspaces z_ρ^{d-r} and x_ρ^r have a positive angle and thus, since $U_r(\rho(\alpha_N)) \rightarrow x_\rho^r$ as $N \rightarrow \infty$ uniformly in N , the angle between z_ρ^{d-r} and $U_r(\rho(\alpha_N))$ is bounded below for all positive big enough N depending only on the representation ρ . Using Lemma 2.5 we deduce that

$$d \left(U_{d-r}(\rho(\alpha_N^{-1})), \rho(\alpha_N^{-1})z_\rho^{d-r} \right) \leq \frac{\sigma_{d-r+1}(\rho(\alpha_N^{-1}))}{\sigma_{d-r}} \frac{1}{\sin \angle (z_\rho^{d-r}, U_r(\rho(\alpha_N)))}. \quad (15)$$

Since the representation is $\{\mathbf{a}_{d-r}\}$ -Anosov, $(\sigma_{d-r+1}/\sigma_{d-r})(\rho(\alpha_N^{-1}))$ is smaller than $\varepsilon_0/2$ for big enough positive N . By hyperconvexity (equation (14)) we know that $(\alpha_N^{-1}x)_\rho^p \oplus (\alpha_N^{-1}y)_\rho^q$ has a definite angle with $(\alpha_N^{-1}z)_\rho^{d-r}$, consequently, by equation (15) we deduce that

$$\angle \left((\alpha_N^{-1}x)_\rho^p \oplus (\alpha_N^{-1}y)_\rho^q, U_{d-r}(\rho(\alpha_N^{-1})) \right) > \varepsilon_0/2.$$

Thus, by Lemma 2.5

$$d \left(\rho(\alpha_N) \left((\alpha_N^{-1}x)_\rho^p \oplus (\alpha_N^{-1}y)_\rho^q \right), U_r(\rho(\alpha_N)) \right) < \frac{\sigma_{r+1}(\rho(\alpha_N))}{\sigma_r} \frac{2}{\varepsilon_0}.$$

This concludes the first step, we can chose $\varepsilon_1 = \delta_{r,\rho}/3$.

We are thus left to verify that, up to possibly shrinking ε_1 and enlarging L_1 , Equation (13) is also verified in the case n for which $\alpha_n^{-1}(x, y, z)$ is ν_0 -far and smaller than a fixed N . Observe that, since the group Γ is finitely generated and N is fixed, we can find C , depending on ρ , such that $d(\alpha_n^{-1}x, \alpha_n^{-1}y) \leq C^n d(x, y)$, and therefore we can find ν_1 depending on N only such that $d(y, x) > \nu_1$. Since furthermore $y_\rho^q \in X_\rho^q(\alpha)$, and thus we have a lower bound on $d(y, z)$, we deduce, up to further shrinking ν_1 , that the triple (x, y, z) is ν_1 -far. The same argument as above let us deduce that there exists ε_2 such that

$$\sin \angle (x_\rho^p \oplus y_\rho^q, z_\rho^{d-r}) > \varepsilon_2.$$

It is then enough to chose L_2 big enough so that $d(z_\rho^{d-r}, U_{d-r}(\rho(\alpha^{-1}))) < \varepsilon_2/2$. The Proposition holds with $L = \max\{L_1, L_2\}$ and $\varepsilon = \min\{\varepsilon_1, \varepsilon_2\}$. \square

Proposition 6.7 combined with Theorem 5.14 yields the following Hausdorff dimension computations.

Corollary 6.9. *Let $\rho : \Gamma \rightarrow \text{PGL}_d(\mathbb{K})$ be $(1, 1, 2)$ -hyperconvex, then*

$$\text{Hff}(\xi^1(\partial\Gamma)) = h_\rho^{a_1}.$$

Corollary 6.10. *Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be $(1, 1, r)$ -hyperconvex. Assume moreover that for every $\gamma \in \Gamma$ one has $\sigma_2(\rho(\gamma)) = \sigma_r(\rho(\gamma))$, then every point of $\partial\Gamma$ is locally conformal for ρ and thus*

$$h_\rho^{a_1} = \mathrm{Hff}(\xi^1(\partial\Gamma)).$$

6.3. Examples: (ir)reducible SL_2 . The easiest examples of hyperconvex representations are induced from representations of $\mathrm{SL}_2(\mathbb{K})$ (see for example Humphreys's book [25] for standard basic facts on the representation theory of SL_2).

Recall that for every $d \in \mathbb{N} - \{0, 1\}$ there is a (unique up to conjugation) irreducible representation $\iota_d : \mathrm{SL}_2(\mathbb{K}) \rightarrow \mathrm{SL}_d(\mathbb{K})$. This representation is given by the action of $\mathrm{SL}_2(\mathbb{K})$ on the symmetric powers $S^{d-1}(\mathbb{K}^2)$, which can be identified with the space of homogenous polynomials on two variables of degree $d - 1$ with coefficients in \mathbb{K} . If we denote by $E_{\mathrm{SL}_2(\mathbb{K})}^*$ the weight space, the representation ι_d has highest weight $\chi_{\iota_d} \in E_{\mathrm{SL}_2(\mathbb{K})}^*$ given by $\chi_{\iota_d}(x) = (d - 1)x$.

Let $\mathcal{F}(S^{d-1}(\mathbb{K}^2))$ denote the full flag space associated to $\mathrm{SL}(S^{d-1}(\mathbb{K}^2))$. The Veronese map $\zeta : \mathbb{P}(\mathbb{K}^2) \rightarrow \mathcal{F}(S^{d-1}(\mathbb{K}^2))$ is defined by

$$\zeta(x) = \{\zeta^k(x)\}_{k=1}^{d-1}$$

where $\zeta^k(\ell)$ is the k -dimensional vector subspace of $S^{d-1}(\mathbb{K}^2)$ consisting of polynomials that have x^{d-k} as a factor. It is easy to check that ζ is ι_d -equivariant and the image of an attractor in $\mathbb{P}(\mathbb{K}^2)$ is an attractor in $\mathcal{F}(S^{d-1}(\mathbb{K}^2))$.

Remark 6.11. Note that for every pair of distinct points $x \neq y$ in $\mathbb{P}(\mathbb{K}^2)$ the flags $\zeta(x)$ and $\zeta(y)$ are in general position, i.e. for every $k \in \llbracket 1, d - 1 \rrbracket$, it holds $\zeta^k(x) \cap \zeta^{d-k}(y) = \{0\}$.

Moreover, using the transitivity of the $\mathrm{SL}_2(\mathbb{K})$ -action on transverse pairs, it is easy to check the following:

Proposition 6.12. *Let $\zeta = \{\zeta^i\}_{i=1}^{d-1}$ be the Veronese embedding of $\mathbb{P}(\mathbb{K}^2)$ into $\mathcal{F}(S_{d-1}(\mathbb{K}^2))$, then for every triple $p + q + r = d$ and pairwise distinct $x, y, z \in \mathbb{P}(\mathbb{K}^2)$ one has*

$$\zeta^p(x) \oplus \zeta^q(y) \oplus \zeta^r(z) = \mathbb{K}^d.$$

Corollary 6.13. *For every convex cocompact⁷ subgroup $\Gamma < \mathrm{SL}_2(\mathbb{K})$, the representation $\iota_d|_\Gamma : \Gamma \rightarrow \mathrm{SL}_d(\mathbb{K})$ is (p, q, r) -hyperconvex for every (p, q, r) such that $r \geq p + q$. The same holds for small deformations.*

We can obtain many more examples of hyperconvex representations by considering direct sums of irreducible representations. A representation $\pi : \mathrm{SL}_2(\mathbb{K}) \rightarrow \mathrm{SL}(V, \mathbb{K})$ decomposes in irreducible modules

$$\pi = \bigoplus_1^k \iota_{d_i},$$

where we have ordered $d_1 \geq \dots \geq d_k$. The highest weight $\chi_\pi \in E_{\mathrm{SL}_2(\mathbb{K})}^*$ is $\chi_\pi(x) = (d_1 - 1)x$. Let us denote by

$$\chi_\pi^{(2)} \geq \dots \geq \chi_\pi^{(\dim V)}$$

the remaining weights in decreasing order.

⁷For non-Archimedean fields \mathbb{K} , in analogy with the Archimedean case, we say that a representation is *convex cocompact* if it is Anosov, as in Definition 3.1.

Definition 6.14. Given $k \in \llbracket 2, \dim V \rrbracket$, we say that π is k -coherent if $\chi_\pi^{(k)} > d_2 - 1$, equivalently if $d_1 > d_2 + 2(k - 1)$.

Observe that a representation π is k -coherent if and only if the representation has a gap of index k and the top k eigenspaces are eigenlines of a diagonalizable element in $\pi(\mathrm{SL}_2(\mathbb{K}))$ and belong to the top irreducible factor. An important example of 2-coherent representations are exterior powers:

Example 6.15. For every $p \in \llbracket 1, d - 1 \rrbracket$ the representation

$$\wedge^p \iota_d : \mathrm{SL}_2(\mathbb{K}) \rightarrow \mathrm{SL}(\wedge^p \mathbb{K}^d)$$

is 2-coherent.

Proof. Considering a diagonalizable element in $\mathrm{SL}_2(\mathbb{K})$ one explicitly checks that the top 3 weights of $\wedge^p \iota_d$ are

$$\begin{aligned} - \chi_{\wedge^p \iota_d} &= d - 1 + \dots + d + 1 - 2p = p(d - p), \\ - \chi_{\wedge^p \iota_d}^{(2)} &= \chi_{\wedge^p \iota_d} - 2, \\ - \chi_{\wedge^p \iota_d}^{(3)} &= \chi_{\wedge^p \iota_d}^{(4)} = \chi_{\wedge^p \iota_d} - 4. \end{aligned}$$

□

Definition 6.14 guarantees some hyperconvexity:

Proposition 6.16. Let $\rho : \Gamma \rightarrow \mathrm{SL}_2(\mathbb{K})$ be convex co-compact. If $\pi : \mathrm{SL}_2(\mathbb{K}) \rightarrow \mathrm{SL}(V, \mathbb{K})$ is k -coherent, then $\pi \circ \rho$ is (p, q, k) -hyperconvex for every p, q with $p + q \leq k$.

Proof. Since $\chi^{(k)} > d_2 - 1$ one has that $\chi^{(k)} > \chi^{(k+1)}$ and thus $\pi \circ \rho$ is $\{\mathbf{a}_k\}$ -Anosov. Coherence implies thus that $\chi^{(l)} > \chi^{(l+1)}$ for every $l \in \llbracket 1, k \rrbracket$ and thus $\pi \circ \rho$ is also $\{\mathbf{a}_p, \mathbf{a}_q\}$ -Anosov since both p and q are smaller than k . The remainder of the statement follows from Lemma 6.12: if N denotes the dimension of V , $\zeta_{d_1}^p : \partial\Gamma \rightarrow \mathcal{G}_p(\mathbb{S}^{d-1}(\mathbb{K}^2)) \subset \mathcal{G}_p(V)$ is the ι_d -equivariant map induced by ζ , and $\xi_\pi^l : \partial\Gamma \rightarrow \mathcal{G}_l(V)$ denotes the boundary map associated to π , we have, for every $l \leq k$, that $\xi_\pi^l = \zeta_{d_1}^l$ and

$$\xi_\pi^{N-l} = \zeta_{d_1}^{d_1-l} \oplus \bigoplus_{i=2}^k \mathbb{S}^{d_i-1}(\mathbb{K}^2).$$

□

In particular Proposition 6.16 can be used to construct example of representations of Kleinian groups satisfying the assumptions of Theorem 5.14.

7. DIFFERENTIABILITY PROPERTIES

7.1. Convergence on pairs and bounds on the Hausdorff dimension. The following result, which follows from Proposition 6.7 is inspired by Guichard [22, Proposition 21], however, Guichard's proof relies heavily on the fact that $\partial\Gamma$ is a circle, and that the representation is (p, q, r) -hyperconvex for every triple p, q, r with $p + q = r$.

Theorem 7.1. Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be (p, q, r) -hyperconvex then for every $(w, y) \in \partial^{(2)}\Gamma$ one has

$$\lim_{(w, y) \rightarrow (x, x)} d(w_\rho^p \oplus y_\rho^q, x_\rho^r) = 0.$$

More precisely there exist constants C, μ such that, if $\{\alpha_i\}$ is a geodesic ray with endpoint x , for every $w, y \in \alpha_i \mathcal{C}_\infty(\alpha_i)$ it holds

$$d(w_\rho^p \oplus y_\rho^q, x_\rho^r) \leq C e^{-\mu i}.$$

Proof. The first claim is a direct consequence of the second, as the sets $\alpha_i \mathcal{C}_\infty(\alpha_i)$ form a fundamental system of neighbourhoods of the point x .

As the representation ρ is (p, q, r) -hyperconvex, and $w, y \in \alpha_i \mathcal{C}_\infty(\alpha_i)$, we deduce from Proposition 6.7 that

$$\sin \angle((\alpha_i^{-1} w)_\rho^p \oplus (\alpha_i^{-1} y)_\rho^q, U_{d-r}(\rho(\alpha_i^{-1}))) > \varepsilon.$$

In particular Lemma 2.5 implies

$$d(w_\rho^p \oplus y_\rho^q, U_r(\alpha_i)) \leq \frac{\sigma_{r+1}(\rho(\alpha_i))}{\sigma_r} \frac{1}{\varepsilon} \leq \frac{C_1}{\varepsilon} e^{-\mu_1 i}.$$

Where C_1, μ_1 are the constants provided by the fact that ρ is $\{\mathbf{a}_r\}$ -Anosov. The result now follows, via triangular inequality, from Lemma 3.6, which guarantees that

$$d(x_\rho^r, U_r(\alpha_i)) \leq C_2 e^{-\mu_2 i}.$$

□

The following easy converse is useful for applications:

Proposition 7.2. *Consider $p, q, r \in \llbracket 1, d-1 \rrbracket$ with $p+q \leq r$. If $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ is $\{\mathbf{a}_p, \mathbf{a}_q, \mathbf{a}_r\}$ -Anosov and for every $x \in \partial\Gamma$ one has*

$$\lim_{(w,y) \rightarrow (x,x)} d(w_\rho^p \oplus y_\rho^q, x_\rho^r) = 0, \quad (16)$$

then ρ is (p, q, r) -hyperconvex.

Proof. Since ρ is $\{\mathbf{a}_p, \mathbf{a}_q\}$ -Anosov and $p+q \leq r \leq d-1$, for every pair of distinct points w, y the sum $w_\rho^p + y_\rho^q$ is direct. Since ρ is $\{\mathbf{a}_r\}$ -Anosov there is a lower bound on $\sin \angle(x_\rho^r, z_\rho^{d-r})$ if x, z are the endpoints of a geodesic through the origin. Combining this fact with (16) we can find ε, δ such that

$$(x_\rho^p \oplus y_\rho^q) \cap z_\rho^{d-r} = \{0\}$$

for every triple with $d(x, y) < \varepsilon$ and $d(x, z) > \delta > \varepsilon$. Any triple in $\partial^{(3)}\Gamma$ can be transformed in such a triple by an element of Γ and thus the claim follows. □

Using the stereographic projection (see Definition 6.4) combined with Theorem 7.1 it is possible to deduce the following estimate on Hausdorff dimension:

Proposition 7.3. *Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be $(1, 1, r)$ -hyperconvex, then*

$$\mathrm{Hff}(\xi^1(\partial\Gamma)) \leq \mathrm{Hff}(\mathbb{P}(\mathbb{K}^r)).$$

Proof. We first claim that if $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ is $(1, 1, r)$ -hyperconvex, then for every x we can find a point z an open neighbourhood \mathcal{U}_x of x in $\xi^1(\partial\Gamma)$ such that the stereographic projection $\pi_{z,\rho}$ is Lipschitz on \mathcal{U}_x . Indeed as ρ is $\{\mathbf{a}_r\}$ -Anosov, we can choose z so that the subspaces x_ρ^r and z_ρ^{d-r} make a definite angle. The claim is then a consequence of Theorem 7.1: Indeed, it implies we can find an open neighbourhood \mathcal{U}_x of x such that for every pair $w, y \in \mathcal{U}_x$ the angle that $w_\rho^1 \oplus y_\rho^1$ makes with z_ρ^{d-r} is bigger than a fixed constant. This is enough to guarantee that the stereographic projection doesn't distort distances too much.

In particular, as Lipschitz maps preserve the Hausdorff dimension, it follows that $\text{Hff}(\mathcal{U}_x) \leq \text{Hff}(\mathbb{P}(\mathbb{K}^r))$. Since the Hausdorff dimension of a compact set is the maximum of the Hausdorff dimensions of the sets in a finite open cover, the result follows. \square

7.2. When $\partial\Gamma$ is a manifold and $\mathbb{K} = \mathbb{R}$. A classical result of Benoist [2] states that if a word hyperbolic group of projective transformations divides a convex set, then the boundary of this set has to be C^1 . These, together with Hitchin representations, have become the paradigm of Zariski-dense projective Anosov representations whose limit set is a regular manifold. The purpose of this section is to provide new examples of such phenomena. Sharper results of similar nature have recently been obtained independently by Zhang-Zimmer [45].

We begin by observing that Theorem 7.1 has the following interesting consequence.

Proposition 7.4. *Let $\rho : \Gamma \rightarrow \text{PGL}_d(\mathbb{R})$ be a $(1, 1, r)$ -hyperconvex representation and assume that $\partial\Gamma$ is topologically a sphere of dimension $r - 1$, then $\xi_\rho^1(\partial\Gamma)$ is a C^1 manifold with $\mathbb{T}_{x_\rho^1} \xi_\rho^1(\partial\Gamma) = \mathbb{T}_{x_\rho^1} \mathbb{P}(x_\rho^r)$.*

Proof. Theorem 7.1 implies that the set $\xi_\rho^1(\partial\Gamma)$ is differentiable at x_ρ^1 with tangent space $\mathbb{T}_{x_\rho^1} \mathbb{P}(x_\rho^r)$. The continuity of $x \mapsto x_\rho^r$ completes the proof. \square

Proposition 7.4 can be applied to many different situations to produce interesting examples through the *represent and deform* method, we now explain how this works in a specific situation. Denote by

$$S^k : \text{PGL}_{d+1}(\mathbb{R}) \rightarrow \text{PGL}(S^k(\mathbb{R}^{d+1}))$$

the k -symmetric power.

Note that in $\text{PGL}_{d+1}(\mathbb{K})$ a $(1, 1, d)$ -hyperconvex representation is a projective Anosov representation ρ such that for each triple $(x, y, z) \in \partial^{(2)}\Gamma$ the sum $x_\rho^1 + y_\rho^1 + z_\rho^1$ is direct.

Proposition 7.5. *Let $\rho : \Gamma \rightarrow \text{PGL}_{d+1}(\mathbb{R})$ be a $(1, 1, d)$ -hyperconvex representation and assume that there exist $c > 0, \mu > 1$ such that, for every $\gamma \in \Gamma$,*

$$\frac{\sigma_1(\rho(\gamma))\sigma_d(\rho(\gamma))}{\sigma_2(\rho(\gamma))^2} > ce^{\mu|\gamma|}. \quad (17)$$

Then the composition

$$S^k \circ \rho : \Gamma \rightarrow \text{PGL}(S^k(\mathbb{R}^{d+1}))$$

is $(1, 1, d)$ -hyperconvex.

Proof. We endow $S^k(\mathbb{R}^{d+1})$ with the norm induced by our choice of norm on \mathbb{R}^{d+1} . For this choice, and for every $g \in \text{PGL}_{d+1}(\mathbb{R})$, the semi-homotecy ratios of $S^k g$ are just the products of k -tuples of semihomotecy ratios of g . Assumption (17) then gives that for all γ apart from possibly finitely many exceptions

$$\begin{aligned} - \mathbf{a}_1(\nu(S^k \rho(\gamma))) &= \mathbf{a}_1(\nu(\rho(\gamma))), \\ - \mathbf{a}_d(\nu(S^k \rho(\gamma))) &= \min\{\mathbf{a}_d(\nu(\rho(\gamma))), \log \frac{\sigma_1(\rho(\gamma))\sigma_d(\rho(\gamma))}{\sigma_2(\rho(\gamma))^2}\} \end{aligned}$$

Since ρ is $\{\mathbf{a}_1, \mathbf{a}_d\}$ -Anosov, we deduce from Definition 3.1 that $S^k \rho$ is also $\{\mathbf{a}_1, \mathbf{a}_d\}$ -Anosov.

Observe that the map S^k is equivariant with respect to the map between the partial flags

$$S^k : \{\text{line} \subset \text{hyperplane}\} \rightarrow \{\text{line} \subset d\text{-dimensional subspace}\}$$

defined by

$$S^k(l, H) = (l^{\odot k}, l^{\odot k-1} \odot H).$$

Here we denote by \odot the symmetric tensors.

It is immediate to verify that Assumption (17) also implies that $S^k \circ \xi$ sends attractors to attractors, therefore, by continuity of $S^k \circ \xi$, we have, for every $x \in \partial\Gamma$,

$$S^k(x_\rho^1, x_\rho^d) = (x_{S^k\rho}^1, x_{S^k\rho}^d).$$

Finally, the convergence property (Theorem 7.1) for ρ , together with the differentiability of $S^2 : \mathbb{P}(\mathbb{R}^{d+1}) \rightarrow \mathbb{P}(S^2(\mathbb{R}^{d+1}))$ implies that

$$\lim_{(w,y) \rightarrow (x,x)} \not\propto (w_{S^2\rho}^1 \oplus y_{S^2\rho}^1, x_{S^2\rho}^d) = 0.$$

Proposition 7.2 yields the result. \square

As a direct corollary we get:

Corollary 7.6. *If $\rho : \Gamma \rightarrow \text{PSO}(d, 1)$ is cocompact, every small deformation*

$$\eta : \Gamma \rightarrow \text{PGL}(S^k(\mathbb{R}^{d+1}))$$

of $S^k\rho$ is $(1, 1, d)$ -hyperconvex. Any such η will have a C^1 -sphere as limit set in $\mathbb{P}(S^k(\mathbb{R}^{d+1}))$.

Applying Johnson-Millson's [26] bending technique we obtain the announced Zariski dense subgroups whose limit set is a C^1 sphere:

Corollary 7.7. *There exists a Zariski dense subgroup $\Gamma < \text{PGL}(S^2(\mathbb{R}^{d+1}))$ whose limit set is a C^1 sphere.*

Proof. Let M be a d -dimensional closed hyperbolic manifold that has a totally geodesic, co-dimension one, closed submanifold N . The inclusion $\Gamma = \pi_1 M \subset \text{SO}(d, 1) \rightarrow \text{SL}(S^2(\mathbb{R}^{d+1}))$ satisfies the hypothesis of Proposition 7.5. Without loss of generality we can assume that $\pi_1 N \subset \text{SO}(d-1, 1)$. Observe that the centralizer of

$$S^2(\text{SO}(d-1, 1)) \subset \text{SL}(S^2(\mathbb{R}^{d+1}))$$

is non-trivial and strictly contains that of $S^2(\text{SO}(d, 1))$: as an $S^2(\text{SO}(d, 1))$ -module, $S^2(\mathbb{R}^{d+1})$ splits as a direct sum of an irreducible representation (usually denoted $\mathbb{S}_{[2]}(\mathbb{R}^{d,1})$) and a trivial representation, its centralizer is thus reduced to \mathbb{R}^* . The decomposition as a $S^2(\text{SO}(d-1, 1))$ -module splits as the sum $\mathbb{S}_{[2]}(\mathbb{R}^{d-1,1}) \oplus \mathbb{R}^{d-1,1} \oplus \mathbb{R}^2$ where the action on the second factor is the standard action, while the action on \mathbb{R}^2 is trivial. In particular the centralizer of $S^2(\text{SO}(n-1, 1))$ is $\text{GL}(2, \mathbb{R}) \times \mathbb{R}^*$. By bending the representation along N with a nontrivial element in $\text{GL}(2, \mathbb{R})$ which doesn't leave invariant the factor \mathbb{R} , we obtain the desired representation. \square

8. EXAMPLES OF LOCALLY CONFORMAL REPRESENTATIONS

The purpose of this section is to discuss some of the many examples in which restricting the Zariski closure of a representation to a non-split real form of $\text{SL}_d(\mathbb{K})$ gives room for $(1, 1, p)$ hyperconvex representations for which we can also guarantee that the second gap p_2 is strictly bigger than 2.

8.1. Hyperconvex representations in $\mathrm{PU}(1, d)$ and $\mathrm{PSp}(1, d)$. The first interesting setting in which Theorem 5.14 applies for large classes of representations is given by considering representations in the rank one groups $\mathrm{PU}(1, d)$ or $\mathrm{PSp}(1, d)$. To unify the treatment we will write $\mathrm{PO}_{\mathbb{K}}(1, d)$ for either $\mathrm{PU}(1, d)$ if $\mathbb{K} = \mathbb{C}$ or $\mathrm{PSp}(1, d)$ if $\mathbb{K} = \mathbb{H}$ and regard $\mathrm{PO}_{\mathbb{K}}(1, d)$ as a subgroup of $\mathrm{PGL}(d + 1, \mathbb{K})$.

Remark 8.1. Unfortunately, as \mathbb{H} is non-commutative, we don't have the setup of Section 2.2.1 at our disposal (as the exterior algebra over a non-commutative field is not well defined), however the issue can be easily solved by considering $\mathrm{SL}(d + 1, \mathbb{H})$ as a subgroup of $\mathrm{SL}(2d + 2, \mathbb{C})$. Given an element $g \in \mathrm{SL}(d + 1, \mathbb{H})$ we denote by $g^{\mathbb{C}}$ the corresponding element in $\mathrm{SL}(2d + 2, \mathbb{C})$; it is then immediate to verify that we can choose a Cartan decomposition of $g^{\mathbb{C}}$ so that, for every p , the subspace $U_{2p}(g^{\mathbb{C}})$ is a quaternionic vector space, and we thus set $U_p(g) := U_{2p}(g^{\mathbb{C}})$. Similarly we say that a sequence $(\alpha_i)_{i \in \mathbb{Z}}$ in $\mathrm{SL}(d + 1, \mathbb{H})$ is p -dominated if $(\alpha_i^{\mathbb{C}})_{i \in \mathbb{Z}}$ is $2p$ -dominated in $\mathrm{SL}(2d + 2, \mathbb{C})$, and that a representation $\rho : \Gamma \rightarrow \mathrm{SL}(d + 1, \mathbb{H})$ is (p, q, r) -hyperconvex if the induced representation $\rho : \Gamma \rightarrow \mathrm{SL}(2d + 2, \mathbb{C})$ is $(2p, 2q, 2r)$ -hyperconvex. With this at hand it is easy to verify that Theorem 5.14 holds for representations with values in $\mathrm{SL}(d + 1, \mathbb{H})$.

Recall that $\mathrm{PO}_{\mathbb{K}}(1, d)$ has rank one, therefore we have at our disposal a good notion of convex co-compactness: a representation $\rho : \Gamma \rightarrow \mathrm{PO}_{\mathbb{K}}(1, d)$ is convex co-compact if and only if there is a convex $\rho(\Gamma)$ -invariant subspace of $\mathbb{H}_{\mathbb{K}}^d$ whose quotient is compact. The induced representation $\rho : \Gamma \rightarrow \mathrm{PGL}(d + 1, \mathbb{K})$ is $\{\mathbf{a}_1\}$ -Anosov if and only if ρ is convex co-compact, see for example Guichard-W. [23, Section 6.1].

Observe that $\mathrm{PO}_{\mathbb{K}}(1, d)$ preserves the closed codimension 1 submanifold $\partial\mathbb{H}_{\mathbb{K}}^d \subset \mathbb{P}(\mathbb{K}^{d+1})$, furthermore one has the following.

Lemma 8.2. *For every $g \in \mathrm{PO}_{\mathbb{K}}(1, d)$, we have $U_1(g) \in \partial\mathbb{H}_{\mathbb{K}}^d \subset \mathbb{P}(\mathbb{K}^{d+1})$ and $U_d(g) = U_1(g)^{\perp}$, where the orthogonal is defined with respect to the Hermitian form defining the group $\mathrm{PO}_{\mathbb{K}}(1, d)$.*

In particular, considering for every point $x \in \partial\mathbb{H}_{\mathbb{K}}^d$ the subspace $x^{\perp} \subset T_x\partial\mathbb{H}_{\mathbb{K}}^d$, one obtains a non-integrable distribution that has (real) codimension 1 if $\mathbb{K} = \mathbb{C}$ and 3 if $\mathbb{K} = \mathbb{H}$. In the complex case this is the standard contact structure on the sphere. We will refer to this distribution also in the quaternionic case as the *generalized contact distribution*. Given a distinct pair $x, y \in \partial\Gamma$ we will denote by $\mathcal{C}_{x,y}$ the intersection $\mathbb{P}(\langle x, y \rangle) \cap \partial\mathbb{H}_{\mathbb{K}}^d$. Of course if \mathbb{K} is \mathbb{C} then $\mathcal{C}_{x,y}$ is a circle, while if $\mathbb{K} = \mathbb{H}$ it is a 3-sphere. In the complex case the sets $\mathcal{C}_{x,y}$ are often referred to as *chains*, and their geometry was extensively studied by Cartan. The incidence geometry of chains (and of suitable generalizations) played an important role in Burger-Iozzi [9] and P. [36].

With these definition at hand we can rephrase our main results in the rank 1 setting:

Proposition 8.3. *A convex cocompact action $\rho : \Gamma \rightarrow \mathrm{PO}_{\mathbb{K}}(1, d)$ is $(1, 1, d)$ -hyperconvex if and only if for every distinct pair $x, y \in \partial\Gamma$, the chain $\mathcal{C}_{x_{\rho}^1, y_{\rho}^1}$ intersects $\xi(\partial\Gamma)$ only in x_{ρ}^1, y_{ρ}^1 . In this case $\mathrm{LC}(\rho) = \xi(\partial\Gamma)$, and $\xi(\partial\Gamma)$ is tangent to the generalized contact distribution.*

Proof. The first statement follows directly from the definitions: for every triple x, y, z the sum $x_{\rho}^1 + y_{\rho}^1 + z_{\rho}^1$ is direct if and only if z_{ρ}^1 doesn't belong to $\mathcal{C}_{x_{\rho}^1, y_{\rho}^1}$. The

second statement follows then from Proposition 6.7, and the last is a consequence of Theorem 7.1. \square

There are many interesting examples of representations satisfying the assumption of Proposition 8.3, a natural class of examples can be obtained deforming totally real embeddings. The following is a direct consequence of Proposition 8.3:

Lemma 8.4. *Let $\Gamma < \mathrm{PO}_{\mathbb{R}}(1, d)$ be a convex cocompact subgroup and let $\rho : \Gamma \rightarrow \mathrm{PO}_{\mathbb{K}}(1, d)$ be obtained extending the coefficients. Then ρ is $(1, 1, d)$ -hyperconvex.*

Corollary 8.5. *Every $\{\mathbf{a}_1\}$ -Anosov representation $\beta : \Gamma \rightarrow \mathrm{PO}_{\mathbb{K}}(1, d)$ sufficiently close to a totally real representation ρ is $(1, 1, d)$ -hyperconvex. In particular for each such representation*

$$\dim_{\mathrm{Hff}}(\xi(\partial\Gamma)) = h_{\beta}^{\mathbf{a}_1} \leq (d-1) \dim \mathbb{K}.$$

Proof. The first statement is a direct consequence of Propositions 6.2 and 8.3. Furthermore we know that for every element $g \in \mathrm{PO}_{\mathbb{K}}(1, d)$, we have $p_2(g) = d$, and hence every point in $\partial\Gamma$ is locally conformal for β . Theorem 5.14 then applies and gives the second statement. \square

Another class of examples was studied by Dufloux in his thesis [15, 16]. He says that a Schottky subgroup $\Gamma < \mathrm{PU}(1, d)$ generated by a symmetric set W is *well positioned* if, for every $w \in W$ there is an open subsets $B(w) \subset \partial\mathbb{H}_{\mathbb{C}}^d$ such that

- the closures $\overline{B(w)}$ are pairwise disjoint;
- $w(\partial\mathbb{H}_{\mathbb{C}}^d \setminus B(w^{-1})) \subset B(w)$;
- no chain passes through three of these open subsets $B(w)$.

Similarly one can define well positioned Schottky subgroups of $\mathrm{PSp}(1, d)$ replacing chains with quaternionic three spheres (recall that in $\partial\mathbb{H}_{\mathbb{H}}^d$ any pair of points uniquely determines a 3 sphere, the boundary of a totally geodesic copy of $\mathbb{H}_{\mathbb{H}}^1$). We will denote also these subspaces of $\partial\mathbb{H}_{\mathbb{H}}^d$ *chains* for notational ease.

Arguments analogue to the ones presented in [16, Section 7.2] imply that well positioned Schottky groups are hyperconvex representations:

Proposition 8.6. *Let $\rho : \Gamma \rightarrow \mathrm{PO}_{\mathbb{K}}(1, d)$ be a well positioned Schottky subgroup. Then $\rho : \Gamma \rightarrow \mathrm{SL}_{d+1}(\mathbb{K})$ is $(1, 1, d)$ -hyperconvex. Furthermore $\mathrm{LC}(\rho) = \xi(\partial\Gamma)$.*

Proof. Observe that since $\mathrm{PO}_{\mathbb{K}}(1, d)$ is a rank one group, $p_2(\alpha)$ doesn't depend on i . Furthermore, as soon as the sequence $\{\alpha_i\}_{i=1}^{\infty}$ forms a geodesic ray, the sequence is d dominated by a classical ping pong argument, and it follows from Lemma 8.2 that $E_{p_2}^{\rho}(x) = x^{\perp} \subset \mathbb{P}(\mathbb{K}^{d+1})$.

In order to verify that every point $x \in \partial\Gamma$ is locally conformal, we need to check that there exists a constant c such that $\angle(\xi(y) \oplus \xi(z), U_1(\rho(\alpha^{-1}))) > c$ for all $y, z \in X_{\infty}(\alpha)$. Since Γ is a well positioned Schottky group, we can choose δ_{ρ} as the smallest distance between two sets $B(w)$. Let w_{α} be the first letter of α . It follows from Lemma 3.6 that if $|\alpha|$ is big enough $U_1(\rho(\alpha^{-1})) \in B(w_{\alpha}^{-1})$ and $X_{\infty}(\alpha) \subset \bigcup_{s \neq w_{\alpha}} B(s)$: by construction $\mathcal{C}_{\infty}(\alpha) \subset \bigcup_{s \neq w_{\alpha}} B(s)$ and the intersection of the $\delta_{\rho}/2$ neighbourhood of $\bigcup_{s \neq w_{\alpha}} B(s)$ with the image of the boundary map is already contained in $\bigcup_{s \neq w_{\alpha}} B(s)$.

Since the chain $\mathcal{C}_{y,z}$ through y_{ρ}^1 and z_{ρ}^1 is the intersection of $\partial\mathbb{H}_{\mathbb{K}}^d$ with $\mathbb{P}(y_{\rho}^1 \oplus z_{\rho}^1)$, and, by assumption, $\mathcal{C}_{y,z}$ doesn't intersect the open subset $B(w_{\alpha}) \subset \partial\mathbb{H}_{\mathbb{K}}^d$, the result

follows. The fact that the representation $\rho : \Gamma \rightarrow \mathrm{SL}_{d+1}(\mathbb{K})$ is $(1, 1, d)$ -hyperconvex is a consequence of Theorem 7.1. \square

Corollary 8.7 (cfr. [15, Corollary 43]). *Let $\rho : \Gamma \rightarrow \mathrm{PO}_{\mathbb{K}}(1, d)$ be a well positioned Schottky subgroup. Then*

$$\mathrm{Hff}(\xi_{\rho}^1(\partial\Gamma)) = h_{\rho}^{a_1}.$$

Proof. If $\mathbb{K} = \mathbb{C}$ this follows directly from Theorem 5.14. For $\mathbb{K} = \mathbb{H}$ it is enough to observe that in the construction of the measure μ^{a_1} performed in Section 5.3 we never used the commutativity of the field \mathbb{K} . \square

We conclude the discussion on convex cocompact subgroups of $\mathrm{PO}_{\mathbb{K}}(1, d)$ by showing that the set of $(1, 1, d)$ -hyperconvex representations is, in general, not closed within the space of projective Anosov representations. We will prove in Proposition 9.3, that, instead, $(1, 1, 2)$ -hyperconvex representations of fundamental groups of surfaces are closed in the space of Anosov representations. Denote by \mathbb{F}_2 the free group on two generators.

Proposition 8.8. *There exists a continuous path of $\{\mathbf{a}_1\}$ -Anosov representations $\rho_t : \mathbb{F}_2 \rightarrow \mathrm{PU}(1, d)$ such that ρ_0 is $(1, 1, d)$ -hyperconvex and ρ_1 is not $(1, 1, d)$ -hyperconvex.*

Proof. As $\mathrm{PU}(1, d)$ has rank 1, for every 4-tuple (a^+, a^-, b^+, b^-) of pairwise distinct points in $\partial\mathbb{H}_{\mathbb{C}}^d$ we can find elements $a, b \in \mathrm{PU}(1, d)$ with prescribed attractive and repulsive fixed points and with translation length big enough so that the group generated by a, b is free and convex cocompact on $\mathbb{H}_{\mathbb{C}}^d$: this follows from a classical ping pong argument. Furthermore, if $(a_t^+, a_t^-, b_t^+, b_t^-)$ vary continuously in t we can also arrange for the elements a_t, b_t to vary continuously in t ; in this way we can define a continuous path $\rho_t : \mathbb{F}_2 \rightarrow \mathrm{PU}(1, d)$ of $\{\mathbf{a}_1\}$ -Anosov representation.

Our claim follows if we choose a_0, b_0 contained in $\mathrm{PO}(1, d)$ (so that the representation is $(1, 1, d)$ -hyperconvex by Lemma 8.4), and $(a_1^+, a_1^-, b_1^+, b_1^-)$ so that (a_1^+, a_1^-, b_1^+) belong to a single chain, but b_1^- doesn't. In this case the representation ρ_1 is clearly not $(1, 1, d)$ -hyperconvex as the sum $\xi(a^+) + \xi(a^-) + \xi(b^+)$ is not direct. \square

8.2. Locally conformal representations in $\mathrm{SO}(p, q)$. We now turn our attention to the group $\mathrm{SO}(p, q)$. Every semi-simple element $g \in \mathrm{SO}(p, q)$ has $|p - q|$ eigenvalues equal to 1. In this subsection, considering suitable exterior representations of $\mathrm{SO}(p, q)$ we will produce examples of hyperconvex representations for which every point is locally conformal, and thus Corollary 6.10 applies. For these representations, the Hausdorff dimension of the limit set computes the critical exponent for the first simple root.

The following generalization of Labourie's property (H) [31, Section 7.1.4] guarantees that a suitable exterior power is hyperconvex:

Proposition 8.9. *Let $\rho : \Gamma \rightarrow \mathrm{SO}(p, q)$ be $\{\mathbf{a}_{p-1}, \mathbf{a}_p\}$ -Anosov (here $p \leq q$). Then $\wedge^p \rho : \Gamma \rightarrow \mathrm{PGL}(\wedge^p(\mathbb{R}^{p,q}))$ is $\{\mathbf{a}_1, \mathbf{a}_{q-p+1}\}$ -Anosov. It is $(1, 1, q - p + 1)$ -hyperconvex if and only if for every $x, y, z \in \partial\Gamma$ pairwise distinct, the sum*

$$x_{\rho}^p + (z_{\rho}^p \cap y_{\rho}^{q+1}) + y_{\rho}^p$$

is direct. In this case every point in $\partial\Gamma$ is locally conformal.

Proof. Observe that the singular values of an element $g \in \mathrm{SO}(p, q) \subset \mathrm{SL}_{p+q}(\mathbb{R})$ have the form $\sigma_1(g) \geq \dots \sigma_p(g) \geq 1 = \dots = 1 \geq \sigma_p(g)^{-1} \geq \dots \sigma_1(g)^{-1}$, where 1 has multiplicity at least $q - p$ (higher if $\sigma_p(g) = 1$). If $\rho : \Gamma \rightarrow \mathrm{SO}(p, q)$ is $\{\mathfrak{a}_{p-1}, \mathfrak{a}_p\}$ -Anosov, then, for every γ with $|\gamma|$ big enough, it holds $\sigma_{p-1}(\rho(\gamma)) > \sigma_p(\rho(\gamma)) > 1$, hence in particular

$$\begin{aligned} \sigma_1(\wedge^p \rho(\gamma)) &= \sigma_1(\rho(\gamma)) \dots \sigma_p(\rho(\gamma)), \\ \sigma_2(\wedge^p \rho(\gamma)) &= \sigma_{q-p+1}(\wedge^p \rho(\gamma)) = \frac{\sigma_1(\wedge^p \rho(\gamma))}{\sigma_p(\rho(\gamma))} \text{ and} \\ \sigma_{q-p+2}(\wedge^p \rho(\gamma)) &= \max \left\{ \frac{\sigma_1(\wedge^p \rho(\gamma))}{\sigma_{p-1}(\rho(\gamma))}, \frac{\sigma_1(\wedge^p \rho(\gamma))}{\sigma_p(\rho(\gamma))^2} \right\}, \end{aligned}$$

which implies that $\wedge^p \rho$ is $\{\mathfrak{a}_1, \mathfrak{a}_{q-p+1}\}$ -Anosov.

Denote by $\mathcal{F}_{p-1, q}(\mathbb{R}^{p, q})$ the partial flag manifold consisting of pairs of $(p-1, q)$ -dimensional isotropic subspaces and consider the map

$$\begin{aligned} L : \mathcal{F}_{p-1, p}(\mathbb{R}^{p, q}) &\rightarrow \mathcal{F}_{1, q-p+1}(\wedge^p \mathbb{R}^{p, q}) \\ (P, Q) &\mapsto (\wedge^p(Q), \wedge^{p-1}(P) \wedge Q^\perp) \end{aligned}$$

where the orthogonal is considered with respect to the bilinear form defining the group $\mathrm{SO}(p, q)$. The map L is clearly equivariant with the homomorphism $\wedge^p : \mathrm{SO}(p, q) \rightarrow \mathrm{SL}(\wedge^p(\mathbb{R}^{p, q}))$; furthermore, if $g \in \mathrm{SO}(p, q)$ is $\mathcal{F}_{p-1, q}(\mathbb{R}^{p, q})$ -proximal, namely g has an attractive fixedpoint g^+ in $\mathcal{F}_{p-1, q}(\mathbb{R}^{p, q})$, then $L(g^+) = (\wedge^p g)^+$. Thus if $(\xi^{p-1}, \xi^p) : \partial\Gamma \rightarrow \mathcal{F}_{p-1, q}(\mathbb{R}^{p, q})$ denote the boundary maps associated to $\rho : \Gamma \rightarrow \mathrm{SO}(p, q)$, the boundary maps associated to $\wedge^p \rho$ have the form $L \circ (\xi^{p-1}, \xi^p)$.

Let N denote the dimension of $\wedge^p(\mathbb{R}^{p, q})$. In order to check if the representation $\wedge^p \rho$ is $(1, 1, q-p+1)$ -hyperconvex, it is enough to verify that for every distinct triple $x, y, z \in \partial\Gamma$, the subspace $x_{\wedge^p}^1 + z_{\wedge^p}^1$ intersects transversely $y_{\wedge^p}^{N-q+p-1}$, or, equivalently, the image of $x_{\wedge^p}^1 + z_{\wedge^p}^1$ in $\wedge^p \mathbb{R}^{p, q} / y_{\wedge^p}^{N-q+p-1}$ is two dimensional.

Recall that if $\rho : \Gamma \rightarrow \mathrm{SO}(p, q)$ is \mathfrak{a}_p -Anosov, then for every distinct pair $(x, y) \in \partial\Gamma^2$ it holds $x_\rho^p \oplus y_\rho^q = \mathbb{R}^d$, furthermore we can interpret any other point z_ρ^p as a linear map $z_\rho^p : x_\rho^p \rightarrow y_\rho^q$. With this notation the condition that the sum $x_\rho^p + (z_\rho^p \cap y_\rho^{q+1}) + y_\rho^p$ is direct is equivalent to requiring that

$$z_\rho^p(x_\rho^p \cap y_\rho^{q+1}) \cap y_\rho^p = \{0\}.$$

Let us then choose a basis $\{b_1 \dots b_p\}$ of x_ρ^p such that $\{b_1, \dots, b_{p-1}\}$ forms a basis of x_ρ^{p-1} and $b_p = x_\rho^p \cap y_\rho^{q+1}$, then we have that a basis of z^p is given by $c_i = b_i + z_\rho^p(b_i)$. Furthermore the only term of the explicit expression of $c_1 \wedge \dots \wedge c_p$ that might not belong to $y_{\wedge^p}^{N-q+p-1}$ is $b_1 \wedge \dots \wedge b_{p-1} \wedge c_p$. This last vector doesn't belong to $y_{\wedge^p}^{N-q+p-1} + x_{\wedge^p}^1$ if and only if $z_\rho^p(x_\rho^p \cap y_\rho^{q+1}) \cap y_\rho^p = \{0\}$. \square

Proposition 8.10. *Assume that there are convex cocompact representations $\rho_1 : \Gamma \rightarrow \mathrm{SO}(1, k)$, $\rho_2 : \Gamma \rightarrow \mathrm{SO}(1, l)$ such that ρ_1 strictly dominates ρ_2 , namely there exists constants c, μ such that $\sigma_1(\rho_1(\gamma)) > c\sigma_1(\rho_2(\gamma))^\mu$. Then the representation $\rho := \rho_1 \oplus \dots \oplus \rho_1 \oplus \rho_2 : \Gamma \rightarrow \mathrm{SO}(p, (p-1)k + l + s)$ satisfies the hypothesis of Proposition 8.9.*

Proof. The representation ρ is \mathfrak{a}_p -Anosov as ρ_2 is convex cocompact, and is \mathfrak{a}_{p-1} -Anosov as ρ_1 strictly dominates ρ_2 . Explicitly writing down the boundary map ξ^p associated to ρ in term of the boundary maps $\xi_1^1 : \Gamma \rightarrow \partial\mathbb{H}_{\mathbb{R}}^k$, $\xi_2^1 : \Gamma \rightarrow \partial\mathbb{H}_{\mathbb{R}}^l$

associated to ρ_1, ρ_2 one verifies that $z_\rho^p(x_\rho^p \cap y_\rho^{q+1}) \cap y_\rho^p \cong z_{\rho_2}^1 \cap y_{\rho_2}^1$ and the latter intersection is empty as the representation is Anosov. \square

Danciger-Gueritaud-Kassel [13, Proposition 1.8] gave an explicit construction of convex cocompact actions ρ_1, ρ_2 on $\mathbb{H}_\mathbb{R}^S$ of the the group Γ generated by reflections in the faces of a 4-dimensional regular right-angled 120-cell, such that ρ_1 strictly dominates ρ_2 and therefore Proposition 8.10 applies. In this case the boundary $\partial\Gamma$ is a 3 sphere. It is also easy to construct representations satisfying the assumption of Proposition 8.10 when the group Γ is free, and in this case it one can deform the representation $\rho : \mathbb{F}_n \rightarrow \mathrm{SO}(p, q)$ to obtain a Zariski dense representation whose image under \wedge^p is locally conformal. We also expect that many more convex cocompact subgroups in rank one have the same property, and it is probably possible to give further examples of situations in which Proposition 8.9 applies for more complicated groups, as, for example, hyperbolic Coxeter groups.

The same argument as in the proof of Proposition 8.9 gives the following

Proposition 8.11. *Let $\rho : \Gamma \rightarrow \mathrm{SL}_d(\mathbb{K})$ be $\{\mathbf{a}_{p-1}, \mathbf{a}_p, \mathbf{a}_s\}$ -Anosov. Assume that*

(i) *there exist constants c, μ such that*

$$\frac{\sigma_{p-1}(\rho(\gamma))\sigma_s(\rho(\gamma))}{\sigma_p(\rho(\gamma))\sigma_{p+1}(\rho(\gamma))} > ce^{\mu|\gamma|},$$

(ii) *for every $x, y, z \in \partial\Gamma$ pairwise distinct, the sum*

$$x_\rho^p + (z_\rho^p \cap y_\rho^{d-p+1}) + y_\rho^{d-s}$$

is direct,

then $\wedge^p \rho$ is $(1, 1, s - p + 1)$ -hyperconvex.

Observe that the first condition, which guarantees that the map

$$\begin{aligned} L : \mathcal{F}_{p-1, p, s}(\mathbb{K}^d) &\rightarrow \mathcal{F}_{1, s-p+1}(\wedge^p \mathbb{K}^d) \\ (P, Q, R) &\mapsto (\wedge^p(Q), \wedge^{p-1}(P) \wedge R) \end{aligned}$$

is proximal, is automatic if $s = p + 1$.

9. FUNDAMENTAL GROUPS OF SURFACES

Let us denote by Γ_S a word-hyperbolic group such that⁸ $\partial\Gamma_S$ is homeomorphic to \mathbb{S}^1 . One has the following direct consequence of Proposition 7.4 and Corollary 6.9.

Corollary 9.1. *Let $\rho : \Gamma_S \rightarrow \mathrm{PGL}_d(\mathbb{R})$ be $(1, 1, 2)$ -hyperconvex, then $h_\rho^{a_1} = 1$.*

9.1. Weak irreducibility and closedness. A projective Anosov representation $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ is *weakly irreducible* if the image of its boundary map is not contained in a proper subspace of $\mathbb{P}(\mathbb{K}^d)$. Clearly if ρ is irreducible, then ρ is weakly irreducible, but it is possible to construct examples of weakly irreducible Anosov representations with non reductive image.

The assumption of weak irreducibility can be used to study properties of the stereographic projection $\pi_{z, \rho}$ defined in Definition 6.4.

⁸A celebrated Theorem of Gabai [18] states that a hyperbolic group Γ_S such that $\partial\Gamma_S$ is a circle is virtually the fundamental group of a connected, closed genus ≥ 2 surface. We will not use this fact.

Lemma 9.2. *Let $\rho : \Gamma \rightarrow \mathrm{PGL}_d(\mathbb{K})$ be $\{\mathbf{a}_1, \mathbf{a}_p\}$ -Anosov. If the stereographic projection $\pi_{z,\rho} : \partial\Gamma \setminus \{z\} \rightarrow \mathbb{P}(\mathbb{K}^d)$ collapses an open set $U \subset \partial\Gamma$, then $\pi_{z,\rho}$ is constant. In particular the representation ρ is not weakly irreducible.*

Proof. Indeed, as fixed points of attractive elements are dense in $\partial\Gamma$ we can find $\gamma \in \Gamma$ with $\gamma^+ \in U$. Up to shrinking U we can assume that $\gamma \cdot U \subset U$. Let $V \subset \mathbb{K}^d$ be the smallest subspace containing $\xi(t)$ for every t in U . As $\pi_{z,\rho}|_U$ is constant, the subspace V is proper, furthermore $\rho(\gamma)V = V$, since if $\xi(x_1), \dots, \xi(x_k)$ is a basis of V then $\xi(\gamma x_1), \dots, \xi(\gamma x_k)$ are also linearly independent vectors contained in V . In particular, for every n , $\pi_{z,\rho}(\gamma^{-n}U)$ is constant. As the union of the sets of the form $\gamma^{-n}U$ is the complement of a point in $\partial\Gamma$, the first result follows by continuity of $\pi_{z,\rho}$.

If the map $\pi_{z,\rho}$ is constant then, for every $x \in \partial\Gamma - \{z\}$, the image of the boundary map is contained in the proper subspace $x_\rho^1 + z_\rho^{d-r}$, hence the representation is not weakly irreducible. \square

Lemma 9.2 is particularly useful to analyze properties of $(1, 1, 2)$ -representations of groups Γ_S . The following argument is very similar to Labourie [31, Proposition 8.3].

Proposition 9.3. *The space of real weakly irreducible $\{\mathbf{a}_1, \mathbf{a}_2\}$ -Anosov representations of Γ_S that are not $(1, 1, 2)$ -hyperconvex is open.*

Proof. Let $\rho : \Gamma_S \rightarrow \mathrm{PGL}_d(\mathbb{R})$ be $\{\mathbf{a}_1, \mathbf{a}_2\}$ -Anosov and not $(1, 1, 2)$ -hyperconvex. By definition, there exists a triple of pairwise distinct points $x, y, z \in \partial\Gamma_S$ such that

$$(x_\rho^1 \oplus y_\rho^1) \cap z_\rho^{d-2} \neq 0, \quad (18)$$

and thus the stereographic projection $\pi_{z,\rho}$ is not injective.

Note that $\mathbb{P}(\mathbb{R}^d/z_\rho^{d-2})$ is topologically a circle. Therefore the stereographic projection $\pi_{z,\rho}$ is a map from an interval with a point removed to a circle that:

- does not collapse intervals,
- is not injective.

One can therefore, using the intermediate value theorem, find an interval $I \subset \partial\Gamma_S \setminus \{z\}$ and a point $w \in \partial\Gamma_S \setminus (\{z\} \cup I)$ such that $\pi_{z,\rho}(w)$ belongs to the interior of $\pi_{z,\rho}(I)$.

This last property will hold for any map close enough to $\pi_{z,\rho}$, in particular for the stereographic projection $\pi_{z,\eta}$ for some η close to ρ . Thus, $\pi_{z,\eta}$ is not injective and hence η is not $(1, 1, 2)$ -hyperconvex, as desired. \square

Recall from Definition 6.14 that a reducible representation $\pi : \mathrm{SL}_2(\mathbb{K}) \rightarrow \mathrm{SL}_d(\mathbb{K})$ is k -coherent if it has a gap of index k and its highest k weights belong to the same irreducible factor. Combining results from previous sections one has the following.

Corollary 9.4. *Let $\pi : \mathrm{SL}_2(\mathbb{R}) \rightarrow \mathrm{SL}_d(\mathbb{R})$ be a 2-coherent representation and $\rho : \Gamma_S \rightarrow \mathrm{PSL}_2(\mathbb{R})$ be co-compact, then any deformation η of $\pi\rho$ among weakly irreducible $\{\mathbf{a}_1, \mathbf{a}_2\}$ -Anosov representations into $\mathrm{PSL}_d(\mathbb{R})$ is $(1, 1, 2)$ -hyperconvex. In particular, verifies:*

- has C^1 -limit set in $\mathbb{P}(\mathbb{R}^d)$,
- the exponential growth rate $h_\eta^{\mathbf{a}_1} = 1$.

Proof. Proposition 6.16 states that $\pi\rho$ is $(1, 1, 2)$ -hyperconvex. Proposition 6.2 states hyperconvexity is an open property and, since $\partial\Gamma_S$ is topologically a circle, Proposition 9.3 implies that $(1, 1, 2)$ -hyperconvex is closed among weakly irreducible $\{\mathbf{a}_1, \mathbf{a}_2\}$ -Anosov representations. The remaining statements follow from Proposition 7.4 and Corollary 6.9 for $\mathbb{K} = \mathbb{R}$. \square

This result can be useful to distinguish some components of weakly irreducible Anosov representations (similar bounds on the number of connected components of Anosov representations were obtained with different techniques by Stecker-Treib [41, Corollary 8.2]).

9.2. The Hitchin component of $\mathrm{PSL}_d(\mathbb{R})$. Let S be a closed connected oriented surface of genus ≥ 2 . The *Hitchin component* of $\mathrm{PSL}_d(\mathbb{R})$ is a connected component of the character variety $\mathfrak{X}(\pi_1 S, \mathrm{PSL}_d(\mathbb{R}))$ that contains a *Fuchsian representation*, i.e. a representation that factors as

$$\pi_1 S \rightarrow \mathrm{PSL}_2(\mathbb{R}) \xrightarrow{\iota_d} \mathrm{PSL}_d(\mathbb{R}),$$

where the first arrow is a the choice of a hyperbolic metric on S . Such a connected component is usually denoted by $\mathcal{H}_d(S)$ and an element $\rho \in \mathcal{H}_d(S)$ is called a *Hitchin representation*.

Recall from Labourie [31] that a map $\xi : \partial\Gamma \rightarrow \mathcal{F}(\mathbb{R}^d)$ satisfies Property (H) if for every triple of distinct points x, y, z and every integer k one has

$$\xi^{k+1}(y) + (\xi^{k+1}(z) \cap \xi^{n-k}(x)) + \xi^{n-k-2}(x) = \mathbb{R}^d.$$

One has the following central result by Labourie [31].

Theorem 9.5 (Labourie [31]). *Every Hitchin representation $\rho : \Gamma \rightarrow \mathrm{PSL}_d(\mathbb{R})$ is (p, q, r) -hyperconvex, for every triple with $p + q = r$. The equivariant boundary map $\xi : \partial\Gamma \rightarrow \mathcal{F}(\mathbb{R}^d)$ has property (H).*

Thus, one concludes the following for deformations of the exterior powers.

Proposition 9.6. *Let $\rho \in \mathcal{H}_d(S)$ and consider any $k \in \llbracket 1, d-1 \rrbracket$. Then any weakly irreducible $\{\mathbf{a}_1, \mathbf{a}_2\}$ -Anosov representation $\eta : \pi_1 S \rightarrow \mathrm{PSL}(\wedge^k \mathbb{R}^d)$ connected by weakly irreducible $\{\mathbf{a}_1, \mathbf{a}_2\}$ -Anosov representations to $\wedge^k \rho$ is $(1, 1, 2)$ -hyperconvex and consequently verifies:*

- has C^1 -limit set in $\mathbb{P}(\wedge^k \mathbb{R}^d)$,
- the exponential growth rate $h_\eta^{\mathbf{a}_1} = 1$.

Proof. Observe that for every s , the representation $\wedge^s \rho$ is $\{\mathbf{a}_1, \mathbf{a}_2\}$ -Anosov, furthermore Proposition 8.11 ensures that $\wedge^s \rho$ is $(1, 1, 2)$ -hyperconvex as the boundary curve satisfies Property (H) (for $k = s - 1$). The result then follows from Proposition 7.4, Corollary 6.9 and Corollary 9.4. \square

When no deformation is applied one recovers the following result from Potrie-S. [35, Theorem B].

Theorem 9.7 (Potrie-S. [35]). *For every $\rho \in \mathcal{H}_d(S)$ and every $k \in \llbracket 1, d-1 \rrbracket$ one has $h_\rho^{\mathbf{a}_k} = 1$.*

9.3. Hitchin representations in other groups. More generally, let $G_{\mathbb{R}}$ be a simple real-split Lie group. These have been classified, i.e. up to finite coverings $G_{\mathbb{R}}$ is a group in the following list: $\mathrm{PSL}_d(\mathbb{R})$, $\mathrm{PSp}(2n, \mathbb{R})$, $\mathrm{SO}(n, n+1)$ $\mathrm{SO}(n, n)$, or it is the split real forms of the exceptional groups $F_{4, \mathrm{split}}$, $G_{2, \mathrm{split}}$, $E_{6, \mathrm{split}}$, $E_{7, \mathrm{split}}$ and $E_{8, \mathrm{split}}$.

The work of Kostant [29] provides a subalgebra $\iota_{\mathfrak{g}_{\mathbb{R}}} : \mathfrak{sl}_2(\mathbb{R}) \rightarrow \mathfrak{g}_{\mathbb{R}}$, unique up to conjugation and called *the principal \mathfrak{sl}_2* , such that the centralizer of $\iota_{\mathfrak{g}_{\mathbb{R}}}\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ has minimal dimension. Denote by $\iota_{G_{\mathbb{R}}} : \mathrm{PSL}_2(\mathbb{R}) \rightarrow G_{\mathbb{R}}$ the induced morphism. For example, $\iota_{\mathrm{PSL}_d(\mathbb{R})} = \iota_d$ is the (unique up to conjugation) irreducible representation of $\mathrm{SL}_2(\mathbb{R})$ in \mathbb{R}^d defined in Subsection 6.3.

Let S be a closed connected genus ≥ 2 surface. The *Hitchin component* of $G_{\mathbb{R}}$ is the connected component of the character variety $\mathfrak{X}(\pi_1 S, G_{\mathbb{R}})$ that contains a *Fuchsian representation*, i.e. a representation that factors as

$$\pi_1 S \rightarrow \mathrm{PSL}_2(\mathbb{R}) \xrightarrow{\iota_{G_{\mathbb{R}}}} G_{\mathbb{R}},$$

where the first arrow is a the choice of a hyperbolic metric on S . We will denote this connected component by $\mathcal{H}(S, G_{\mathbb{R}})$ and an element $\rho \in \mathcal{H}(S, G_{\mathbb{R}})$ is called a *Hitchin representation*.

Remark 9.8 (Canonical inclusions). By construction, one sees that the irreducible representation $\iota_d : \mathrm{SL}_2(\mathbb{R}) \rightarrow \mathrm{SL}_d(\mathbb{R})$ factors, depending on the parity of d , as

$$\begin{aligned} \mathrm{SL}_2(\mathbb{R}) &\xrightarrow{\iota_{\mathrm{Sp}(2n, \mathbb{R})}} \mathrm{Sp}(2n, \mathbb{R}) \rightarrow \mathrm{SL}_{2n}(\mathbb{R}), \\ \mathrm{SL}_2(\mathbb{R}) &\xrightarrow{\iota_{\mathrm{SO}(n, n+1)}} \mathrm{SO}(n, n+1) \rightarrow \mathrm{SL}_{2n+1}(\mathbb{R}), \\ \mathrm{SL}_2(\mathbb{R}) &\xrightarrow{G_{2, \mathbb{R}}} G_{2, \mathbb{R}} \rightarrow \mathrm{SO}(3, 4) \rightarrow \mathrm{SL}_7(\mathbb{R}), \end{aligned}$$

where, in each case, the first arrows is the principal inclusion $\iota_{G_{\mathbb{R}}}$. Thus

$$\begin{aligned} \mathcal{H}(S, \mathrm{PSp}(2n, \mathbb{R})) &\subset \mathcal{H}_{2n}(S), \\ \mathcal{H}(S, \mathrm{PSO}(n, n+1)) &\subset \mathcal{H}_{2n+1}(S), \\ \mathcal{H}(S, \mathbb{P}G_{2, \mathbb{R}}) &\subset \mathcal{H}(S, \mathrm{PSO}(3, 4)) \subset \mathcal{H}_7(S). \end{aligned}$$

On the other hand if we consider the embedding $\mathrm{SO}(n-1, n) \subset \mathrm{SO}(n, n)$ as the stabilizer of a positive definite line. The morphism $\iota_{\mathrm{SO}(n, n)}$ is the composition of $\iota_{\mathrm{SO}(n-1, n)}$ with such inclusion. Hence the induced action of $\iota_{\mathrm{SO}(n, n)}$ on $\mathbb{R}^{n, n}$ decomposes in $\mathrm{SL}_2(\mathbb{R})$ -irreducible modules as

$$\iota_{2n-1} \oplus \iota_1,$$

and in particular $\mathcal{H}(S, \mathrm{PSO}(n, n))$ is not a subset of a $\mathrm{PSL}_{2n}(\mathbb{R})$ -Hitchin component.

It is known to experts that every Hitchin representation is Anosov with respect to the minimal parabolic of G , see for example Fock-Goncharov [17].

Recall that the simple roots of the group $\mathrm{PSO}(n, n)$ are given by $\{\mathbf{a}_1, \dots, \mathbf{a}_{n-1}, \mathbf{b}_n\}$ where, as above, $\mathbf{a}_i(x) = x_i - x_{i+1}$ and \mathbf{b}_n is defined by

$$\mathbf{b}_n(x) = x_{n-1} + x_n.$$

Thus every representation $\rho \in \mathcal{H}(S, \mathrm{PSO}(n, n))$, when considered as a representation in $\mathrm{SL}_{2n}(\mathbb{R})$ under the canonical inclusion, is $\{\mathbf{a}_p\}$ -Anosov for every $p \leq n-1$.

Furthermore it is easy to check that the n -th exterior power

$$\wedge^n : \mathrm{PSO}(n, n) \rightarrow \mathrm{PSL}(\wedge^n \mathbb{R}^{2n})$$

splits as the direct sum of two irreducible $\mathrm{PSO}(n, n)$ -modules, which have respectively \mathfrak{a}_{n-1} and \mathfrak{b}_n as first root (see for example Danciger-Zhang [14]). In particular we obtain the following result, independently announced by Labourie [30].

Theorem 9.9. *For every $\rho \in \mathcal{H}(S, \mathrm{PSO}(n, n))$ and every $p \leq n - 2$ the exterior power $\wedge^p \rho$ is $(1, 1, 2)$ -hyperconvex, and the same holds for each one of the two irreducible submodules of $\wedge^n \rho$. Thus the associated limit curve of ρ on the p -Grassmannian for $p \leq n - 2$, as well as each one of the two limit curves in the n -Grassmannian, is C^1 and one has $h_{\mathfrak{a}}^p = 1$ for every simple root \mathfrak{a} .*

Proof. Considering a diagonalizable element in $\mathrm{SL}_2(\mathbb{R})$ as in the proof of Lemma 6.15 we obtain that $\wedge^k(\iota_{2n-1} \oplus \iota_1)$ is 2-coherent for every $k \in \llbracket 1, n - 2 \rrbracket$. Similarly a direct computation shows that the 5 highest weights of $\wedge^n(\iota_{2n-1} \oplus \iota_1)$ are

$$\begin{aligned} - \chi_{\wedge^n(\iota_{2n-1} \oplus \iota_1)}^{(2)} &= \chi_{\wedge^n(\iota_{2n-1} \oplus \iota_1)}^{(2)} = 2n + \dots + 2 = n(n + 1), \\ - \chi_{\wedge^n(\iota_{2n-1} \oplus \iota_1)}^{(3)} &= \chi_{\wedge^n(\iota_{2n-1} \oplus \iota_1)}^{(4)} = 2n + \dots + 4 = n(n + 1) - 2 \\ - \chi_{\wedge^n(\iota_{2n-1} \oplus \iota_1)}^{(5)} &= 2n + \dots + 6 = n(n + 1) - 4 \end{aligned}$$

and each of the first four weights appears with multiplicity one in each irreducible $\mathrm{SO}(n, n)$ -submodules of $\wedge^n \mathbb{R}^{2n}$. We deduce that the restriction of the representation $\wedge^n(\iota_{2n-1} \oplus \iota_1)$ to each of the two submodules is also $(1, 1, 2)$ -hyperconvex. The result is then a consequence of Corollary 9.4 together with the classification of Zariski closures due to Guichard [21]. \square

Remark 9.10. Danciger-Zhang [14] recently proved that when a representation $\rho \in \mathcal{H}(S, \mathrm{PSO}(n, n))$ is regarded as a representation in $\mathrm{PSL}_{2n}(\mathbb{R})$, it is, instead, never $\{\mathfrak{a}_n\}$ -Anosov and the limit curve in the $n - 1$ -Grassmannian is never C^1 .

REFERENCES

- [1] B. Bárány, M. Hochman, and A. Rapaport. Hausdorff dimension of planar self-affine sets and measures. <https://arxiv.org/abs/1712.07353>, 2017. (Cited on page 2.)
- [2] Y. Benoist. Convexes divisibles I. In *Algebraic groups and arithmetic*, pages 339–374. Tata Inst. Fund. Res., 2004. (Cited on page 36.)
- [3] J. Bochi and N. Gourmelon. Some characterizations of domination. *Math. Z.*, 263:221–231, 2009. (Cited on pages 6 and 12.)
- [4] J. Bochi, R. Potrie, and A. Sambarino. Anosov Representations and dominated splittings. To appear in *J. Eur. Math. Soc.*, <https://www.emis-ph.org/journals/forthcoming.php?jrn=jems>, 2017. (Cited on pages 2, 3, 6, 7, 9, 10, 12, 13, 14, 16, 17, 18, 19, and 20.)
- [5] M. Bourdon. Structure conforme au bord et flot géodésique d’un $\mathrm{CAT}(-1)$ -espace. *Enseign. Math. (2)*, 41:63–102, 1995. (Cited on page 2.)
- [6] M. Bridgeman, R. Canary, F. Labourie, and A. Sambarino. The pressure metric for Anosov representations. *Geom. Funct. Anal.*, 25(4):1089–1179, 2015. (Cited on page 3.)
- [7] M.R. Bridson and A. Haefliger. *Metric spaces of non-positive curvature*, volume 319. Springer-Verlag, Berlin, 1999. (Cited on pages 14 and 15.)
- [8] F. Bruhat and J. Tits. Groupes réductifs sur un corps local, i. données radicielles valuées. *Publ. Math. de l’I.H.E.S.*, 41:5–251, 1972. (Cited on page 8.)
- [9] M. Burger and A. Iozzi. A measurable cartan theorem and applications to deformation rigidity in complex hyperbolic geometry. *Pure Appl. Math. Q.*, 41:181–202, 2008. Special Issue: In honor of Grigory Margulis. (Cited on page 38.)
- [10] J. Chen and Y. Pesin. Dimension of non-conformal repellers: a survey. *Nonlinearity*, 23(4):93–114, 2010. (Cited on page 2.)
- [11] M. Coornaert, T. Delzant, and A. Papadopoulos. *Géométrie et théorie des groupes: Les groupes hyperboliques de Gromov*. Springer-Verlag, Berlin, 1990. (Cited on pages 14 and 15.)

- [12] S. Crovisier and R. Potrie. Introduction to partially hyperbolic dynamics. In *School on Dynamical Systems*, Trieste, 2015. ICTP. (Cited on page 12.)
- [13] J. Danciger, F. Guéritaud, and F. Kassel. Proper affine actions for right-angled coxeter groups. <https://arxiv.org/abs/1804.03132>. (Cited on page 42.)
- [14] J. Danciger and T. Zhang. Affine actions with hitchin linear part. <https://arxiv.org/abs/1812.03930>. (Cited on page 46.)
- [15] L. Dufloux. Dimension de Hausdorff des ensembles limites. (Cited on pages 39 and 40.)
- [16] L. Dufloux. Hausdorff dimension of limit sets. *Geom. Dedic.*, 191:1–35, 2017. (Cited on pages 2 and 39.)
- [17] V. Fock and A. Goncharov. Moduli spaces of local systems and higher teichmüller theory. *Publ. Math. de l'I.H.E.S.*, 103(1):1–211, 2006. (Cited on page 45.)
- [18] D. Gabay. Convergence groups are fuchsian groups. *Ann. of Math.*, 136:447–510, 1992. (Cited on page 42.)
- [19] O. Glorieux, D. Monclair, and N. Tholozan. Hausdorff dimension of limit set for projective anosov groups. In preparation, 2018. (Cited on pages 3 and 20.)
- [20] F. Guéritaud, O. Guichard, F. Kassel, and A. Wienhard. Anosov representations and proper actions. *Geom. & Top.*, 21:485–584, 2017. (Cited on pages 2, 3, and 16.)
- [21] O. Guichard. Personal communication. (Cited on page 46.)
- [22] O. Guichard. Composantes de Hitchin et représentations hyperconvexes de groupes de surface. *J. Differential Geom.*, 80:391–431, 2008. (Cited on page 34.)
- [23] O. Guichard and A. Wienhard. Anosov representations: domains of discontinuity and applications. *Invent. Math.*, 190:357–438, 2012. (Cited on pages 2, 3, 16, 19, and 38.)
- [24] P. Haïssinsky. Géométrie quasiconforme, analyse au bord des espaces métriques hyperboliques et rigidités. (Cited on page 28.)
- [25] J. Humphreys. *Introduction to Lie algebras and representation theory*. Springer – Verlag, 1972. (Cited on page 33.)
- [26] D. Johnson and J. Millson. Deformation spaces associated to compact hyperbolic manifolds. In R. Howe, editor, *Discrete groups in geometry and analysis*, volume 67 of *Progress in Mathematics*, pages 48–106. Birkhäuser, Boston, MA., 1987. (Cited on page 37.)
- [27] M. Kapovich, B. Leeb, and J. Porti. A Morse Lemma for quasigeodesics in symmetric spaces and euclidean buildings. <https://128.84.21.199/abs/1411.4176v1>, 2014. (Cited on pages 2, 16, and 18.)
- [28] M. Kapovich, B. Leeb, and J. Porti. Morse actions of discrete groups on symmetric space. <http://front.math.ucdavis.edu/1403.7671>, 2014. (Cited on pages 2 and 3.)
- [29] Bertram Kostant. The principal three-dimensional subgroup and the Betti numbers of a complex simple Lie group. *Amer. J. Math.*, 81:973–1032, 1959. (Cited on page 45.)
- [30] F. Labourie. Entropy and affine actions for surfaces groups. <https://arxiv.org/abs/1908.00599>. (Cited on page 46.)
- [31] F. Labourie. Anosov Flows, Surface Groups and Curves in Projective Space. *Invent. Math.*, 165:51–114, 2006. (Cited on pages 2, 3, 16, 19, 29, 40, 43, and 44.)
- [32] F. Ledrappier and L.-S. Young. The metric entropy of diffeomorphisms: Part I: Characterization of measures satisfying Pesin’s entropy formula. *Ann. of Math.*, 122(3):509–539, 1985. (Cited on page 2.)
- [33] D. Lind and R. Marcus. *An introduction to symbolic dynamics and coding*. Cambridge University Press, Cambridge, 1995. (Cited on page 15.)
- [34] G. Margulis. *Discrete subgroups of semisimple Lie groups*. Springer Verlag, 1991. (Cited on page 13.)
- [35] R. Potrie and A. Sambarino. Eigenvalues and entropy of a Hitchin representation. *Invent. Math.*, 209:885–925, 2017. (Cited on pages 3, 5, 6, and 44.)
- [36] B. Pozzetti. Maximal representations of complex hyperbolic lattices into $su(m, n)$. *GAF*, 41:1290–1332, 2015. (Cited on page 38.)
- [37] J.-F. Quint. Cônes limites de sous-groupes discrets des groupes réductifs sur un corps local. *Transformation groups*, 7(3):247–266, 2002. (Cited on page 8.)
- [38] J.-F. Quint. Divergence exponentielle des sous-groupes discrets en rang supérieur. *Comment. Math. Helv.*, 77:563–608, 2002. (Cited on pages 7 and 8.)
- [39] J.-F. Quint. Mesures de Patterson-Sullivan en rang supérieur. *Geom. funct. anal.*, 12:776–809, 2002. (Cited on pages 25, 26, and 29.)

- [40] M. Sambarino. A (short) survey on dominated splittings. In J.A. de la Peña, J.A. López-Mimbela, M. Nakamura, and J. Petean, editors, *Mathematical congress of the Americas*, volume 656 of *Contemporary Mathematics*, pages 149–183, 2016. (Cited on page 13.)
- [41] F. Stecker and N. Treib. Domains of discontinuity in oriented flag manifolds. <https://arxiv.org/abs/1806.04459>, 2018. (Cited on page 44.)
- [42] D. Sullivan. The density at infinity of a discrete group of hyperbolic motions. *Publ. Math. de l’I.H.E.S.*, 50:171–202, 1979. (Cited on pages 1, 26, and 28.)
- [43] P. Tukia. Convergence groups and Gromov’s hyperbolic spaces. *New Zealand J. Math.*, 23, 1994. (Cited on page 31.)
- [44] C. Yue. The ergodic theory of discrete isometry groups on manifolds of variable negative curvature. *Trans. of the A.M.S.*, 348(12):4965–5005, 1996. (Cited on page 2.)
- [45] T. Zhang and A. Zimmer. Regularity of limit sets of Anosov representations. In preparation. (Cited on pages 5 and 36.)

Beatrice Pozzetti
 Ruprecht-Karls Universität Heidelberg
 Mathematisches Institut, Im Neuenheimer Feld 205, 69120 Heidelberg, Germany
 pozzetti@mathi.uni-heidelberg.de

Andrés Sambarino
 Sorbonne Université
 IMJ-PRG (CNRS UMR 7586)
 4 place Jussieu 75005 Paris France
 andres.sambarino@imj-prg.fr

Anna Wienhard
 Ruprecht-Karls Universität Heidelberg
 Mathematisches Institut, Im Neuenheimer Feld 205, 69120 Heidelberg, Germany
 HITS gGmbH, Heidelberg Institute for Theoretical Studies
 Schloss-Wolfsbrunnenweg 35, 69118 Heidelberg, Germany
 wienhard@uni-heidelberg.de