



HAL
open science

Permanent grassland classifications predict agronomic and environmental characteristics well, but not ecological characteristics

Geoffrey Mesbahi, Alice Michelot-Antalik, Jérémie Goulnik, Sylvain Plantureux

► To cite this version:

Geoffrey Mesbahi, Alice Michelot-Antalik, Jérémie Goulnik, Sylvain Plantureux. Permanent grassland classifications predict agronomic and environmental characteristics well, but not ecological characteristics. *Ecological Indicators*, 2020, 110, pp.105956. 10.1016/j.ecolind.2019.105956 . hal-02392277

HAL Id: hal-02392277

<https://hal.science/hal-02392277>

Submitted on 3 Dec 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Title

2 Permanent grassland classifications predict agronomic and environmental characteristics well,
3 but not ecological characteristics

4

5

6 Authors

7 Geoffrey Mesbahi ^{a,b,*}, Alice Michelot-Antalik ^a, Jérémie Goulnik ^a, Sylvain Plantureux ^a

8

9 Affiliation

10

11 a: Université de Lorraine, Inra, LAE, F-54000 Nancy, France

12 b: Parc Naturel Régional des Vosges du Nord, 67290 La Petite Pierre, France

13 * corresponding author: geoffrey.mesbahi@gmail.com

14

15

16 Declaration of interest

17 None

18

19

20 Colour

21 Figure 1, Figure 2 and Figure 3 require colour.

22

23 Abstract

24 Permanent grasslands produce highly diverse ecosystem goods and services, which need to be
25 easily assessed by decision makers. Naturalists and agronomists classify grasslands in
26 different ways to predict ecological, agronomic and environmental characteristics of the
27 grasslands. However, few studies have compared the prediction abilities of these different
28 classifications using the same botanical relevés, and none has explored the utility of
29 combining classifications. In this study, we attributed a grassland class from each of three
30 classifications (phytosociological, agronomic and functional) to 250 permanent grasslands in
31 north-eastern France to predict 16 characteristics: nine ecological, three agronomic and four
32 environmental. We used statistical model selection to identify the classification or
33 combination of classifications that best predicted each characteristic. Our results showed great
34 prediction ability of agronomic classification, which created the best models for predicting
35 agronomic (yield) and environmental (management, elevation) characteristics. We also
36 identified a strong prediction ability of combining two or all three classifications to predict
37 seven other grassland characteristics. However, grassland classifications did not predict most
38 of the ecological characteristics well. We can assume that phytosociological classification,
39 despite its mainstream use, predicts grassland characteristics less well than agronomic
40 classification. We recommend combining grassland classifications to improve rapid
41 prediction abilities. This study provides new knowledge useful for developing grassland
42 classifications which meet the needs of agronomists and naturalists.

43 Key words: phytosociology; agricultural type; functional type; management; biodiversity;
44 typology

45

46 1 Introduction

47 Worldwide, permanent grassland is the main terrestrial ecosystem, covering 26% of the world
48 land area (FAO, 2019) and 28% of UE total utilised agricultural area in 2013 (Eurostat, 2017).
49 Permanent grasslands provide diverse ecosystem goods and services of global importance,
50 such as forage production, species habitat, pollination, water purification, flood prevention
51 and mitigation of global warming through carbon storage (D'Ottavio et al., 2017). Provision
52 of these services is threatened by four main dangers: disappearance of grasslands (cultivation,
53 degradation), underuse (abandonment), agricultural intensification and climate change (Biró
54 et al., 2013; Muller et al., 1998; WallisDeVries et al., 2002; Young et al., 2005).

55 Conservation of grasslands and associated goods and services requires strong policy, but
56 grasslands differ and do not need equal protection. Indeed, agronomic and ecological services
57 provided by grasslands are complex consequences of management choices, environmental
58 constraints and botanical composition (Dumont et al., 2018; Michaud et al., 2012). Decision-
59 makers need simple tools such as grassland classifications to assess or predict the levels of
60 grassland services provided. At local scales, farmers and agronomists mainly need to predict
61 forage yield and quality, while naturalists are interested in ecological values of grasslands. At
62 regional, national and international scales, policy makers and administrators need to assess
63 trade-offs between economics and the environment, and to design best policies. To solve these
64 issues, naturalists and agronomists have developed different grassland classifications, using
65 divergent methods. These classifications simplify prediction of grassland characteristics,
66 which depend on the viewpoint and method of study.

67 Naturalists have used phytosociology since the early 20th century to classify vegetation
68 communities and assess habitat conservation. The development of phytosociological

69 classification required botanical relevés of all vascular species and a coefficient of abundance-
70 dominance for each species of each relevé (Braun-Blanquet, 1964). Phytosociology is the
71 mainstream method for classifying vegetation communities as it can be applied to all
72 ecosystems worldwide (e.g. Cheng et al., 2013; Rodríguez-Rojo et al., 2017, 2001; Setubal
73 and Boldrini, 2012), vegetation classes are arranged into a hierarchical system, and their
74 names follow scientific rules (Dengler et al., 2008). Once developed, the use of
75 phytosociological classifications requires only a complete botanical relevé and less than an
76 hour in each field. Phytosociology can also be used to predict agronomic characteristics, such
77 as forage quality, and environmental characteristics by using indicator species (Petrovic et al.,
78 2013); however, coefficients of abundance-dominance correlate only weakly with each
79 species' percentage of total grassland biomass (i.e. relative biomass), which can skew
80 assessment of agronomic characteristics (Daget and Poissonet, 1971; Pittarello et al., 2018).
81 Naturalists use phytosociology frequently worldwide; in the European Union, it is used to
82 assess habitat conservation and define priority habitats to protect faunal and floral
83 biodiversity.

84 Agronomists have developed agronomic classifications since the 1950s to quickly assess
85 forage yield and quality, as well as impacts of practices on forage production and the
86 environment (Michaud et al., 2013). Complete knowledge of botanical diversity is needed to
87 develop a classification because effects of each plant species on animal production depend on
88 its prevalence in an animal's diet (Diquélou et al., 2003), but classification approaches may
89 vary. Agronomic classifications provide accurate information about agronomic performances
90 of grasslands but are closely tied to the areas for which they were developed. Once developed,
91 agronomic classifications can often be used without botanical knowledge, but they do require
92 knowledge about agronomic practices, which needs to be collected when the classifications
93 are developed. Doing so requires time, but can begin a discussion between farmers and
94 agricultural advisors. Moreover, agronomic classifications can be used to predict ecological
95 characteristics such as species richness and pollinator value (Hulin et al., 2011; Launay et al.,
96 2011), but not all classifications have been developed for this purpose. Several agronomic
97 classifications have been developed, especially in western Europe, and they are used by a
98 wide range of decision makers, including agricultural advisors and farmers (Michaud et al.,
99 2013).

100 More recently, during the 1990s, ecologists attempted to explain the functioning of grassland
101 ecosystems using functional traits of plants (e.g. Lavorel and Garnier, 2002; Mouillot et al.,
102 2013; Violle et al., 2007). A functional trait is "any morphological, physiological or
103 phenological feature measurable at the individual level, ... without reference to the
104 environment or any other level of organization" (Violle et al., 2007). Agronomists applied this
105 knowledge to permanent grasslands to develop functional classifications, which are used to
106 predict grassland agronomic characteristics such as forage quality, earliness and management
107 flexibility. Dominant grass (Cruz et al., 2010; Durante et al., 2012) and forb species have been
108 classified by their functional traits, which are then used to classify grasslands. Functional
109 ecology has brought a fresh perspective to ecology and agronomy, improved prediction of
110 ecosystem goods and services, such as fodder production or carbon and nitrogen cycling
111 (Carol Adair et al., 2018; Lavorel, 2013), and been particularly useful for characterising
112 community response to environmental change (Lavorel and Garnier, 2002; Nock et al., 2016).
113 Unlike agronomic classifications, functional classifications are not restricted to the areas for
114 which they were developed (Cruz et al., 2010) and, once developed, do not require great
115 botanical knowledge: the relative biomass of the grass and forb species classified is sufficient
116 to classify the grassland and predict its agronomic characteristics. Developing functional
117 classifications, however, requires knowledge about functional traits of plant species. These

118 classifications are already used in France to help farmers manage grasslands (Carrère et al.,
119 2012).

120 The first objective of our study was to compare how phytosociological, agronomic and
121 functional classifications assess grassland characteristics, as they use different methods and
122 have different objectives. Phytosociological classification's main aim is to describe botanical
123 associations, while agronomic classification focuses on predicting forage yield and quality. In
124 comparison, functional classification predicts mechanisms (e.g. strategy for acquiring
125 nutrients, development of plant structure) that express themselves when exposed to
126 environmental variation. We hypothesised that each classification could predict certain
127 characteristics well: phytosociological for ecological characteristics, agronomic for agronomic
128 characteristics and functional for agronomic and environmental characteristics.

129 Classifications are used to predict grassland ecosystem goods and services, but usually only
130 one classification approach is used to do so. However, differences in classification objectives
131 lead to divergent methods for constructing classifications, which can cause classifications'
132 predictions of grassland characteristics to differ despite having similar prediction potentials.
133 From this perspective, the second objective of this study was to investigate prediction
134 capacities of combined grassland classifications. To our knowledge, few studies have
135 combined classification approaches. Carrère et al. (2012) combined phytosociological class,
136 agronomic management and functional class to study potential evolution from one grassland
137 class to another (e.g. due to modifying fertilisation or earliness of first use). Macedo et al.
138 (2010) compared the species prediction abilities of three classifications of dune plants:
139 phytosociological, dominant-strategy (Grime's CSR theory) and functional. They found that
140 these classifications predicted plant species similarly but differed in their predictions of plant
141 response to disturbance. However, Macedo et al. (2010) predicted ecological characteristics of
142 dune communities only, and neither of these two studies assessed the increase in prediction
143 accuracy provided by combinations of classifications. We hypothesised that although each
144 classification has strengths and weaknesses, combined classifications would create a new
145 classification that was more polyvalent and accurate at predicting grassland characteristics.

146 Using a sample of 250 permanent grasslands, we determined a phytosociological, agronomic
147 and functional class for each. We analysed the prediction accuracy of each of the three
148 classifications and explored whether combining two or all three of them improved prediction
149 of each of 16 ecological, agronomic and environmental characteristics. Finally, we
150 determined whether one classification alone or a combination of classifications could predict
151 all 16 of the characteristics well.

152

153 2 Materials and methods

154 2.1 Site


155 The dataset consisted of 250 grasslands in the Vosges Mountains (north-eastern France,
156 7000+ km²) (Fig. 1). Their elevations vary greatly (170-1424 m a.s.l.), as does their geology:
157 from limestone and sandstone in the north to plutonic volcanic rock in the south. The climate
158 is under oceanic and semi-continental influences, and can be polar at mountain summits.
159 According to climate normals (1981-2010), mean monthly temperatures vary from -4 to
160 +12°C, and mean annual precipitation varies greatly (700-2300 mm) under complex
161 latitudinal, longitudinal and elevational gradients (Ferrez et al., 2017). Finally, the Vosges
162 Mountains cover a large gradient of agricultural practices: 26 % of the grasslands were
163 exclusively mown, 38% were exclusively grazed, and nitrogen fertilisation ranged from 0-277

164 kg nitrogen/ha (69% organic (manure applied or directly deposited during grazing) and 31%
165 synthetic, by mass).

166 2.2 Attribution of grassland classes

167 Three studies were performed from 2001-2013 (Bayeur et al., 2013; Collectif, 2006; Launay
168 et al., 2011) of 250 permanent grasslands on commercial farms (Fig. 1). They focused on the
169 main homogeneous vegetation community of each grassland in areas 1000-10,000 m² in size
170 to avoid effects of paddock size on botanical diversity. One botanical relevé per grassland was
171 performed in the spring (peak biomass production), identifying all vascular botanical species
172 and their relative biomass (%) in 25 circles of 0.2 m² randomly placed in the homogeneous
173 vegetation community. Finally, homogeneous vegetation community were prospected to
174 search for species too rare to be found in the 25 samples.

175 353 species were observed in 250 grassland homogeneous vegetation communities. The mean
176 (\pm SD) number of species per community was 37.6 (\pm 8.7) [min-max = 16-65]. Thirty species
177 formed 80% of the total biomass, with a dominance of red fescue *Festuca rubra* L. (14.9%),
178 Yorkshire fog *Holcus lanatus* L. (9.1%), perennial ryegrass *Lolium perenne* L. (7.4%) and
179 sweet vernal grass *Anthoxanthum odoratum* L. (5.6%). 11 threatened or endangered species
180 were identified (seven vulnerable, two endangered and two critically endangered) according
181 to the IUCN Red List, distributed in 30 grasslands. We used this dataset to attribute an
182 agronomic, functional and phytosociological class to each of the 250 grasslands.



183
184 Figure 1. Location of the 250 botanical relevés performed in the Vosges Mountains, France,
185 from 2001-2013 (Bayeur et al., 2013; Collectif, 2006; Launay et al., 2011). Coordinate
186 system: RGF93-Lambert93

187

188

189 2.2.1 Agronomic class attribution

190 Previous studies identified 25 agronomic classes for homogenous grassland based on species
191 presence and relative biomass (Bayeur et al., 2013; Collectif, 2006). To this end, they
192 performed multivariate analyses of grassland composition: Correspondence Analysis (CA)
193 and Hierarchical Cluster Analysis (HCA) of species presence and Principal Component
194 Analysis (PCA) and HCA of relative biomass. Thus, by combining CA and PCA, they
195 attributed one agronomic class to each grassland based on the presence and relative biomass
196 of its botanical species (Table S1). The number of grasslands per agronomic class varied from
197 4-24 (Table 1).

198 This classification is called “agronomic” because its main aim is not to predict botanical
199 associations, but to use these associations to predict forage yield and quality.

200 2.2.2 Functional class attribution

201 Cruz et al. (2010) attributed one class (A, B, b, C and D) to 38 of the most common French
202 grasses (Poaceae), based on a combination of functional traits for leaves (dry matter content,
203 specific leaf area, longevity, resistance to breakage) and plants (flowering date, maximum
204 height) (Table S2). Then, Cruz et al. (2010) attributed functional classes to grasslands
205 according to the percentage of each grass class in them.

206 We adapted their method of functional class attribution to reduce the number of grassland
207 classes containing only one grassland. If one grass class represented more than 60% of
208 grassland grasses, the grassland was defined as that class (e.g. 76% of grasses of class A, 3%
209 of B and 21% of C yielded grassland class A), as proposed by Cruz et al. (2010). Otherwise,
210 grassland was defined by the two main grass classes, in decreasing order (e.g. 28% of A, 30%
211 of B, 10% of b and 32% of C yielded grassland class CB).

212 We then attributed one functional class to each permanent grassland based on the Cruz et al.
213 (2010) grass classifications and the relative biomass of these grasses in each grassland. Thus,
214 we attributed 20 functional classes, with 1-41 grasslands per class (Table 1). The mean (\pm SD)
215 percentage of Poaceae in relevés was $64.8 \pm 13.5\%$ [min-max = 24.0-96.7%], while that of
216 other families was $35.2 \pm 13.5\%$ [3.3-76.0%].

217 2.2.3 Phytosociological class attribution

218 Ferrez et al. (2017) developed a phytosociological classification for the entire Vosges
219 Mountain range using 1628 relevés performed from 1993-2015 following the Braun-Blanquet
220 method. They identified 35 phytosociological classes and selected the 22 of them that had the
221 highest appearance rates and most important agronomic and ecological roles in the Vosges
222 Mountains (Table S3). These 22 classes are based on 550 grassland relevés on commercial
223 farms.

224 Normally, phytosociological classes are attributed directly in the field. For this study,
225 however, we had to attribute phytosociological classes to the 250 grasslands *a posteriori*,
226 from a dataset of existing botanical relevés. Therefore, we developed a new key to determine
227 phytosociological classes. We used a statistical regression tree to design the new key, using
228 the complete set of 550 relevés (Ferrez et al. 2017) and the mvpart package of R software v.
229 3.4.2 (R Core Team, 2019; Therneau and Atkinson, 2014). The key, based on 48 botanical
230 genera, determines phytosociological classes easily. We verified the key’s percentage of good
231 classification for the dataset (550 grasslands) using a confusion matrix. According to the
232 confusion matrix, the key had an agreement of 81%.

233 Using this key and the presence of botanical genus that we observed in each grassland during
 234 the field sampling from 2001-2013, we attributed a phytosociological class to each of the 250
 235 studied grasslands. All of the following analyses were performed for these 250 grasslands.

236 Thus, we attributed 17 phytosociological classes to the 250 grasslands, with 3-61 grasslands
 237 per class (Table 1). As 22 classes could have been attributed at most, five classes were not
 238 observed in the set of grasslands.

239

240 Table 1. Number of grasslands (n = 250) attributed to 17 phytosociological classes, 25
 241 agronomic classes and 20 functional classes when attributing one phytosociological, one
 242 agronomic and one functional class to each grassland (Table S1, S2 and S3).

Phytosociological classes		Agronomic classes				Functional classes	
Phy_12	61	BV_02	24	VN_09	14	C	41
Phy_05	25	BV_01	17	VN_07	13	A	40
Phy_01	24	BV_10	13	VN_12	11	AC	38
Phy_02	21	BV_06	12	VN_13	11	CA	31
Phy_11	20	BV_07	11	VN_05	9	AB	26
Phy_20	20	BV_08	11	VN_08	9	Ab	15
Phy_07	14	BV_03	10	VN_14	9	B	10
Phy_09	12	BV_04	10	VN_01	8	BA	8
Phy_25	9	BV_09	10	VN_02	8	Cb	8
Phy_04	8	BV_11	6	VN_06	8	CB	6
Phy_06	8	BV_05	5	VN_03	6	CD	6
Phy_08	8			VN_11	6	BC	5
Phy_16	6			VN_10	5	Bb	4
Phy_18	5			VN_04	4	bB	3
Phy_29	3					DC	3
Phy_30	3					bC	2
Phy_31	3					AE	1
						b	1
						bA	1
						E	1

243

244

245 2.3 Grassland characteristics

246 We chose to analyse ecological, agronomic and environmental characteristics of grasslands.
247 Some of them were ecosystem services, which are benefits that people obtain from
248 ecosystems (Millennium Ecosystem Assessment, 2005); however, this definition is still
249 debated, and goods can be separated from services (Fisher et al., 2009). In addition, goods and
250 services can be delivered by both the ecosystem and human practices; for example, grassland
251 yield can be increased by applying synthetic fertilisers. Because we assessed a variety of
252 goods and services delivered by the ecosystem and by human practices, we referred to them
253 collectively as “grassland characteristics”.

254 We selected nine ecological characteristics to assess taxonomic diversity (species richness,
255 family richness, Shannon heterogeneity index, Simpson diversity index), functional diversity
256 (life form abundance, pollinator value, richness and weighted mean of the Ellenberg root
257 system depth index) and patrimonial functions (oligotrophic species richness, which can be
258 used as a proxy for species of high ecological value, threaten by the homogenization and
259 intensification of agricultural management). We selected three agronomic characteristics to
260 assess actual forage quantity (yield), potential forage quality and quantity (pastoral value) and
261 phenology (fodder earliness). Finally, we selected four environmental characteristics to assess
262 the sensitivity of classification to natural and human factors (Ellenberg humidity and fertility
263 indices, elevation and management). We used a variety of methods (e.g. calculation, farmer
264 interviews) to determine the 16 characteristics (Table 2, S4 and S5).

265 The botanical relevés of 250 grasslands and interviews of 52 farmers were performed in
266 previous studies, which were used to develop the agronomic classification (Bayeur et al.,
267 2013; Collectif, 2006; Launay et al., 2011). Botanical relevés allowed the collection of species
268 composition and relative biomass, which we entered into the e-FLORA-sys program
269 (Plantureux and Amiaud, 2010, 2009). This program calculated information at the vegetation
270 community scale: all ecological characteristics, as well as pastoral values and life form
271 diversity came from direct observation and calculation by e-FLORA-sys. Using a
272 questionnaire to avoid bias, farmers were interviewed to obtain information about yield and
273 grassland management. Knowledge from the interviews were extracted to deduce grassland
274 management (“mown”, “grazed” or a “mixture” of both) and estimate yield from the number
275 and mass of bales and from livestock stocking rates. On 14 grasslands, vegetation was
276 sampled and dried to measured yields without interviews. Finally, grassland elevation was
277 collected using BD ALTI dataset and QGIS software (IGN, 2016; QGIS Development Team,
278 2019).

279

280

281 Table 2. Methods for determining the 16 characteristics studied for permanent grasslands, in
282 ecological, agronomic and environmental categories.

Characteristic		Determination method
Ecological	Species richness	Number of botanical species observed in the grassland
	Oligotrophic species richness	Number of botanical species observed in the grassland with an Ellenberg nitrogen index of 1-3
	Family richness	Number of botanical families observed in the grassland
	Shannon heterogeneity index	$-\sum(Y_k \times \log_2 Y_k)$, where Y_k is the relative biomass of species k ($0 < Y_k \leq 1$)
	Simpson diversity index	$1 - (\sum Y_k^2)$, where Y_k is the relative biomass of species k ($0 < Y_k \leq 1$)
	Pollinator value	Relative biomass (% of total biomass) of entomophilous botanical species observed in the grassland
	Ellenberg root system depth index richness	Number of different Ellenberg root system depth indices observed in the grassland
	Ellenberg root system depth index weighted mean	$\sum(Y_k \times \text{individual root index})$, where Y_k is the relative biomass of species k ($0 < Y_k \leq 1$)
	Life form abundance	Relative biomass (% of total biomass) of each life form: grasses (Cyperaceae, Juncaceae, Liliaceae, Poaceae), legumes (Fabaceae) and forbs
Agronomic	Yield	Estimated from the number of bales and livestock stocking rate, or measured in the field
	Pastoral value	$\sum(Y_k \times \text{individual pastoral value})$, where Y_k is the relative biomass of species k ($0 < Y_k \leq 100$). For each species, pastoral value is estimated by expert opinion from potential yield, nutritive value, appetite and digestibility (Daget and Poissonet, 1971)
	Fodder earliness	Relative biomass (% of total biomass) of early, middle and late species, according to Cruz et al. (2010) and Theau et al. (2017)
Environmental	Ellenberg humidity index	$\sum(Y_k \times \text{individual humidity index})$, where Y_k is the relative biomass of species k ($0 < Y_k \leq 100$)
	Ellenberg fertility index	$\sum(Y_k \times \text{individual fertility index})$, where Y_k is the relative biomass of species k ($0 < Y_k \leq 100$)
	Elevation	Mean elevation of the grassland
	Management	“Mown”, “grazed”, or a “mixture” of both, obtained from farmer interviews

285 2.4 Statistical analyses

286 We first identified the classification or combination of classifications that predicted each
287 characteristic the best. We then assessed the quality of the best models to identify the most
288 polyvalent classification or combination of classifications for all 16 characteristics. The
289 specific method used depended on the grassland characteristic.

290 2.4.1 Modelling each characteristic with classifications

291 We used statistical model selection to calculate the probability that each classification alone
292 (phytosociological (P), agronomic (A) or functional (F)), or a combination of classifications,
293 yielded the best model, using R software. We modelled each characteristic as a function of
294 each classification or combinations of classifications. To this end, we generated a model
295 containing the main effects of all the classifications (A+P+F), and we then derived all
296 possible submodels, including an intercept-only model (null model) (Burnham and Anderson,
297 2002; Grueber et al., 2011). We then repeated this methodology from a model containing the
298 interactions of all the classifications (A×P×F). Thus, a set of 12 models could be created,
299 based on classifications alone and combinations of classifications. The sets of models
300 generated to predict each characteristic are presented in Table S6.

301 Combining agronomic and phytosociological classifications yielded 116 combinations (e.g.
302 agronomic class BV_01 with phytosociological class Phy_05 for the same grassland).
303 Similarly, we identified 136 combinations of agronomic and functional classifications, 93
304 combinations of phytosociological and functional classifications, and 202 combinations of all
305 three classifications. The large number of combinations of classifications demonstrated that
306 classifications differed in how they classified grasslands.

307 We modelled each continuous variable (species richness, oligotrophic species richness, family
308 richness, Shannon heterogeneity and Simpson diversity indices, pollinator value, Ellenberg
309 root system depth richness and root system depth index weighted mean, yield, pastoral value,
310 Ellenberg humidity and fertility indices and elevation) using generalised linear models from
311 the gamma family. We modelled each characteristics by 12 models.

312 We also fitted life form abundance and fodder earliness using generalised linear models.
313 However, as we used only one model for life form (grass, legume and forb) and one model for
314 earliness (early, middle and late), models were created from the binomial family. Here again,
315 we modelled each characteristics by 12 models.

316 Grassland management was a categorical variable, so we created multinomial logit models
317 using the mlogit package (Croissant, 2015). As logit models do not support interactions
318 (Croissant, 2015), we fitted only eight models to grassland management.

319 2.4.2 Identification of the best model

320 After creating the models, we selected the best one for each characteristic. Except for
321 grassland management models, the best model was that with the lowest second-order Akaike
322 Information Criterion (AICc), calculated using the MuMIn package and R software (Barton,
323 2018). Grassland management models were ranked according to the Akaike Information
324 Criterion (AIC) itself, calculated using the qpcR package and R software (Spiess, 2014),
325 because AICc cannot be calculated for logit models (Barton, 2018).

326 During model selection, an Akaike weight (hereafter, “weight”) is attributed to each model
327 tested. This weight is the probability that a given model is the best, and the sum of the weights
328 of all models tested equals 1 (Symonds and Moussalli, 2011). For each analysis, we compared

329 the weight of each model to those of other models: the closer a model's weight is to 1, the
330 better the model is. If only one model had a high weight (> 0.8), we assumed it was the best,
331 but if several models had similar weight, we examined them more closely. We verified the
332 quality of the model selection by ensuring that the null model had a weight close to zero
333 (Symonds and Moussalli, 2011).

334 2.4.3 Identification of a reliable polyvalent classification

335 Best model do not mean good model: the best among poor quality models remain a poor
336 quality model. Moreover, the best model for a given characteristic may not be the best model
337 for multiple characteristics together. To solve these two issues, we calculated the quality of
338 each best model. First, model quality informed about the accuracy of each characteristics
339 model. Then we could compare the accuracy of models selected for several characteristics: if
340 a model would be selected as best model for several characteristics and its accuracy would be
341 sufficient, we could assume it is a polyvalent and reliable model.

342 AIC and AICc indicate model quality, but as they are relative to the dataset for each
343 characteristic we could not use them to compare best models for multiple characteristics
344 (Burnham and Anderson, 2002). We used each best model to predict all 16 characteristics and
345 calculated a goodness of fit among the predictions using pseudo- R^2 , because R^2 cannot be
346 calculated for generalised linear models (Tjur, 2009).

347 For generalised linear models for the gamma family, we calculated Cox and Snell pseudo- R^2
348 (Cox and Snell, 1989), which is the easiest to use because it expresses R^2 in terms of log-
349 likelihood (Tjur, 2009). Because the Cox and Snell pseudo- R^2 is not optimal for generalised
350 linear models from the binomial family or for multinomial logit models (Cox and Snell,
351 1989), we calculated McFadden's pseudo- R^2 for these models, which is also based on the
352 likelihood of the model (Hoetker, 2007). We used the DescTools package and R software to
353 calculate pseudo- R^2 (Signorell, 2017). Finally, we calculated generalised variance-inflation
354 factors to check for collinearity among models, using the car package (Fox and Sanford,
355 2011). We assumed that models with pseudo- $R^2 > 0.5$ had a satisfactory goodness of fit.

356

357 3 Results

358 3.1 Identification of the best classification

359 The model with the highest weight for 15 of the 16 grassland characteristics contained one or
360 more classifications, as root system depth richness was predicted best by the null model (Fig.
361 2). Among these 15 models, four of them had a weak weight < 0.8 and required a closer look
362 to identified best models: Shannon heterogeneity index (0.66), pollinator value (0.52),
363 Ellenberg root system index weighted mean (0.73) and Ellenberg fertility index (0.59).

364 One classification alone was sufficient to predict four characteristics the best. Three
365 characteristics were predicted best by agronomic classification alone (A): yield (agronomic),
366 grassland management (environmental) and elevation (environmental). One characteristic was
367 predicted best by phytosociological classification alone (P): species richness (ecological).

368 For two characteristics, the model with the highest weight contained only one classification,
369 but the weight was not high enough to select it as the best model. For Shannon heterogeneity
370 index, phytosociological classification (P) alone had the highest weight (0.66), but it did not
371 differ enough from the second highest weight (0.25), which was for the main effects of
372 phytosociological and functional classifications (P+F). Similarly, for pollinator value, weights

373 of agronomic classification (A) and the null model (0.52 and 0.35, respectively) were too
374 similar to identify one best model.

375 The main effects of two or three classifications predicted six characteristics the best:
376 agronomic and phytosociological (A+P) for family richness; agronomic and functional (A+F)
377 for Ellenberg humidity index; phytosociological and functional (P+F) for Simpson
378 heterogeneity index and pastoral value; and all three classifications (A+P+F) for fodder
379 earliness. Ellenberg root system index weighted mean highest weight was associated to the
380 P+F model, but was <0.8. However, the second highest weight was different enough to
381 assume Ellenberg root system index weighted mean was best model by the main effects of
382 phytosociological and functional classifications.

383 On the other hand, weights of the A+P+F and A+P models for the Ellenberg fertility index
384 were too similar (0.59 and 0.41, respectively), we could not identify which one was best.
385 Finally, two best models contained an interaction: that between agronomic and
386 phytosociological classifications (A×P) for life form abundance, and between
387 phytosociological and functional classifications (P×F) for oligotrophic species richness.


388 The highest weights equalled 1.0 for seven characteristics, 0.7-1.0 for six characteristics, and
389 less than 0.7 for the Shannon heterogeneity index, pollinator value and Ellenberg fertility
390 index. Null model weights equalled zero, except for those for pollinator value (0.35) and root
391 system depth richness (1.0).

392 3.2 Identification of a reliable and polyvalent classification

393 The goodness of fit (pseudo-R²) varied greatly among the best models for characteristics
394 (0.20-0.88) (Fig. 2). The goodness of fit of the model for the Ellenberg root system index
395 weighted mean equalled zero, due to the selection of the null model. Models for agronomic
396 and environmental characteristics had high weights and/or quality. In contrast, most models
397 for ecological characteristics had low quality, except for life form abundance, oligotrophic
398 species richness and, to a lesser extent, family richness (pseudo-R² = 0.87, 0.83 and 0.54,
399 respectively). We identified ten grassland characteristics whose models had goodness of fit >
400 0.5 (Fig. 3): oligotrophic species richness, family richness and life form abundance
401 (ecological); yield, pastoral value and fodder earliness (agronomic); and Ellenberg indices for
402 fertility and humidity, elevation and management (environmental). Unlike these ten grassland
403 characteristics, the other six, all ecological, had best models of low quality despite having
404 high weights: species richness, Shannon heterogeneity index, Simpson diversity index,
405 pollinator value and richness and weighted mean of Ellenberg root system depth


406 We could not identify one polyvalent and reliable classification or combination of
407 classifications that could predict ecological, agronomic and environmental characteristics
408 well. Only agronomic classification could predict characteristics the best when alone, with
409 high quality models. Moreover, most of the characteristics predicted well were predicted
410 totally or partially by agronomic classification. The combination of phytosociological and
411 functional classifications predicted oligotrophic species richness and pastoral value well.

412



413

414 Figure 2. Akaike weights and goodness of fit (pseudo-R²) of the 16 grassland characteristics
 415 modelled using agronomic (A), phytosociological (P) or functional (F) classification, the main
 416 effects of two or all three of them (A+P, A+F, P+F and A+P+F), their interactions (A×P,
 417 A×F, P×F and A×P×F) and a null model (Null). The term “n.c.” (“not considered”) indicates
 418 characteristics that could not be modelled with interactions. The goodness of fit shown is that
 419 of the model with the highest weight for each grassland characteristic.



420

421 Figure 3. Representation of which phytosociological, agronomic and functional classifications
 422 (or combinations of them) were needed to predict ten grassland characteristics well (goodness
 423 of fit > 0.5). Red, green and blue text identify ecological, agronomic and environmental
 424 characteristics, respectively.

425

426 4 Discussion

427 4.1 Prediction of grassland characteristics

428 4.1.1 Quality of prediction models

429 Four of the six characteristics whose models had goodness of fit < 0.5 involved species or
 430 trait richness and equitability. This poor prediction of ecological characteristics is not
 431 surprising for agronomic and functional classifications, which were not designed to predict
 432 them and rely mainly on relative biomass rather than presence of species. The best models for
 433 each of these characteristics are built in part on phytosociological classification, except those
 434 for pollinator value and Ellenberg root system index weighted mean. However, the poor
 435 quality of these ecological characteristics highlights the lack of a relationship between
 436 ecological characteristics and phytosociological classification, which is more surprising.
 437 While grasslands in the same phytosociological class have species in common (a
 438 “characteristic combination”), they also have some different species among them, which gives
 439 them different values of species richness and may help explain our result. Bias in relevés
 440 could also be a reason: Chytrý (2001) showed that botanists increase the area sampled in
 441 species-poor grasslands to increase the number of species, which could bias attribution of
 442 class and calculation of richness.

443 Despite its common use and importance for estimating total biodiversity, species richness is
 444 not a key characteristic to predict because of its high variability in similar grasslands
 445 (Brunbjerg et al., 2018). These limits to biodiversity measurements have led researchers and
 446 decision-makers to use diversity indices such as the Shannon heterogeneity index and

447 Simpson diversity index (Bello et al., 2010; Mauchamp et al., 2014). In our study, we could
448 not identify models that predicted diversity indices well, but ecological indices were predicted
449 best by phytosociological classification alone or a combination of phytosociological and
450 functional classifications. These indices provide different types of information: the Shannon
451 heterogeneity index describes species equitability, while the Simpson diversity index
452 describes both species equitability and richness. One way to predict grassland plant diversity
453 well could be to use disturbance intensity: Vujnovic et al. (2002) showed that intermediate
454 disturbance by grazing, trampling or soil management maximised grassland biodiversity,
455 while Rodríguez-Rojo et al. (2017) observed that management intensity had more influence
456 on grassland biodiversity than management practices. However, biodiversity measurement,
457 through species richness or ecological indices, is not necessary a useful indicator. Indeed,
458 these measurements do not take into account vegetation characteristics as proportion of rare
459 species, which is not correlated to total richness (Pykälä et al., 2005). Assessment of
460 biodiversity should be implemented with proxy for species of high ecological value like
461 oligotrophic species richness (Michaud et al., 2012; Muller, 2002). In the present study,
462 oligotrophic species richness was predicted best by the combination of phytosociological and
463 functional classifications.

464 Among ecological characteristics, only pollinator value and the weighted mean of Ellenberg
465 root system depth needed species relative biomass in their calculations. However, they are
466 based on traits (pollinator attractiveness traits and root system depth, respectively) not
467 included in the functional classification studied nor in other classifications. Agronomic
468 classification has been used to estimate pollinator value (Bayeur et al., 2013; Hulin et al.,
469 2011; Launay et al., 2011). Our study confirms that agronomic classification predicts
470 pollinator value the best, but its low model quality (weight = 0.52, pseudo-R² = 0.20) raises
471 questions about the reliability of the prediction. Beyond our study, phytosociological
472 classification may help predict pollinator species or traits, which are particularly related to
473 flower diversity (Hegland and Totland, 2005; Warzecha et al., 2017). However,
474 entomophilous botanical species relative biomass assesses a potential pollinator value at
475 grassland scale. To assess a real pollinator service, flowering time and cutting/grazing dates
476 should be studied because they express the real abundance of flowers (Kleijn et al., 2001).
477 The landscape also influences pollinator diversity and abundance strongly, but it is not
478 included in grassland classifications (Sutter et al., 2017). Finally, our modelling of root
479 system depth is similar to the study of Oram et al. (2018), which found that mean root system
480 depth was related to other characteristics but root system depth richness was not.

481 4.1.2 Grassland characteristics predicted by one classification alone

482 Three grassland characteristics – elevation, management and yield – were predicted best by
483 agronomic classification alone, which was the only classification that could predict grassland
484 characteristics well by itself (Fig. 3). Management and elevation predict species composition
485 well (Rodríguez-Rojo et al., 2017; van Oijen et al., 2018), and agronomic classification
486 predicted management and elevation well with high weights and model qualities. We can
487 assume that botanical associations developed from taxonomic relative biomass are good
488 indicators of elevation and might be indicators of climate more generally.

489 Prediction of yield is crucial, especially for farmers. As agronomic classifications are
490 developed mainly to predict agronomic characteristics, we expected a strong relationship
491 between yield and agronomic classification. Functional classifications are also developed to
492 predict grassland yields, and this ability was confirmed by Lavorel and Garnier (2002), who
493 identified strong relationships between nutrient-response traits and traits determining yields.
494 Plasticity in trait expression (at the species scale) and trait co-occurrence (at the community

495 scale) of grasses and forbs need to be better understood to improve predictions of functional
496 classification (Roscher et al., 2018). Yield can be estimated easily using Ellenberg fertility
497 index (Hill and Carey, 1997) or by more complex models based on functional traits, life form
498 abundance (Michaud et al., 2015) and phytosociological classes (Magiera et al., 2017).
499 Finally, field measurement also estimates grassland yield well but requires more time and
500 equipment than using grassland classifications or Ellenberg indices (Ni, 2004).

501 4.1.3 Combining classifications: a solution?

502 To our knowledge, our study is the first to compare three classification approaches using the
503 same relevés. Carrère et al. (2012) created tool based on phytosociological, agronomic and
504 functional classifications but did not compare the quality of predictions provided by each
505 classification to those of the combination of two or three classifications. Macedo et al. (2010)
506 compared the prediction abilities of three classifications but focused only on predicting
507 ecological characteristics of dune communities.

508 In our study, seven of the characteristics studied were predicted better by combinations of
509 classifications than by one classification alone (Fig. 3): oligotrophic species richness, family
510 richness and life form abundance (ecological), pastoral value and fodder earliness
511 (agronomic) and Ellenberg humidity and fertility indices (environmental). Notably, among the
512 goodness of fit >0.8 , three out of four models required combination of classifications: life
513 form abundance (0.87), oligotrophic species richness (0.83) and Ellenberg fertility index
514 (0.8). Moreover, despite its weak prediction capacities when used alone, phytosociological
515 classification is part of these three best models.

516 Both fertility and humidity indices were predicted well (high weight and goodness of fit) by
517 combining classifications. Humidity and fertility indices have been linked to elevation (de
518 Almeida Campos Cordeiro and Neri, 2019; Pittarello et al., 2018; Sevruk, 1997) but also to
519 taxonomic (Critchley et al., 2002; Melts et al., 2018; Wang et al., 2007) and functional (Čop
520 et al., 2009; Schellberg, 1999; Suding et al., 2005) diversity. We also used the fertility index
521 to calculate oligotrophic species richness. However, fertility needed all three classifications to
522 be predicted well, while oligotrophic species richness needed only phytosociological and
523 functional classifications and was predicted better. This difference could have been due to the
524 method used: the fertility index is based on relative biomass, unlike oligotrophic species
525 richness. Ellenberg fertility index is influenced by fertilizer application, but is also dependant
526 of soil properties like the cation exchange capacity. Thus, the prediction of Ellenberg fertility
527 index and oligotrophic species richness can bring information about presence of species of
528 high ecological value, but cannot be used to assess the impact of fertilization at scales larger
529 than grassland community.

530 Life form abundance, as well as root system depth, was not included in previous
531 classifications. The latter was not predicted well, but the life form abundance was predicted
532 well by an interaction between classifications ($A \times P$). This result is particularly interesting
533 because it shows that grassland classifications can predict characteristics beyond those for
534 which they were developed. Moreover, family richness was predicted best by a similar
535 combination of classifications: the main effects of agronomic and phytosociological
536 classifications ($A+P$). These characteristics are related by the method of calculation. Indeed,
537 life forms are based on botanical families: legumes belong to Fabaceae, while grasses belong
538 to Cyperaceae, Juncaceae, Liliaceae and Poaceae. This result could highlight a correlation
539 between richness and relative biomass of families. Grassland management influences life
540 form diversity strongly: mowing increases grass and forb relative biomass, while grazing
541 selects rosettes and creeping plants (Gaujour et al., 2012; Lanta et al., 2009). However,

542 intensification also selects rosettes and creeping plants (Gaujour et al., 2012), and livestock
543 type also has a strong influence on life form diversity (Tóth et al., 2018). These previous
544 studies show that knowledge about grassland management could improve prediction of life
545 form abundance; however, Craine et al. (2001) identified correlations between leaf traits and
546 life form but no correlation between intensification and life form. The prediction of life form
547 abundance can be useful for agronomists and naturalists. Indeed, grassland with highest life
548 form diversity produce higher yields (Bullock et al., 2006), and allow heavier livestock live
549 weight than grassland with poor life form diversity (Grace et al., 2019). In the present study,
550 the goodness of fit of life form abundance (0.87) was higher than goodness of fit of yield
551 (0.51) and fodder quality (0.60). Finally, life form abundance also determines the resistance of
552 botanical species to drought: life form of weak abundance have more probabilities to
553 disappear (Tilman and Haddi, 1992). Thus, life form abundance could be an important
554 characteristic for naturalist who want to assess the risk of extinction of grassland botanical
555 species.

556 It was particularly surprising that phytosociological and functional classifications combined
557 (P+F) predicted pastoral value the best, because i) yield, part of the pastoral value equation, is
558 predicted well by agronomic classification, and ii) calculating the pastoral value requires the
559 pastoral value and relative biomass of each species, which is provided by agronomic
560 classification (Daget and Poissonet, 1971). Functional traits (Tasset et al., 2019) and
561 phytosociology (Petrovic et al., 2013) can be used to predict forage quality, but our result
562 shows that combining them improves predictions. Phytosociology alone may not be sufficient
563 to predict pastoral value due to the inclusion of different facies in the same phytosociological
564 class (Bagella and Roggero, 2004). Prediction of pastoral value could be improved by using
565 Ellenberg fertility index and environmental indicators such as elevation and slope (Bagella
566 and Roggero, 2004; Pittarello et al., 2018).

567 Forage earliness is key information in grassland management, because peak forage quality is
568 related to plant phenology (Pontes et al., 2007). Moreover, coupled with cutting or grazing
569 dates, earliness could also be used to assess pollination service (Kleijn et al., 2001). Climate is
570 the main factor that influences flowering time: temperature (Kudernatsch et al., 2008) and, to
571 a lesser extent, precipitation (Hovenden et al., 2008) create annual variability. We used degree
572 days to be free from annual temperature variability. Generally, alpine vegetation needs fewer
573 heat units to initiate flowering (Heide, 1994), which could create a fodder earliness gradient
574 according to elevation. Management, expressed indirectly through botanical composition,
575 creates differences in earliness of French grasslands, but differences are also perceptible
576 among species from the same community (Ansquer et al., 2009). We conclude that the three
577 classifications are essential to predict fodder earliness: while the functional classification is
578 based in part on grass earliness, phytosociological and agronomic classifications could
579 provide information about other species.

580 4.2 Limits of the study and perspectives

581 It may be possible to extrapolate results of this study. From the viewpoint of environmental
582 conditions, our study focused on 250 permanent grasslands of the French Vosges Mountains
583 but covered wide gradients of climates, elevations, soils and agricultural practices. From the
584 viewpoint of vegetation, we found 17 phytosociological classes out of the 35 identified by
585 botanists in the Vosges Mountains (Ferrez et al., 2017). However, these 17 classes are the
586 most representative of commercial farms according to expert opinion and can be found in
587 many other lowlands and low-elevation mountains in western Europe. Finally, the functional
588 classification studied can also be used in a wide range of grasslands, as it is representative of
589 western-European grassland Poaceae (Cruz et al., 2010).

590 All grassland classifications are based in part on arbitrary choices, either of the classification
591 criteria (here, taxonomic richness, taxonomic relative biomass, or functional relative biomass)
592 or the number of grassland classes. The method used to attribute functional classes provided
593 quick and easy classification, but it can also be restrictive and attribute different functional
594 classes to similar grasslands. This method also creates several functional classes containing
595 few grasslands: seven classes contained less than five grasslands. Moreover, the functional
596 classification focused only on Poaceae, even though other families represent a relatively large
597 percentage of each grassland (mean \pm SD = $35 \pm 14\%$), which could lead to lower accuracy in
598 measuring response traits if Poaceae respond differently to disturbances than other plant
599 families. A new functional classification based not only on Poaceae but also on
600 dicotyledonous species may improve prediction abilities of functional classification. Finally,
601 including new classification approaches should also improve knowledge about predicting
602 grassland characteristics. Phytosociology diverges into two approaches: Braun-Blanquet and,
603 more recently, numerical (Dengler et al. 2008). In the latter, relevés are grouped into classes
604 using statistics, based on presence/absence (as in the Braun-Blanquet approach) or relative
605 biomass (as in agronomic classifications). Classes are not phytosociological classes from the
606 literature but are instead free from the subjective concept of the vegetation unit.

607 Of the 16 characteristics studied, five were calculated using Ellenberg indices, whose use may
608 raise some questions. Indeed, some indices are estimated rather than measured, species
609 response to the environment depends on location and plant stage, and the strength of
610 correlations between indices and field measurements diverges among studies (Diekmann,
611 2003; Schaffers and Sýkora, 2000). Nevertheless, Ellenberg weighted mean indices are
612 considered reliable (Diekmann, 2003).

613 In a more extensive study, we could have analysed the ability of grassland classifications to
614 predict other characteristics. Classifications have been used to predict organoleptic
615 characteristics of animal products (Diquélou et al., 2003) and soil carbon stocks (Hulin et al.,
616 2011), but other services could be studied, such as maintenance of genetic diversity,
617 regulation of water flows and landscape heritage (D'Ottavio et al., 2017). Moreover, a more
618 complete dataset of grassland characteristics could improve statistical analyses: the lack of
619 homogeneity of grasslands per class might induce statistical bias, and it is important to assess
620 the quality of new models on an external dataset.

621 Our study raises questions about the use of phytosociological classification alone to identify
622 habitats of interest in Europe. Indeed, we demonstrated that ecological characteristics remain
623 difficult to predict and, above all, that phytosociological classification alone does not predict
624 grassland characteristics well. However, identification of ecological characteristics could lead
625 to conservation of key grasslands, while identification of agronomic characteristics could
626 improve distribution of economic incentives for ecologically valuable grasslands.
627 Characteristics can be predicted more accurately by combining classifications, as in this
628 study, but also by using other methods. Indicator species or life traits such as life forms or
629 flower colour are also easy to use but might be less accurate or more time consuming (Arnold
630 et al., 2009; Ellenberg et al., 1992). Finally, models can consider soil and weather variability,
631 but such models are often difficult for non-specialists to use and do not predict agronomic and
632 ecological characteristics at the same time (van Oijen et al., 2018).

633

634 5 Conclusion

635 Our study focused on the ability of three classification approaches to predict 16 grassland
636 characteristics. Our results show that agronomic classification has great potential to predict

637 three agronomic and environmental characteristics, which demonstrates that it meets the
638 needs of farmers and agronomists well. Although phytosociological classification is designed
639 to assess habitat conservation, it failed to predict the studied ecological characteristics well.
640 Functional classification appears promising, but it was developed more recently and may
641 suffer from knowledge gaps. As ecological characteristics remain especially difficult to
642 predict, naturalists and policy makers should use tools other than phytosociological
643 classification alone to predict them. Indicator species or life traits such as life forms or flower
644 colour are also easy to use but might be more time consuming than a phytosociological relevé.
645 To identify trade-offs between economics and ecology, it may be important to combine
646 grassland classifications: our results show that combinations of phytosociological, agronomic
647 and/or functional classification could accurately predict seven ecological, agronomic and
648 environmental characteristics. Finally, models may improve prediction of characteristics: they
649 can consider soil and weather variability but are often difficult for non-specialists to use and
650 do not predict agronomic and ecological characteristics at the same time.

651

652 Acknowledgments

653 The authors acknowledge the Northern Vosges Regional Nature Park, the Regional Natural
654 Park Ballons des Vosges and the Franche-Comté National Botanical Conservatory for their
655 contribution to the botanical relevés and to the development of agronomic and
656 phytosociological classifications. The authors also gratefully thank MéliSSa Berthet, Christof
657 Neumann and Jean Villerd for their statistical support. This project was funded by the
658 European Regional Development Fund, the Agence de l'eau Rhin-Meuse, the Fonds National
659 d'Aménagement et de Développement du Territoire Massif des Vosges and the Région
660 Grand-Est.

661

662 References

- 663 Ansquer, P., Al Haj Khaled, R., Cruz P, P., Theau, J.P., Therond, O., Duru, M., 2009.
664 Characterizing and predicting plant phenology in species-rich grasslands. *Grass and*
665 *Forage Science* 64, 57–70. <https://doi.org/10.1111/j.1365-2494.2008.00670.x>
- 666 Arnold, S.E.J., Le Comber, S.C., Chittka, L., 2009. Flower color phenology in European
667 grassland and woodland habitats, through the eyes of pollinators. *Israel Journal of*
668 *Plant Sciences* 57, 211–230. <https://doi.org/10.1560/IJPS.57.3.211>
- 669 Bagella, S., Roggero, P.P., 2004. Integrating phytosociological and agronomic analysis to
670 support the sustainable management of Mediterranean grasslands. *Fitosociologia* 41,
671 101–107.
- 672 Barton, K., 2018. MuMIn: Multi-Model Inference.
- 673 Bayeur, C., Kleiber, F., L'Hospitalier, M., Loridat, F., Plantureux, S., 2013. Typologie des
674 prairies permanentes Vosges du Nord et Vosges Mosellanes : guide technique 2013.
- 675 Bello, F. de, Lavergne, S., Meynard, C.N., Lepš, J., Thuiller, W., 2010. The partitioning of
676 diversity: showing Theseus a way out of the labyrinth. *Journal of Vegetation Science*
677 21, 992–1000.
- 678 Biró, M., Czúcz, B., Horváth, F., Révész, A., Csatári, B., Molnár, Z., 2013. Drivers of
679 grassland loss in Hungary during the post-socialist transformation (1987–1999).
680 *Landscape Ecology* 28, 789–803. <https://doi.org/10.1007/s10980-012-9818-0>
- 681 Braun-Blanquet, J., 1964. *Pflanzensoziologie*. Springer Vienna, Vienna.
682 <https://doi.org/10.1007/978-3-7091-8110-2>

683 Brunbjerg, A.K., Bruun, H.H., Dalby, L., Fløjgaard, C., Frøslev, T.G., Høye, T.T., Goldberg,
684 I., Læssøe, T., Hansen, M.D.D., Brøndum, L., Skipper, L., Fog, K., Ejrnæs, R., 2018.
685 Vascular plant species richness and bioindication predict multi-taxon species richness.
686 *Methods in Ecology and Evolution* 9, 2372–2382. [https://doi.org/10.1111/2041-](https://doi.org/10.1111/2041-210X.13087)
687 [210X.13087](https://doi.org/10.1111/2041-210X.13087)

688 Bullock, J.M., Pywell, R.F., Walker, K.J., 2006. Long-term enhancement of agricultural
689 production by restoration of biodiversity: Restoring biodiversity enhances production.
690 *Journal of Applied Ecology* 44, 6–12. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2664.2006.01252.x)
691 [2664.2006.01252.x](https://doi.org/10.1111/j.1365-2664.2006.01252.x)

692 Burnham, K.P., Anderson, D.R., 2002. Model selection and multimodel inference: a practical
693 information-theoretic approach, 2nd ed. ed. Springer, New York.

694 Carol Adair, E., Hooper, D.U., Paquette, A., Hungate, B.A., 2018. Ecosystem context
695 illuminates conflicting roles of plant diversity in carbon storage. *Ecology Letters* 21,
696 1604–1619. <https://doi.org/10.1111/ele.13145>

697 Carrère, P., Seytre, L., Piquet, M., Landrieux, J., Rivière, J., Chabalier, C., Orth, D., 2012. A
698 multifunctional typology of grassland in AOP-certified dairy systems in the Massif
699 Central combining an agronomic and ecological approach. *Fourrages* 209, 9–21.

700 Cheng, Y., Kamijo, T., Tsubo, M., Nakamura, T., 2013. Phytosociology of Hulunbeier
701 grassland vegetation in Inner Mongolia, China. *Phytocoenologia* 43, 41–51.
702 <https://doi.org/10.1127/0340-269X/2013/0043-0540>

703 Chytrý, M., 2001. Phytosociological data give biased estimates of species richness. *Journal of*
704 *Vegetation Science* 12, 441–444. <https://doi.org/10.1111/j.1654-1103.2001.tb00190.x>

705 Collectif, 2006. Le Massif Vosgien : Typologie des Prairies Naturelles.

706 Čop, J., Vidrih, M., Hacin, J., 2009. Influence of cutting regime and fertilizer application on
707 the botanical composition, yield and nutritive value of herbage of wet grasslands in
708 Central Europe: Management of wet grasslands. *Grass and Forage Science* 64, 454–
709 465. <https://doi.org/10.1111/j.1365-2494.2009.00713.x>

710 Cox, D.R., Snell, E.J., 1989. *The Analysis of Binary Data*, 2nd ed., Chapman and Hall. ed.
711 London.

712 Craine, J.M., Froehle, J., Tilman, D.G., Wedin, D.A., Chapin, III, F.S., 2001. The
713 relationships among root and leaf traits of 76 grassland species and relative abundance
714 along fertility and disturbance gradients. *Oikos* 93, 274–285.
715 <https://doi.org/10.1034/j.1600-0706.2001.930210.x>

716 Critchley, C.N.R., Chambers, B.J., Fowbert, J.A., Sanderson, R.A., Bhogal, A., Rose, S.C.,
717 2002. Association between lowland grassland plant communities and soil properties.
718 *Biological Conservation* 105, 199–215.

719 Croissant, Y., 2015. Multinomial logit model.

720 Cruz, P., Theau, J.P., Lecloux, E., Jouany, C., Duru, M., 2010. Functional typology of
721 perennial forage grasses : a classification based on several characteristics. *Fourrages*
722 201, 11–17.

723 Daget, P., Poissonet, J., 1971. Une méthode d'analyse phytosociologique des prairies.
724 *Annales Agronomiques* 22, 5–41.

725 de Almeida Campos Cordeiro, A., Neri, A.V., 2019. Spatial patterns along an elevation
726 gradient in high altitude grasslands, Brazil. *Nordic Journal of Botany* 37.
727 <https://doi.org/10.1111/njb.02065>

728 Dengler, J., Chytrý, M., Ewald, J., 2008. Phytosociology, in: *Encyclopedia of Ecology*.
729 Elsevier, pp. 2767–2779. <https://doi.org/10.1016/B978-008045405-4.00533-4>

730 Diekmann, M., 2003. Species indicator values as an important tool in applied plant ecology –
731 a review. *Basic and Applied Ecology* 4, 493–506. [https://doi.org/10.1078/1439-1791-](https://doi.org/10.1078/1439-1791-00185)
732 [00185](https://doi.org/10.1078/1439-1791-00185)

733 Diquélou, S., Leconte, D., Simon, J.C., 2003. Diversité floristique des prairies permanentes de
734 Basse Normandie (synthèse des travaux antérieurs). *Fourrages* 173, 3–22.

735 D’Ottavio, P., Francioni, M., Trozzo, L., Sedić, E., Budimir, K., Avanzolini, P., Trombetta,
736 M.F., Porqueddu, C., Santilocchi, R., Toderi, M., 2017. Trends and approaches in the
737 analysis of ecosystem services provided by grazing systems: A review. *Grass and*
738 *Forage Science* 73, 15–25. <https://doi.org/10.1111/gfs.12299>

739 Dumont, B., Ryschawy, J., Duru, M., Benoit, M., Chatellier, V., Delaby, L., Donnars, C.,
740 Dupraz, P., Lemauviel-Lavenant, S., Méda, B., Vollet, D., Sabatier, R., 2018. Review:
741 Associations among goods, impacts and ecosystem services provided by livestock
742 farming. *animal* 1–12. <https://doi.org/10.1017/S1751731118002586>

743 Durante, M., Lezana, L., Massa, E., Figari, M., Lezama, F., Jaurena, M., Cruz, P., 2012. A
744 first attempt to classify in functional groups grasses of Entre Rios (Argentina) and
745 Uruguay. Presented at the International Symposium on Integrated Crop-Livestock
746 Systems, Porto Alegre, Brasil, p. 4.

747 Ellenberg, H., Weber, H.E., Düll, R., Wirth, V., Werner, W., Paulissen, D., 1992. Zeigerwerte
748 von Pflanzen in Mitteleuropa, *Scripta Geobotanica*.

749 Eurostat, 2017. Permanent grassland: number of farms and areas by agricultural size of farm
750 (UAA) and size of permanent grassland area.

751 FAO, 2019. FAOSTAT - Land Cover [WWW Document]. Food and Agriculture
752 Organization of the United Nations\$. URL <http://www.fao.org/faostat/en/#data/LC>
753 (accessed 7.29.19).

754 Ferrez, Y., Cholet, J., Dor, J.-C., Dupont, F., Froehlicher, T., Giovannacci, L., Hennequin, C.,
755 Laumond, J.-S., L’Hospitalier, M., Nguéfacq, J., Simler, N., Voirin, M., 2017. Guide
756 phytosociologique des prairies du massif des Vosges et du Jura alsacien. Autechaux,
757 France.

758 Fisher, B., Turner, R.K., Morling, P., 2009. Defining and classifying ecosystem services for
759 decision making. *Ecological Economics* 68, 643–653.
760 <https://doi.org/10.1016/j.ecolecon.2008.09.014>

761 Fox, J., Sanford, W., 2011. *An R Companion to Applied Regression*, Second Edition.

762 Gaujour, E., Amiaud, B., Mignolet, C., Plantureux, S., 2012. Factors and processes affecting
763 plant biodiversity in permanent grasslands. A review. *Agronomy for Sustainable*
764 *Development* 32, 133–160. <https://doi.org/10.1007/s13593-011-0015-3>

765 Grace, C., Lynch, M.B., Sheridan, H., Lott, S., Fritch, R., Boland, T.M., 2019. Grazing
766 multispecies swards improves ewe and lamb performance. *Animal* 13, 1721–1729.
767 <https://doi.org/10.1017/S1751731118003245>

768 Grueber, C.E., Nakagawa, S., Laws, R.J., Jamieson, I.G., 2011. Multimodel inference in
769 ecology and evolution: challenges and solutions: Multimodel inference. *Journal of*
770 *Evolutionary Biology* 24, 699–711. <https://doi.org/10.1111/j.1420-9101.2010.02210.x>

771 Hegland, S.J., Totland, Ø., 2005. Relationships between species’ floral traits and pollinator
772 visitation in a temperate grassland. *Oecologia* 145, 586–594.
773 <https://doi.org/10.1007/s00442-005-0165-6>

774 Heide, O.M., 1994. Control of flowering and reproduction in temperate grasses. *New*
775 *Phytologist* 128, 347–362. <https://doi.org/10.1111/j.1469-8137.1994.tb04019.x>

776 Hill, M.O., Carey, P.D., 1997. Prediction of yield in the Rothamsted Park Grass Experiment
777 by Ellenberg indicator values. *Journal of Vegetation Science* 8, 579–586.
778 <https://doi.org/10.2307/3237210>

779 Hoetker, G., 2007. The use of logit and probit models in strategic management research:
780 Critical issues. *Strategic Management Journal* 28, 331–343.
781 <https://doi.org/10.1002/smj.582>

782 Hovenden, M.J., Wills, K.E., Vander Schoor, J.K., Williams, A.L., Newton, P.C.D., 2008.
783 Flowering phenology in a species-rich temperate grassland is sensitive to warming but
784 not elevated CO₂. *New Phytologist* 178, 815–822. [https://doi.org/10.1111/j.1469-](https://doi.org/10.1111/j.1469-8137.2008.02419.x)
785 [8137.2008.02419.x](https://doi.org/10.1111/j.1469-8137.2008.02419.x)

786 Hulin, S., Carrère, P., Chabalier, C., Farruggia, A., Landrieaux, J., Orth, D., Piquet, M.,
787 Rivière, J., Seytre, L., 2011. Typologie multifonctionnelle des prairies Outil 1 -
788 Typologie multifonctionnelle des prairies. Niveau 2.

789 IGN, 2016. BD ALTI®. Institut National de l'information Géographique et forestière, é,
790 France.

791 Kleijn, D., Berendse, F., Smit, R., Gilissen, N., 2001. Agri-environment schemes do not
792 effectively protect biodiversity in Dutch agricultural landscapes. *Nature* 413, 723–725.
793 <https://doi.org/10.1038/35099540>

794 Kudernatsch, T., Fischer, A., Bernhardt-Römermann, M., Abs, C., 2008. Short-term effects of
795 temperature enhancement on growth and reproduction of alpine grassland species.
796 *Basic and Applied Ecology* 9, 263–274. <https://doi.org/10.1016/j.baae.2007.02.005>

797 Lanta, V., Doležal, J., Lantová, P., Kelíšek, J., Mudrák, O., 2009. Effects of pasture
798 management and fertilizer regimes on botanical changes in species-rich mountain
799 calcareous grassland in Central Europe. *Grass and Forage Science* 64, 443–453.
800 <https://doi.org/10.1111/j.1365-2494.2009.00709.x>

801 Launay, F., Baumont, R., Plantureux, S., Farrié, J.-P., Michaud, A., Pottier, E., 2011. Prairies
802 permanentes Des références pour valoriser leur diversité. Institut de l'Élevage, Paris.

803 Lavorel, S., 2013. Plant functional effects on ecosystem services. *Journal of Ecology* 101, 4–
804 8. <https://doi.org/10.1111/1365-2745.12031>

805 Lavorel, S., Garnier, É., 2002. Predicting changes in community composition and ecosystem
806 functioning from plant traits: revisiting the Holy Grail. *Functional ecology* 16, 545–
807 556.

808 Macedo, J.A., Alves, P., Lomba, Â., Vicente, J., Henriques, R., Granja, H., Honrado, J., 2010.
809 On the interest of plant functional classifications to study community—level effects of
810 increased disturbance on coastal dune vegetation. *Acta Botanica Gallica* 157, 305–
811 315. <https://doi.org/10.1080/12538078.2010.10516208>

812 Magiera, A., Feilhauer, H., Waldhardt, R., Wiesmair, M., Otte, A., 2017. Modelling biomass
813 of mountainous grasslands by including a species composition map. *Ecological*
814 *Indicators* 78, 8–18. <https://doi.org/10.1016/j.ecolind.2017.02.039>

815 Mauchamp, L., Mouly, A., Badot, P.-M., Gillet, F., 2014. Impact of management type and
816 intensity on multiple facets of grassland biodiversity in the French Jura Mountains.
817 *Applied Vegetation Science* 17, 645–657. <https://doi.org/10.1111/avsc.12116>

818 Melts, I., Lanno, K., Sammuli, M., Uchida, K., Heinsoo, K., Kull, T., Laanisto, L., 2018.
819 Fertilising semi-natural grasslands may cause long-term negative effects on both
820 biodiversity and ecosystem stability. *Journal of Applied Ecology* 55, 1951–1955.
821 <https://doi.org/10.1111/1365-2664.13129>

822 Michaud, A., Carrère, P., Farruggia, A., Jeangros, B., Orth, D., Pauthenet, Y., Plantureux, S.,
823 2013. Building grassland typologies in order to evaluate their potential to provide
824 agro-environmental services. *Fourrages* 213, 35–44.

825 Michaud, A., Plantureux, S., Amiaud, B., Carrère, P., Cruz, P., Duru, M., Dury, B., Farruggia,
826 A., Fiorelli, J.L., Kerneis, E., Baumont, R., 2012. Identification of the environmental
827 factors which drive the botanical and functional composition of permanent grasslands.
828 *The Journal of Agricultural Science* 150, 219–236.
829 <https://doi.org/10.1017/S0021859611000530>

830 Michaud, A., Plantureux, S., Pottier, E., Baumont, R., 2015. Links between functional
831 composition, biomass production and forage quality in permanent grasslands over a

832 broad gradient of conditions. *The Journal of Agricultural Science* 153, 891–906.
833 <https://doi.org/10.1017/S0021859614000653>

834 Millennium Ecosystem Assessment (Ed.), 2005. *Ecosystems and human well-being:*
835 *synthesis*. Island Press, Washington, DC.

836 Mouillot, D., Graham, N.A.J., Villéger, S., Mason, N.W.H., Bellwood, D.R., 2013. A
837 functional approach reveals community responses to disturbances. *Trends in Ecology*
838 *& Evolution* 28, 167–177. <https://doi.org/10.1016/j.tree.2012.10.004>

839 Muller, S., 2002. Diversity of management practices required to ensure conservation of rare
840 and locally threatened plant species in grasslands: a case study at a regional scale
841 (Lorraine, France). *Biodiversity and Conservation* 11, 1173–1184.
842 <https://doi.org/10.1023/A:1016049605021>

843 Muller, S., Dutoit, T., Alard, D., Gréville, F., 1998. Restoration and Rehabilitation of
844 Species-Rich Grassland Ecosystems in France: a Review. *Restoration ecology* 6, 94–
845 101.

846 Ni, J., 2004. Estimating Net Primary Productivity of Grasslands from Field Biomass
847 Measurements in Temperate Northern China. *Plant Ecology* 174, 217–234.

848 Nock, C.A., Vogt, R.J., Beisner, B.E., 2016. Functional Traits, in: ELS. John Wiley & Sons,
849 Ltd, Chichester, UK, pp. 1–8. <https://doi.org/10.1002/9780470015902.a0026282>

850 Oram, N.J., Ravenek, J.M., Barry, K.E., Weigelt, A., Chen, H., Gessler, A., Gockele, A., de
851 Kroon, H., van der Paauw, J.W., Scherer-Lorenzen, M., Smit-Tiekstra, A., van
852 Ruijven, J., Mommer, L., 2018. Below-ground complementarity effects in a grassland
853 biodiversity experiment are related to deep-rooting species. *Journal of Ecology* 106,
854 265–277. <https://doi.org/10.1111/1365-2745.12877>

855 Petrovic, M., Acic, S., Zornic, V., Andjelkovic, B., Dajic-Stevanovic, Z., Babic, S., 2013.
856 Evaluation of quality of semi-natural grasslands of central Serbia upon
857 phytosociological and numerical analysis. *Biotechnology in Animal Husbandry* 29,
858 363–371. <https://doi.org/10.2298/BAH1302363P>

859 Pittarello, M., Lonati, M., Gorlier, A., Perotti, E., Probo, M., Lombardi, G., 2018. Plant
860 diversity and pastoral value in alpine pastures are maximized at different nutrient
861 indicator values. *Ecological Indicators* 85, 518–524.
862 <https://doi.org/10.1016/j.ecolind.2017.10.064>

863 Pontes, L.S., Carrere, P., Andueza, D., Louault, F., Soussana, J.F., 2007. Seasonal
864 productivity and nutritive value of temperate grasses found in semi-natural pastures in
865 Europe: responses to cutting frequency and N supply. *Grass and Forage science* 62,
866 485–496.

867 Pykälä, J., Luoto, M., Heikkinen, R.K., Kontula, T., 2005. Plant species richness and
868 persistence of rare plants in abandoned semi-natural grasslands in northern Europe.
869 *Basic and Applied Ecology* 6, 25–33. <https://doi.org/10.1016/j.baae.2004.10.002>

870 QGIS Development Team, 2019. QGIS Geographic Information System. Open Source
871 Geospatial Foundation Project.

872 R Core Team, 2019. R: A language and environment for statistical computing. R Foundation
873 for Statistical Computing, Vienna, Austria.

874 Rodríguez-Rojo, M.P., Jiménez-Alfaro, B., Jandt, U., Bruelheide, H., Rodwell, J.S.,
875 Schaminée, J.H.J., Perrin, P.M., Kaçki, Z., Willner, W., Fernández-González, F.,
876 Chytrý, M., 2017. Diversity of lowland hay meadows and pastures in Western and
877 Central Europe. *Applied Vegetation Science* 20, 702–719.
878 <https://doi.org/10.1111/avsc.12326>

879 Rodríguez-Rojo, M.P., Sánchez-Mata, D., Gavilan, R.G., Rivas-Martínez, S., Barbour, M.G.,
880 2001. Typology and ecology of Californian serpentine annual grasslands. *Journal of*
881 *Vegetation Science* 12, 687–698.

- 882 Roscher, C., Gubsch, M., Lipowsky, A., Schumacher, J., Weigelt, A., Buchmann, N., Schulze,
883 E.-D., Schmid, B., 2018. Trait means, trait plasticity and trait differences to other
884 species jointly explain species performances in grasslands of varying diversity. *Oikos*
885 127, 865–865. <https://doi.org/10.1111/oik.04815>
- 886 Schaffers, A.P., Sýkora, K.V., 2000. Reliability of Ellenberg indicator values for moisture,
887 nitrogen and soil reaction: a comparison with field measurements. *Journal of*
888 *Vegetation Science* 11, 225–244. <https://doi.org/10.2307/3236802>
- 889 Schellberg, J., 1999. Long-term effects of fertilizer on soil nutrient concentration, yield,
890 forage quality and floristic composition of a hay meadow in the Eifel mountains,
891 Germany. *Grass and Forage Science* 54, 195–207.
- 892 Setubal, R.B., Boldrini, I.I., 2012. Phytosociology and natural subtropical grassland
893 communities on a granitic hill in southern Brazil. *Rodriguésia* 63, 513–524.
894 <https://doi.org/10.1590/S2175-78602012000300003>
- 895 Sevruck, B., 1997. Regional Dependency of Precipitation-Altitude Relationship in the Swiss
896 Alps, in: Diaz, H.F., Beniston, M., Bradley, R.S. (Eds.), *Climatic Change at High*
897 *Elevation Sites*. Springer Netherlands, Dordrecht, pp. 123–137.
898 https://doi.org/10.1007/978-94-015-8905-5_7
- 899 Signorell, A., 2017. DescTools: Tools for Descriptive Statistics.
- 900 Spiess, A.-N., 2014. Modelling and analysis of real-time PCR data.
- 901 Suding, K.N., Collins, S.L., Gough, L., Clark, C., Cleland, E.E., Gross, K.L., Milchunas,
902 D.G., Pennings, S., 2005. Functional- and abundance-based mechanisms explain
903 diversity loss due to N fertilization. *Proceedings of the National Academy of Sciences*
904 102, 4387–4392. <https://doi.org/10.1073/pnas.0408648102>
- 905 Sutter, L., Jeanneret, P., Bartual, A.M., Bocci, G., Albrecht, M., 2017. Enhancing plant
906 diversity in agricultural landscapes promotes both rare bees and dominant crop-
907 pollinating bees through complementary increase in key floral resources. *Journal of*
908 *Applied Ecology* 54, 1856–1864. <https://doi.org/10.1111/1365-2664.12907>
- 909 Symonds, M.R.E., Moussalli, A., 2011. A brief guide to model selection, multimodel
910 inference and model averaging in behavioural ecology using Akaike's information
911 criterion. *Behavioral Ecology and Sociobiology* 65, 13–21.
912 <https://doi.org/10.1007/s00265-010-1037-6>
- 913 Tasset, E., Boulanger, T., Diquélou, S., Laîné, P., Lemauiel-Lavenant, S., 2019. Plant trait to
914 fodder quality relationships at both species and community levels in wet grasslands.
915 *Ecological Indicators* 97, 389–397. <https://doi.org/10.1016/j.ecolind.2018.10.035>
- 916 Theau, J.P., Pauthenet, Y., Cruz, P., 2017. Assessing the diversity and usage value of
917 permanent grasslands with a classification system based on non-grass species.
918 *Fourrages* 232, 321–329.
- 919 Therneau, T.M., Atkinson, B., 2014. mvpart.
- 920 Tilman, D., Haddi, A.E., 1992. Drought and Biodiversity in Grasslands. *Oecologia* 89, 257–
921 264.
- 922 Tjur, T., 2009. Coefficients of Determination in Logistic Regression Models—A New
923 Proposal: The Coefficient of Discrimination. *The American Statistician* 63, 366–372.
924 <https://doi.org/10.1198/tast.2009.08210>
- 925 Tóth, E., Deák, B., Valkó, O., Kelemen, A., Migléc, T., Tóthmérész, B., Török, P., 2018.
926 Livestock Type is More Crucial Than Grazing Intensity: Traditional Cattle and Sheep
927 Grazing in Short-Grass Steppes: Traditional Cattle and Sheep Grazing in Short-Grass
928 Steppes. *Land Degradation & Development* 29, 231–239.
929 <https://doi.org/10.1002/ldr.2514>

- 930 van Oijen, M., Bellocchi, G., Höglind, M., 2018. Effects of Climate Change on Grassland
931 Biodiversity and Productivity: The Need for a Diversity of Models. *Agronomy* 8, 14.
932 <https://doi.org/10.3390/agronomy8020014>
- 933 Violle, C., Navas, M.-L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., Garnier, E., 2007.
934 Let the concept of trait be functional! *Oikos* 116, 882–892.
935 <https://doi.org/10.1111/j.2007.0030-1299.15559.x>
- 936 Vujnovic, K., Wein, R.W., Dale, M.R.T., 2002. Predicting plant species diversity in response
937 to disturbance magnitude in grassland remnants of central Alberta. *Canadian Journal*
938 *of Botany* 80, 504–511. <https://doi.org/10.1139/b02-032>
- 939 WallisDeVries, M.F., Poschlod, P., Willems, J.H., 2002. Challenges for the conservation of
940 calcareous grasslands in northwestern Europe: integrating the requirements of flora
941 and fauna. *Biological Conservation* 104, 264–273.
- 942 Wang, C.T., Long, R.J., Wang, Q.J., Ding, L.M., Wang, M.P., 2007. Effects of altitude on
943 plant-species diversity and productivity in an alpine meadow, Qinghai–Tibetan
944 plateau. *Australian Journal of Botany* 55, 110–117. <https://doi.org/10.1071/BT04070>
- 945 Warzecha, D., Diekötter, T., Wolters, V., Jauker, F., 2017. Attractiveness of wildflower
946 mixtures for wild bees and hoverflies depends on some key plant species. *Insect*
947 *Conservation and Diversity* 11, 32–41. <https://doi.org/10.1111/icad.12264>
- 948 Young, J., Watt, A., Nowicki, P., Alard, D., Clitherow, J., Henle, K., Johnson, R., Laczko, E.,
949 McCracken, D., Matouch, S., Niemela, J., Richards, C., 2005. Towards sustainable
950 land use: identifying and managing the conflicts between human activities and
951 biodiversity conservation in Europe. *Biodiversity and Conservation* 14, 1641–1661.
952 <https://doi.org/10.1007/s10531-004-0536-z>
953