



HAL
open science

Plasma Dihydroceramides Are Diabetes Susceptibility Biomarker Candidates in Mice and Humans

Leonore Wigger, Céline Cruciani-Guglielmacci, Anthony Nicolas, Jessica Denom, Neïké Fernandez, Frédéric Fumeron, Pedro Marques-Vidal, Alain Ktorza, Werner Kramer, Anke Schulte, et al.

► **To cite this version:**

Leonore Wigger, Céline Cruciani-Guglielmacci, Anthony Nicolas, Jessica Denom, Neïké Fernandez, et al.. Plasma Dihydroceramides Are Diabetes Susceptibility Biomarker Candidates in Mice and Humans. *Cell Reports*, 2017, 18 (9), pp.2269–2279. 10.1016/j.celrep.2017.02.019 . hal-02392734

HAL Id: hal-02392734

<https://hal.science/hal-02392734>

Submitted on 16 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DR. PER HARALD JONSON (Orcid ID : 0000-0003-2423-359X)

MR. MRIDUL JOHARI (Orcid ID : 0000-0002-3549-558X)

Article type : Short Communications

Novel mutations in DNAJB6 cause LGMD1D and distal myopathy in French families

Authors

Per H. Jonson^{1*}, PhD; Johanna Palmio^{2*}, MD, PhD; Mridul Johari¹, MSci; Sini Penttilä², MSci; Anni Evilä¹, PhD; Isabelle Nelson³, PhD; Gisèle Bonne³, PhD; Nicolas Wiart⁴, MSci; Vincent Meyer⁴, PhD; Anne Boland⁴, PhD; Jean-François Deleuze⁴, PhD; Cécile Masson⁵, PhD; Tanya Stojkovic³, MD, PhD; Françoise Chapon⁶, MD, PhD; Norma B. Romero⁷, MD, PhD; Guilhem Solé⁸, MD; Xavier Ferrer⁸, MD, PhD; Ana Ferreira⁹, MD, PhD; Peter Hackman¹, PhD; Isabelle Richard¹⁰, PhD; Bjarne Udd¹¹, MD, PhD

Full address:

1. PHJ, MJ, AE, PH: Folkhälsan Institute of Genetics and University of Helsinki, Medicum, Finland.
2. JP, SP: Neuromuscular Research Center, Tampere University Hospital and University of Tampere, Finland.
3. IN, GB, TS: Sorbonne Universités, UPMC Univ Paris 06, INSERM UMRS 974, Center of Research in Myology, Institut de Myologie, F-75013, Paris, France.
4. NW, VM, AB & J-F D: Centre National de Recherche en Génomique Humaine (CNRGH), CEA, Evry, France.
5. CM: Bioinformatics Core Facility, Université Paris Descartes - Structure Fédérative de Recherche Necker, INSERM US24/CNRS UMS3633, INSERM UMR 1163, Institut Imagine, Paris, France.
6. FC: Université de Normandie, INSERM U1075, Neuromuscular Competence Center, CHU Caen, Caen, France CHU Caen, France.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/ene.13598

This article is protected by copyright. All rights reserved.

7. NBR: Unit of Neuromuscular Morphology, Institute of Myology, UPMC Paris 6, INSERM UMRS 974, Pitié-Salpêtrière Hospital, Paris, France.
8. GS, XF: Neuromuscular Reference Center, CHU Bordeaux, Bordeaux, France.
9. AF: Unité de Biologie Fonctionnelle et Adaptative, Université Paris Diderot/CNRS, France; Reference Center for Neuromuscular Disorders, Pitié-Salpêtrière Hospital, AP-HP, France.
10. IR: Généthon INSERM, U951, INTEGRARE research unit, University Paris–Saclay, Evry, F-91002, France.
11. BU: Neuromuscular Research Center, Tampere University Hospital and University of Tampere, Finland; Folkhälsan Institute of Genetics and University of Helsinki, Medicum, Finland; Department of Neurology, Vaasa Central Hospital, Vaasa, Finland.

Phone, fax, e-mail of corresponding author: Phone +358-3-3116111, fax: +358-3-35516164,

e-mail: johanna.palmio@staff.uta.fi

Corresponding author: Johanna Palmio

Running title: DNAJB6 myopathy in French families

Keywords: Neuromuscular diseases < NEUROLOGICAL DISORDERS, Myopathies < Neuromuscular diseases < NEUROLOGICAL DISORDERS, Genetic and inherited disorders < NEUROLOGICAL DISORDERS

Conflict of interest: The authors declare no conflict of interest.

*Equal contribution

Abstract

Background and purpose: To determine the genetic background of unknown muscular dystrophy in five French families.

Methods: Twelve patients with limb girdle muscular dystrophy or distal myopathy were clinically evaluated. Gene mutations were identified using targeted exon sequencing and mutated DNAJB6 was tested *in vitro*.

Results: Five patients presented with distal lower limb weakness while others had proximal presentation with variable rate of progression starting at the median age of 34.6 years. Two novel mutations (c.284A>T, p.Asn95Ile, 2 families and c.293_295delATG, p.Asp98del, 1 family) as well as the previously reported c.279C>G (p.Phe93Leu, 2 families) mutation in *DNAJB6* were identified. All showed a reduced capacity to prevent protein aggregation.

This article is protected by copyright. All rights reserved.

Conclusions: The mutational and phenotypical spectrum of *DNAJB6*-caused muscle disease is larger than previously reported, including also dysphagia. The originally reported c.279C>G (p.Phe93Leu) mutation is now identified in four different populations and appears to be a mutational hotspot. Our report confirms that some *DNAJB6* mutations cause distal-onset myopathy and hence *DNAJB6* defects should be considered broadly in dominant muscular dystrophy families.

Introduction

Autosomal dominant limb-girdle muscular dystrophy type 1D (LGMD1D, OMIM 603511) caused by *DNAJB6* mutations [1,2] has so far been identified in Europe [1,3,4], Asia [5-8] and the US [1,2,9,10]. All reported mutations affect *DNAJB6*'s G/F-domain, encoded by exon 5. Classical LGMD1D is late-onset with slow progression, and most patients remain ambulatory at high age. However, some *DNAJB6* mutations can cause a much earlier onset and severe progression [3,5], and some may present with a more distal phenotype. Cardiac involvement has not been observed, but the severe p.Phe91Ile and p.Phe91Leu mutations lead to respiratory failure [3]. Muscle biopsies show myofibrillar protein accumulations and later autophagic rimmed vacuolar dystrophic pathology indicating secondary abnormal autophagic induction. The accumulations commonly contain sarcomeric Z-disc proteins such as α B-crystallin, myotilin and desmin. The rimmed vacuoles stain for the autophagic markers TDP-43, LC3 and SQSTM1, but not for lysosomal marker LAMP2, suggesting problems with autophagosome maturation and lysosome fusion [11].

We report the identification of *DNAJB6* mutations in five French families. Two novel mutations (c.284A>T; p.Asn95Ile, c.293_295delATG; p.Asp98del), in addition to the previously reported c.279C>G (p.Phe93Leu), were identified. Functional studies of the novel mutations confirm reduced anti-aggregation capacity.

Methods

Twelve patients from five French families with distal myopathy or LGMD were evaluated (Figure 1A). DNA from families 1 and 2 were sequenced using MYOcap [12] and families 3-5 with exome sequencing.

Histochemical analyses were performed on cryosections using standard methods. The anti-aggregation activity of *DNAJB6* was tested using a filter trap method [1]. Further experimental details in Supplementary methods.

The study was approved by the IRB of Tampere University Hospital. All participants provided appropriate consent and the study was conducted according to the Helsinki Declaration.

Results

Clinical details and muscle investigations of the patients in Table 1. The mean age of onset was 38.5 years (range 16-55). Five patients presented with distal weakness, mainly with difficulties walking on toes, nine with proximal lower limb weakness. Three had dysphagia. CK levels were normal or slightly elevated. Electromyography findings in all patients studied (5 patients) were consistent with myopathy with occasional pseudomyotonic or myotonic discharges in three patients. Fatty degenerative involvement was observed in the hamstrings and calf muscles on muscle imaging (Figure 1B). Two patients experienced dyspnoea; none of the others had cardiac or respiratory symptoms. Muscle biopsy was available from nine patients and rimmed vacuolar pathology was a constant finding (Figure 1C).

Targeted high-throughput sequencing revealed a novel c.284A>T (p.Asn95Ile) mutation in the probands in families F1 and F2. Sanger sequencing of remaining family members confirmed the segregation. The families had no known relation, but a common haplotype showed that they are distantly related (supplement table 1).

Exome sequencing of F3:II:2 revealed a novel c.293_295delATG (p.Asp98del) mutation in *DNAJB6*. It was confirmed by Sanger sequencing and segregated in the affected. Families 4 and 5 had a typical LGMD1D phenotype with the previously reported c.279C>G (p.Phe93Leu) mutation.

Filter-trap analysis of the two novel mutation together with two earlier described Finnish mutations showed significant loss of anti-aggregation capacity (Figure 1D).

Discussion

We present the first French families with *DNAJB6* mutations. *DNAJB6* mutations were first related to LGMD1D, but this and other recent studies have shown a wider phenotype and variable severity compared to the most frequent mutation, p.Phe93Leu [1,2,6,13-15].

The novel mutations caused a larger variation in phenotype, both in presenting symptoms and disease severity. Family F1 (p.Asn95Ile) and patient F3:II:2 (p.Asp98del) presented with distal lower limb weakness, but this was not entirely consistent with mutation. Thus, we could not confirm the earlier suggestion of a genotype – phenotype correlation that C-terminal mutations in the G/F-domain correlate with a more distal phenotype [4], although some correlation cannot be excluded. Dysphagia is not uncommon and was present in three of our patients. Although *DNAJB6* is a reported cardiomyopathy susceptibility gene [16], cardiomyopathy has not been observed in any patient so far, and severe respiratory insufficiency seems very rare [17].

All described mutations affect the G/F-domain of *DNAJB6* clearly indicating an important function. The C-terminal of *DNAJB6* is alternatively spliced giving two isoforms with distinct cellular

This article is protected by copyright. All rights reserved.

localisation. Mutations in exon 5 will affect both isoforms, but the cytoplasmic b is the pathogenic isoform in both zebrafish [1] and mouse [18]. This is probably due to the cellular localisation as muscle phenotypes apparently arise from defect chaperonal processing of cytoplasmic proteins.

DNAJB6 interacts [1] with proteins in the chaperone-assisted selective autophagy (CASA) pathway [19]. Other CASA proteins also have similar aggregation preventing effects [20,21] and mutations in the CASA proteins HSPB8 [22] and BAG3 [23] are known to cause neuromuscular disease. DNAJB6 has recently been shown to be important for stress granule handling via TDP-43 [24], and colocalization of DNAJB6 with the stress granule protein TIA1 in protein accumulations can be observed in patient muscle (Supplementary figure 1). DNAJB6 is also important for handling of prions in yeast [25]. The ability of DNAJB6 to recognise specific protein conformers can prevent aggregation of these proteins; and when impaired this will lead to defect handling of misfolded proteins and aggregations.

Our findings broaden the mutational spectrum of DNAJB6 with two novel mutations. Further, as p.Phe93Leu is now identified in four populations it emerges as a mutational hotspot. Our report confirms that *DNAJB6* mutations may cause distal-onset myopathy and hence that *DNAJB6* defects should be considered also in distal myopathy patients with weakness starting in the posterior compartment.

Acknowledgements

Lydia Sagath & Helena Luque for technical assistance and Dr. Anna Vihola for providing immunofluorescent images.

This work was supported by the France Génomique National infrastructure, funded as part of the « Investissements d’Avenir » program managed by the Agence Nationale pour la Recherche (ANR-10-INBS-09) and the Fondation Maladies Rares within the Myocapture sequencing project.

Conflicts of interest: no

References

1. Sarparanta J, Jonson PH, Golzio C, *et al.* Mutations affecting the cytoplasmic functions of the co-chaperone DNAJB6 cause limb-girdle muscular dystrophy. *Nat Genet* 2012; **44**: 450-455.
2. Harms MB, Sommerville RB, Allred P, *et al.* Exome sequencing reveals DNAJB6 mutations in dominantly-inherited myopathy. *Ann Neurol* 2012; **71**: 407-416.
3. Palmio J, Jonson PH, Evilä A, *et al.* Novel mutations in *DNAJB6* gene cause a very severe early-onset limb-girdle muscular dystrophy 1D disease. *Neuromuscul Disord* 2015; **25**: 835-842.
4. Ruggieri A, Brancati F, Zanotti S, *et al.* Complete loss of the DNAJB6 G/F domain and novel missense mutations cause distal-onset DNAJB6 myopathy. *Acta Neuropathol Commun* 2015; **3**: 44.
5. Nam TS, Li W, Heo SH, *et al.* A novel mutation in DNAJB6, p.(Phe91Leu), in childhood-onset LGMD1D with a severe phenotype. *Neuromuscul Disord* 2015; **25**: 843-851.
6. Sato T, Hayashi YK, Oya Y, *et al.* DNAJB6 myopathy in an Asian cohort and cytoplasmic/nuclear inclusions. *Neuromuscul Disord* 2013; **23**: 269-276.
7. Monies D, Alhindi HN, Almuhaizea MA, *et al.* A first-line diagnostic assay for limb-girdle muscular dystrophy and other myopathies. *Hum Genomics* 2016; **10**: 32.
8. Tsai PC, Tsai YS, Soong BW, *et al.* A novel *DNAJB6* mutation causes dominantly inherited distal-onset myopathy and compromises DNAJB6 function. *Clin Genet* 2017; .
9. Suarez-Cedeno G, Winder T, Milone M. DNAJB6 myopathy: a vacuolar myopathy with childhood onset. *Muscle Nerve* 2014; **49**: 607-610.
10. Couthouis J, Raphael AR, Siskind C, *et al.* Exome sequencing identifies a *DNAJB6* mutation in a family with dominantly-inherited limb-girdle muscular dystrophy. *Neuromuscul Disord* 2014; **24**: 431-435.
11. Sandell S, Huovinen S, Palmio J, *et al.* Diagnostically important muscle pathology in DNAJB6 mutated LGMD1D. *Acta Neuropathol Commun* 2016; **4**: 9.
12. Evilä A, Palmio J, Vihola A, *et al.* Targeted next-generation sequencing reveals novel *TTN* mutations causing recessive distal titinopathy. *Mol Neurobiol* 2016; .
13. Sandell S, Huovinen S, Sarparanta J, *et al.* The enigma of 7q36 linked autosomal dominant limb girdle muscular dystrophy. *J Neurol Neurosurg Psychiatry* 2010; **81**: 834-839.
14. Hackman P, Sandell S, Sarparanta J, *et al.* Four new Finnish families with LGMD1D; refinement of the clinical phenotype and the linked 7q36 locus. *Neuromuscul Disord* 2011; **21**: 338-344.
15. Sandell SM, Mahjneh I, Palmio J, Tasca G, Ricci E, Udd BA. 'Pathognomonic' muscle imaging findings in DNAJB6 mutated LGMD1D. *Eur J Neurol* 2013; **20**: 1553-1559.

16. Ding Y, Long PA, Bos JM, *et al.* A modifier screen identifies DNAJB6 as a cardiomyopathy susceptibility gene. *JCI Insight* 2016; **1**: e88797.
17. Ruggieri A, Saredi S, Zanotti S, Pasanisi MB, Maggi L, Mora M. DNAJB6 Myopathies: Focused Review on an Emerging and Expanding Group of Myopathies. *Front Mol Biosci* 2016; **3**: 63.
18. Bengoechea R, Pittman SK, Tuck EP, True HL, Weihl CC. Myofibrillar disruption and RNA-binding protein aggregation in a mouse model of limb-girdle muscular dystrophy 1D. *Hum Mol Genet* 2015; **24**: 6588-6602.
19. Arndt V, Dick N, Tawo R, *et al.* Chaperone-assisted selective autophagy is essential for muscle maintenance. *Curr Biol* 2010; **20**: 143-148.
20. Carra S, Sivilotti M, Chávez Zobel AT, Lambert H, Landry J. HspB8, a small heat shock protein mutated in human neuromuscular disorders, has *in vivo* chaperone activity in cultured cells. *Hum Mol Genet* 2005; **14**: 1659-1669.
21. Hishiya A, Salman MN, Carra S, Kampinga HH, Takayama S. BAG3 directly interacts with mutated α B-crystallin to suppress its aggregation and toxicity. *PLoS One* 2011; **6**: e16828.
22. Ghaoui R, Palmio J, Brewer J, *et al.* Mutations in HSPB8 causing a new phenotype of distal myopathy and motor neuropathy. *Neurology* 2016; **86**: 391-398.
23. Selcen D, Muntoni F, Burton BK, *et al.* Mutation in BAG3 causes severe dominant childhood muscular dystrophy. *Ann Neurol* 2009; **65**: 83-89.
24. Stein KC, Bengoechea R, Harms MB, Weihl CC, True HL. Myopathy-causing mutations in an HSP40 chaperone disrupt processing of specific client conformers. *J Biol Chem* 2014; **289**: 21120-21130.
25. Reidy M, Sharma R, Roberts BL, Masison DC. Human J-protein DnaJB6b cures a subset of *Saccharomyces cerevisiae* prions and selectively blocks assembly of structurally related amyloids. *J Biol Chem* 2016; **291**: 4035-4047.

Supplementary material references (for online only E-extra publication)

26. Gordon CT, Petit F, Kroisel PM, *et al.* Mutations in endothelin 1 cause recessive auriculocondylar syndrome and dominant isolated question-mark ears. *Am J Hum Genet* 2013; **93**: 1118-1125.
27. Hasholt L, Abell K, Nørremølle A, Nellesmann C, Fenger K, Sørensen SA. Antisense downregulation of mutant huntingtin in a cell model. *J Gene Med* 2003; **5**: 528-538.

Legends

Figure 1. Pedigree and muscle findings of the patients with novel mutations: (A) Pedigree of the families. (B) Rimmed and subsarcolemmal non-rimmed vacuolation and some myofibrillar aggregation are seen in F1 III:1. Staining with antibodies against myotilin, α B-crystallin and desmin shows strong expression in the myofibrillar lesions with aggregates. In patient F3 II:2 the findings are similar although more advanced dystrophic with increased fat and fibrosis. (C) Fatty degenerative changes are most severe in the distal lower limb muscles, soleus and medial gastrocnemius in patient F1 III:1. Less severe involvement is present in semimembranosus on the left. The posterior compartment of the calf is fatty degenerated in patient F2 III:1 with asymmetrical involvement of biceps femoris, semimembranosus and adductor magnus. Patient's F3 II:2 muscle MRI was performed at the time when she was already bedridden and consistently all muscles are severely affected. (D) Both novel mutations showed significantly impaired anti-aggregation activity (Mann-Whitney U-test). Aggregation score was calculated as a ratio of aggregated/soluble in the induced to the uninduced cells.

Table 1 Clinical data of the patients

| Patients | Age at last exam/ Sex | Age of onset | First symptoms | Muscle weakness (MRC, the Medical Research Council Scale) | Ambulation | Creatine kinase levels | Muscle histology | Cardiac / Respiratory symptoms |
|-----------------|------------------------------|---------------------|---|--|----------------------|-------------------------------|--|--|
| F1 II:2 | 85/M | 55 | Difficulty walking on toes | Distal and proximal upper and lower limbs, dysphagia | Wheelchair (aged 85) | Normal | Rimmed vacuoles | No/ No |
| F1 II:4 | 75/M | 40 | Difficulty walking on toes | Lower limbs; tibialis (0), gastrocnemius (2), quadriceps (4), hamstrings (3) | One cane | 2 X UNL | Rimmed vacuoles | No/ No |
| F1 III:1 | 64/F | 50-54 | Difficulty walking on toes | Distal lower limbs | Ambulant | Normal | Rimmed vacuoles, myofibrillar aggregates | No/ No |
| F1 III:3 | 65/M | 54 | Difficulty walking on toes | Mild distal lower limbs; gastrocnemius (4) | Ambulant | Normal | Not done | No/ No |
| F2 III:1 | 60/M | 40 | Left lower limb weakness, difficulty climbing stairs | Scapular (4), axial (4), hyperlordosis, pelvic (3), unable to walk toes/heels | One cane 50m | Not available | Dystrophy, rimmed vacuoles | No/ No |
| F2 III:4 | 54/M | 36 | Left lower limb weakness difficulty climbing stairs, frequent falls | Abdominal, pelvic and thigh (2), gastrocnemius (3+), tibialis (2), peroneus (3/4-), triceps/biceps (3) | Two canes 50m | 2 X UNL | Dystrophy, rimmed vacuoles | No/ exertional dyspnoea Echo: Left EF 63% / restrictive pulmonary dysfunction |

| | | | | | | | | |
|-----------------|------|----|------------------------------|---|---|---------------|---------------------|--|
| F3 II:2 | 71/F | 16 | Distal lower limb weakness | Severe distal and proximal lower/upper limbs; dysphagia, cricopharyngeal myotomy; Achilles tendon surgery | Wheelchair (aged 49), bedridden (aged 65) | Normal | Rimmed vacuoles | No/Dyspnoea Echo: aortic dilatation, no cardiomyopathy / restrictive pulmonary dysfunction, VC 52 % |
| F3 III:1 | 46/M | 20 | Proximal lower limb weakness | Lower limbs (0-1), deltoid (3-), biceps (3-), triceps (2), wrist fl/ext. (4/3-), finger fl/ext (4/2) | Wheelchair | Not available | Rimmed vacuoles | No/ No Echo: normal / FVC 4,1 l (90 %) |
| F4 II:2 | 86/F | 44 | Waddling gait | Proximal and distal limbs; dysphagia | Wheelchair (aged 70) | Normal | Not available | Cardiac infarction aged 79/ No |
| F4 III:1 | 48/M | 40 | Difficulty climbing stairs | Severe in lower limbs; moderate in axial and proximal upper limbs | Rollator and wheelchair | 2 x UNL | Not available | No/ No Echo: left ventricular hypertrophy |
| F5 I:2 | 62/F | 40 | Difficulty climbing stairs | Proximal and distal lower limbs, mild scapular winging | One cane 20m | Normal | Unspecific myopathy | Not available |
| F5 II:1 | 40/M | 25 | Difficulty climbing stairs | Proximal > distal lower limbs | Ambulant | 6 x UNL | Unspecific myopathy | No/ No |

Echo, echocardiogram; FVC, forced vital capacity; UNL, upper normal limit; VC, vital capacity

