



HAL
open science

Generalized description of spectral incoherent solitons

Gang Xu, Josselin Garnier, Matteo Conforti, Antonio Picozzi

► **To cite this version:**

Gang Xu, Josselin Garnier, Matteo Conforti, Antonio Picozzi. Generalized description of spectral incoherent solitons. *Optics Letters*, 2014, 39 (14), pp.4192. 10.1364/OL.39.004192 . hal-02393796

HAL Id: hal-02393796

<https://hal.science/hal-02393796>

Submitted on 4 Dec 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Generalized description of spectral incoherent solitons

Gang Xu,^{1,*} Josselin Garnier,² Matteo Conforti,³ Antonio Picozzi¹

¹Laboratoire Interdisciplinaire Carnot de Bourgogne (ICB), CNRS - University of Burgundy, Dijon, France

²Laboratoire de Probabilités et Modèles Aléatoires, University Paris Diderot, 75205 Paris Cedex 13, France

³Laboratoire de Physique des Lasers, Atomes et Molécules (PhLAM), CNRS - University of Lille 1, Villeneuve d'Ascq, France

*Corresponding author: gang.xu@u-bourgogne.fr

Compiled May 16, 2014

Spectral incoherent solitons (SIS) refer to an incoherent solitonic structure that can only be identified in the frequency domain and which is sustained by noninstantaneous Raman-like nonlinearities. We provide a generalized formulation of SISs by considering the generalized nonlinear Schrödinger equation. We show that nonlinear dispersive effects (e.g., self-steepening) dramatically affect the structure of SISs, in particular by significantly decelerating their spectral velocity in frequency space. The wave turbulence description of the problem reveals an underlying self-organization process: The soliton trajectory in frequency space is selected in such a way that it allows the system to self-organize into a stationary SIS state. © 2014 Optical Society of America

OCIS codes: 190.4370, 190.5650, 030.1640

The propagation of partially coherent nonlinear optical waves is a topic of growing interest in different fields of investigations, such as, e.g., wave propagation in homogeneous [1–4] or periodic media [5], waveguides [6], cavity systems [7–11], supercontinuum generation (SC) [12–14], shock waves [15], or nonlinear interferometry [16]. In analogy with kinetic gas theory, an incoherent wave is expected to exhibit a nonequilibrium process of thermalization toward thermodynamic equilibrium [17]. Besides wave thermalization, a partially coherent wave can self-organize into incoherent solitons (IS), which thus constitute genuine nonequilibrium stable states of the random field. Originally observed in photorefractive crystals [18], ISs have become a bloomy area of research, as illustrated by important achievements, such as, e.g., the modulational instability of incoherent waves [19, 20], the existence of ISs with spatial or temporal nonlocal nonlinearities [21–23].

An IS of a fundamentally different nature has been identified by exploiting the noninstantaneous property of the nonlinear Raman response in optical fibers [25–27]. The incoherent wave associated to this soliton exhibits fluctuations that are statistically stationary in time, so that it does not exhibit a confinement in the temporal domain, but exclusively in the frequency domain. For this reason, it has been termed ‘spectral incoherent soliton’ (SIS). Some experimental signatures of the generation of SISs have been obtained through the study of SC generation in photonic crystal fibers [27, 28].

Our aim in this Letter is to report a nontrivial generalization of SIS structures. We consider the 1D generalized nonlinear Schrödinger equation (GNLSE) which accounts, in particular, for the self-steepening (or shock) term, and a frequency dependence of the nonlinear Kerr coefficient [29–31]. Nonlinear dispersive effects are shown to dramatically affect the dynamics of the incoherent wave. Starting from the GNLSE, we derive a generalized kinetic equation (GKE) which does not admit incoher-

ent soliton solutions: When considered as initial condition, the standard SIS exhibits a significant deceleration in frequency space as well as a significant reduction of its power. However, the underlying soliton structure is unveiled by a change of variables, which reveals an unexpected process of self-organization: The specific form of the soliton trajectory in frequency space is selected in such a way that it allows the system to self-organize into a stationary SIS state. In this way, the spectral deceleration of SIS is obtained in explicit analytical form and in quantitative agreement with the simulations of GNLSE, without adjustable parameters. Besides its fundamental interest, this generalized formulation of SISs can shed new light on the spectral dynamics of SC generation [27, 28].

The starting point is the GNLSE ~~which is usually~~ written in the frequency domain [29–31]

$$\partial_z \tilde{\psi}_\omega(z) = ik(\omega)\tilde{\psi}_\omega - \alpha\tilde{\psi}_\omega(z) + i\Gamma(\omega)\mathcal{F}\{\psi R * |\psi|^2\} \quad (1)$$

where \mathcal{F} is the Fourier operator [$\tilde{\psi}_\omega(z) = \mathcal{F}\{\psi(z, t)\} = \int \psi(z, t) \exp(i\omega t) dt$], while $*$ denotes the temporal convolution product between the intensity $|\psi|^2(z, t)$ and the response function, which includes both the instantaneous (Kerr) contribution and the noninstantaneous (Raman) contribution, $R(t) = (1 - f_R)\delta(t) + f_R h_R(t)$, where $h_R(t) = H(t) \frac{1+\eta^2}{\eta\tau_R} \sin(\eta t/\tau_R) \exp(-t/\tau_R)$, $H(t)$ being the Heaviside function [$\tau_R = 32\text{fs}$, $\eta = \tau_R/\tau_1$, $f_R \simeq 0.18$]. The Taylor expansion of the dispersion relation reads $k(\omega) = \sum_{j \geq 2} \beta_j \omega^j / j!$, β_j being the j -th dispersion coefficient. The function $\Gamma(\omega) = (1 + \omega/\omega_0)\gamma(\omega_0 + \omega)$ accounts for the self-steepening effect (1st factor), as well as the frequency dependence of the nonlinear coefficient that arises from the frequency variation of the effective area and corresponding effective index (2nd factor), ω_0 being the carrier frequency of the wave. The parameter α accounts for the losses during propagation. If $\alpha = 0$, Eq.(1) conserves the ‘number of photons’, $N = \int |\tilde{\psi}(\omega, z)|^2 / \Gamma(\omega) d\omega$. Note that by setting

$\Gamma(\omega) = \gamma_0$ where $\gamma_0 = \gamma(\omega_0)$, Eq.(1) recovers the conventional NLS equation.

It is important to recall that the typical spectral width of SISs is of the same order than the Raman gain spectral bandwidth ($\sim 13\text{THz}$). To study SISs we thus need to avoid spectral broadening effects due to SC-like phenomena. In the following we thus consider the spectral dynamics of SISs far from a zero-dispersion frequency. Note that, as revealed by the theory [Eq.(2)], linear dispersive effects do not play any role in the dynamics of SISs, as previously discussed in several works [24,26,27]. This also reveals that, in spite of their apparent similarities, the phenomenon of spectral deceleration of SISs discussed here is of a fundamental different nature than that discussed in the framework of Raman-induced self-frequency shift of optical solitons [28,31,32], which are inherently *temporally localized coherent entities*.

We report in Fig. 1 (left column) the evolution of the spectrum of the incoherent wave obtained by integrating numerically the GNLSE (1). As usual, the initial condition is a partially coherent wave with Gaussian spectrum and random spectral phases, superimposed on a white noise spectral background [24,27]. Neglecting nonlinear dispersive effects, i.e., $\Gamma(\omega) = \gamma_0$, the initial incoherent wave evolves into a conventional SIS, which propagates with a constant spectral velocity V_0 , and thus virtually crosses the zero-frequency component of the optical field, i.e., $\omega = -\omega_0$ (see Fig. 1a-d). Self-steepening and nonlinear dispersion regularize this ‘unphysical’ negative frequency propagation [33] through a significant spectral deceleration of the SIS, as well as a significant reduction of its power $N = \int |\tilde{\psi}_\omega|^2 d\omega$ (Fig. 1). Note that the frequency reached by the SIS $\omega \sim -180\tau_0^{-1}$ can correspond to a physical wavelength $\lambda \sim 3\mu\text{m}$ (for a source at $\lambda_0 \sim 1.06\mu\text{m}$), where GNLSE (1) can still be considered as an accurate model. Also note that propagation losses have been neglected, so as to highlight the fact that the decrease of power is solely due to nonlinear dispersion effects. In this example we considered a linear frequency variation $\gamma(\omega_0 + \omega) = \gamma_0(1 + q\omega)$, with $q = \gamma_0^{-1} \partial_\omega \gamma(\omega)|_{\omega_0}$.

To describe these novel properties of SISs, we resort to a statistical description of the random wave based on the wave turbulence theory [1,24]. Since we are looking for a soliton behavior, we neglect losses, $\alpha = 0$ in Eq.(1). In the weakly nonlinear regime, we obtain a closure of the hierarchy of moments equations from Eq.(1), in which the averaged spectrum of the wave $\langle \tilde{\psi}_{\omega+\Omega/2} \tilde{\psi}_{\omega-\Omega/2}^* \rangle = n_\omega(z) \delta(\Omega)$ is governed by the following GKE (details on derivation will be given elsewhere):

$$\partial_z n_\omega = \frac{n_\omega}{\pi} \int G(\omega, \omega') n_{\omega'} d\omega', \quad (2)$$

where $G(\omega, \omega') = \Gamma(\omega)g(\omega - \omega')$, and $g(\omega) = \Im[\tilde{R}(\omega)]$ is the imaginary part of the Fourier transform of $R(t)$. Note that GKE (2) accounts for nonlinear dispersion effects, but not for linear dispersion effects (although linear dispersion is known to be essential for the establishment of a




Fig. 1. Numerical simulations of the GNLSE (1) (left column (a)-(c)) and GKE (2) (right column (b)-(d)), for $p = q = 0$ ($\Gamma(\omega) = \gamma_0$) (1st row, (a)-(b)), and for $p = \omega_0^{-1}, q = 0.1\omega_0^{-1}$ (2nd row, (c)-(d)). (e)-(f) Corresponding spectral profiles obtained by simulations of GNLSE (1) (gray line), and GKE (2) (red line), at $z = 200L_{nl}$ and $z = 600L_{nl}$. The vertical dashed-line denotes the physical zero frequency component, i.e., $\omega = -\omega_0$. [$\omega_0/(2\pi) = 282\text{THz}$ ($\lambda_0 = 1.06\mu\text{m}$), $\eta = 2.6$, $\tau_0 = \sqrt{\beta_2 L_{nl}/2}$, $L_{nl} = 1/(\gamma_0 P)$, where P is the average intensity of the incoherent wave.]

weakly nonlinear regime, which in turn leads to a closure of the moments hierarchy [1]). Simulations of GKE (2) are in quantitative agreement with GNLSE without adjustable parameters, as remarkably illustrated in Fig. 1.

Because of nonlinear dispersive effects, $\Gamma(\omega) \neq \gamma_0$, the integral in Eq.(2) does not reduce to a convolution. For this reason, the GKE no longer conserves the power $N = \int n_\omega d\omega$, and thus does not admit soliton solutions [34]. However, to unveil the underlying soliton dynamics, let us consider the variable change $\tilde{n}_\omega(z) = n_\omega(z)/\Gamma(\omega)$, so that GKE (2) can be recast into the symmetric form

$$\partial_z \tilde{n}_\omega = \frac{1}{\pi} \int \tilde{n}_\omega \Gamma(\omega) g(\omega - \omega') \Gamma(\omega') \tilde{n}_{\omega'} d\omega'. \quad (3)$$

This modified GKE (MGKE) conserves the power $\tilde{N} = \int \tilde{n}_\omega d\omega$ and can thus exhibit soliton solutions. However, as revealed by the simulations in Fig. 1, such soliton solutions are not conventional, in the sense that they do not propagate with a constant velocity in frequency space. We thus look for a stationary soliton solution in a ‘non-inertial’ reference frame in frequency space, i.e., a *reference frame travelling with a non-stationary (z -dependent) spectral velocity* $V(z) = \partial_z \Omega^s(z)$: $\Omega =$

$\omega - \Omega^s(z)$. The analysis will reveal that the soliton trajectory, $\Omega^s(z)$, is selected in such a way that it allows the system to self-organize into a stationary SIS state. In the non-inertial reference frame, the MGKE (3) reads

$$\partial_z \tilde{n}_\Omega - \partial_z \Omega^s(z) \partial_\Omega \tilde{n}_\Omega = \frac{\tilde{n}_\Omega}{\pi} \int g(\Omega - \Omega') \tilde{n}_{\Omega'} \times \Gamma[\Omega + \Omega^s(z)] \Gamma[\Omega' + \Omega^s(z)] d\Omega', \quad (4)$$

where $\Omega' = \omega' - \Omega^s(z)$. In general, one cannot decouple Eq.(4) into two independent equations for $\tilde{n}_\Omega(z)$, and for the soliton trajectory $\Omega^s(z)$. However, the SIS profile and $g(\omega)$ vary on a frequency scale much smaller than $\Gamma(\omega)$. We thus introduce the multi-scale expansion $\tilde{n}_\Omega(z) = \varepsilon^{-1} m_{\Omega/\varepsilon}(z)$, $g(\omega) = \varepsilon^{-1} \tilde{g}(\omega/\varepsilon)$, and look for a solution of Eq.(4) in the form, $m_{\bar{\Omega}}(z) = m_{\bar{\Omega}}^s + \varepsilon \delta m_{\bar{\Omega}}(z)$, where $m_{\bar{\Omega}}^s$ is the stationary soliton profile in the non-inertial reference frame and $\delta m_{\bar{\Omega}}(z)$ a residual radiation ($\bar{\Omega} = \Omega/\varepsilon$). To first order in ε , Eq.(4) then decouples into two independent equations for the soliton profile, $\tilde{n}_\Omega^s = \varepsilon^{-1} m_{\Omega/\varepsilon}^s$, and the soliton trajectory:

$$\partial_z \Omega^s(z) = -V_0 \Gamma^2(\Omega^s(z))/\gamma_0^2, \quad (5)$$

$$V_0 \partial_\Omega \tilde{n}_\Omega^s = \frac{\gamma_0^2}{\pi} \tilde{n}_\Omega^s \int g(\Omega - \Omega') \tilde{n}_{\Omega'}^s d\Omega'. \quad (6)$$

Here, by introducing $V_0(> 0)$, we have recast Eq.(6) into the usual stationary equation for a conventional SIS with $\Gamma(\omega) = \gamma_0$ that propagates with the *constant velocity* V_0 . This reveals a remarkable result: Within the new variable $\tilde{n}_\omega = n_\omega/\Gamma(\omega)$, the SIS spectral profile is almost identical to the conventional SIS spectral profile for $\Gamma(\omega) = \gamma_0$, while the corresponding soliton trajectory, $\Omega^s(z)$, satisfies the differential Eq.(5). Note that the multi-scale expansion also readily gives the power decay

$$N(z) = \tilde{N} \Gamma[\Omega^s(z)], \quad (7)$$

where $\tilde{N} = \int \tilde{n}_\omega^s d\omega$ is the conserved power of MGKE (3) and $\Omega^s(z)$ is obtained in explicit form in Eq.(8). The power decay (7) is found in quantitative agreement with simulations, as illustrated in Fig. 2b.

The soliton trajectory can be obtained analytically under the assumption of a linear frequency dependence, $\gamma(\omega_0 + \omega) = \gamma_0(1 + q\omega)$. In this case (5) takes the form $\int_0^{-\Omega^s} (1 - qu)^{-2} (1 - pu)^{-2} du = V_0 z$, with $p = 1/\omega_0$ and $\Omega^s(z = 0) = 0$. The solution can be written as a function of z vs Ω^s :

$$z(\Omega^s) = \frac{1}{V_0(q-p)^3} \left[2pq \log \left(\frac{1 + q\Omega^s}{1 + p\Omega^s} \right) - Q_{qp}(\Omega^s) \right] \quad (8)$$

where $Q_{qp}(\Omega^s) = (q-p)[qp(q+p)\Omega^s + q^2 + p^2]\Omega^s / [(1 + q\Omega^s)(1 + p\Omega^s)]$. Note that, by neglecting nonlinear dispersive effects, i.e., $\Gamma(\omega) = \gamma_0$ (i.e., $p = q = 0$), Eq.(8) reduces to the linear trajectory, $\Omega^s(z) = -V_0 z$, inherent to conventional SISs [24, 25, 27]. Conversely, whenever




Fig. 2. Soliton trajectory Ω^s vs z plotted from Eq.(8), in which V_0 is given by Eq.(10): $q = p = 0$, $\Omega^s(z) = -V_0 z$ (green); $q = 0, p = \omega_0^{-1}$ (red); $q = 0.2p, p = \omega_0^{-1}$ (dark); $q = -0.2p, p = \omega_0^{-1}$ (blue). The corresponding crosses on the curves refer to the simulations of MGKE (3): a quantitative agreement is obtained without adjustable parameters. The inset (a) compare the simulations of the MGKE (3) (green), GNLSE (1) (gray), with the analytical SIS solution Eq.(9-10), in \log_{10} -scale. The superposition of 5 spectral profiles at $z = 200, 400, 600, 800, 1000$ (in units of L_{nl}) has been plotted to stress the stationary character of the SIS solution obtained by simulation of MGKE (3). The inset (b) shows the power decrease, $N = \int |\psi_\omega|^2 d\omega$ vs z , in the GNLSE (1) (red), and the theoretical prediction in Eq.(7) (dashed blue). [$\omega_0/(2\pi) = 282\text{THz}$, $\eta = 1$].

$\Gamma(\omega) \neq \gamma_0$, $\Omega^s(z)$ asymptotically reaches $-\omega_0$ for large propagation lengths z (see Fig. 2), which regularizes the unphysical negative frequency propagation predicted by the linear trajectory, $\Omega^s(z) = -V_0 z$.

An analytical expression of SIS solution with velocity V_0 was obtained in Ref. [25] through a generalization of the particular solution originally derived in [35] in the limit $n_0/n_m \ll 1$, where n_m and n_0 respectively denote the soliton peak and the level of the background noise. Note that, so far this SIS solution has not been compared with numerical simulations. The solution reads

$$\tilde{n}_\Omega^s = n_0 \exp \left[\log \left(\frac{n_m}{n_0} \right) \frac{H(\Omega)}{g_i \omega_i} \right], \quad (9)$$

where $g_i = [|\partial_\omega g(0)| H(0)]^{1/2}$ denotes the typical amount of gain of $g(\omega)$, $\omega_i = [|\partial_\omega g(0)| H(0)]^{1/2}$ its typical spectral bandwidth, and $H(\Omega) = -\int_\Omega^\infty g(\omega) d\omega$. The SIS propagates with the constant velocity

$$V_0 = \frac{\gamma_0^2}{\pi} \frac{\int (\tilde{n}_\Omega^s - n_0) d\Omega}{\int \log(\tilde{n}_\Omega^s/n_0) d\Omega} \int H(\Omega) d\Omega. \quad (10)$$

We have compared the expression of the soliton trajectory (8) as well as the corresponding SIS solution (9-10) with the results of the numerical simulations of the




Fig. 3. Transition to a discrete behavior of generalized SISs in the presence of a small noise background, $n_0/n_m^0 = 10^{-7}$, n_m^0 being the initial spectral peak amplitude: Simulations of the GNLSE (1) (a), and GKE (2) (b) [$\omega_0/(2\pi) = 282\text{THz}$, $\eta = 2.6$, $p = \omega_0^{-1}$, $q = 0.1\omega_0^{-1}$].

MGKE (3) and NLS Eq.(1). A quantitative agreement has been obtained without adjustable parameters, as remarkably illustrated in Fig. 2. In the limit of a small SIS spectral shift, the trajectory recovers the straight line, $\Omega^s(z) = -V_0z$ of conventional SISs, where V_0 is given by (10). In the limit $q = 0$ ($\Gamma(\omega) = 1 + p\omega$), the SIS trajectory takes the simple form $\Omega^s(z) = -V_0z/(1 + V_0z/\omega_0)$ (red line in Fig. 2). A positive (negative) value of q then decelerates (accelerates) the spectral shift of the SIS (see dark and blue lines in Fig. 2). Finally, the plot of the analytical SIS solution (9-10) has been found in quantitative agreement with simulations of MGKE (3) and GNLSE (1), as illustrated in Fig. 2a.

We have provided a generalized wave turbulence formulation of incoherent optical waves and SISs accounting for nonlinear dispersive effects. Note that, as the spectral noise background decreases, the generalized SIS exhibits a transition to a discrete behavior (see Fig. 3), in a way similar to conventional SISs [26], a feature that will be the subject of future investigations.

References

1. J. Garnier, M. Lisak, A. Picozzi, J. Opt. Soc. Am. B. **29**, 2229 (2012).
2. T. Hansson, D. Anderson, M. Lisak, V.E. Semenov, U. Österberg, J. Opt. Soc. Am. B **25**, 1780 (2008).
3. A. Piskarskas, V. Pyragaitė, and A. Stabinis, Phys. Rev. A **82**, 053817 (2010).
4. J. Laurie, U. Bortolozzo, S. Nazarenko, S. Residori, Phys. Rep. **514**, 121 (2012).
5. Y. Silberberg, Y. Lahini, Y. Bromberg, E. Small, R. Morandotti, Phys. Rev. Lett. **102**, 233904 (2009).
6. P. Aschieri, J. Garnier, C. Michel, V. Doya, A. Picozzi, Phys. Rev. A **83**, 033838 (2011).
7. C. Conti, M. Leonetti, A. Fratalocchi, L. Angelani, G. Ruocco, Phys. Rev. Lett. **101**, 143901 (2008).
8. S. Babin, D. Churkin, A. Ismagulov, S. Kablukov, E. Podivilov, J. Opt. Soc. Am. B **24**, 1729 (2007).
9. E. Turitsyna, S. Smirnov, S. Sugavanam, N. Tarasov, X. Shu, S. Babin, E. Podivilov, D. Churkin, G. Falkovich, S. Turitsyn, Nat. Photon. **7**, 783 (2013).
10. R. Weill, B. Fischer, O. Gat, Phys. Rev. Lett. **104**, 173901 (2010).
11. A. Schwache, F. Mitschke, Phys. Rev. E **55**, 7720 (1997).
12. B. Barviau, B. Kibler, A. Picozzi, Phys. Rev. A **79**, 063840 (2009).
13. B. Barviau, J. Garnier, G. Xu, B. Kibler, G. Millot, A. Picozzi, Phys. Rev. A **87**, 035803 (2013).
14. M. Erkintalo, M. Surakka, J. Turunen, A.T. Friberg, G. Genty, Opt. Lett. **37**, 169 (2012).
15. J. Garnier, G. Xu, S. Trillo, A. Picozzi, Phys. Rev. Lett. **111**, 113902 (2013).
16. S. Derevyanko, E. Small, Phys. Rev. A **85**, 053816 (2012).
17. P. Suret, S. Randoux, H.R. Jauslin, A. Picozzi, Phys. Rev. Lett. **104**, 054101 (2010).
18. M. Mitchell, Z. Chen, M. Shih, and M. Segev, Phys. Rev. Lett. **77**, 490 (1996).
19. M. Soljacic, M. Segev, T. Coskun, D.N. Christodoulides, A. Vishwanath, Phys. Rev. Lett. **84**, 467 (2000).
20. B. Kibler, C. Michel, J. Garnier, A. Picozzi, Opt. Lett. **37**, 2472 (2012).
21. C. Rotschild, T. Schwartz, O. Cohen, and M. Segev, Nat. Photon. **2**, 371 (2008).
22. A. Picozzi, J. Garnier, Phys. Rev. Lett. **107**, 233901 (2011).
23. C. Michel, B. Kibler, J. Garnier, A. Picozzi, Phys. Rev. A **86**, 041801(R) (2012).
24. A. Picozzi, S. Pitois, G. Millot, Phys. Rev. Lett. **101**, 093901 (2008).
25. J. Garnier, A. Picozzi, Phys. Rev. A **81**, 033831 (2010).
26. C. Michel, B. Kibler, A. Picozzi, Phys. Rev. A **83**, 023806 (2011).
27. B. Kibler, C. Michel, A. Kudlinski, B. Barviau, G. Millot, A. Picozzi, Phys. Rev. E **84**, 066605 (2011).
28. G.P. Agrawal, *Nonlinear Fiber Optics*, 5th ed. (Academic, 2012).
29. K. J. Blow, D. Wood, IEEE J. Quantum Electron. **25**, 2665–2673 (1989).
30. J. Laegsgaard, Opt. Express **15**, 16110 (2007).
31. J. Travers, M. Frosz, J. Dudley, “Nonlinear fibre optics overview,” in J. Dudley, J.R. Taylor, *Supercontinuum Generation in Optical Fibers*, p. 32 (Cambridge University Press, Cambridge, 2010).
32. A.A. Voronin, A.M. Zheltikov, M. Aleksei, Opt. Lett. **33**, 1723 (2008).
33. M. Conforti, A. Marini, T.X. Tran, D. Faccio, F. Biancalana, Opt. Express **21**, 31239 (2013).
34. S.L. Musher, A.M. Rubenchik, V.E. Zakharov, Phys. Reports **252**, 177 (1995).
35. C. Montes, Phys. Rev. A **20**, 1081 (1979).

References

1. J. Garnier, M. Lisak, A. Picozzi, "Toward a wave turbulence formulation of statistical nonlinear optics," *J. Opt. Soc. Am. B* **29**, 2229 (2012).
2. T. Hansson, D. Anderson, M. Lisak, V.E. Semenov, U. Österberg, "Propagation of partially coherent light beams with parabolic intensity distribution in noninstantaneous nonlinear Kerr media," *J. Opt. Soc. Am. B* **25**, 1780 (2008).
3. A. Piskarskas, V. Pyragaitė, and A. Stabinis, "Generation of coherent waves by frequency up-conversion and downconversion of incoherent light," *Phys. Rev. A* **82**, 053817 (2010).
4. J. Laurie, U. Bortolozzo, S. Nazarenko, S. Residori, "One-dimensional optical wave turbulence: Experiment and theory," *Phys. Rep.* **514**, 121 (2012).
5. Y. Silberberg, Y. Lahini, Y. Bromberg, E. Small, R. Morandotti, "Universal Correlations in a Nonlinear Periodic 1D System," *Phys. Rev. Lett.* **102**, 233904 (2009).
6. P. Aschieri, J. Garnier, C. Michel, V. Doya, A. Picozzi, "Condensation and thermalization of classical optical waves in a waveguide," *Phys. Rev. A* **83**, 033838 (2011).
7. C. Conti, M. Leonetti, A. Fratallocchi, L. Angelani, G. Ruocco, "Condensation in Disordered Lasers: Theory, 3D+1 Simulations, and Experiments," *Phys. Rev. Lett.* **101**, 143901 (2008).
8. S. Babin, D. Churkin, A. Ismagulov, S. Kablukov, E. Podivilov, "Four-wave-mixing-induced turbulent spectral broadening in a long Raman fiber laser," *J. Opt. Soc. Am. B* **24**, 1729 (2007).
9. E. Turitsyna, S. Smirnov, S. Sugavanam, N. Tarasov, X. Shu, S. Babin, E. Podivilov, D. Churkin, G. Falkovich, S. Turitsyn, "The laminar-turbulent transition in a fibre laser," *Nat. Photon.* **7**, 783 (2013).
10. R. Weill, B. Fischer, O. Gat, "Light-Mode Condensation in Actively-Mode-Locked Lasers," *Phys. Rev. Lett.* **104**, 173901 (2010).
11. A. Schwache, F. Mitschke, "Properties of an optical soliton gas," *Phys. Rev. E* **55**, 7720 (1997).
12. B. Barviau, B. Kibler, A. Picozzi, "Wave-turbulence approach of supercontinuum generation: Influence of self-steepening and higher-order dispersion," *Phys. Rev. A* **79**, 063840 (2009).
13. B. Barviau, J. Garnier, G. Xu, B. Kibler, G. Millot, A. Picozzi, "Truncated thermalization of incoherent optical waves through supercontinuum generation in photonic crystal fibers," *Phys. Rev. A* **87**, 035803 (2013).
14. M. Erkintalo, M. Surakka, J. Turunen, A.T. Friberg, G. Genty, "Coherent-mode representation of supercontinuum," *Opt. Lett.* **37**, 169 (2012).
15. J. Garnier, G. Xu, S. Trillo, A. Picozzi, "Incoherent dispersive shocks in the spectral evolution of random waves," *Phys. Rev. Lett.* **111**, 113902 (2013).
16. S. Derevyanko, E. Small, "Nonlinear propagation of an optical speckle field," *Phys. Rev. A* **85**, 053816 (2012).
17. P. Suret, S. Randoux, H.R. Jauslin, A. Picozzi, "Anomalous thermalization of nonlinear wave systems," *Phys. Rev. Lett.* **104**, 054101 (2010).
18. M. Mitchell, Z. Chen, M. Shih, and M. Segev, "Self-trapping of partially spatially incoherent light," *Phys. Rev. Lett.* **77**, 490 (1996).
19. M. Soljacic, M. Segev, T. Coskun, D.N. Christodoulides, A. Vishwanath, "Modulation Instability of Incoherent Beams in Noninstantaneous Nonlinear Media," *Phys. Rev. Lett.* **84**, 467 (2000).
20. B. Kibler, C. Michel, J. Garnier, A. Picozzi, "Temporal dynamics of incoherent waves in noninstantaneous response nonlinear Kerr media," *Opt. Lett.* **37**, 2472 (2012).
21. C. Rotschild, T. Schwartz, O. Cohen, and M. Segev, "Incoherent spatial solitons in effectively-instantaneous nonlocal nonlinear media," *Nat. Photon.* **2**, 371 (2008).
22. A. Picozzi, J. Garnier, "Incoherent soliton turbulence in nonlocal nonlinear media," *Phys. Rev. Lett.* **107**, 233901 (2011).
23. C. Michel, B. Kibler, J. Garnier, A. Picozzi, "Temporal incoherent solitons supported by a defocusing nonlinearity with anomalous dispersion," *Phys. Rev. A* **86**, 041801(R) (2012).
24. A. Picozzi, S. Pitois, G. Millot, "Spectral incoherent solitons: a localized soliton behavior in the frequency domain," *Phys. Rev. Lett.* **101**, 093901 (2008).
25. J. Garnier, A. Picozzi, Unified kinetic formulation of incoherent waves propagating in nonlinear media with noninstantaneous response," *Phys. Rev. A* **81**, 033831 (2010).
26. C. Michel, B. Kibler, A. Picozzi, "Discrete spectral incoherent solitons in nonlinear media with noninstantaneous response," *Phys. Rev. A* **83**, 023806 (2011).
27. B. Kibler, C. Michel, A. Kudlinski, B. Barviau, G. Millot, A. Picozzi, "Emergence of spectral incoherent solitons through supercontinuum generation in photonic crystal fiber," *Phys. Rev. E* **84**, 066605 (2011).
28. G.P. Agrawal, *Nonlinear Fiber Optics*, 5th ed. (Academic, 2012).
29. K. J. Blow, D. Wood, "Theoretical description of transient stimulated Raman scattering in optical fibers," *IEEE J. Quantum Electron.* **25**, 2665–2673 (1989).
30. J. Laegsgaard, "Mode profile dispersion in the generalized nonlinear Schrödinger equation," *Opt. Express* **15**, 16110 (2007).
31. J. Travers, M. Frosz, J. Dudley, "Nonlinear fibre optics overview," in J. Dudley, J.R. Taylor, *Supercontinuum Generation in Optical Fibers*, p. 32 (Cambridge University Press, Cambridge, 2010).
32. A.A. Voronin, A.M. Zheltikov, M. Aleksei, "Soliton self-frequency shift decelerated by self-steepening," *Opt. Lett.* **33**, 1723 (2008).
33. M. Conforti, A. Marini, T.X. Tran, D. Faccio, F. Biancalana, "Interaction between optical fields and their conjugates in nonlinear media" *Opt. Express* **21**, 31239 (2013).
34. S.L. Musher, A.M. Rubenchik, V.E. Zakharov, "Weak Langmuir turbulence," *Phys. Reports* **252**, 177 (1995).
35. C. Montes, "Photon soliton and fine structure due to nonlinear Compton scattering," *Phys. Rev. A* **20**, 1081 (1979).