

#### A phase I/II clinical trial of autologous myoblast transplantation in facioscapulohumeral muscular dystrophy

Sabrina Sacconi, Jean-Thomas Vilquin, Serge Colson, Jonathan Bocquet, Jacques Darcourt, Jérémy Garcia, Véronique Tanant, Emmanuelle Salort-Campana, Pierre Carlier, Jean-Pierre Marolleau, et al.

#### ▶ To cite this version:

Sabrina Sacconi, Jean-Thomas Vilquin, Serge Colson, Jonathan Bocquet, Jacques Darcourt, et al.. A phase I/II clinical trial of autologous myoblast transplantation in facioscapulohumeral muscular dystrophy. Myology 2016, Mar 2016, Lyon, France. hal-02394504

#### HAL Id: hal-02394504 https://hal.science/hal-02394504v1

Submitted on 4 Dec 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.









Les Programmes Hospitaliers de Recherche Clinique (PHRC)

# A phase I/II clinical trial of autologous myoblast transplantation in facioscapulohumeral muscular dystrophy

Sabrina Sacconi<sup>1,2</sup>, Jean-Thomas Vilquin<sup>3,4,5</sup>, Serge Colson<sup>2,6</sup>, Jonathan Bocquet<sup>7</sup>, Jacques Darcourt<sup>2,8</sup>, Jérémy Garcia<sup>1</sup>, Véronique Tanant<sup>1</sup>, Emmanuelle Salort-Campana<sup>9</sup>, Pierre Carlier<sup>5,10</sup>, Jean-Pierre Marolleau<sup>11</sup>, Daniel Skuk<sup>12</sup>, Jérôme Larghéro<sup>13</sup>, Claude Desnuelle<sup>1,2</sup>

<sup>1</sup>Neuromuscular Disease specialized center, Pasteur 2 University Hospital - IRCAN, CNRS UMR 7284/INSERM U1081; <sup>2</sup>University of Nice-Sophia Antipolis; <sup>3</sup>Sorbonne Universités UPMC Univ Paris 06, Inserm UMRS974, CNRS FRE3617; <sup>4</sup>Myology Research Center, Paris; <sup>5</sup>Pitié Salpêtrière Hospital, Paris; <sup>6</sup>University of Toulon, Laboratory of Human Motricity, Education and Health, Nice; <sup>7</sup>Clinique St Jean, Toulon; <sup>8</sup>TIRO Laboratory, UMRE 4320, iBEB, DSV, CEA, Nice; <sup>9</sup>Neuromuscular Diseases and ALS specialized center, La Timone University Hospital, Aix-Marseille University, Marseille; <sup>10</sup>AIM-CEA, NMR Laboratory, CEA, I2BM, MIRCen, Paris; <sup>11</sup>EA4666, LNPC, UPJV - Department of Clinical Hematology and Cell Therapy, Amiens University Medical Center, Amiens; <sup>12</sup>Neurosciences Division-Human Genetics, CHUQ Research Center-CHUL, Quebec; <sup>13</sup>Cell Therapy Unit, Inserm, CIC Biotherapies, U1160, University Institute of Hematology, APHP Saint Louis Hospital, University Paris Diderot, Sorbonne Paris Cité, Paris.










## Introduction

Facioscapulohumeral muscular dystrophy type 1 (FSHD1) is one of the most frequent adult myopathies (1/20.000), with selective involvement of specific groups of muscles : facial, scapular fixator, anterior foreleg muscles, abdominal and humeral muscles.

Vastus lateralis (VL) is usually spared clinically until late stages of the disease, and myoblasts grown from VL have similar behaviour in vivo and in vitro than myoblasts from control patients



# Purpose of the study

Primary endpoints: Safety, feasability

- Feasability of cell preparation
- Safety of intramuscular injections of cells
- Clinical and biological tolerance of cell transplantation

#### -> Proposal:

Transplantation of autologous myoblast from spared muscle (VL) into an affected muscle as the Tibialis anterior (TA) muscle could locally improve the muscle's regenerative capacities.

Secondary endpoints : Follow-up of muscle strength and resistance to fatigue over 2 years

- Mechanical testing of strength and resistance to fatigue
- Surface electromyography, MRI and FDG fixation by PET-Scan

#### Specific inclusion criteria

- Men and women aged 18-65.

- Clinical manifestations of DMFSH confirmed by molecular diagnosis (D4Z4 repeats).

- Lack of clinical deficit in at least one VL muscle (assessed by score at knee extension, MRC = 5), and absence of adipo-fibrotic invasion (assessed by MRI).

- Motor deficiency of at least one anterior leg (assessed by score at foot dorsiflexion, MRC < 4) and fatty infiltration in at least one TA muscle (assessed by MRI).


-> Three groups of at least three patients selected in a sequential fashion.

#### Clinical parameters

Informations collected repeatedly from early (D0) to late phases (1mo). **Clinical monitoring:** overall wellness, heart rate and pressure, fever, cutaneous status, pain, redness, edema...


### Cell cultures





1-2g of VL muscle harvested under local anaesthesia in Nice Hospital and shipped to the Saint Louis Cell Therapy Laboratory in Paris. Within 24h, biopsies minced, digested, filtered and cells grown in a myogenic proprietary medium. Cells settled and expanded on days 8, 11, 14, 17, 20/21. Phenotypical characterization (CD56 NCAM), viability, microbiological controls, endotoxins assessed according to classical procedures.

#### Autologous cell transplantations



800 million cells containing >50% CD56+ cells concentrated in an isotonic saline buffer.

Injection of cells suspended in 10ml, into half of TA muscle (surface: approx. 20  $\text{cm}^2$ ; volume: approx. 40  $\text{cm}^3$ ).

3 groups of patients : 3 injection modalities with increased densities of needle trajectories per surface unit :

**Biological monitoring:** blood formulation, sedimentation, inflammation (CRP), CPK, myoglobinemia, myoglobinuria, creatininemia, ions, calcemia, phosphoremia, transaminases, transferases.

#### MMT testing and Electromyography

**Global manual testing** performed at time of inclusion, then at 15 d, 21 d, 1, 2, 3, 4, 5, 6, 12 months. Measurement of dorsal and plantar flexion of ankles, extension and flexion of knees, abduction and antepulsion of shoulders, extension and flexion of arms.

Maximal voluntary contraction (MVC) of ankle dorsiflexor muscles were measured with an ankle dynamometer (isometric strength).

Fatiguing exercise consisted in a maximal isometric dorsiflexion lasting 30 s. A fatigue index (FI) was determined.

Muscle activity of TA muscle was recorded by bipolar sEMG electrodes (10 mm diameter, 20 mm inter-electrode distance).

**Common peroneal nerve stimulation** was induced with a constant-current stimulator to evoke the electrophysiological and associated mechanical responses. The peak-to-peak amplitudes of the three M<sub>max</sub> responses of the TA muscle and the **maximal amplitude** of the three mechanical twitches (Pt) elicited by single stimuli at rest were averaged. The root mean square (RMS) of the sEMG recordings of the TA muscle was analyzed over a 500ms period around the peak force.

#### NMR imagery

At inclusion, absence of fatty infiltration in one VL (donor muscle) and the presence of abnormal fatty infiltration in one TA (recipient muscle) are documented without Gadolinium.

Graded semi-quantitative analysis expressing the ratio fat signal / skeletal muscle signal obtained by NMR imaging using Gadolinium contrast agent and **T1 ponderation** done before, 1, 3 and 6 months after implantation to evaluate the inflammatory reaction and the evolution of the volume of fatty infiltration (at middle and lower parts of the thigh and of the leg).


Methodologies, products and clinical protocols validated by regulatory agencies.

Description of patients

#### **Patient characteristics**

Patient	Age at inclusion	Duration of disease at inclusion	Number of D4Z4 repeats	Global Sumscore at inclusion (/80)	Tibialis MMT score at inclusion	Quadrcieps MMT score at inclusion
FSHD1	54	9	6	62,99	2	5
FSHD2	51	16	7	64,66	3+	5
FSHD3	49	16	6	65,68	2-	5
FSHD4	44	23	7	61,33	1	5
FSHD5	60	10	7	62,65	2+	5
FSHD6	48	32	7	70,00	3-	5
FSHD7	52	15	7	65,33	1+	5
FSHD8	59	2	7	74,33	3-	5
FSHD9	64	15	6	73,33	4+	5
FSHD10	61	20	8	65,69	3-	5
FSHD11	59	8	9	63,33	2	5
Mean +/- SD	55 +/- 6,4	15 +/- 8,1		66,3 +/- 4,3		

#### **Clinical outcomes**



-> Group 1 : 64 sites, interspace 5 mm – Group 2 : 100 sites, interspace 4 mm – Group 3: 189 sites, interspace 3mm.

#### Results of Cell cultures

Cell culture feasibility									
Patient	Bio	psy	Culture	Final number			Microbiologic		
	Weight (g)	Initial number of cells (10 <sup>6</sup> )	duration (days)	of cells (millions)	Final CD56 (%)	Viability (%)	al controls / Endotoxins	Feasibility	
FSHD1	1,7	1,58	22	1238	96,2	93,4	Neg / Neg	Good	
FSHD2	1,9	0,95	20	1937	98,2	92,5	Neg / Neg	Good	
FSHD3	1,9	0,68	20	993	94,1	95,6	Neg / Neg	Good	
FSHD4	1,1	0,71	21	999	92	93,2	Neg / Neg	Good	
FSHD5	1,9	0,9	22	1357	71,3	92,5	Neg / Neg	Good	
FSHD6	1,4	2,55	21	2037	50,05	91,45	Neg / Neg	Good	
FSHD7	1,7	0,85	21	1767	63,6	98,1	Neg / Neg	Good	
FSHD8	1,1	0,28	21	1081	25,3 *	97	Neg / Not Done	Not grafted / Not reached	
FSHD9	1,2	2,09	21	2000	18,3 *	96,4	Neg / Not Done	Not grafted / Not reached	
FSHD10	1,6	1,26	28	804,5	95,94	95,7	Neg / Neg	Good	
FSHD11	2	0,38	20	1855	97,63	97,8	Neg / Neg	Good	

Inter-individual variabilities may be important. CD56+ cut-off reached 9 times out of 11 within three weeks. In two cases, biopsies looked fibro-adipocytic, purity was not reached. Muscle biopsy were then extemporaneously qualified using histology.

## MMT and Electromyography

Mean differential MMT in each methodological series										
		Serie 1			Serie 2		Serie 3			
	∆Treated	∆Untreated	Global ΔSumscore	∆Treated	∆Untreated	Global ΔSumscore	∆Treated	∆Untreated	Global ∆Sumscore	
M1	-19,97%	-6,79%	-17,51%	6,31%	0,00%	-0,94%	23,71%	0,00%	-1,88%	
M3	-13,31%	-2,59%	-11,48%	17,31%	0,00%	-1,28%	23,71%	0,00%	-2,24%	
M6	-16,62%	-9,38%	-10,16%	17,31%	0,00%	-2,32%	19,79%	0,00%	-1,74%	
M9	-22,12%	-9,38%	-8,09%	17,31%	-1,33%	-2,35%	18,21%	-8,27%	0,23%	
M12	-15,51%	-9,38%	-7,65%	28,64%	-1,33%	-1,86%	15,44%	-13,37%	-2,07%	

## **PET-Scan analysis**

<sup>18</sup>F-fluorodéoxyglucose (<sup>18</sup>FDG) accumulates in some tissues as a function of their metabolic activity, especially in brain, myocardium and skeletal muscle upon exercise.

Measurements have been done at the level of TA upon standardized exercise before implantation, then 3 and 6 months later to quantify the volumes of metabolically active muscle tissue.

#### Acknowledgments

The authors wish to thank the patients and their families for their motivation, together with the several collaborators involved in the set up of this study at multiple levels.

	ent	_	- 1	N)	00	g/L	. at	Z	a, 1m)		
FSHD1	Y (< 50 mm)	Ν	Ν	Ν	Ν	N	Ν	Y (clinically not significant)	Ν	Y	Good
FSHD2	ND	ND	N	Ν	Ν	N	Ν	Ν	Ν	Y	Good
FSHD3	Y (< 50 mm)	ND	Ν	Ν	Ν	Ν	Ν	Ν	Ν	Y	Good
FSHD4	Ν	Ν	Ν	Ν	Ν	N	Ν	Ν	Ν	Y	Good
FSHD5	Ν	ND	Ν	Ν	Ν	Ν	Ν	Ν	Ν	Y	Good
FSHD6	Ν	ND	N	Ν	Ν	Ν	Ν	Ν	Ν	Y	Good
FSHD7	Ν	Ν	Ν	Ν	Ν	Ν	Ν	Y (CRP 160,2)	AN	Y	Questionable
FSHD10	Ν	Ν	Ν	Ν	Ν	Ν	Ν	Y (clinically not significant)	Ν	Y	Good
FSHD11	Ν	ND	Ν	Ν	Ν	Ν	Ν	Ν	Ν	Y	Good

#### **PET-Scan evolution**

No significant changes were noted over the follow-up period.

The maximal voluntary isometric strength of the dorsiflexor muscles remained constant during the follow-up period. The **fatigue index** of the dorsiflexor muscles of the experimental leg tend to decrease over the 12 months whereas the one from the control leg fluctuated.

The neuromuscular transmission was preserved after the cell therapy. The mechanical twitch response of the dorsiflexor muscles of the experimental leg slightly increased over the follow-up period.

Taken together, and considering that no changes were observed on the control leg, these results lead us to conclude that the **cell therapy may have positively affected** the experimental leg.

# **Discussion and Conclusions**

Cell productions were feasible but the quality of the initial muscle biopsy is important. Cell administrations were feasible and clinically well tolerated by all patients but one. The control of local cell distribution may be improved by echographic monitoring. Results show slight increases in twitch response and slight decrease in fatigue in the 3rd group. The combination of cell type (myoblasts) and of the modality (dense multisite injections) may have positively affected the TA muscle, BUT: No clinically significant gain of function perceived by FSHD patients, and no significant changes were noted at MRI and PET-Scan. The local FSHD1 degenerated muscle environment may be detrimental to the stability of the fibers or of the niches, and muscle regeneration may have been inefficient, too transitory, aborted or too unstable.

The slight muscle strength increase may not be clinically significant for FSHD patients, but may improve the quality of life of patients with more advanced muscle loss (e.g. DMD) patients) in other indications.