



**HAL**  
open science

## World Scientists' Warning of a Climate Emergency

William J Ripple, Christopher Wolf, Thomas M Newsome, Phoebe Barnard,  
William R Moomaw, Philippe Grandcolas

► **To cite this version:**

William J Ripple, Christopher Wolf, Thomas M Newsome, Phoebe Barnard, William R Moomaw, et al.. World Scientists' Warning of a Climate Emergency. Bioscience, In press, 10.1093/biosci/biz088/5610806 . hal-02397151

**HAL Id: hal-02397151**

**<https://hal.science/hal-02397151>**

Submitted on 28 Apr 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# World Scientists' Warning of a Climate Emergency

WILLIAM J. RIPPLE, CHRISTOPHER WOLF, THOMAS M. NEWSOME, PHOEBE BARNARD, WILLIAM R. MOOMAW, AND 11,258 SCIENTIST SIGNATORIES FROM 153 COUNTRIES (LIST IN SUPPLEMENTAL FILE S1)

**Scientists have a moral obligation** to clearly warn humanity of any catastrophic threat and to “tell it like it is.” On the basis of this obligation and the graphical indicators presented below, we declare, with more than 11,000 scientist signatories from around the world, clearly and unequivocally that planet Earth is facing a climate emergency.

Exactly 40 years ago, scientists from 50 nations met at the First World Climate Conference (in Geneva 1979) and agreed that alarming trends for climate change made it urgently necessary to act. Since then, similar alarms have been made through the 1992 Rio Summit, the 1997 Kyoto Protocol, and the 2015 Paris Agreement, as well as scores of other global assemblies and scientists' explicit warnings of insufficient progress (Ripple et al. 2017). Yet greenhouse gas (GHG) emissions are still rapidly rising, with increasingly damaging effects on the Earth's climate. An immense increase of scale in endeavors to conserve our biosphere is needed to avoid untold suffering due to the climate crisis (IPCC 2018).

Most public discussions on climate change are based on global surface temperature only, an inadequate measure to capture the breadth of human activities and the real dangers stemming from a warming planet (Briggs et al. 2015). Policymakers and the public now urgently need access to a set of indicators that convey the effects of human activities on GHG emissions and the consequent impacts on climate, our environment, and society. Building on prior work (see supplemental file S2), we present a suite of graphical vital signs of climate change over the last 40 years for human activities that can affect GHG emissions and change the climate (figure 1), as well

as actual climatic impacts (figure 2). We use only relevant data sets that are clear, understandable, systematically collected for at least the last 5 years, and updated at least annually.


The climate crisis is closely linked to excessive consumption of the wealthy lifestyle. The most affluent countries are mainly responsible for the historical GHG emissions and generally have the greatest per capita emissions (table S1). In the present article, we show general patterns, mostly at the global scale, because there are many climate efforts that involve individual regions and countries. Our vital signs are designed to be useful to the public, policymakers, the business community, and those working to implement the Paris climate agreement, the United Nations' Sustainable Development Goals, and the Aichi Biodiversity Targets.

Profoundly troubling signs from human activities include sustained increases in both human and ruminant livestock populations, per capita meat production, world gross domestic product, global tree cover loss, fossil fuel consumption, the number of air passengers carried, carbon dioxide (CO<sub>2</sub>) emissions, and per capita CO<sub>2</sub> emissions since 2000 (figure 1, supplemental file S2). Encouraging signs include decreases in global fertility (birth) rates (figure 1b), decelerated forest loss in the Brazilian Amazon (figure 1g), increases in the consumption of solar and wind power (figure 1h), institutional fossil fuel divestment of more than US\$7 trillion (figure 1j), and the proportion of GHG emissions covered by carbon pricing (figure 1m). However, the decline in human fertility rates has substantially slowed during the last 20 years (figure 1b), and the pace of

forest loss in Brazil's Amazon has now started to increase again (figure 1g). Consumption of solar and wind energy has increased 373% per decade, but in 2018, it was still 28 times smaller than fossil fuel consumption (combined gas, coal, oil; figure 1h). As of 2018, approximately 14.0% of global GHG emissions were covered by carbon pricing (figure 1m), but the global emissions-weighted average price per tonne of carbon dioxide was only around US\$15.25 (figure 1n). A much higher carbon fee price is needed (IPCC 2018, section 2.5.2.1). Annual fossil fuel subsidies to energy companies have been fluctuating, and because of a recent spike, they were greater than US\$400 billion in 2018 (figure 1o).

Especially disturbing are concurrent trends in the vital signs of climatic impacts (figure 2, supplemental file S2). Three abundant atmospheric GHGs (CO<sub>2</sub>, methane, and nitrous oxide) continue to increase (see figure S1 for ominous 2019 spike in CO<sub>2</sub>), as does global surface temperature (figure 2a–2d). Globally, ice has been rapidly disappearing, evidenced by declining trends in minimum summer Arctic sea ice, Greenland and Antarctic ice sheets, and glacier thickness worldwide (figure 2e–2h). Ocean heat content, ocean acidity, sea level, area burned in the United States, and extreme weather and associated damage costs have all been trending upward (figure 2i–2n). Climate change is predicted to greatly affect marine, freshwater, and terrestrial life, from plankton and corals to fishes and forests (IPCC 2018, 2019). These issues highlight the urgent need for action.

Despite 40 years of global climate negotiations, with few exceptions, we have generally conducted business




**Figure 1.** Change in global human activities from 1979 to the present. These indicators are linked at least in part to climate change. In panel (f), annual tree cover loss may be for any reason (e.g., wildfire, harvest within tree plantations, or conversion of forests to agricultural land). Forest gain is not involved in the calculation of tree cover loss. In panel (h), hydroelectricity and nuclear energy are shown in figure S2. The rates shown in panels are the percentage changes per decade across the entire range of the time series. The annual data are shown using gray points. The black lines are local regression smooth trend lines. Abbreviation: Gt oe per year, gigatonnes of oil equivalent per year. Sources and additional details about each variable are provided in supplemental file S2, including table S2.

as usual and have largely failed to address this predicament (figure 1). The climate crisis has arrived and is accelerating faster than most scientists expected (figure 2, IPCC 2018). It is

more severe than anticipated, threatening natural ecosystems and the fate of humanity (IPCC 2019). Especially worrisome are potential irreversible climate tipping points and nature's

reinforcing feedbacks (atmospheric, marine, and terrestrial) that could lead to a catastrophic “hothouse Earth,” well beyond the control of humans (Steffen et al. 2018). These climate



**Figure 2. Climatic response time series from 1979 to the present.** The rates shown in the panels are the decadal change rates for the entire ranges of the time series. These rates are in percentage terms, except for the interval variables (d, f, g, h, i, k), where additive changes are reported instead. For ocean acidity (pH), the percentage rate is based on the change in hydrogen ion activity,  $a_{H^+}$  (where lower pH values represent greater acidity). The annual data are shown using gray points. The black lines are local regression smooth trend lines. Sources and additional details about each variable are provided in supplemental file S2, including table S3.

chain reactions could cause significant disruptions to ecosystems, society, and economies, potentially making large areas of Earth uninhabitable.

To secure a sustainable future, we must change how we live, in ways that

improve the vital signs summarized by our graphs. Economic and population growth are among the most important drivers of increases in  $CO_2$  emissions from fossil fuel combustion (Pachauri et al. 2014, Bongaarts and

O'Neill 2018); therefore, we need bold and drastic transformations regarding economic and population policies. We suggest six critical and interrelated steps (in no particular order) that governments, businesses, and the rest of

humanity can take to lessen the worst effects of climate change. These are important steps but are not the only actions needed or possible (Pachauri et al. 2014, IPCC 2018, 2019).

### Energy

The world must quickly implement massive energy efficiency and conservation practices and must replace fossil fuels with low-carbon renewables (figure 1h) and other cleaner sources of energy if safe for people and the environment (figure S2). We should leave remaining stocks of fossil fuels in the ground (see the timelines in IPCC 2018) and should carefully pursue effective negative emissions using technology such as carbon extraction from the source and capture from the air and especially by enhancing natural systems (see “Nature” section). Wealthier countries need to support poorer nations in transitioning away from fossil fuels. We must swiftly eliminate subsidies for fossil fuels (figure 1o) and use effective and fair policies for steadily escalating carbon prices to restrain their use.

### Short-lived pollutants

We need to promptly reduce the emissions of short-lived climate pollutants, including methane (figure 2b), black carbon (soot), and hydrofluorocarbons (HFCs). Doing this could slow climate feedback loops and potentially reduce the short-term warming trend by more than 50% over the next few decades while saving millions of lives and increasing crop yields due to reduced air pollution (Shindell et al. 2017). The 2016 Kigali amendment to phase down HFCs is welcomed.

### Nature

We must protect and restore Earth’s ecosystems. Phytoplankton, coral reefs, forests, savannas, grasslands, wetlands, peatlands, soils, mangroves, and sea grasses contribute greatly to sequestration of atmospheric CO<sub>2</sub>. Marine and terrestrial plants, animals, and microorganisms play significant roles in carbon and nutrient cycling and storage.

We need to quickly curtail habitat and biodiversity loss (figure 1f–1g), protecting the remaining primary and intact forests, especially those with high carbon stores and other forests with the capacity to rapidly sequester carbon (proforestation), while increasing reforestation and afforestation where appropriate at enormous scales. Although available land may be limiting in places, up to a third of emissions reductions needed by 2030 for the Paris agreement (less than 2°C) could be obtained with these natural climate solutions (Griscom et al. 2017).

### Food

Eating mostly plant-based foods while reducing the global consumption of animal products (figure 1c–d), especially ruminant livestock (Ripple et al. 2014), can improve human health and significantly lower GHG emissions (including methane in the “Short-lived pollutants” step). Moreover, this will free up croplands for growing much-needed human plant food instead of livestock feed, while releasing some grazing land to support natural climate solutions (see “Nature” section). Cropping practices such as minimum tillage that increase soil carbon are vitally important. We need to drastically reduce the enormous amount of food waste around the world.

### Economy

Excessive extraction of materials and overexploitation of ecosystems, driven by economic growth, must be quickly curtailed to maintain long-term sustainability of the biosphere. We need a carbon-free economy that explicitly addresses human dependence on the biosphere and policies that guide economic decisions accordingly. Our goals need to shift from GDP growth and the pursuit of affluence toward sustaining ecosystems and improving human well-being by prioritizing basic needs and reducing inequality.

### Population

Still increasing by roughly 80 million people per year, or more than 200,000 per day (figure 1a–b), the world

population must be stabilized—and, ideally, gradually reduced—within a framework that ensures social integrity. There are proven and effective policies that strengthen human rights while lowering fertility rates and lessening the impacts of population growth on GHG emissions and biodiversity loss. These policies make family-planning services available to all people, remove barriers to their access and achieve full gender equity, including primary and secondary education as a global norm for all, especially girls and young women (Bongaarts and O’Neill 2018).

### Conclusions

Mitigating and adapting to climate change while honoring the diversity of humans entails major transformations in the ways our global society functions and interacts with natural ecosystems. We are encouraged by a recent surge of concern. Governmental bodies are making climate emergency declarations. Schoolchildren are striking. Ecocide lawsuits are proceeding in the courts. Grassroots citizen movements are demanding change, and many countries, states and provinces, cities, and businesses are responding.

As the Alliance of World Scientists, we stand ready to assist decision-makers in a just transition to a sustainable and equitable future. We urge widespread use of vital signs, which will better allow policymakers, the private sector, and the public to understand the magnitude of this crisis, track progress, and realign priorities for alleviating climate change. The good news is that such transformative change, with social and economic justice for all, promises far greater human well-being than does business as usual. We believe that the prospects will be greatest if decision-makers and all of humanity promptly respond to this warning and declaration of a climate emergency and act to sustain life on planet Earth, our only home.

### Contributing reviewers

Franz Baumann, Ferdinando Boero, Doug Boucher, Stephen Briggs, Peter

Carter, Rick Cavicchioli, Milton Cole, Eileen Crist, Dominick A. DellaSala, Paul Ehrlich, Iñaki Garcia-De-Cortazar, Daniel Gilfillan, Alison Green, Tom Green, Jillian Gregg, Paul Grogan, John Guillebaud, John Harte, Nick Houtman, Charles Kennel, Christopher Martius, Frederico Mestre, Jennie Miller, David Pengelley, Chris Rapley, Klaus Rohde, Phil Sollins, Sabrina Speich, David Victor, Henrik Wahren, and Roger Worthington.

### Funding

The Worthy Garden Club furnished partial funding for this project.

### Project website

To view the Alliance of World Scientists website or to sign this article, go to <https://scientistswarning.forestry.oregonstate.edu>.

### Supplemental material

Supplemental data are available at *BIOSCI* online. A list of the signatories appears in supplemental file S1.

### References cited

- Briggs S, Kennel CF, Victor DG. 2015. Planetary vital signs. *Nature Climate Change* 5: 969.
- Bongaarts J, O'Neill BC. 2018. Global warming policy: Is population left out in the cold? *Science* 361: 650–652.
- Griscom BW, et al. 2017. Natural climate solutions. *Proceedings of the National Academy of Sciences* 114: 11645–11650.
- [IPCC] Intergovernmental Panel on Climate Change. 2018. *Global Warming of 1.5°C: An IPCC Special Report*. IPCC.
- [IPCC] Intergovernmental Panel on Climate Change. 2019. *Climate Change and Land*. IPCC.
- Pachauri RK, et al. 2014. *Climate Change 2014: Synthesis Report*. Intergovernmental Panel on Climate Change.
- Ripple WJ, Smith P, Haberl H, Montzka SA, McAlpine C, Boucher DH. 2014. Ruminants, climate change, and climate policy. *Nature Climate Change* 4: 2–5.
- Ripple WJ, Wolf C, Newsome TM, Galetti M, Alamgir M, Crist E, Mahmoud MI, Laurance WF. 2017. World scientists' warning to humanity: A second notice. *BioScience* 67: 1026–1028.
- Shindell D, Borgford-Parnell N, Brauer M, Haines A, Kuylenstierna J, Leonard S, Ramanathan V, Ravishankara A, Amann M, Srivastava L. 2017. A climate policy pathway for near- and long-term benefits. *Science* 356: 493–494.
- Steffen W, et al. 2018. Trajectories of the Earth System in the Anthropocene. *Proceedings*

of the National Academy of Sciences 115: 8252–8259.

*William J. Ripple (bill.ripple@oregonstate.edu) and Christopher Wolf (christopher.wolf@oregonstate.edu) are affiliated with the Department of Forest Ecosystems and Society at Oregon State University, in Corvallis and contributed equally to the work. Thomas M. Newsome is affiliated with the School of Life and Environmental Sciences at The University of Sydney, in Sydney, New South Wales, Australia. Phoebe Barnard is affiliated with the Conservation Biology Institute, in Corvallis, Oregon, and with the African Climate and Development Initiative, at the University of Cape Town, in Cape Town, South Africa. William R. Moomaw is affiliated with The Fletcher School and the Global Development and Environment Institute, at Tufts University, in Medford, Massachusetts.*

doi:10.1093/biosci/biz088