

Implementation of various approaches to study the prevalence, incidence and progression of disseminated neoplasia in mussel stocks

Erika A.V. Burioli, Suzanne Trancart, Alexis Simon, I. Bernard, Maud Charles, Elise Oden, Nicolas Bierne, Maryline Houssin

To cite this version:

Erika A.V. Burioli, Suzanne Trancart, Alexis Simon, I. Bernard, Maud Charles, et al.. Implementation of various approaches to study the prevalence, incidence and progression of disseminated neoplasia in mussel stocks. Journal of Invertebrate Pathology, 2019, 168, pp.107271. 10.1016/j.jip.2019.107271. hal-02404031

HAL Id: hal-02404031 <https://hal.science/hal-02404031>

Submitted on 25 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Implementation of various approaches to study the prevalence, incidence 1

and progression of disseminated neoplasia in mussel stocks 2

Burioli E.A.V1*, Trancart S.¹ , Simon A.² , Bernard I.³ , Charles M.¹ , Oden E.¹ , Bierne N.² , Houssin M.1,4 3

4*Corresponding author: <u>erikaastrid.burioli@gmail.com</u>, LABÉO, 1 Route de Rosel, 14280 St Contest, France

5¹ LABÉO, 1 Route de Rosel, 14280 St Contest, France

6² Institut des Sciences de l'Évolution UMR5554, CNRS-IRD-EPHE-UM, Route de Mende, 34090 Montpellier, France 7

3 Eurêka Modélisation, 13 rue de Kermarquer, Lézardrieux, France 8

4 Université de Caen Normandie, UMR BOREA, MNHN, UPMC, UCBN, CNRS-7208, IRD-207, Esplanade de 9 10la paix, 14000 Caen, France

11

Abstract 12

13Marine mussel production is of substantial economic interest in numerous coastal areas worldwide, 14 making crucial the study of pathologies that affect them. Disseminated neoplasia (DN) has recently 15been suggested to be linked to blue mussel, Mytilus edulis, mortality outbreaks observed in France 16 since 2014, although the evidence remains indirect. In order to improve DN detection and monitoring, 17we compared the sensitivity of four diagnostic tools, namely haemocytology, histology, flow 18 cytometry, and genetics. Haemocytological examination gave the best results in sensitivity and had the 19advantage of being non-invasive, allowing disease progression to be followed in affected mussels. 20Using this approach, we showed that DN progression is usually slow, and we provide evidence of 21 remission events. We observed a high diversity of forms and mitotic features of neoplastic cells 22located in the vesicular connective tissue but rarely in the haemolymph. Circulating cells occur as four 23 main types but are homogenous in morphology and DNA content within a single individual. 24Polyploidy proved very high, from 8N to 18N. Genetic analysis of haemolymph DNA showed that a *Mytilus trossulus* genetic signal was associated with almost all the DN cases here diagnosed by 25 26haemocytological examination, regardless of the DN type. This result corroborates DN is a 27transmissible cancer that first originated in a *M. trossulus* host and subsequently crossed into *M*. 28edulis. No pre-neoplastic conditions were detectable. The prevalence of the disease was quite low,

29which, together with the low morbidity observed in the lab, suggest DN is unlikely to be the direct 30 cause of mortality outbreaks in France.

31

32Keywords: Blue mussel, Genetic chimerism, Genomic abnormalities, Haemolymph cytology, 33 Neoplasia, Trossulus

34

351. Introduction

36Disseminated neoplasia (DN) has been reported in several species of bivalves belonging to various families, including *Cardiidae*, *Myidae*, *Mytilidae*, *Ostreidae*, *Tellinidae*, and *Veneridae* (Farley, 1969a; 37 38Bayne, 1976; Balouet et al., 1978; Auffret and Poder, 1986; Elston et al., 1992; Barber, 2004; Carballal 39et al., 2015), with a world-wide distribution. Within the *Mytilidae* family, the disease has been described in blue (*Mytilus edulis*), Mediterranean (*Mytilus galloprovincialis*), and Pacific (*Mytilus* 40 *trossulus*) mussels (Farley, 1969b; Green and Alderman, 1983; Moore et al., 1991; Elston et al., 1992; 41 42Galimany and Sunila, 2008; Carella et al., 2013; Gombač et al., 2013). The DN condition is 43 characterised by the presence of circulating anomalous enlarged basophilic cells (Lowe and Moore, 1978; Green and Alderman, 1983) and by their infiltration of hemal spaces of all organs, with the loss 44 45of vesicular connective tissue in severe cases. These round-shaped cells display enlarged 46heteromorphic and sometimes multiple nuclei with prominent nucleoli, and high nucleus to cell 47 diameter ratio (Farley and Spark, 1970; Lowe and Moore, 1978; Mix et al., 1979). Anomalies in DNA 48 content and ploidy in neoplastic bivalve cells have been reported in many studies (Lowe and Moore, 1978; Elston et al., 1990; González-Tizón et al., 2000; da Silva et al., 2005; AboElkhair et al., 2009; 49 50Le Grand et al., 2010; Díaz et al., 2013; Vassilenko and Baldwin, 2014; Carella et al., 2017).

51Various aetiological origins have been suggested. Exposure to pollutants and sublethal levels of 52biotoxins were suspected to be involved in DN development (Lowe and Moore, 1978; Reinisch et al., 1984; Landsberg, 1996) but was not confirmed in subsequent studies (Green and Alderman, 1983). On 53 54the other hand, the transmission of the disease between mollusc individuals including mussels, through 55 cohabitation or haemolymph injection, has been proven (Elston et al., 1988; Twomey and Mulcahy, 1988; Kent et al., 1991; House, 1997; Weinberg et al., 1997), indicating that an infective agent, or the 56 57 neoplastic cells themselves, could be involved. DN transmission has been effected by injection of 58 neoplastic whole cell preparations, but when cell-free 0.45 μ m filtered preparations were used, 59 conclusions by different researchers on the success of the disease transmission were not unanimous 60(McLaughlin et al., 1992; House et al., 1998; Collins and Mulcahy, 2003).

61 Reverse transcriptase activity has been reported in DN-affected bivalves (House et al., 1998; Romalde 62et al., 2007; AboElkhair et al., 2009) and some studies have highlighted the presence of a retrovirus-63like agent that could readily produce the chromosomal aberrations associated with DN and possesses a 64 reverse transcriptase (Oprandy et al., 1981; Oprandy and Chang, 1983; Romalde et al., 2007). 65However, in most cases, no infectious agent was confirmed and AboElkhair et al. (2012) did not 66 provide clear evidence of retrovirus involvement in the disease. Recent studies (Metzger et al., 2015; Metzger et al., 2016) showed that DN can sometimes be a contagious form of neoplasia. Transmissible 67 68 cancers, evidenced by intra- or inter-species genetic chimerism, have recently been identified in 69 various marine bivalves (Metzger et al., 2016), including mussels. Riquet et al. (2017) identified 70 genetic chimerism in some French *M. edulis* individuals that carried *M. trossulus* alleles at diagnostic 71 markers even though this species is not present in France, suggesting that this could be related to the 72 contagious cancer described by Metzger et al. (2016).

73Mussel farming is of great economic importance in several European countries. French production is 74 represented by two species, M. edulis and M. galloprovincialis, which are able to hybridise in several 75 areas where they coexist (Bierne et al., 2003). Production in France amounts to around 75.000 tons 76each year (FAO, 2016). However, since 2014, mass mortality events (90-100%) have occurred in France, beginning in the Pertuis Charentais (Nouvelle-Aquitaine, France) (Béchemin et al., 2015) and, 77 78since 2016, in other farming areas in Brittany (France) (Bernard and Allain, 2017, Bernard et al., 2017; 792018). The causes of these mortality events are still unresolved. Some studies hypothesized the 80involvement of *Vibrio* species (Ben Cheikh et al., 2016, 2017; Oden et al., 2016), probably coupled 81 with stressful environmental conditions (Polsenaere et al., 2017). Benabdelmouna and Ledu (2016)

82associated abnormal outbreaks of mussel mass mortality with poor cytogenetic quality, particularly in 83terms of ploidy variations in circulating cells. In a second study, Benabdelmouna et al. (2018) linked 84 these genomic abnormalities to ongoing neoplastic processes. DN was also suspected to be involved in 85these events after the finding of a high prevalence of DN in a *M. edulis* collection from Lannion 86(France) affected by mortality in October 2016 (Burioli et al., 2017). Massive mortalities of 87 economically important bivalve stocks have occasionally been linked to high prevalence of DN in 88other parts of the world (Frierman and Andrews, 1976; Elston et al., 1992; McGladdery et al., 2001; 89 Villalba, et al., 2001).

90We studied the prevalence of DN in various mussel stocks, some of them regularly affected by 91 mortality events. Because we wanted to maintain our live mussel specimens to follow the incidence 92 and progression of the disease in the lab, we tested the sensitivity of two non-invasive methods, based 93on cytology and flow cytometry of haemolymph. On sub-samples, we compared the results of these 94two methods with histological observation and genetic analysis of haemolymph DNA to diagnose the 95DN state. Finally, we used cytology to follow DN progression in five mussel groups over 4 to 7 96months.

97

282. Materials and Methods

2.1. Sampling 99

100 Eighteen batches of mussels were collected during sampling campaigns conducted between July 2017 101 and June 2018. A total of 2,188 mussels were collected from six farming sites and a wild population 102(Table 1 and Figure 1). We also reported the origin areas where mussel spat was collected from the 103 wild in the prior year before being moved to their final growing sites. At time of sampling, individuals 104 from 12 batches were more than 1 year in age, while the others were aged approximately 1 year or less 105(the age of wild individuals is unknown). Sampled mussels were farmed on the intertidal zone (Agon-106 Coutainville, St Brieuc, Daoulas, and Pénestin) or in suspension in deep waters (Lannion and 107 Camaret). Wild mussels (Roscanvel) were collected on the intertidal zone. Sixteen batches were 108 previously identified as mainly *M. edulis* and two as mainly *M. galloprovincialis* according to 109 morphological characters and by genotyping the repetitive region of foot protein 1 in 10 individuals 110 per batch according to Inoue et al. (1995).

111A monitoring program for mussel mortality was conducted in Northern Brittany during 2017 and 2018 112(Bernard et al., 2017, 2018), in the sites of St Brieuc, Lannion, and Daoulas. The station of Daoulas 113was affected by heavy mortality between March and June 2017 ($\geq 65\%$ cumulative mortality rate of 114the period) and during May 2018 (about 30%). In Lannion, the phenomenon was weaker but lasted at 115least until July 2017 with \geq 35% cumulative mortality. Unfortunately, this population was not 116 monitored thereafter. No anomalous mortality was observed in St Brieuc. The sites of Roscanvel, 117 Camaret, and Pénestin were not included in this monitoring program, thus no official information on 118 putative mortality is available for these populations. However, the farmer at Camaret reported 119 cumulative mortality ranging between 60% and 80% in 2017. In Normandy, according to the French 120network for the monitoring of mollusc health REPAMO (https://wwz.ifremer.fr/sante_mollusques/), a 121 mortality event started in Agon-Coutainville in June 2018.

122

2.2. Evaluation of the various diagnostic methods 123

124 Because we wanted to maintain mussels alive with the aim to study the incidence of DN and the 125 progression of the disease, we tested the non-invasiveness of regular haemolymph drawing by 126 comparing the mortality of sampled and naïve individuals. Mortality was monitored daily for 2 months 127in four mussel batches (N=791) (see Table 1). Cumulative mortality rates were compared for 128individuals from which 100 µL of haemolymph was drawn every 3 weeks and those not subjected to 129haemolymph drawing (naïve) (see Table 2). Once the mussel batches arrived at the LABÉO 130laboratory, the groups used for this trial were subdivided and held in 10-L tanks, a maximum of 100 131 individuals per tank, in natural sea water maintained at 15° C, under oxygenation conditions. Complete 132 water changes were done on alternate days.

133Haemocytological observation was conducted on all batches on their arrival at the laboratory. The 134 sensitivities of haemocytological observation and flow cytometry were then compared and 135 accompanied by histological observation and genetic analysis in some batches (see Table 1). A total of 136331 individuals underwent haemolymph drawing to allow both cytological observation and flow 137 cytometry. We used a neoplastic individual from Batch 2 to implement the flow cytometry protocol. 138All individuals from the Batches 10, 12, and 13 ($N=241$) were simultaneously subjected to both 139 analyses on their arrival; flow cytometry was not immediately carried out for Batch 11, but on 140 survivors in February alongside haemocytological observation; for Batches 2 and 5, flow cytometry 141was carried out on randomly sampled individuals. Histological observation is an invasive method. 142 Thus, when the batch was kept in tank for a long period of time in order to study the DN incidence and 143 disease progression in affected mussels (Batches 2, 4, 5, and 8), histology was carried out on each 144 moribund individual (along with haemocytological observation) and on half of the survivors at the end 145 of the monitoring period (402 individuals in total). Genetic analysis was conducted on 114 individuals 146also subjected to haemocytological observation, flow cytometry, and histology. The various protocols 147 used are reported below.

2.3. Haemocytological examination 148

149The mussels were anaesthetised by bathing them for 30 min in a solution containing 50 g/L of 150 magnesium chloride dissolved in 2:1 distilled water:sea water (Suquet et al., 2009). Haemolymph (40 151 µl) was withdrawn from the adductor muscle with a 1-mL sterile syringe fitted with a 27-gauge needle 152 and refrigerated to avoid aggregation. After addition of 160 μ L of artificial seawater, the haemolymph 153 solution was "cytospun" (Shandon™ Cytospin™ 4, Thermo Scientific™; 10 min, 800 rpm) on coated 154 slides (Shandon[™] Cytoslides[™], Thermo Scientific[™]), fixed, stained with May-Grünwald Giemsa, 155 and observed under light microscopy (100X, 500X).

156The entire area of the cell spot on the cytoslide was scanned in all samples to search for neoplastic 157 cells, which are characterised by an enlarged diameter, elevated nucleus/cell diameter ratio (NCR), and 158high basophily. Four developmental stages of DN were differentiated according to the proportion of 159 neoplastic cells in haemolymph: "non-diseased" when no neoplastic cells were observed, "low 160 intensity" (less than 20% of neoplastic cells), "moderate intensity" (20% to 75%), and "high intensity" 161 (more than 75%).

162In order to determine if a pre-neoplastic stage could be detected in the individuals before the 163 appearance of the first typical neoplastic cells, the various haemocyte populations were then also 164 characterised in terms of morphology and relative proportions. The mussel conditions showing 165 atypical haemocytograms in terms relative percentage of cell populations according to statistical 166 analysis were considered to be affected by "haemocytic disorders". The cell and nucleus diameters of 167each haemocyte type were then measured. Cell and nuclear diameters were determined in 50 cells per 168cell type (when there were sufficient numbers), in 10 mussels/batch. The disorder-affected individuals 169were initially excluded from this data collection carried out to determine the normal condition. 170Nevertheless, in order to better characterise these anomalies, 50 cells per cell type in 1 to 3 atypical 171 mussels per batch were measured. Finally, neoplastic cells were described in the same way for all 172 affected mussels, with 1 to 30 neoplastic cells measured depending on the number of anomalous cells 173in the haemolymph.

174The total haemocyte count was performed directly from cytological stained slides made from 30 175 individuals in each of the 18 batches, excluding initially the individuals affected by DN and 176 "haemocytic disorders". Haemocytes were counted in four random microscope fields at 500X 177 magnification. Haemocyte densities per millilitre of haemolymph were estimated as follows:

179Then, total haemocyte count was also determined in mussels affected by neoplasia and "haemocytic 180disorders".

2.4. Flow cytometry analysis 181

182When flow cytometry analysis was conducted on mussel individuals at the same time as cytological 183analysis, a supplementary volume of 100 µL haemolymph was drawn and placed in a 1.5-mL 184 microtube; 500 µL of ethanol 100% was added to fix the cells, prevent aggregation, and increase cell 185 permeability. This suspension was stored at -20 $^{\circ}$ C until the next steps of the process. First, each 186aliquot of haemolymph was centrifuged at $600x$ g for 5 min and the pellet was resuspended in 500 µL 187 sterile artificial seawater (1 L of reverse osmosis purified water, 23 g of sodium chloride, 1.49 g of 188 potassium chloride, 0.3 g of calcium chloride, and 1.23 g of magnesium sulfate heptahydrate) for 15 189 min at room temperature. A second centrifugation at 600 x g for 5 min was carried out and the pellet 190was treated in 500 µL of propidium iodide (PI) solution: PI (Life technologies) 0.05 mg/mL; RNase A 191(Life technologies) 0.02 mg/mL; PBS (Eurobio 1X). Cells were stored for 45 min in the dark at room 192temperature. Haemocytes were then analysed by flow cytometry using a Cytoflex (Beckman Coulter) 193 equipped with a laser providing 50 mW at wavelength 488 nm. For each individual, 10,000 cells were 194 analysed at a medium flow rate $(30 \mu L/min)$ to collect multiparametric data. The measurement of 195DNA contents in haemolymph cells was based on the properties of PI, which intercalate into nucleic 196 acid molecules. When PI binds to nucleic acids its fluorescence intensity is proportional to the DNA 197 cell content, thus allowing the study of cell ploidy (Krishan, 1975). Because some cell populations had 198 extraordinarily high ploidy levels, each sample was run twice with two different gain settings. In order 199to remove doublets from single cells, the IPBLUE-height *versus* IPBLUE-area dot-plot was used and a 200 region comprising single cells was drawn on this representation (adapted from da Silva et al., 2005), 201doublets being excluded on the right of the selected area. The cells included in this region were then 202 plotted on a histogram in linear scale, used to calculate the percentage of cell populations according to 203their DNA content. In particular, the percentages of cells in the normal diploid G_0/G_1 and $S/G_2/M$ 204 phases were estimated according to Delaporte et al. (2008) and Benabdelmouna and Ledu (2016).

2.5. Histological examination 205

206A standard section of the mussel body, \sim 5 mm thick, containing the digestive gland, gut, mantle, gills, 207 and foot was fixed for 48 hrs in Davidson's fixative. Fixed tissues were then processed for routine 208histological examination and 4-µm thick sections were stained with Harris's haematoxylin and eosin 209(H&E). Histological sections were examined under a light microscope for the presence of abnormal 210 cells related to neoplasia.

2.6. Genetics 211

212DNA was extracted from the haemolymph using a NucleoMag 96 Tissue kit (Macherey-Nagel) and a 213KingFisherTM Flex (ThermoFisher Scientific). Following the study of Riquet et al. (2017) we used 214SNP markers diagnostic between *M. trossulus* and *M. edulis/M. galloprovincialis*. We used 12 SNPs 215(supplementary Table S2) described in Simon et al. (2018). These markers were identified as being 216diagnostic (fixed for alternative alleles in *M. trossulus* and *M. edulis/M. galloprovincialis*) in the 217 dataset of Fraïsse et al. (2016) and were subsequently verified to remain diagnostic with larger sample 218 sizes (Simon et al., unpublished). Non-hybrid individuals can nonetheless occasionally be 219heterozygous at one or two of the 12 loci because incomplete lineage sorting and introgression is 220 pervasive in the *M. edulis* complex of species. Genotyping was subcontracted to LGC-genomics and 221 performed with the KASPTM chemistry (Kompetitive Allele Specific PCR, Semagn et al., 2014). The 222 results are a combination of two fluorescence values, one for allele $1 (X)$, and another for allele $2 (Y)$. 223Following Cuenca et al. (2013) we transformed the data to produce a single measure of the relative 224 fluorescence of the two alleles, from 0 when the fluorescence of allele 1 dominates, to 1 when the 225 fluorescence of allele 2 (*i.e.* the *trossulus* allele) dominates, using the following formula: $y' = Y/(X+Y)$. 226The fluorescence of the *trossulus* allele was averaged over the 12 loci to obtain a compound 227 multilocus estimate.

2.7. Determination of DN prevalence, incidence and progression 228

229The occurrence of DN in mussels was described in terms of prevalence, defined as the number of 230 mussels presenting anomalous cells in 100 individuals investigated by cytological examination. We 231determined the initial DN prevalence in each of the 18 batches and the developmental stage of the 232 disease in affected animals.

233The incidence of the disease, defined as the number of new cases observed in survivors between two 234 consecutive haemolymph drawings expressed in percentage, and its progression in affected individuals 235 were evaluated in five batches (2, 4, 5, 8, and 11). Individuals were maintained in six different 10-L 236tanks (two tanks were used for Batch 11), in natural sea water maintained at around 15° C, under 237 oxygenation conditions. Complete water changes were done on alternate days. Haemolymph drawings 238 and haemocytological observations were carried out on all specimens, each month, for 4 to 7 months 239 according to the batch (Table 1). In individuals that developed DN during the monitoring, a 240 retrospective observation of haemocytological slides was carried out in search of haemocytogram 241 modifications or changes in cell morphology that could represent a pre-neoplastic stage.

2.8. Statistical analysis 242

243At the end of the trial for evaluation of the invasiveness of haemolymph drawing, the significance of 244the differences in mortality rate between the groups was evaluated using a chi-square test. Data from 245 morphological characterisation (cell diameter, nucleus diameter and NCR), relative percentage of cell 246 populations and total haemocyte count, determined by haemocytological observation, were analysed 247 statistically to determine the significance of the inter-individual and inter-site differences. An 248Anderson-Darling test for normality was initially performed. If the data were found to follow a normal 249 distribution, they were analysed using analysis of variance (ANOVA) followed by Tukey's post-hoc 250test on mean values. A Kruskal-Wallis test, followed by the ad-hoc multiple pair comparison test 251Kruskalmc (package pgirmess), were used on results requiring nonparametric analysis. Individuals 252 showing significant differences were considered to be affected by haemocytic disorders.

253Potential correlation between haemocytic anomalies and minority cells was evaluated by the 254 calculation of the Pearson correlation coefficient. The same inferential statistic tests were used to 255 compare the percentage of cells in the various peaks of ploidy during flow cytometry. Significance of 256the differences in the initial DN prevalence observed between the farming techniques (with and 257 without tide exposure) and between the two age categories (>1 year and \leq 1 year) was estimated 258through a chi-square test.

259Statistical analysis was performed using R software, version 3.3.1. Results with a p-value <0.05 were 260 considered statistically significant.

3. Results 262

3.1. Haemolymph drawing is non-invasive 263

264At the end of the trial to assess the invasiveness of the haemolymph drawing (2 months duration), 265 cumulative mortality in the different batches ranged between 12.0% and 38.2% (Table 2). No 266 significant differences were observed between control individuals and the individuals for which 267haemolymph was drawn in each batch. The haemolymph drawing method was, thus, confirmed to be 268non-invasive.

269

3.2. Haemocytological observations 270

271In DN-unaffected individuals, cytological examination confirmed the presence of two main cellular 272types, eosinophilic granulocytes (Figure 2A) and agranulocytes (Figure 2B and 2C). Granulocytes 273 were characterised by the presence of abundant brick-red granules in a beige cytoplasm, which varied 274in number and size. The nuclei were typically spherical and occasionally ovoid and eccentric. A small 275 number of granulocytes were observed to contain two or more nuclei (Figure 2D). Lucent vesicles, 276 consistent with phagosomes, were sometimes observed in these cells. The cytoplasm of agranulocytes 277 appeared bluish and no granules were present. However, this population appeared more 278heterogeneous, in terms of size, nucleus/cell diameter ratio (NCR), basophily, and nucleus shape. We 279 observed highly basophilic small cells (Figure $2B$) with a very high NCR and larger cells, 280 characterised by a light-blue cytoplasm and a lower NCR with some cells showing a U-shaped nucleus 281 (Figure 2C). In some individuals, very pale red or blue vesicle coloration was observed in a few cells. 282The results of the morphometric analysis carried out on cell and nucleus diameters and the NCR 283 values are reported in Table 3. Data collected from granulocytes showed a normal distribution while 284those from agranulocytes did not. According to both parametric (ANOVA) and nonparametric 285(Kruskal-Wallis) tests, no significant differences were observed among inter-individual and inter-site 286 measurements. From the first haemolymph drawing, we observed the presence of a few basophilic

287 granulocytes (Figure $2E$), smaller than eosinophilic granulocytes, characterised by numerous small 288blue granules in the cytoplasm in about 1.90% mussels (Table 4). When present, these cells 289 represented less than 1% of the total haemocyte population. They were consistently present in the 290batches from Daoulas (Batches 10, 14, and 15). Brown cells (Figure $2F$) containing small iridescent 291brown granules were also observed in about 10.83% of mussels. Their presence varied from batch to 292batch (Table 4) and were observed in 0% to 89% of individuals, depending on the site and the 293 sampling date, and increased relative to the number of bacteria in haemolymph, reaching a proportion 294of 6% of the circulating haemocytes in some individuals from Batch 9. Agranulocytes showed 295 aggregation capability with eosinophilic granulocytes accumulated around the aggregates. In several 296 cases, we observed "rosetting" where the central cell was a mitotic cell surrounded by agranulocytes. 297Phagocytosed cells were sometimes observed inside granulocytes.

298The investigation on the morphology of neoplastic cells was conducted on all the DN-affected mussels 299detected during the present study. They appeared enlarged, highly basophilic, with prominent nucleoli. 300The scatterplot for cell diameter and NCR ($Figure 3$) showed that neoplastic cells were separated from 301normal haemocytes, even if an overlap area existed with agranulocytes. Three distinct neoplastic 302 populations emerged from the analyses. The Kruskal-Wallis test showed a significant difference $303(p<0.05)$ in cell and nucleus diameters and for NCR among the three groups. These characteristics 304 corresponded only in part with the A and B types previously described (Lowe and Moore, 1978, Green 305 and Alderman, 1983; Carella et al., 2013). The most frequent cell type (approx. 40% of diseased 306 individuals), which we termed P-type (Figure 4A), had a lower NCR, a pleomorphic (usually lobed or 307 reniform) nucleus with many nucleolus-like structures and, typically, a larger cytoplasm diameter 308when compared with the second type cells. A rarer sub-type with smaller cells and round nucleus 309named Ps-type (approx. 4% of individuals) was also observed in a few individuals. The second most 310 prevalent cell type (approx. 32% of individuals), U-type (Figure 4B), showed a regular outline, scant 311 cytoplasm, an ovoid-shaped hyperchromatic nucleus, and one (occasionally 2) nucleolus. The third 312cell type, S-type, occurred less frequently (approx. 17% of individuals). It was represented by smaller 313 cells, with a diameter comparable to large agranulocytes, but highly more basophilic and with at least

314 one prominent nucleolus (Figure 4C). We used new terms to avoid any confusion with previous 315 descriptions: P-cells (pluri-nucleolus), and the smaller sub-type Ps-cells, U-cells (unique nucleolus), 316and S-cell (small cells). Their morphometric characteristics are summarised in Table 3. Most 317 individuals were affected by only one of the four neoplastic cell types but in few individuals $(7%)$, 318both types P and U were present. Very occasionally, binucleated and multinucleated neoplastic cells 319 were observed but only in P-type affected individuals. Both cell types contained cytoplasmic vacuoles, 320but they were particularly abundant in the P-type. Mitoses were very rare, but few multipolar atypical 321mitoses were seen (Figure 4D).

322

323

324

325 The total haemocyte count in normal mussels was estimated as a mean of 4.4 (\pm 1.3) x10⁶ cells/mL but, 326in the individuals sampled in February, haemocyte count was significantly lower, mean 9.8 (\pm 1.2) $327x10⁵$ cells/mL (ANOVA, p<0.05), in the three sampled sites, without significant inter-individual 328 variations. The proportion of granulocytes was mean 64 (\pm 4.3) %. Mitoses, easily observable at the 329 prophase and metaphase stages, were observed in most individuals and concerned both granulocytes 330and hyalinocytes.

331Neoplastic cells were initially observed in 59 of the 1791 specimens analysed by haemocytological 332 observation (3.3%). About 59% of the 59 specimens were lightly affected, 19% moderately, and 22% 333 heavily. The neoplastic cell-type was not associated with a specific disease stage. The total haemocyte 334 count in DN-affected individuals was not significantly higher. Two other haemocytic disorders were 335 observed. Their frequencies among the batches are reported in Table 4. "Granulocytosis" consisted in 336the significant proliferation (p <0.05) of eosinophilic granulocytes reaching >90% of the circulating 337haemocytes and a total mean haemocyte count $>8 \times 10^6$ cells/mL. No relevant differences in typical 338 cell morphology were observed. Conversely, in some individuals we observed a predominance of 339 agranulocytes and, in some specimens, a nearly total absence of granulocytes ("granulocytopenia")

340with a significant decrease of total mean haemocyte count (2.8 x 10^6 cells/mL). No significant 341 differences in the morphometric values were observed when compared with the agranulocytes of 342unaffected mussels. However, aggregates were nearly absent on these smears as also observed in S-343type neoplasia.

344We calculated a positive correlation between the basophilic granulocytes and granulocytopenia 345 frequencies (Pearson, 0.5127, p=0.02) but no correlation was detected between neoplasia frequencies 346 and other variables.

3.3 Histological observations 347

348Neoplastic cells were easily identifiable during histological observation. They appeared as enlarged, 349basophilic, agranular cells, with a high NCR and one or more nucleolus-like inclusions. Among the 350402 specimens examined histologically, the result was not interpretable in three samples due to 351 degradation. Disseminated neoplasia was observed in 65 individuals. All five morphological types of 352 atypical haemocytes previously described by Carella et al. (2013) were identified. In particular, in P-353type neoplasia (according to haemocytological observation), we observed a high heteromorphism of 354 nuclei and numerous multinucleated cells (Figure 5). The DN advanced condition was characterised 355by the infiltration and intense invasion of the vesicular connective tissue by enlarged, basophilic 356neoplastic cells. Atypical cells were also present in muscle and gills, where they induced enlargement 357 of the vessels. Generally, a high mitotic rate was observed in the neoplastic compartment, usually 358 represented by atypical mitoses. In less advanced stages, the foci of neoplastic cells were present 359 around the digestive tubules, the stomach or the intestine and, less intensely, in the vesicular 360 connective tissue. No correlation between DN stage and neoplastic cell types was observed. In 361addition, two types of inflammatory disorders were observed in about 20% of the individuals not 362 affected by DN. The first consisted of the local infiltration of connective tissue by the normal 363inflammatory component (granulocytes and agranulocytes). The second corresponded to the 364 granulocytomas previously described by Lowe & Moore (1979). The frequency of these lesions did 365not differ significantly among the batches and were not correlated with DN prevalence.

3673.4. Flow cytometry

368A total of 331 haemolymph samples were individually analysed by flow cytometry and 277 were 369 retained for their suitable quality. In particular, part of the samples from Batches 12 and 13 were 370 excluded. These two samplings were conducted just before the spawning period, usually occurring in 371March, and mature spermatozoids contaminated the haemolymph sample during drawing (as observed 372in haemocytology). In mussels diagnosed as asymptomatic for DN by haemocytological observation, 373we observed a main peak in the cytometry histogram representing the dominant cell population and 374 constituting the G_0/G_1 compartment (diploid) of the normal cell cycle, and a second peak attributable 375to the G_2M (4N) phase (Figure 6A). However, some mussels presented more complex patterns. Twenty 376 specimens from Daoulas and few specimens from other sites (Table 5), not affected by DN in 377haemocytology, displayed a second peak situated between the normal 2N and 4N peaks. The ploidy 378level of this anomalous peak was $2.92(\pm 0.29)$ N, ranging between 2.56 and 3.54N and represent 37933.12(±10.02)% of the circulating cells. No haemocytic disorder, increased phagocytosis, or gamete 380 presence was observed during haemocytological and histological observations in these mussels.

381

382Eight mussels showed profiles consistent with DN. All presented a cell population with an 383 extraordinarily high ploidy, in addition to the normal G_0/G_1 and G_2/M peaks (6B, 6C). The DNA 384 content of the neoplastic peak ranged from 4.18 to 9.9x the diploid DNA content, corresponding to 3858.36 to 18.8N. The neoplastic cell population within an individual was characterised by a low 386 dispersion of the value of ploidy, as confirmed by a low coefficient of variation $\leq 6\%$. Only one sample 387 from St. Brieuc had two aneuploid peaks corresponding to 9.4 and 11.64N (Figure 6C). The 388 percentages of each cell population in neoplastic mussels are reported in Table 6. There was no 389 correlation between the degree of ploidy and the disease stage. On the contrary, a trendline between 390ploidy and mean nucleus diameter of neoplastic cells, determined during haemocytological 391 observation, emerged (Figure 7).

393In order to estimate the cell replication activity, we computed the fraction of cells in the $S-G_2/M$ phase. 394We never observed a second peak on the cytometry histogram of the DN-affected individuals that 395 could represent the G_2/M phase of the neoplastic cell population, even with the lower amplifier gain. 396Among the mussels with no neoplastic disorder according to haemocytological observation, we 397 excluded those bearing the anomalous "nearly triploid" peak for the statistical analysis. The 398 individuals from Daoulas showed a significantly higher mean percentage of cells in the S-G₂/M phase 399 compared with the other sites (Kruskal-Wallis, K= 100.57, df=5, p < 2.2e-16), representing 7.45(±3.56)% *vs* 2.51(±1.92)%, respectively. Moreover, in 19.54% of the individuals from Daoulas, 400 401 more than 10% of circulating haemocytes were in the S-G2/M phases. Only one individual from Agon 402 exceeded slightly the 10% threshold (10.39%), considered by Elston et al. (1990) to be the upper limit 403to consider the mussel unaffected by genomic abnormalities. No hypodiploid peaks were observed.

3.5. Genetic analysis 404

405 Genetic analysis of haemolymph DNA was conducted with 12 diagnostic SNPs between *M. trossulus* 406and *M. edulis/M. galloprovincialis* in a subsample of 114 mussels, 11 of which DN-positive with 407haemocytology: a sample of 100 mussels from Batch 10, 6 mussels from Batch 2 (St Brieuc, 408including 3 DN-positive specimens), 6 mussels from Batch 5 (Lannion, including 5 DN-positive 409 specimens) and 2 mussels from Batch 8 and 11 (Camaret) were analysed. (Unfortunately, 410haemocytology and histology found only one specimen from Daoulas to be lightly affected.) The 411 compound multilocus estimates of the *trossulus* allele fluorescence are presented in Figure 8 and the 41212 single locus fluorescence values are provided in supplementary Figure S1. We found six samples 413with evidence of amplification of *trossulus* alleles (Figure 8): two specimens diagnosed as heavily 414 affected, three specimens diagnosed as moderately affected, and one lightly affected. Thus, 4 of the 415lightly affected and one of the four moderately affected specimens were negative for genetic 416 chimerism. The four types of neoplasia were detected positive by genetic analysis.

417

3.6. Sensitivity of the various diagnostic methods 418

419In our study, 115 individuals tested DN-positive by at least one of the four diagnostic methods. Among 420the 572 mussels showing interpretable results with at least two diagnostic techniques, results from 12 421individuals were not concordant among the techniques. The individual results of the four diagnostic 422 methods are detailed in supplementary Table S1. The sensitivities of histology and haemocytological 423 observation for DN-detection were compared in 399 mussels and results were in accordance in 42499.25% of cases.. All heavily and moderately-affected mussels diagnosed by haemocytology were 425 positive in histological observation, but two lightly affected individuals in haemocytology were not 426 detected by histology. On the contrary, in a mussel diagnosed as negative by haemocytology, a few 427 neoplastic cells were present around the stomach in histology. Flow cytometry, when compared with 428haemocytological observation of 277 haemolymph samples, positively diagnosed all the heavily and 429 moderately-diseased mussels while the lightly-diseased individuals showed profiles consistent with 430non-diseased individuals. Nevertheless, the cytometry histogram of one individual, alternatively 431 investigated only by haemocytology and diagnosed as non-diseased, displayed the presence of a few 432hyperploid cells compatible with neoplasia. The sensitivity of flow cytometry appeared significantly 433lower than haemocytological observation, with concordant results in 98.19% of samples. Finally, 434 according to the genetic analysis carried out on 114 mussels, genetic chimerism was detected in all the 435heavily-neoplastic stages, in 75% of the moderately-affected mussels, and in 20% of the lightly 436 neoplastic specimens. None of the non-diseased mussels was found to present this genetic anomaly. 437 Considering the present results, haemocytological observation was chosen for the monitoring of the 438 disease progression.

3.7. Prevalence and monitoring of the disease progression 439

According to haemocytological results, disseminated neoplasia was detected in all sites, except in 440 441Pénestin and Roscanvel (Table 4). Initial prevalence ranged between 0% and 25.45%, with a mean of 4423.29%. Neoplastic animals were found consistently in Lannion, Camaret, and Agon-Coutainville. The 443chi-square test revealed that the DN prevalence was higher in the groups with higher mortality $444(\chi^{2(1,0.05)} = 53.16$, p-value=3.07x10⁻¹³). The highest prevalences were observed in Batches 1, 3, 8, and 17; 445however, DN was absent or prevalence was low in Batches 15 and 18, respectively, which also had

446increased mortality. Additionally, DN prevalence in Batches 4, 5, 7, and 12 was in the average range 447but there was no increased mortality. Both species, *M. edulis* and *M. galloprovincialis*, sampled in 448 Lannion on $26th$ July and $1st$ August 2017, respectively, had comparable prevalence. Significant 449 differences (χ 2, p<0.05) were observed between both farming technique and animal age, with higher 450 prevalence in older mussels and those not exposed to tide. In all sites, we reported simultaneously the 451 presence of various neoplasia types.

452According to haemocytological observation, a total of 75 individuals of the 627 monitored were 453diagnosed with DN during the monitoring. Twenty-two of the monitored individuals were positive 454upon arrival and 53 developed the disease during the subsequent months. Incidence of DN between 455two consecutive haemolymph samplings is represented in Figure 9. Interestingly, in all the batches, we 456 observed an increase of incidence during the first 9 weeks followed by a decrease. In Batch 11, 457 monitored for 7 months, only two new cases appeared after the fourth month. There were fewer 458diseased individuals at the end of the monitoring in two batches (2 and 11) that had lower initial 459 prevalences $(\chi^2 (4, 0.05)=14.23; \text{ p-value}=0.006)$).

460In the individuals that developed DN, we did not observe previous changes in the haemocytogram or 461 in cell morphology before the occurrence of the first neoplastic cells. We represented the evolution of 462the disease in individuals acquiring the disease in Figure $10A$. Remission was observed in 12% of 463 diseased individuals (9/75), one previously diagnosed as high intensity stage, and concerned both P 464 and U-type neoplasia. Remission was characterized by a rapid decrease of the number of neoplastic 465 circulating cells some of which appeared picnotic or necrotic (until clearance in 3-6 weeks). The 466 remission stage was confirmed in five individuals analysed by histology; in three the remission was 467 complete and, in two, few neoplastic cells were still present in the connective tissue. No anomaly was 468 detectable by flow cytometry in the three mussels showing a total remission but, in one, the remission 469was partial and some hyperploid cells were still present. On the contrary, genetic analyses detected the 470 presence of *trossulus* chimerism in the single individual analysed by this method and otherwise 471 presenting an apparent total remission. If we consider only individuals tested negative at the first 472 haemolymph drawing (n=53), the heavy intensity stage was usually reached 6-9 weeks after the first

473 neoplastic cells appeared. However, in a few individuals, the course of the disease was more acute: in 474six individuals the process lasted 3 weeks and in two others, after having been diagnosed negative 475 during the last haemolymph drawing, the haemocytological observations carried out at their death 10 476 and 12 days later revealed an advanced stage of the disease. Interestingly, in all individuals the same 477 neoplastic cell type was observed during all the course of the disease. Except for individuals 478 undergoing remission processes, DN-affected mussels that reached the high intensity stage usually 479 died between about 2-4 weeks later. Moreover, all the individuals diagnosed as high intensity at their 480 arrival died within the first month.

481Mortality of individuals with no DN cells occurred in all the batches, especially between the $6th$ and 482the $9th$ weeks (Figure 10B) and some batches experienced high losses (48% for Batch 2). In most of 483these cases, moribund individuals analysed by histology showed haemocytic infiltration and 484 granulomas associated with bacterial invasion.

4854. Discussion

486We studied the haemolymph of 1791 mussels Mytilus sp. using cytological analysis. Haemocytes 487 constitute the cellular component of haemolymph and their morphological classification was a 488 prerequisite to further exploring the DN development. However, controversy over their 489 characterisation remains. As in previous studies (Rasmussen et al., 1985; Carballal et al., 1997), we 490 classified the circulating haemocytes of both *M. edulis* and *M. galloprovincialis* specimens in two 491 morphologically distinct cell types: granulocytes and agranulocytes, depending on the presence of 492 cytoplasmic granules. The distinction between two types of granulocytes suggested by Pipe (1990), 493some with small or large granules, was not univocal in our samples because granule dimension varied 494in a single cell. Carballal et al. (1997) observed bi/multi-nucleated granulocytes in a mussel parasitized 495by trematodes. The presence of basophilic granulocytes appeared to be very limited; they were 496 observed in less than 2% of individuals and represented only few cells, but were reported from nearly 497all sampling sites. Le Foll et al. (2010) reported a higher proportion of basophilic granulocytes in 498haemolymph (20-25%). We did not observe the granulocytes containing both types of granules 499(eosinophilic and basophilic) described by Carballal et al. (1997). These differences could be

500 attributed to different physiological or environmental conditions among the studies. We observed the 501 small and large agranulocytes that Moore and Lowe (1977) suggested considering as two sub-502 populations. Brown cells were observed in the haemolymph of some individuals, corresponding 503 morphologically to the brown cells described in mussel tissues by Moore and Lowe (1977) and 504 Carballal et al. (1997). We observed that their presence in the haemolymph was linked to the invasion 505of bacteria and the number of cells was particularly high in the batch from Lannion, sampled on $20th$ 506July during a mortality event. Granular haemocytes were the major constituent of the circulating cell 507 populations. The haemocytograms from mussels of all origins showed a similar pattern with no 508 significant differences except for the two batches sampled in February, both with a lower total 509haemocyte count. A seasonal evolution in total haemocyte count was previously observed in bivalves, 510linked to water temperature variations or gonadal development (Soudant et al., 2004; da Silva et al., 5112008; Flye-Sainte-Marie et al., 2009; Perrigault et al., 2011) but also to the presence of pathogens 512(Parisi et al., 2008; Ciacci et al., 2009) or pollution (Auffret et al., 2006). Granulocytosis and 513 granulocytopenia were also observed in approximately 4% and 8% of individuals, respectively, but 514 further studies are needed to understand their causes and the mechanisms leading to these conditions. 515In fact, haemocyte populations might vary greatly with environmental and physiological conditions or 516in presence of pollutants and pathogens (Pipe and Coles, 1995; Soudant at al., 2004; da Silva et al., 5172008; Donaghy and Volety, 2011). By quantifying the mitotic activity of haemolymph cells, Mayrand 518et al. (2005) showed that blue mussels exposed to a stressor were able to control the number of 519 circulating haemocytes by adjusting the mitotic activity of haemocytes. Contrary to reports by Moore 520and Lowe (1977) in *M. edulis* and Matozzo et al. (2008) in *Ruditapes decussatus*, we did not observe 521 mitotic figures in the small agranular cells but mitoses were observed in granulocytes and, more rarely, 522in the large agranular cells, questioning the function as stem cells attributed to small agranulocytes. 523The haematopoiesis in bivalves is still a debated topic, principally due to the lack of specific markers. 524 There are several theories on bivalve haemocyte renewal and maturation (see review Pila et al., 2016). 525A definitive haematopoietic site has yet to be identified.

526Our first goal was to identify the best method to diagnose and follow the evolution of DN. We showed 527that drawing haemolymph samples was non-destructive and individuals can be repeatedly sampled 528 without affecting survival, as also demonstrated in *Mya arenaria* (Brown et al., 1978; Cooper et al., 5291982). This finding is relevant for future studies on mussel haemolymph. We showed that the 530 sensitivity of haemocytological observation appears to be appropriate for DN diagnosis, comparable to 531 that of histology in early stages of the disease. The relative accuracy of these two approaches is most 532likely a function of the relative amount of tissue that is processed and examined. Haemocytological 533 observation detected one neoplastic cell among about 150,000 fixed on the slide. Flow cytometry and 534 genetics appeared to be reasonable diagnostic methods but might not be sufficiently sensitive in early 535 stages of the disease. However, genetic analysis is the only method that can diagnose neoplasia as a 536trossulus transmissible cancer. Although the fluorescence signal is likely too low for lightly infected 537 specimens, it remains possible that some neoplasms were not the *trossulus* transmissible cancer.

538Neoplastic cells were easily identifiable by haemocytology and histology because of their peculiar 539 morphology compared to normal haemocytes. They were characterised by a larger diameter, higher 540nuclear-cytoplasm ratio, prominent nucleoli and marked basophilia. Histological observation revealed 541that, in tissues, the neoplastic cell population within a specimen was heteromorphic, confirming the 542 observations of Carella et al. (2013). In contrast, the population circulating in the haemolymph 543 appeared to be relatively homogeneous. Among mussels with DN, four types of anomalous cells were 544 observed in the haemolymph (P-type, Ps-type, U-type, and S-type) but usually only one type was 545 detected in a single individual. These cell-types clearly differed in size, NCR, nucleus shape, number 546 of nucleoli, and basophilia. The P-type was the most frequent form of neoplasia. Previous studies, 547 conducted on other bivalve species, described only two neoplastic circulating cell-types, type A and 548type B (Farley and Sparks, 1970; Lowe and Moore, 1978; Green and Alderman, 1983) that correspond 549only partially to P-type and U-type of our study. We did not find a link between cell type and disease 550 stage, thus it is unlikely that the various cell types correspond to a continuum of differentiation. Cells 551 undergoing mitosis and plurinucleated cells were common in tissues, as reported in previous studies 552(Mix et al., 1979; Carella et al., 2013) but were rare among the circulating cells.

553Neoplastic cells were also easily distinguishable by flow cytometry due to their high DNA content, 554 ranging between about 8 and 18 times the normal haploid DNA content, depending on the individual. 555 Clearly, a relation between the nucleus size (measured by haemocytological observation) and ploidy 556 was established allowing a predictive estimation of the DNA content. The low coefficient of variation 557in fluorescence values among the neoplastic cells of a single mussel showed that the pattern of DNA 558 content of this population within the host was uniform with a comparable ploidy in all the circulating 559anomalous cells. Only one individual presented simultaneously two neoplastic cell populations, 560 distinct for their ploidy but not enough to be distinguishable morphologically. The high DNA content 561of neoplastic cells in mussels was mentioned in previous reports, even when different degrees of 562 ploidy were reported. Interestingly, we observed the absence of correlation between the disease stage 563and ploidy value. This conclusion contrasted with the results of Benabdelmouna et al. (2018) in 564 mussels and Le Grand et al. (2010) in cockles, who suggested an evolutive continuum of ploidies 565during disease development. However, these two studies did not use genetic analysis, which might 566 have determined if they were describing the transmissible neoplasia we studied. Our results are more 567 consistent with the observations of neoplasia in *M. trossulus* by Moore et al. (1991) and Elston et al. 568(1990), which were likely transmissible neoplasias (Metzger, 2016). These authors showed that 569neoplasia exists in two forms in this host, pentaploid and tetraploid, representing distinct pathogenetic 570 processes rather than sequential stages of a single pathogenesis. In our study concerning *M. edulis* 571 hosts, the definition of disease forms according to ploidy was not so conclusive as a high inter-572 individual variability existed. Unfortunately, all previous flow cytometry analyses carried out on 573 mussel DN, except the study of Benabdelmouna et al. (2018), concerned *M. trossulus* hosts and 574therefore the descriptions of the ploidy patterns are limited for *M. edulis* and M. galloprovincialis. In 575 our samples, the number of circulating neoplastic cells in the S/G2M phases was very low, to such an 576 extent that no corresponding peak was graphically distinguishable, whereasElston et al. (1990) and 577 Moore et al (1991) in *M. trossulus*, described a cell population in the S/G2M phases. Our observations 578 in flow cytometry confirmed the low mitotic rate detected by haemocytological observation.

579The genetic analysis shows that DN of the French mussels we studied is a *trossulus* transmissible 580 cancer, as proposed by Riquet et al. (2017). Trossulus alleles in haemolymph DNA of *M. edulis* hosts 581 were amplified regardless of the type of neoplasia. Although the four DN types might correspond to 582 different lineages of transmissible cancer, our results with 12 diagnostic genetic markers show they all 583share a *M. trossulus* origin. Further investigations will be required to determine if the different types 584 are sublineages of the same emergence or independent emergences. Of interesting note, no *trossulus* 585 individuals have been identified previously in France. Questions remain regarding how and when this 586neoplasm colonized *M. edulis* mussels in France from its M. trossulus population of origin: (i) 587 naturally by stepping stone propagation from *M. trossulus* populations, and it has been missed until 588 now or (ii) by recent anthropogenic induced migration.

589Disseminated neoplasia in mussels was widespread among the sampling sites and was found to affect 590both *M. edulis* and *M. galloprovincialis*. No geographic distribution of the various neoplasia types 591 emerged. Interestingly, older individuals and stocks not subjected to tide (farmed in suspension) were 592 most affected, suggesting that they are promoting factors. It is not clear if it is time of exposure that 593 increases the probability of infection or if adult stages are more vulnerable and if the culture technique 594 in suspension is a particular source of physiological stress for these species. Seasonality in prevalence 595 was difficult to assess because it was not possible to sample the same batch over a longer time period 596 due to the short farming cycle of mussels.

597We detected disseminated neoplasia in locations affected by higher mortality as well as in mortality598 free sites (Bernard et al., 2017, 2018; REPAMO [https://wwz.ifremer.fr/sante_mollusques\)](https://wwz.ifremer.fr/sante_mollusques), and DN 599 was not detected in the Daoulas site in May 2018 during an event of high mortality. This site regularly 600has higher than normal mortality but DN was rare (only one positive of the 300 mussels tested). 601 However, 23% of the individuals from this batch showed a supplementary population of nearly triploid 602 cells during flow cytometry analysis while no anomalies were observed in haemocytological 603 observation. A similar condition with a second slightly hyperdiploid peak was observed in M. *trossulus* by Elston et al. (1990), who suggested it to be a distinct normal cell population. In the same 604 605Daoulas batch, other 20% of individuals presented with more than 10% of circulating haemocytes in

606the S-G2/M phases. This kind of genomic abnormality, the origin of which is unknown but may be 607 promoted by contaminants, was linked to mortalities by Benabdelmouna and Ledu (2016). 608Nevertheless, the prevalence of DN was above the mean value of 3.29% in four of the six batches 609 regularly affected by high mortality, but was also far lower than the mortality rate recorded. According 610to these observations, it was not possible to assign the cause of mortality to DN alone.

611Our efforts to anticipate the diagnosis of the disease by identifying early indicators were not 612 conclusive. No correlation between neoplasia and other variables (haemocytogram values, 613inflammatory lesions such as granulocytomas and haemocytic infiltration, and early genomic 614abnormalities) was demonstrated. Presently, the disease seems to be diagnosable only when 615 characteristic neoplastic cells appear. However, with the evidence of the genetic peculiarity of 616 neoplastic cells, a molecular test that is rapid, sensitive and specific for DN is likely to be developed 617in the near future. According to the results of our monitoring, carried out by haemocytological 618 observation of mussels maintained under controlled and stable conditions, disease incidence was not 619 constant over time but followed the same trend in all the batches, reaching a maximum after 2 months 620 and then decreasing. Various interpretations are possible: i) the infection occurred in the natural 621 environment before the collection and was not transmitted under the laboratory conditions, even if at 622least one DN-affected mussel was always present in each tank; ii) the sensitivity to DN varies among 623 individuals and some of them, the survivors, are resistant; iii) physiological conditions are 624 determinants for the development of the disease. Assuming the first hypothesis is supported, it would 625 mean that we were not able to detect the first stage of infection for months and suggests the existence 626of a quiescent state. Disease progression in affected mussels appeared usually rather slow to reach the 627 complete replacement of normal haemocytes by neoplastic cells, followed by death about 1 month 628later. Disease progression is, however, poorly generalisable. In fact, in few cases, we observed a rapid 629 and total invasion of haemolymph by neoplastic cells in less than 2 weeks, probably promoted by 630 stress factors or due to a more aggressive DN form. Unfortunately, because our monitored batches 631 suffered mortality not attributable to DN, we were not able to evaluate consistently whether 632 differences exist between the various forms of neoplasia. Finally, remission of DN was observed and

633this proved to be quite frequent (12%), regardless of the neoplasia type. A higher remission rate (20%) 634 was observed in *M. trossulus* (Elston et al., 1988), confirming that it is common in the course of DN 635 infection. The rapid decrease of the number of circulating anomalous cells during remission was also 636 observed in tissues. Interestingly, the DNA of neoplastic cells was still detectable after complete 637 remission by genetic analysis. Two assumptions can be made i) *trossulus* DNA resulting from the 638 destruction of neoplastic cells is still present, free or within phagosomes, ii) *trossulus* cells are still 639 present in the haemolymph but are undistinguishable from host haemocytes at this stage.

640Presently, it is not possible to conclude that mussel mortality events in northern France are attributable 641 to DN. We observed neoplasia in batches not affected by mortality and, on the contrary, the absence of 642 neoplasia in batches suffering these events. Prevalence was higher than the mean prevalence in four of 643the six batches sampled during mortality events; however, during the monitoring in laboratory, we 644 observed a contemporaneity between an increase in mortality in DN-unaffected individuals and an 645 increase in DN incidence. This may indicate that mussels are developing a particular sensitivity to 646 pathogens and the promotion of DN development. DN is particularly interesting due to its peculiarity 647among neoplasms of being directly transmissible from individual to individual and across species 648(here from *M. trossulus* to *M. edulis*). The infection route remains to be determined.

5. Conclusions 649

650In the present study, we obtained information on the circulating cells of the blue mussel and on 651 disseminated neoplasia. We observed that DN is a widespread disease in mussels in northern France, 652 although prevalence is low. Furthermore, various types of DN coexist in the same site. Our genetic 653 studies showed that *trossulus* DNA amplification was associated with most of the DN cases in *Mytilus* 654 edulis, regardless the type. Interestingly, we observed a high diversity of forms and high mitotic rate of 655 neoplastic cells located in the vesicular connective tissue and not in haemolymph, which raised the 656 question of the proliferation site of these cells.

657 Some observations raised new questions:

658We described the normal cellular composition of the haemolymph of blue mussels and detected that 659 some variations occur. Further studies are needed to understand the origin, the consequences, and the 660 function of these alterations.

We detected *trossulus* DNA in the haemolymph of mussels with DN, consistent with a transmissible 661 662 cancer; however, the route of infection and the invasive mechanism are still unknown.

663We demonstrated that haemocytological examination is a good method to diagnose DN and monitor 664the disease; however, it is not sufficient to understand the details of DN pathogenesis, and a 665 comparison between the neoplastic cells in tissues and those circulating in the haemolymph is 666necessary, along with the development of specific cellular markers.

667We detected genomic anomalies, different from DN, that may be linked with host mortality and was 668 already described in previous studies associated with mortality events; however, the cause of these 669anomalies and the real link with mortalities is still unclear;

670We produced data suggesting that DN is not alone responsible for the mortality outbreaks in farmed 671 mussel stocks in northern France; however, the real impact of DN on an entire cycle of production 672 should be evaluated. Because of the economic relevance of mussel farming in Europe, all these topics 673 should be the subject of further studies, aiming to contribute to reduction of mortality events.

674

Acknowledgements 675

676We express our gratitude to the farmers for supplying mussel samples. This work received financial 677 support from the "Conseil Régional de Normandie". We thank the Comité Régional de la 678Conchyliculture de Bretagne Nord (CRC). A. S. and N. B. acknowledge funding by Montpellier 679Université d'Excellence (BLUECANCER project) and Agence Nationale de la Recherche (TRANSCAN project, ANR-18-CE35-0009). 680

681

References 682

683AboElkhair, M., Iwamoto, T., Clark, K.F., McKenna, P., Siah, A., Greenwood, S.J., Berthe, F.C.J., Casey, J.W., Cepica, A., 2012. Lack of detection of a putative retrovirus associated with haemic 684

neoplasia in the soft-shell clam *Mya arenaria*. J. Invertebr. Pathol., 109: 97-104. doi: 10.1016/j.jip.2011.10.008. 685 686

687AboElkhair, M., Siah, A., Clark, K.F., McKenna, P., Pariseau, J., Greenwood, S.J., Berthe, F.C.J., Cepica, A., 2009. Reverse transcriptase activity associated with haemic neoplasia in the softshell clam *Mya arenaria*. Dis. Aquat. Org., 84: 57-63. doi: 10.3354/dao02038. 688 689

690Auffret, M., Poder, M., 1986. Sarcomatous lesion in the cockle *Cerastoderma edule*. II. Electron microscopy study. Aquaculture, 58: 9-15. doi: 10.1016/0044-8486(86)90152-3. 691

692 Auffret, M., Rousseau, S., Boutet, I., Tanguy, A., Baron, J., Moraga, D., Duchemin, M.B., 2006. A

multiparametric approach for monitoring immunotoxic responses in mussels from contaminated 693

sites in Western Mediterranea. Ecotoxicology and Environmental Safety 63(3): 393-405. doi: 694

10.1016/j.ecoenv.2005.10.016. 695

696Balouet, G., Poder, M., Cahour, A., Auffret, M., 1978. Proliferative hemocytic condition in European flat oysters (*Ostrea edulis*) from Breton coasts: A 6-year survey. J. Invert. Pathol., 48(2): 208- 215. doi: 10.1016/0022-2011(86)90125-4. 697 698

699Barber, B.J., 2004. Neoplastic diseases of commercially important marine bivalves. Aquat. Living Resour. 17, 449-466. doi: 10.1051/alr:2004052. 700

701Bayne, B.L., 1976. Marine mussels, Their Ecology and Physiology. The University Press, Cambridge, MA, 506 pp. 702

703Béchemin, C., Soletchnik, P., Polsenaere, P., Le Moine, O., Pernet, F., Protat, M., Fuhrman, M., Quéré,

C., Goulitquer, S., Corporeau, C., Lapègue, S., Travers, A., Morga, B., Garrigues, M., Garcia C., 704

Haffner, P., Dubreuil, C., Faury, N., Baillon, L., Baud, J.-P., Renault, T., 2015. Episodes de 705

mortalité massive de moules bleues observés en 2014 dans le Pertuis charentais. Bull. 706

épidémiol., santé anim. aliment., 67: 6-9. 707

708Ben Cheikh, Y., Travers, M. A., Morga, B., Godfrin, Y., Rioult, D., Le Foll, F., 2016. First evidence for a *Vibrio* strain pathogenic to *Mytilus edulis* altering hemocyte immune capacities. Developmental and Comparative Immunology, 57: 107-119. doi: 10.1016/j.dci.2015.12.014. 709 710

Ben Cheikh, Y., Travers, M. A., Le Foll, F., 2017. [Infection dynamics of a](https://www.sciencedirect.com/science/article/pii/S1050464817305636) *V. splendidus* strain 711 pathogenic to *Mytilus edulis*: In vivo and in [vitro interactions with hemocytes.](https://www.sciencedirect.com/science/article/pii/S1050464817305636) Developmental and Comparative Immunology, 70: 515-523. doi: 10.1016/j.fsi.2017.09.047. 712 713

Benabdelmouna, A., Ledu, C., 2016. The mass mortality of blue mussels (*Mytilus* spp.) from the 714 Atlantic coast of France is associated with heavy genomic abnormalities as evidenced by flow cytometry. J. Invertebr. Pathol., 138: 30-38. doi: 10/1016/j.jip.2016.06.001. 715 716

717 Benabdelmouna, A., Saunier, A., Ledu, C., Travers, M.A., Morga, B., 2018. Genomic abnormalities

affecting mussels (Mytilus edulis-galloprovincialis) in France are related to ongoing neoplastic 718

processes, evidenced by dual flow cytometry and cell monolayer analyses. J. Invertebr. Pathol., 719

157: 45-52. doi: 10.1016/j.jip.2018.08.003. 720

721Bernard, I., Allain, G., 2017. Mortalités des moules en Bretagne nord : bilan des connaissances. Rapp. tech. CRC Bretagne nord. 37 pp. 722

723Bernard, I., Allain, G., Charles, M., Villalba, A., Burioli, E., Houssin, M., Deydier, Y., 2017. Investigation on blue mussel high mortality in France: a phenomenon with high spatio-temporal variability but no conclusive cause. AMMR 2017, 2nd International Symposium on the Advances in Marine Mussel Research, 7th & 8th of September 2017, Sète (France). 724 725 726

727 Bernard, I., Charles, M., Allain, G., Burioli, E., Villalba, A., Foll, F. L., Deydier, Y., Houssin, M., 2018. Bilan de l'observatoire des mortalités de moules en Bretagne Nord pour la saison 2016- 2017 et premiers éléments sur les organismes pathogènes présents. Rapp. tech. CRC Bretagne Nord. 26 pp. 728 729 730

731Bierne, N., Borsa, P., Daguin, C., Jollivet, D., Viard, F., Bonhomme, F., David, P., 2003. Introgression patterns in the mosaic hybrid zone between *Mytilus edulis* and *M. galloprovincialis*. Mol. Ecol., 12(2): 447-461. doi: 10.1046/j.1365-294X.2003.01730.x. 732 733

734 Brown, R.S., Wolke, R.E., Saila, S.B., Brown, C.W., 1978. Prevalence of neoplasia in 10 New England populations of the soft-shell clam (*Mya arenaria*). Proc. Conf. on Aquatic Pollutants and Biologic Effects with Emphasis on Neoplasia. Ann. N.Y. Acad. Sci., 298: 522-534. doi: 10.1111/j.1749-6632.1977.tb19287.x. 735 736 737

738Burioli, E.A.V., Charles, M., Bernard, I., Pitel, P.H., Houssin, M., 2017. First description of disseminated neoplasia in *Mytilus edulis* in Northern Brittany (France) and development of a rapid diagnostic tool. AMMR 2017, 2nd International Symposium on the Advances in Marine Mussel Research, 7th & 8th of September 2017, Sète (France). doi: 10.13140/RG.2.2.10832.20488. 739 740 741 742

743 Carballal, M.J., Barber B.J., Iglesias, D., Villalba, A., 2015. Neoplastic diseases of marine bivalves. J. Invertebr. Pathol., 131: 83-106. doi: 10.1016/j.jip.2015.06.004. 744

745 Carballal, M.J., López, M.C., Azevedo, C., Villalba, A., 1997. Hemolymph cell types of the mussel *Mytilus galloprovincialis*. Dis. Aquat. Org., 29: 127-135. 746

747 Carella, F., De Vico G., Landini, G., 2017. Nuclear morphometry and ploidy of normal and neoplastic

haemocytes in mussels. PLoS ONE 12(3): e0173219. doi: 10.1371/journal.pone.0173219. 748

749 Carella, F., Figueras, A., Novoa, B., De Vico, G., 2013. Cytomorphology and PCNA expression

pattern in bivalves *Mytilus galloprovincialis* and *Cerastoderma edule* with haemic neoplasia. 750

Dis. Aquat. Org., 105: 81-87. doi: 10.3354/dao02612. 751

752 Ciacci, C., Citterio, B., Betti, M., Canonico, B., Roch, P., Canesi, L., 2009. Functional differential

immune responses of *Mytilus galloprovincialis* to bacterial challenge. Comp. Biochem. Physiol. 753

Biochem. Mol. Biol, 153(4): 365-71. doi: 10.1016/j.cbpb.2009.04.007. 754

755Collins, C.M., Mulcahy, M.F., 2003. Cell-free transmission of a haemic neoplasm in the cockle *Cerastoderma edule*. Dis. Aquat. Organ., 54(1): 61-67. doi: 10.3354/dao054061. 756

757 Cooper, K.R., Brown, R.S., Chang, P.W., 1982. Accuracy of blood cytological screening techniques for the diagnosis of a possible hematopoietic neoplasm in the bivalve mollusc, *Mya arenaria*. J. Invertebr. Pathol., 39: 281-289. 758 759

760Cuenca, J., Aleza, P., Navarro, L., Ollitrault, P., 2013. Assignment of SNP allelic configuration in polyploids using competitive allele-specific PCR: application to citrus triploid progeny. Ann. Bot., 111(4), 731–742. doi: 10.1093/aob/mct032. 761 762 763

764da Silva, P.C., Comesaña, P., Fuentes, J., Villalba, A., 2008. Variability of haemocyte and haemolymph parameters in European flat oyster *Ostrea edulis* families obtained from brood stocks of different geographical origins and relation with infection by the protozoan *Bonamia ostreae*. Fish Shellfish Immunol., 24: 551-563. doi: 10.1016/j.fsi.2007.11.003. 765 766 767

768da Silva, P.M., Soudant, P., Carballal, M.J., Lambert, C., Villalba, A., 2005. Flow cytometric DNA content analysis of neoplastic cells in haemolymph of the cockle *Cerastoderma edule*. Dis. Aquat. Organ. 67, 133-139. doi: 0.3354/dao067133. 769 770

771Delaporte, M., Synard, S., Pariseau, J., McKenna, P., Tremblay, R., Davidson, J., Berthe, F.C., 2008. Assessment of haemic neoplasia in different soft-shell clam *Mya arenaria* populations from eastern Canada by flow cytometry. J. Invertebr. Pathol. 98, 190-197. doi : 10.1016/j.jip.2007.12.005. 772 773 774

775Díaz, S., Villalba, A., Insua, A., Soudant, P., Fernandez-Tajes, J., Mendez, J., Carballal, M.J., 2013. Disseminated neoplasia causes changes in ploidy and apoptosis frequency in cockles *Cerastoderma edule*. J. Invertebr. Pathol. 113, 214-219. doi: 10.1016/j.jip.2013.03.010. 776 777

778Donaghy, L., Volety, A.K., 2011. Functional and metabolic characterization of hemocytes of the green mussel, *Perna viridis*: *in vitro* impacts of temperature. Fish Shellfish Immunol., 31: 808-814. doi: 10.1016/j.fsi.2011.07.018. 779 780

781Elston, R.A., Drum, A.S., Allen, S.K.Jr., 1990. Progressive development of circulating polyploid cells in *Mytilus* with hemic neoplasia. Dis. Aquat. Org., 8: 51-59. 782

783Elston, R.A., Kent M.L., Drum, A.S., 1988. Transmission of hemic neoplasia in the bay mussel,

Mytilus edulis, using whole cells and cells homogenate. Dev. Comp. Immunol., 12: 719-727. doi: 10.1016/0145-305X(88)90047-X. 784 785

786Elston, R.A., Moore, J.D., Brooks, K., 1992. Disseminated neoplasia of bivalve molluscs. Rev. Aquat. Sci. 6, 405-466. 787

788FAO, Food and Agriculture Organisation of the United Nations (2016). Fisheries and Aquaculture Department, Dataset Global Aquaculture production. On-Line Query. 789

790Farley, C.A., 1969a. Probable neoplastic disease of the hematopoietic system in oysters, *Crassostrea virginica* and *Crassostrea gigas*. Natl. Cancer I. Monogr. 31: 541-555. 791

792Farley, C.A., 1969b. Sarcomatoid proliferative disease in a wild population of blue mussels (Mytilus *edulis*). Natl. Cancer I. Monogr., 43: 509-516. 793

794 Farley, C.A., Sparks, A.K., 1970. Proliferative diseases of hemocytes, endothelial cells, and connective tissue cells in mollusks. Bibl. Haematol. 36: 610-617. 795

796Flye-Sainte-Marie, J., Soudant, P., Lambert, C., Le Goïc, N., Goncalvez, M., Travers, M.A., Paillard, C., Jean, F., 2009. Variability of the hemocyte parameters of *Ruditapes philippinarum* in the field during an annual cycle. J. Exp. Marine Biol. Ecol., 377: 1-11. doi: 10.1013/j.embe.2009.06.003. 797 798 799

800Fraïsse, C., Belkhir, K., Welch, J.J., Bierne, N., 2016. Local interspecies introgression is the main cause of extreme levels of intraspecific differentiation in mussels. Mol. Ecol., 25(1), 269–286. doi: 10.1111/mec.13299. 801 802

803Frierman, E.M., Andrews, J.D., 1976. Occurrence of haematopoietic system in oysters, *Crassostrea virginica* and *Crassostrea gigas*. J. Natl. Cancer I., 56: 319-324. 10.1093/jnci/56.2.319. 804

Galimany, E., Sunila, I., 2008. Several cases of disseminated neoplasia in mussels *Mytilus edulis* (L.) 805 in western long island sound. J. Shellfish Res. 27(5), 1201-1207. doi: 10.2983/0730-8000- 27.5.1201. 806 807

Gombač, M., Sitar, R., Pogačnik, M., Arzul, I., Jenčič, V., 2013. Haemocytic neoplasia in 808 Mediterranean mussels (*Mytilus galloprovincialis*) in the Slovene Adriatic Sea. Mar. Freshwater Behav. Physiol., 46(2): 135-143. doi: 10.1080/10236244.2013.782736. 809 810

811González-Tizón, A.M., Martínez-Lage, A., Rego, I., Ausió, J., Méndez J., 2000. DNA content, karyotypes, and chromosomal location of 18S-5.8S-28S ribosomal loci in some species of bivalve molluscs from the Pacific Canadian coast. Genome, 43(6): 1065-1072. doi: 10.1139/g00-089. 812 813 814

815Green, M., Alderman, D.J., 1983. Neoplasia in Mytilus edulis L. from United Kingdom waters. Aquaculture, 30(1-4):1-10. doi: 10.1016/0044-8486(83)90146-1. 816

817 House, M.L., 1997. Transmission of disseminated neoplasia in the soft-shell clam, *Mya arenaria*. PhD thesis, Oregon State University, Corvallis. 818

819House, M.L., Kim, C.H., Reno, P.W., 1998. Soft shell clams Mya arenaria with disseminated neoplasia demonstrate reverse transcriptase activity. Dis. Aquat. Org., 34(3): 187-192. doi: 10.3354/dao034187. 820 821

822Inoue, K., Waite, J.H., Matsuoka, M., Odo, S., Harayama, S., 1995. Interspecific variations in adhesive protein sequences of *Mytilus edulis*, *M. galloprovincialis*, and *M. trossulus*. Biol. Bull. 189(3):370-375. doi: 10.2307/1542155. 823 824

825 Kent, M.L., Wilkinson, M.T., Drum, A.S., Elston, R.A., 1991. Failure of transmission of hemic neoplasia of bay mussels, *Mytilus trossulus*, to other bivalves. J. Invertebr. Pathol., 57: 435-436. 826

827 Krishan, A., 1975. Rapid cytofluorometric analysis of mammalian cell cycle by propidium iodide staining. J. Cell. Biol., 66:188–193.doi: 10.1083/jcb.66.1.188. 828

829Landsberg, J.H., 1996. Neoplasia and biotoxins in bivalves: is there a connexion? J. Shellfish Res. 15, 203-230. 830

831Le Foll, F., Rioult, D., Boussa, S., Pasquier, J., Dagher, Z., Leboulenger, F., 2010. Characterisation of Mytilus edulis hemocyte subpopulations by single cell time-lapse motility imaging. Fish Shellfish Immunol., 28(2):372-86. doi: 10.1016/j.fsi.2009.11.011. 832 833

834Le Grand, F., Kraffe, E., de Montaudouin, X., Villalba, A., Marty, Y., Soudant, P., 2010. Prevalence,

intensity, and aneuploidy patterns of disseminated neoplasia in cockles (*Cerastoderma edule*) 835

from Arcachon Bay: seasonal variation and position in sediment. J. Invertebr. Pathol. 104, 110- 836

118. doi: 10.1016/j.jip.2010.01.011. 837

838Lowe, D.M., Moore, M.N., 1978. Cytology and quantitative cytochemistry of a proliferative atypical hemocytic condition in *Mytilus edulis* (Bivalvia, Mollusca). J. Natl. Cancer I., 60(6):1455-1459. doi: 10.1093/jnci/60.6.1455. 839 840

841Lowe, D.M., Moore, M.N., 1979. The cytology and occurrence of granulocytomas in mussels. Mar. Pollut. Bull., 10: 137-141. doi: 10.1016/0025-326X(79)90081-X. 842

843Matozzo, V., Marin, M.G., Cima, F., Ballarin, L., 2008. First evidence of cell division in circulating haemocytes from the Manila clam *Tapes philippinarum*. Cell Biol. Intern., 32: 865-868. doi: 10.1016/j.cellbi.2008.03.008. 844 845

846Mayrand, E., St-Jean, S.D., Courtenay, S.C., 2005. Haemocyte responses of blue mussels (Mytilus edulis L.) transferred from a contaminated site to a reference site: can the immune system recuperate? Aquaculture Res., 36: 962-971. doi: 10.1111/j.1365-2109.2005.01302.x. 847 848

849McGladdery, S.E., Reinisch, C.L., MacCallum, G.S., Stephens, R.E., Walker, C.L., Davidson, J.T., 2001. Haemic neoplasia in soft-shell clams *Mya arenaria*: recent outbreaks in Atlantic Canada and discovery of a p53 gene homologue associated with the condition. Bull. Aquat. Assoc. Can., 101, 19-26. 850 851 852

853McLaughlin, S.M., Farley, C.A., Hetrick, F.M., 1992. Transmission studies of sarcoma in the soft-shell clam, *Mya arenaria*. In Vivo, 6: 367-370. 854

855Metzger, M.J., Reinisch, C., Sherry, J., Goff, S.P., 2015. Horizontal transmission of clonal cancer cells causes leukemia in soft-shell clams. Cell, 161: 255-263. doi: 10.1016/j.cell.2015.02.042. 856

857Metzger, M.J., Villalba, A., Carballal, M.J., Iglesias, D., Sherry, J., Reinisch, C., Muttray, A.F.,

Baldwin, S.A., Goff, S.P., 2016. Widespread transmission of independent cancer lineages within 858

multiple bivalve species. Nature, 534: 705–709. doi: 10.1038/nature18599. 859

860Mix, M.C., Hawkes, J.W., Sparks, A.K., 1979. Observations of the ultrastructure of large cells associated with putative neoplastic disorders of mussels, *Mytilus edulis*, from Yaquina Bay, 861

Oregon. J. Invertebr. Pathol., 34: 41-56. doi: 10.1016/0022-2011(79)90052-1. 862

863Moore, J.D., Elston, R.A., Drum, A.S., Wilkinson, M.T., 1991. Alternate pathogenesis of systemic neoplasia in the bivalve mollusc *Mytilus*. J. Invertebr. Pathol., 58: 231-243. doi: 10.1016/0022- 2011(91)90067-Z. 864 865

866Moore, M.N., Lowe, D.M., 1977. The cytology and cytochemistry of the hemocytes of Mytilus edulis and their response to experimentally injected carbon particles. J. Invertebr. Pathol., 29: 18-30. doi: 10.1016/0022-2011(77)90167-7. 867 868

869Oden, E., Burioli, E.A.V., Trancart, S, Pitel, P.H., Houssin, M., 2016. Multilocus Sequence Analysis of

Vibrio splendidus related-strains isolated from blue mussel Mytilus sp. during mortality events. 870

Aquaculture, 464: 420-427. doi: 10.1016/j.aquaculture.2016.07.024. 871

872Oprandy, J.J., Chang, P.W., 1983. 5-bromodeoxyuridine induction of hematopoietic neoplasia and retrovirus activation in the soft-shell clam, *Mya arenaria*. J. Invertebr. Pathol., 42: 196-206. doi: 10.1016/0022-2011(83)90062-9. 873 874

875 Oprandy, J.J., Chang, P.W., Pronovost, A.D., Cooper, K.R., Brown, R.S., Yates, V.J., 1981. Isolation of a viral agent causing haematopoietic neoplasia in the soft-shell clam *Mya arenaria*. J. Inverteb. Pathol., 38(1): 45-51. doi: 10.1016/0022-2011(81)90033-1. 876 877

878Parisi, M.G., Li, H., Jouvet, L.B.P., Dyrynda, E.A., Parrinello, N., Cammarata, M., Roch, P., 2008. Differential involvement of mussel hemocyte sub-populations in the clearance of bacteria. Fish Shellfish Immunol., 25: 834-840. doi: 10.1016/j.fsi.2008.09.005. 879 880

881Perrigault, M., Dahl, S.F., Pales Espinosa, E., Gambino, L., Allam, B., 2011. Effects of temperature on hard clam (*Mercenaria mercenaria*) immunity and QPX (Quahog Parasite Unknown) disease development: II. Defence parameters. J. Invertebr. Pathol., 106: 322-332. doi: 10.1016/j.jip.2010.11.004. 882 883 884

885Pila, E.A., Sullivan, J.T., Wu, X.Z., Fang, J., Rudko, S.P., Gordy, M.A., Hanington, P.C., 2016. Haematopoiesis in molluscs: A review of haemocyte development and function in gastropods, cephalopods and bivalves. Dev. Comp. Immunol., 58: 119-128. doi: 10.1016/j.dci.2015.11.010. 886 887

888Pipe, R.K., 1990. Differential binding of lectins to haemocytes of the mussel Mytilus edulis. Cell Tissue Res., 261: 261:268. doi: 10.1007/BF00318667. 889

890Pipe, R.K., Coles, J.A., 1995. Environmental contaminants influencing immune function in marine bivalve molluscs. Fish Shellfish Immunol., 5: 581-595. doi: 10.1016/S1050-4648(95)80043-3. 891

892Polsenaere, P., Soletchnik, P., Le Moine, O., Gohin, F., Robert, S., Pépin, J. F., Stanisière, J. Y., Dumas, F., Béchemin, C., Goulletquer, P., 2017. Potential environmental drivers of a regional blue mussel mass mortality event (winter of 2014, Breton Sound, France). J. Sea Res., 123: 39- 50. doi: 10.1016/j.seares.2017.03.005. 893 894 895

896Rasmussen, L.P.D., Hage, E., Karlog, O., 1985. An Electron Microscope Study of the Circulating Leucocytes of the Marine Mussel, *Mytilus edulis*. J. Invertebr. Pathol., 45: 158-167. doi: 10.1016/0022-2011(85)90005-9. 897 898

899 Reinisch, C.L., Charles, A.M., Stone, A.M., 1984. Epizootic neoplasia in soft shell clams collected from New Bedford Harbor. Hazardous waste, 1(1): 73-81. doi: 10.1089/hzw.1984.1.73. 900

901 Riquet, F., Simon, A., Bierne, N., 2017. Weird genotypes? Don't discard them, transmissible cancer could be an explanation. Evol. Appl., 10: 140-145. doi: 10.1111/eva.12439. 902

903Romalde, J.L., Luz Vilariño, M., Beaz, R., Rodríguez, J.M., Díaz, S., Villalba, A., Carballal, M.J., 2007. Evidence of retroviral etiology for dissaminated neoplasia in cockles (*Cerastoderma edule*), J. Invertebr. Pathol., 94: 95-101. doi: 10.1016/j.jip.2006.09.007. 904 905

906Semagn, K., Babu, R., Hearne, S., Olsen, M., 2014. Single nucleotide polymorphism genotyping using Kompetitive Allele Specific PCR (KASP): overview of the technology and its application in crop improvement. Mol. Breeding, 33:1–14. doi: 10.1007/s11032-013-9917-x. 907 908

909Simon, A., Bierne, N., Welch, J. J., 2018. Coadapted genomes and selection on hybrids: Fisher's geometric model explains a variety of empirical patterns. Evolution Letters, 2(5): 472-498. doi: 10.1002/evl3.66. 910 911

912 Soudant, P., Paillard C., Choquet, G., Lambert, C., Reid, H.I., Marhic, A., Donaghy, L., Birkbeck, T.H., 2004. Impact of season and rearing site on the physiological and immunological parameters of the Manila clam *Venerupis* (=Tapes, =*Ruditapes*) *philippinarum*. Aquaculture; 229: 401-418. doi: 10.1016/S0044-8486(03)00352-1. 913 914 915

916Suquet, M., de Kermoysan, G., Gonzalez Araya, R., Queau, I., Lebrun, L., Le Souchu, P., Mingant, C. (2009). Anesthesia in Pacific oyster *Crassostrea gigas*. Aquat. Living Resour., 22, 29–34. doi: 10.1051/alr/2009006. 917 918

919Twomey, E., Mulcahy, M.F., 1988. Transmission of a sarcoma in the cockle *Cerastoderma edule* (Bivalvia: Mollusca) using cell transplants. Dev. Comp. Immunol., 12:195-200. 920

921Vassilenko, E., Baldwin, S.A., 2014. Using flow cytometry to detect haemic neoplasia in mussels (*Mytilus trossulus*) from the Pacific Coast of Southern British Columbia, Canada. J. Invertebr. Pathol., 117: 68-72. doi: 10.1016/j.jip.2014.02.002. 922 923

924 Villalba, A., Carballal, M.J., López, C., 2001. Disseminated neoplasia and large foci indicating heavy haemocytic infiltration in cockles *Cerastoderma edule* from Galicia (NW Spain). Dis. Aquat. Org. 46: 213-216. doi: 10.3354/dao046213. 925 926

927Weinberg, J.R., Leavitt, D.F., Lancaster, B.A., Capuzzo, J.M., 1997. Experimental field studies with *Mya arenaria* (Bivalvia) on the induction and effect of hematopoietic neoplasia. J. Invertebr.

Pathol., 61: 253-259. doi: 10.1006/jipa.1996.4641.