



HAL
open science

Free (tri)dendriform family algebras

Dominique Manchon, Gao Xing, Yuanyuan Zhang

► **To cite this version:**

Dominique Manchon, Gao Xing, Yuanyuan Zhang. Free (tri)dendriform family algebras. Journal of Algebra, 2020, <10.1016/978-3-030-37030-5>. <hal-02404042>

HAL Id: hal-02404042

<https://hal.science/hal-02404042v1>

Submitted on 21 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

FREE (TRI)DENDRIFORM FAMILY ALGEBRAS

YUANYUAN ZHANG, XING GAO, AND DOMINIQUE MANCHON*

ABSTRACT. In this paper, we first prove that a Rota-Baxter family algebra indexed by a semigroup amounts to an ordinary Rota-Baxter algebra structure on the tensor product with the semigroup algebra. We show that the same phenomenon arises for dendriform and tridendriform family algebras. Then we construct free dendriform family algebras in terms of typed decorated planar binary trees. Finally, we generalize typed decorated rooted trees to typed valently decorated Schröder trees and use them to construct free tridendriform family algebras.

CONTENTS

1. Introduction	1
2. Rota-Baxter family algebras and (tri)dendriform family algebras	3
2.1. Rota-Baxter family algebras	3
2.2. Dendriform family algebras and tridendriform family algebras	4
3. Free dendriform family algebras	8
3.1. Free dendriform algebras	8
3.2. Free dendriform family algebras	9
4. Free tridendriform family algebras	17
4.1. Free tridendriform algebras	17
4.2. Free tridendriform family algebras	17
References	26
5. Appendix	27

1. INTRODUCTION

The concept of Rota-Baxter family algebra is a generalization of Rota-Baxter algebras [18], which was proposed by Guo. It arises naturally in renormalization of quantum field theory ([7, Proposition 9.1] and [19, Theorem 3.7.2]). The free objects were constructed in [28], in which the authors described free commutative Rota-Baxter family algebras, and also described free non-commutative Rota-Baxter family algebras via the method of Gröbner-Shirshov bases.

Dendriform algebras were introduced by Loday [20] in 1995 with motivation from algebraic K-theory. They have been studied quite extensively with connections to several areas in mathematics and physics, including operads [24], homology [9], arithmetics [23], Hopf algebras [21, 25, 26]

⁰* Corresponding author.

Date: November 14, 2019.

2010 *Mathematics Subject Classification.* 16W99, 16S10, 13P10, 08B20,

Key words and phrases. Rota-Baxter algebra, Rota-Baxter family algebra, (tri)dendriform algebra, (tri)dendriform family algebra, typed decorated planar binary trees, typed valently decorated rooted trees.

and quantum field theory [8, 10], in connection with the theory of renormalization of Connes and Kreimer [4, 5, 6].

Some years later after [20], Loday and Ronco introduced the concept of tridendriform algebra in the study of polytopes and Koszul duality [25]. The construction of free (tri)dendriform algebras can be referred to [20, 21, 25]. The free objects play a crucial role in the study of any algebraic structures, such as the construction of free differential algebras in terms of differential monomials, the construction of free Rota-Baxter algebras [15] which is more involved, the free differential Rota-Baxter algebras [16] composing the construction of free differential algebras followed by that of the free Rota-Baxter algebras, and the free integro-differential algebras [12, 14] for analyzing the underlying algebraic structures of boundary problems for linear ordinary differential equations. In [28], the authors proposed the concepts of (tri)dendriform family algebras and they obtained that Rota-Baxter family algebras induce (tri)dendriform family algebras. Besides, they construct free commutative (tri)dendriform family algebras.

It is natural to ask how to use reduced planar rooted trees (also known as Schröder trees) to construct free (tri)dendriform family algebras. Here we exhibit a way to construct free (tri)dendriform family algebras via typed decorated Schröder trees, whose vertices are decorated by elements of a set X and edges are decorated by elements of a semigroup Ω . Typed decorated trees are used by Bruned, Hairer and Zambotti in [2] to give a systematic description of a canonical renormalisation procedure of stochastic PDEs. They also appear [27] in a context of low dimension topology and also appear in [3] for the description of combinatorial species. The concept of typed decorated planar binary trees enables us to construct free dendriform family algebras, and the concept of typed valently decorated Schröder trees will be used to construct free tridendriform family algebras.

The layout of the paper is as follows. In Section 1, we prove that any Rota-Baxter family algebra R on a base ring \mathbf{k} indexed by a semigroup Ω induces an ordinary Rota-Baxter algebra structure on $R \otimes \mathbf{k}\Omega$ (Theorem 2.4) and vice-versa. The semigroup Ω at hand is not necessarily commutative. Then we prove that Rota-Baxter family algebras can induce (tri)dendriform family algebras (Proposition 2.10). Finally, we give the relationship between (tri)dendriform family algebras and (tri)dendriform algebras (Theorem 2.11). Section 3 is devoted to typed decorated planar binary trees. We derive a useful recursive expression for typed decorated planar binary trees endowed with the operations $(\langle_\omega, \rangle_\omega)_{\omega \in \Omega}$ introduced in the third section (Definition 3.7). We first construct dendriform family algebras (Proposition 3.10) in terms of typed decorated planar binary trees, then we prove the freeness of this dendriform family algebra (Theorem 3.12). In Section 4, closely related to Section 3, we first introduce the concept of typed valently decorated Schröder trees. Secondly, we construct tridendriform family algebras (Proposition 4.8) in terms of typed valently decorated Schröder trees, and finally prove the freeness of this tridendriform family algebra (Theorem 4.10).

The related notions of *matching Rota-Baxter algebra*, *matching dendriform algebra* and *matching pre-Lie algebra* are addressed in the recent paper [13]. The main difference is that the parameter at hand runs into a set Ω without any semigroup structure: see Remark 2. 2 (b) therein.

Notation: In this paper, we fix a ring \mathbf{k} and assume that an algebra is a \mathbf{k} -algebra. Denote by Ω a semigroup, unless otherwise specified.

2. ROTA-BAXTER FAMILY ALGEBRAS AND (TRI)DENDRIFORM FAMILY ALGEBRAS

In this section, we first recall the concepts of Rota-Baxter family algebras and (tri)dendriform family algebras. We then give a method to induce Rota-Baxter (resp. (tri)dendriform) algebras from Rota-Baxter (resp. (tri)dendriform) family algebras, and vice-versa.

2.1. Rota-Baxter family algebras. Rota-Baxter algebras (first called Baxter algebras) are originated in the work of the American mathematician Glen E. Baxter [1] in the realm of probability theory.

Definition 2.1. Let λ be a given element in \mathbf{k} . A **Rota-Baxter algebra of weight λ** is a pair (R, P) consisting of an algebra R with a linear operator $P : R \rightarrow R$ that satisfies the Rota-Baxter equation

$$P(a)P(b) = P(P(a)b + aP(b) + \lambda ab), \text{ for } a, b \in R.$$

Then P is called a **Rota-Baxter operator of weight λ** . If further R is commutative, then (R, P) is called a commutative Rota-Baxter algebra of weight λ .

The following is the concept of Rota-Baxter family algebra proposed by Guo, which arises naturally in renormalization of quantum field theory [7, Proposition 9.1], see also [19].

Definition 2.2. ([7, 17]) Let Ω be a semigroup and $\lambda \in \mathbf{k}$ be given. A **Rota-Baxter family of weight λ** on an algebra R is a collection of linear operators $(P_\omega)_{\omega \in \Omega}$ on R such that

$$P_\alpha(a)P_\beta(b) = P_{\alpha\beta}(P_\alpha(a)b + aP_\beta(b) + \lambda ab), \text{ for } a, b \in R \text{ and } \alpha, \beta \in \Omega.$$

Then the pair $(R, (P_\omega)_{\omega \in \Omega})$ is called a **Rota-Baxter family algebra of weight λ** . If further R is commutative, then $(R, (P_\omega)_{\omega \in \Omega})$ is called a commutative Rota-Baxter family algebra of weight λ .

Definition 2.3. Let $(R, (P_\omega)_{\omega \in \Omega})$ and $(R', (P'_\omega)_{\omega \in \Omega})$ be two Rota-Baxter family algebras of weight λ . A map $f : R \rightarrow R'$ is called a **Rota-Baxter family algebra morphism** if f is an algebra homomorphism and $f \circ P_\omega = P'_\omega \circ f$ for each $\omega \in \Omega$.

The following theorem precises the link between Rota-Baxter family algebras and ordinary Rota-Baxter algebras.

Theorem 2.4. $(R \otimes \mathbf{k}\Omega, P)$ is a Rota-Baxter algebra of weight λ , where

$$\begin{aligned} P : R \otimes \mathbf{k}\Omega &\rightarrow R \otimes \mathbf{k}\Omega \\ x \otimes \omega &\mapsto P_\omega(x) \otimes \omega \end{aligned}$$

if and only if $(R, (P_\omega)_{\omega \in \Omega})$ is a Rota-Baxter family algebra of weight λ .

Proof. For $x, y \in R$ and $\alpha, \beta \in \Omega$, we have

$$\begin{aligned} P(x \otimes \alpha)P(y \otimes \beta) &= (P_\alpha(x) \otimes \alpha)(P_\beta(y) \otimes \beta) = (P_\alpha(x)P_\beta(y)) \otimes \alpha\beta \\ &= P_{\alpha\beta}(P_\alpha(x)y + xP_\beta(y) + \lambda xy) \otimes \alpha\beta \\ &= P((P_\alpha(x)y + xP_\beta(y) + \lambda xy) \otimes \alpha\beta) \\ &= P(P_\alpha(x)y \otimes \alpha\beta + xP_\beta(y) \otimes \alpha\beta + \lambda xy \otimes \alpha\beta) \\ &= P((P_\alpha(x) \otimes \alpha)(y \otimes \beta) + (x \otimes \alpha)(P_\beta(y) \otimes \beta) + \lambda(x \otimes \alpha)(y \otimes \beta)) \end{aligned}$$

$$= P(P(x \otimes \alpha)(y \otimes \beta) + (x \otimes \alpha)P(y \otimes \beta) + \lambda(x \otimes \alpha)(y \otimes \beta)),$$

as required. Conversely, we obtain

$$\begin{aligned} P_\alpha(x)P_\beta(y) \otimes \alpha\beta &= P(x \otimes \alpha)P(y \otimes \beta) \\ &= P(P(x \otimes \alpha)(y \otimes \beta) + (x \otimes \alpha)P(y \otimes \beta) + \lambda(x \otimes \alpha)(y \otimes \beta)) \\ &= P\left((P_\alpha(x) \otimes \alpha)(y \otimes \beta) + (x \otimes \alpha)(P_\beta(y) \otimes \beta) + \lambda(x \otimes \alpha)(y \otimes \beta)\right) \\ &= P(P_\alpha(x)y \otimes \alpha\beta + xP_\beta(y) \otimes \alpha\beta + \lambda xy \otimes \alpha\beta) \\ &= P\left((P_\alpha(x)y + xP_\beta(y) + \lambda xy) \otimes \alpha\beta\right) \\ &= P_{\alpha\beta}(P_\alpha(x)y + xP_\beta(y) + \lambda xy) \otimes \alpha\beta. \end{aligned}$$

So we get

$$P_\alpha(x)P_\beta(y) = P_{\alpha\beta}(P_\alpha(x)y + xP_\beta(y) + \lambda xy).$$

In other words, $P \otimes \mathbf{k}\Omega$ is an Ω -graded Rota-Baxter algebra, with ω -homogeneous component equal to $P \otimes \mathbf{k}\omega$ for any $\omega \in \Omega$. Identifying each homogeneous component on both sides of the Rota-Baxter identity leads to the Rota-Baxter family identity. This completes the proof. \square

2.2. Dendriform family algebras and tridendriform family algebras. The concept of dendriform family algebras was proposed in [28], as a generalization of dendriform algebras invented by Loday [20] in the study of algebraic K -theory.

Definition 2.5. [20] A **dendriform algebra** is a \mathbf{k} -module D with two binary operations \langle, \rangle such that for $x, y, z \in D$,

$$\begin{aligned} (x \langle y) \langle z &= x \langle (y \langle z + y \rangle z), \\ (x \rangle y) \langle z &= x \rangle (y \langle z), \\ (x \langle y + x \rangle y) \rangle z &= x \rangle (y \rangle z). \end{aligned}$$

Definition 2.6. [28] Let Ω be a semigroup. A **dendriform family algebra** is a \mathbf{k} -module D with a family of binary operations $(\langle_\omega, \rangle_\omega)_{\omega \in \Omega}$ such that for $x, y, z \in D$ and $\alpha, \beta \in \Omega$,

$$\begin{aligned} (1) \quad & (x \langle_\alpha y) \langle_\beta z = x \langle_{\alpha\beta} (y \langle_\beta z + y \rangle_\alpha z), \\ (2) \quad & (x \rangle_\alpha y) \langle_\beta z = x \rangle_\alpha (y \langle_\beta z), \\ (3) \quad & (x \langle_\beta y + x \rangle_\alpha y) \rangle_{\alpha\beta} z = x \rangle_\alpha (y \rangle_\beta z). \end{aligned}$$

The concept of tridendriform family algebras was also introduced in [28], which is a generalization of tridendriform algebras invented by Loday and Ronco [25] in the study of polytopes and Koszul duality.

Definition 2.7. [25] A **tridendriform algebra** is a \mathbf{k} -module T equipped with three binary operations \langle, \rangle and \cdot such that for $x, y, z \in T$,

$$\begin{aligned} (x \langle y) \langle z &= x \langle (y \langle z + y \rangle z + y \cdot z), \\ (x \rangle y) \langle z &= x \rangle (y \langle z), \\ (x \langle y + x \rangle y + x \cdot y) \rangle z &= x \rangle (y \rangle z), \\ (x \rangle y) \cdot z &= x \rangle (y \cdot z), \end{aligned}$$

$$\begin{aligned}(x < y) \cdot z &= x \cdot (y > z), \\ (x \cdot y) < z &= x \cdot (y < z), \\ (x \cdot y) \cdot z &= x \cdot (y \cdot z).\end{aligned}$$

Definition 2.8. [28] Let Ω be a semigroup. A **tridendriform family algebra** is a \mathbf{k} -module T equipped with a family of binary operations $(\langle_\omega, \rangle_\omega)_{\omega \in \Omega}$ and a binary operation \cdot such that for $x, y, z \in T$ and $\alpha, \beta \in \Omega$,

$$\begin{aligned}(4) \quad & (x \langle_\alpha y) \langle_\beta z = x \langle_{\alpha\beta} (y \langle_\beta z + y \rangle_\alpha z + y \cdot z), \\ (5) \quad & (x \rangle_\alpha y) \langle_\beta z = x \rangle_\alpha (y \langle_\beta z), \\ (6) \quad & (x \langle_\beta y + x \rangle_\alpha y + x \cdot y) \rangle_{\alpha\beta} z = x \rangle_\alpha (y \rangle_\beta z), \\ (7) \quad & (x \rangle_\alpha y) \cdot z = x \rangle_\alpha (y \cdot z), \\ (8) \quad & (x \langle_\alpha y) \cdot z = x \cdot (y \rangle_\alpha z), \\ (9) \quad & (x \cdot y) \langle_\alpha z = x \cdot (y \langle_\alpha z), \\ (10) \quad & (x \cdot y) \cdot z = x \cdot (y \cdot z).\end{aligned}$$

Remark 2.9. When the semigroup Ω is taken to be the trivial monoid with one single element, a dendriform (resp. tridendriform) family algebra is precisely a dendriform (resp. tridendriform) algebra.

Let $\mathfrak{RB}\mathfrak{F}_\lambda$ be the category of Rota-Baxter family algebras of weight λ , and let $\mathfrak{DD}\mathfrak{F}$ (resp. $\mathfrak{DT}\mathfrak{F}$) be the category of dendriform (resp. tridendriform) family algebras. A functor $\varepsilon : \mathfrak{RB}\mathfrak{F}_\lambda \rightarrow \mathfrak{DT}\mathfrak{F}$ has been introduced in [28, Theorem 4.4]. Further we have

Proposition 2.10. *Let Ω be a semigroup.*

(a) *A Rota-Baxter family algebra $(R, (P_\omega)_{\omega \in \Omega})$ of weight λ induces a dendriform family algebra $(R, (\langle_\omega, \rangle_\omega)_{\omega \in \Omega})$, where*

$$x \langle_\omega y := xP_\omega(y) + \lambda xy \text{ and } x \rangle_\omega y := P_\omega(x)y, \text{ for } x, y \in R.$$

(b) *A tridendriform family algebra $(T, (\langle_\omega, \rangle_\omega)_{\omega \in \Omega}, \cdot)$ induces two dendriform family algebras $(T, (\langle'_\omega, \rangle'_\omega)_{\omega \in \Omega})$ and $(T, (\langle''_\omega, \rangle''_\omega)_{\omega \in \Omega})$, where*

$$\begin{aligned}x \langle'_\omega y &:= x \langle_\omega y + x \cdot y \text{ and } x \rangle'_\omega y := x \rangle_\omega y, \\ x \langle''_\omega y &:= x \langle_\omega y \text{ and } x \rangle''_\omega y := x \rangle_\omega y + x \cdot y, \text{ for } x, y \in T.\end{aligned}$$

Proof. (a) For any $x, y, z \in R$ and $\alpha, \beta \in \Omega$,

$$\begin{aligned}(x \langle_\alpha y) \langle_\beta z &= (xP_\alpha(y) + \lambda xy) \langle_\beta z \\ &= (xP_\alpha(y) + \lambda xy)P_\beta(z) + \lambda(xP_\alpha(y) + \lambda xy)z \\ &= xP_\alpha(y)P_\beta(z) + \lambda xyP_\beta(z) + \lambda xP_\alpha(y)z + \lambda^2xyz \\ &= xP_{\alpha\beta}(y)P_\beta(z) + P_\alpha(y)z + \lambda yz + \lambda x(yP_\beta(z) + P_\alpha(y)z + \lambda yz) \\ &= x \langle_{\alpha\beta} (yP_\beta(z) + P_\alpha(y)z + \lambda yz) \\ &= x \langle_{\alpha\beta} (y \langle_\beta z + y \rangle_\alpha z),\end{aligned}$$

$$\begin{aligned}
(x \succ_{\alpha} y) \prec_{\beta} z &= P_{\alpha}(x)y \prec_{\beta} z = P_{\alpha}(x)yP_{\beta}(z) + \lambda P_{\alpha}(x)yz \\
&= P_{\alpha}(x)(yP_{\beta}(z) + \lambda yz) = P_{\alpha}(x)(y \prec_{\beta} z) \\
&= x \succ_{\alpha} (y \prec_{\beta} z), \\
x \succ_{\alpha} (y \succ_{\beta} z) &= x \succ_{\alpha} (P_{\beta}(y)z) = P_{\alpha}(x)(P_{\beta}(y)z) = (P_{\alpha}(x)P_{\beta}(y))z \\
&= P_{\alpha\beta}(P_{\alpha}(x)y + xP_{\beta}(y) + \lambda xy)z \\
&= (P_{\alpha}(x)y + xP_{\beta}(y) + \lambda xy) \succ_{\alpha\beta} z \\
&= (x \succ_{\alpha} y + x \prec_{\beta} y) \succ_{\alpha\beta} z.
\end{aligned}$$

(b) The proof is straightforward. \square

Let $\eta : \mathfrak{RB}\mathfrak{F}_{\lambda} \rightarrow \mathfrak{DD}\mathfrak{F}$ and $\gamma : \mathfrak{DI}\mathfrak{F} \rightarrow \mathfrak{DD}\mathfrak{F}$ be the functors obtained in Proposition 2.10. Then we have the commutative diagram:

$$\begin{array}{ccc}
\mathfrak{RB}\mathfrak{F}_{\lambda} & \xrightarrow{\varepsilon} & \mathfrak{DI}\mathfrak{F} \\
& \searrow \eta & \downarrow \gamma \\
& & \mathfrak{DD}\mathfrak{F}
\end{array}$$

The following result precises the link between (tri)dendriform family algebras and ordinary (tri)dendriform algebras.

Theorem 2.11. (a) $(D \otimes \mathbf{k}\Omega, \prec, \succ)$ is a dendriform algebra, if and only if, $(D, (\prec_{\omega}, \succ_{\omega})_{\omega \in \Omega})$ is a dendriform family algebra, where

$$\begin{aligned}
(x \otimes \alpha) \prec (y \otimes \beta) &:= (x \prec_{\beta} y) \otimes \alpha\beta \\
(x \otimes \alpha) \succ (y \otimes \beta) &:= (x \succ_{\alpha} y) \otimes \alpha\beta, \text{ for } x, y \in D \text{ and } \alpha, \beta \in \Omega.
\end{aligned}$$

(b) $(T \otimes \mathbf{k}\Omega, \prec, \succ, \bullet)$ is a tridendriform algebra, if and only if, $(T, (\prec_{\omega}, \succ_{\omega})_{\omega \in \Omega}, \cdot)$ is a tridendriform family algebra, where

$$\begin{aligned}
(x \otimes \alpha) \prec (y \otimes \beta) &:= (x \prec_{\beta} y) \otimes \alpha\beta, \\
(x \otimes \alpha) \succ (y \otimes \beta) &:= (x \succ_{\alpha} y) \otimes \alpha\beta, \\
(x \otimes \alpha) \bullet (y \otimes \beta) &:= (x \cdot y) \otimes \alpha\beta, \text{ for } x, y \in T \text{ and } \alpha, \beta \in \Omega.
\end{aligned}$$

Proof. We only prove Item (b), as the proof of Item (a) can be deduced from tridendriform case. Since a dendriform family algebra is a tridendriform family algebra which product \cdot is equal to zero. Applying Item (b) to a dendriform family algebra D , we obtain a tridendriform algebra $D \otimes \mathbf{k}\Omega$ which product \cdot is trivial, that is to say a dendriform algebra.

For $x \otimes \alpha, y \otimes \beta, z \otimes \gamma \in T \otimes \mathbf{k}\Omega$, we get

$$\begin{aligned}
((x \otimes \alpha) \prec (y \otimes \beta)) \prec (z \otimes \gamma) &= ((x \prec_{\beta} y) \otimes \alpha\beta) \prec (z \otimes \gamma) \\
&= ((x \prec_{\beta} y) \prec_{\gamma} z) \otimes \alpha\beta\gamma \\
&= (x \prec_{\beta\gamma} (y \prec_{\gamma} z + y \succ_{\beta} z + y \cdot z)) \otimes \alpha\beta\gamma \quad (\text{by Eq. (4)}) \\
&= (x \otimes \alpha) \prec ((y \prec_{\gamma} z + y \succ_{\beta} z + y \cdot z) \otimes \beta\gamma)
\end{aligned}$$

$$\begin{aligned}
&= (x \otimes \alpha) < \left((y <_{\gamma} z) \otimes \beta\gamma + (y >_{\beta} z) \otimes \beta\gamma + (y \cdot z) \otimes \beta\gamma \right) \\
&= (x \otimes \alpha) < \left((y \otimes \beta) < (z \otimes \gamma) + (y \otimes \beta) > (z \otimes \gamma) + (y \otimes \beta) \bullet (z \otimes \gamma) \right), \\
((x \otimes \alpha) > (y \otimes \beta)) < (z \otimes \gamma) &= \left((x >_{\alpha} y) \otimes \alpha\beta \right) < (z \otimes \gamma) = \left((x >_{\alpha} y) <_{\gamma} z \right) \otimes \alpha\beta\gamma \\
&= \left(x >_{\alpha} (y <_{\gamma} z) \right) \otimes \alpha\beta\gamma \quad (\text{by Eq. (5)}) \\
&= (x \otimes \alpha) > \left((y <_{\gamma} z) \otimes \beta\gamma \right) \\
&= (x \otimes \alpha) > \left((y \otimes \beta) < (z \otimes \gamma) \right), \\
(x \otimes \alpha) > \left((y \otimes \beta) > (z \otimes \gamma) \right) &= (x \otimes \alpha) > \left((y >_{\beta} z) \otimes \beta\gamma \right) \\
&= \left(x >_{\alpha} (y >_{\beta} z) \right) \otimes \alpha\beta\gamma \\
&= \left((x <_{\beta} y + x >_{\alpha} y + x \cdot y) >_{\alpha\beta} z \right) \otimes \alpha\beta\gamma \quad (\text{by Eq. (6)}) \\
&= \left((x <_{\beta} y + x >_{\alpha} y + x \cdot y) \otimes \alpha\beta \right) > (z \otimes \gamma) \\
&= \left((x <_{\beta} y) \otimes \alpha\beta + (x >_{\alpha} y) \otimes \alpha\beta + (x \cdot y) \otimes \alpha\beta \right) > (z \otimes \gamma) \\
&= \left((x \otimes \alpha) < (y \otimes \beta) + (x \otimes \alpha) > (y \otimes \beta) + (x \otimes \alpha) \bullet (y \otimes \beta) \right) > (z \otimes \gamma), \\
((x \otimes \alpha) > (y \otimes \beta)) \bullet (z \otimes \gamma) &= \left((x >_{\alpha} y) \otimes \alpha\beta \right) \bullet (z \otimes \gamma) \\
&= \left((x >_{\alpha} y) \cdot z \right) \otimes \alpha\beta\gamma = \left(x >_{\alpha} (y \cdot z) \right) \otimes \alpha\beta\gamma \quad (\text{by Eq. (7)}) \\
&= (x \otimes \alpha) > \left((y \cdot z) \otimes \beta\gamma \right) \\
&= (x \otimes \alpha) > \left((y \otimes \beta) \bullet (z \otimes \gamma) \right), \\
((x \otimes \alpha) < (y \otimes \beta)) \bullet (z \otimes \gamma) &= \left((x <_{\beta} y) \otimes \alpha\beta \right) \bullet (z \otimes \gamma) = \left((x <_{\beta} y) \cdot z \right) \otimes \alpha\beta\gamma \\
&= \left(x \cdot (y >_{\beta} z) \right) \otimes \alpha\beta\gamma \quad (\text{by Eq. (8)}) \\
&= (x \otimes \alpha) \bullet \left((y >_{\beta} z) \otimes \beta\gamma \right) \\
&= (x \otimes \alpha) \bullet \left((y \otimes \beta) > (z \otimes \gamma) \right), \\
((x \otimes \alpha) \bullet (y \otimes \beta)) < (z \otimes \gamma) &= \left((x \cdot y) \otimes \alpha\beta \right) < (z \otimes \gamma) = \left((x \cdot y) <_{\gamma} z \right) \otimes \alpha\beta\gamma \\
&= \left(x \cdot (y <_{\gamma} z) \right) \otimes \alpha\beta\gamma \quad (\text{by Eq. (9)}) \\
&= (x \otimes \alpha) \bullet \left((y <_{\gamma} z) \otimes \beta\gamma \right) \\
&= (x \otimes \alpha) \bullet \left((y \otimes \beta) < (z \otimes \gamma) \right), \\
((x \otimes \alpha) \bullet (y \otimes \beta)) \bullet (z \otimes \gamma) &= \left((x \cdot y) \otimes \alpha\beta \right) \bullet (z \otimes \gamma) = \left((x \cdot y) \cdot z \right) \otimes \alpha\beta\gamma \\
&= \left(x \cdot (y \cdot z) \right) \otimes \alpha\beta\gamma \quad (\text{by Eq. (10)}) \\
&= (x \otimes \alpha) \bullet \left((y \cdot z) \otimes \beta\gamma \right) \\
&= (x \otimes \alpha) \bullet \left((y \otimes \beta) \bullet (z \otimes \gamma) \right).
\end{aligned}$$

Conversely, we notice that the tridendriform structure on $T \otimes \mathbf{k}\Omega$ is Ω -graded, with ω -homogeneous component equal to $T \otimes \mathbf{k}\omega$ for any $\omega \in \Omega$. Identifying each homogeneous component in both sides of the three dendriform axioms leads to tridendriform family axioms on T . This completes the proof. \square

3. FREE DENDRIFORM FAMILY ALGEBRAS

In this section, we construct free dendriform family algebras. For this, let us first briefly recall the construction of free dendriform algebras. For details, see [20, 21, 22, 26].

3.1. Free dendriform algebras. Let X be a set. For $n \geq 0$, let $Y_{n,X}$ be the set of planar binary trees with $n + 1$ leaves and with internal vertices decorated by elements of X . The unique tree with one leaf is denoted by $|$. So we have $Y_{0,X} = \{| \}$. Here are the first few of them.

$$Y_{0,X} = \{| \}, \quad Y_{1,X} = \left\{ \begin{array}{c} x \\ \diagup \quad \diagdown \\ | \quad | \end{array} \mid x \in X \right\}, \quad Y_{2,X} = \left\{ \begin{array}{c} y \\ \diagup \quad \diagdown \\ x \quad | \\ | \quad | \end{array}, \begin{array}{c} y \\ \diagup \quad \diagdown \\ | \quad x \\ | \quad | \end{array} \mid x, y \in X \right\},$$

$$Y_{3,X} = \left\{ \begin{array}{c} z \\ \diagup \quad \diagdown \\ y \quad x \\ | \quad | \end{array}, \begin{array}{c} z \\ \diagup \quad \diagdown \\ x \quad y \\ | \quad | \end{array}, \begin{array}{c} z \\ \diagup \quad \diagdown \\ | \quad x \\ | \quad y \\ | \quad | \end{array}, \begin{array}{c} y \\ \diagup \quad \diagdown \\ z \quad x \\ | \quad | \end{array}, \begin{array}{c} z \\ \diagup \quad \diagdown \\ | \quad x \\ | \quad y \\ | \quad | \end{array} \mid x, y, z \in X \right\}.$$

For $T \in Y_{m,X}$, $U \in Y_{n,X}$ and $x \in X$, the grafting \vee_x of T and U over the vertex x is defined to be the planar binary tree $T \vee_x U \in Y_{m+n+1,X}$ obtained by adding a new vertex decorated by x and joining the roots of T and U to the new vertex.

Remark 3.1. In the graphical representation above, the edge pointing downwards is the root, the upper edges are the leaves. The other edges, joining two internal vertices, are called internal edges.

Example 3.2. Let $T = |$ and $U = \begin{array}{c} y \\ \diagup \quad \diagdown \\ | \quad | \end{array}$ with $y \in X$. Then

$$T \vee_x U = | \vee_x \begin{array}{c} y \\ \diagup \quad \diagdown \\ | \quad | \end{array} = \begin{array}{c} y \\ \diagup \quad \diagdown \\ x \quad | \\ | \quad | \end{array}.$$

Given a planar binary tree $T \in Y_{n,X}$ not equal to $|$, there is a unique decomposition $T = T^l \vee_x T^r$ for some $x \in X$. For example,

$$\begin{array}{c} x \\ \diagup \quad \diagdown \\ | \quad | \end{array} = | \vee_x |, \quad \begin{array}{c} y \\ \diagup \quad \diagdown \\ x \quad | \\ | \quad | \end{array} = \begin{array}{c} y \\ \diagup \quad \diagdown \\ | \quad | \end{array} \vee_x |, \quad \begin{array}{c} y \\ \diagup \quad \diagdown \\ | \quad x \\ | \quad | \end{array} = | \vee_x \begin{array}{c} y \\ \diagup \quad \diagdown \\ | \quad | \end{array}.$$

Let $\text{DD}(X) := \bigoplus_{n \geq 1} \mathbf{k}Y_{n,X}$. Define binary operations $<$ and $>$ on $\text{DD}(X)$ recursively by

- (a) $| > T := T < | := T$ and $| < T := T > | := 0$ for $T \in Y_{n,X}$ with $n \geq 1$.
- (b) For $T = T^l \vee_x T^r$ and $U = U^l \vee_y U^r$, define

$$T < U := T^l \vee_x (T^r < U + T^r > U), \quad T > U := (T < U^l + T > U^l) \vee_y U^r.$$

Let j be the unique linear map from X into $\text{DD}(X)$ defined by $j(x) = \begin{array}{c} x \\ \diagup \quad \diagdown \\ | \quad | \end{array}$ for $x \in X$. The following result is well-known.

Theorem 3.3. [21] *Let X be a set. Then $(\text{DD}(X), <, >)$, together with the map j , is the free dendriform algebra on X .*

3.2. Free dendriform family algebras. In this subsection, we apply typed decorated planar binary trees to construct free dendriform family algebras. For this, let us first recall typed decorated rooted trees studied in [2, 11].

For a rooted tree T , denote by $V(T)$ (resp. $E(T)$) the set of its vertices (resp. edges).

Definition 3.4. [2] Let X and Ω be two sets. An X -decorated Ω -typed (abbreviated typed decorated) rooted tree is a triple $T = (T, \text{dec}, \text{type})$, where

- (a) T is a rooted tree.
- (b) $\text{dec} : V(T) \rightarrow X$ is a map.
- (c) $\text{type} : E(T) \rightarrow \Omega$ is a map.

In other words, vertices of T are decorated by elements of X and edges of T are decorated by elements of Ω .

As we can see from the examples drawn above, the graphical representation of planar binary trees is slightly different from the graphical representation of rooted trees used in [2, 11]: the root and the leaves are now edges rather than vertices. Here the set $V(T)$ must be replaced by the set $IV(T)$ of internal vertices of T , and the set $E(T)$ must be replaced by the set $IE(T)$ of internal edges, i.e. edges which are neither a leaf nor the root. Hence we propose the following definition:

Definition 3.5. Let X and Ω be two sets. An X -decorated Ω -typed planar binary tree is a triple $T = (T, \text{dec}, \text{type})$, where

- (a) T is a planar binary tree.
- (b) $\text{dec} : IV(T) \rightarrow X$ is a map.
- (c) $\text{type} : IE(T) \rightarrow \Omega$ is a map.

Note that in the definition of dendriform family algebras, Ω is a semigroup. However in the following construction of free dendriform family algebras, if the semigroup Ω has no identity element, it will be convenient to consider the monoid $\Omega^1 := \Omega \sqcup \{1\}$ obtained from Ω by adjoining an identity:

$$1\omega := \omega 1 := \omega, \text{ for } \omega \in \Omega \text{ and } 11 := 1.$$

Let X be a set and let Ω be a semigroup. For $n \geq 0$, let $Y_n := Y_{n, X, \Omega}$ be the set of X -decorated Ω^1 -typed planar binary trees with $n + 1$ leaves, such that leaves are decorated by the identity 1 in Ω^1 and internal edges are decorated by elements of Ω . Let us expose some examples for better understanding. For convenience, we omit the decoration 1 in the sequel.

$$Y_0 = \{\emptyset\}, \quad Y_1 = \left\{ \begin{array}{c} x \\ \diagdown \quad \diagup \\ \quad \quad \quad \end{array} \mid x \in X \right\}, \quad Y_2 = \left\{ \begin{array}{c} y \\ \diagdown \quad \diagup \\ \alpha \quad \quad \quad \end{array}, \begin{array}{c} y \\ \diagdown \quad \diagup \\ \quad \quad \alpha \end{array} \mid x, y \in X, \alpha \in \Omega \right\},$$

$$Y_3 = \left\{ \begin{array}{c} z \quad y \\ \diagdown \quad \diagup \\ \beta \quad \alpha \quad x \\ \quad \quad \quad \end{array}, \begin{array}{c} \quad y \quad z \\ \diagdown \quad \diagup \\ x \quad \alpha \quad \beta \\ \quad \quad \quad \end{array}, \begin{array}{c} z \quad \quad y \\ \diagdown \quad \diagup \\ \beta \quad x \quad \alpha \\ \quad \quad \quad \end{array}, \begin{array}{c} \quad y \quad z \\ \diagdown \quad \diagup \\ \alpha \quad \beta \quad x \\ \quad \quad \quad \end{array}, \begin{array}{c} \quad \quad z \quad y \\ \diagdown \quad \diagup \\ x \quad \beta \quad \alpha \\ \quad \quad \quad \end{array}, \dots \mid x, y, z \in X, \alpha, \beta \in \Omega \right\}.$$

For $T \in Y_m, U \in Y_n, x \in X$ and $\alpha, \beta \in \Omega^1$, the grafting $\vee_{x, (\alpha, \beta)}$ of T and U over x and (α, β) is defined to be the tree $T \vee_{x, (\alpha, \beta)} U \in Y_{m+n+1}$, obtained by adding a new vertex decorated by x and joining the roots of T and U with the new vertex via two new edges decorated by α and β respectively. The **depth** $\text{dep}(T)$ of a rooted tree T is the maximal length of linear chains of

vertices from the root to the leaves of the tree. For example,

$$\text{dep}\left(\begin{array}{c} x \\ \diagdown \quad \diagup \\ \end{array}\right) = 1 \quad \text{and} \quad \text{dep}\left(\begin{array}{c} y \\ \diagdown \quad \diagup \\ \alpha \end{array}\right) = 2.$$

Example 3.6. Let $T = \begin{array}{c} x \\ \diagdown \quad \diagup \\ \end{array}$ and $U = \begin{array}{c} y \\ \diagdown \quad \diagup \\ \end{array}$ with $x, y \in X$. For $z \in X$ and $\alpha, \beta \in \Omega$,

$$T \vee_{z, (\alpha, \beta)} U = \begin{array}{c} x \\ \diagdown \quad \diagup \\ \end{array} \vee_{z, (\alpha, \beta)} \begin{array}{c} y \\ \diagdown \quad \diagup \\ \end{array} = \begin{array}{c} x \quad y \\ \diagdown \quad \diagup \\ \alpha \quad \beta \end{array}.$$

Given a typed decorated planar binary tree $T \in Y_n$ not equal to $|$, there is a unique decomposition

$$T = T^l \vee_{x, (\alpha, \beta)} T^r \quad \text{for some } x \in X \text{ and } \alpha, \beta \in \Omega^1.$$

For example,

$$\begin{array}{c} x \\ \diagdown \quad \diagup \\ \end{array} = | \vee_{x, (1, 1)} |, \quad \begin{array}{c} y \\ \diagdown \quad \diagup \\ \alpha \end{array} = \begin{array}{c} y \\ \diagdown \quad \diagup \\ \end{array} \vee_{x, (\alpha, 1)} |, \quad \begin{array}{c} y \\ \diagdown \quad \diagup \\ \alpha \end{array} = | \vee_{x, (1, \alpha)} \begin{array}{c} y \\ \diagdown \quad \diagup \\ \end{array}, \\ \begin{array}{c} y \quad z \\ \diagdown \quad \diagup \\ \alpha \quad \beta \end{array} = \begin{array}{c} y \\ \diagdown \quad \diagup \\ \end{array} \vee_{x, (\alpha, \beta)} \begin{array}{c} z \\ \diagdown \quad \diagup \\ \end{array}.$$

Denote by

$$\text{DD}(X, \Omega) := \bigoplus_{n \geq 1} \mathbf{k}Y_n.$$

Definition 3.7. Let X be a set and let Ω be a semigroup. We define binary operations $(\prec_\omega, \succ_\omega)_{\omega \in \Omega}$ on $\text{DD}(X, \Omega)$ recursively on $\text{dep}(T) + \text{dep}(U)$ by

(a) $| \succ_\omega T := T \prec_\omega | := T$ and $| \prec_\omega T := T \succ_\omega | := 0$ for $\omega \in \Omega$ and $T \in Y_n$ with $n \geq 1$.

(b) For $T = T^l \vee_{x, (\alpha_1, \alpha_2)} T^r$ and $U = U^l \vee_{y, (\beta_1, \beta_2)} U^r$, define

$$(11) \quad T \prec_\omega U := T^l \vee_{x, (\alpha_1, \alpha_2 \omega)} (T^r \prec_\omega U + T^r \succ_{\alpha_2} U),$$

$$(12) \quad T \succ_\omega U := (T \prec_{\beta_1} U^l + T \succ_\omega U^l) \vee_{y, (\omega \beta_1, \beta_2)} U^r, \quad \text{where } \omega \in \Omega.$$

In the following, we employ the convention that

$$(13) \quad | \succ_1 T := T \prec_1 | := T \quad \text{and} \quad | \prec_1 T := T \succ_1 | := 0.$$

Example 3.8. Let $T = \begin{array}{c} x \\ \diagdown \quad \diagup \\ \end{array}$ and $U = \begin{array}{c} y \\ \diagdown \quad \diagup \\ \end{array}$ with $x, y \in X$. For $\omega \in \Omega$, we have

$$\begin{aligned} \begin{array}{c} x \\ \diagdown \quad \diagup \\ \end{array} \prec_\omega \begin{array}{c} y \\ \diagdown \quad \diagup \\ \end{array} &= (| \vee_{x, (1, 1)} |) \prec_\omega \begin{array}{c} y \\ \diagdown \quad \diagup \\ \end{array} \\ &= | \vee_{x, (1, \omega)} \left(| \prec_\omega \begin{array}{c} y \\ \diagdown \quad \diagup \\ \end{array} + | \succ_1 \begin{array}{c} y \\ \diagdown \quad \diagup \\ \end{array} \right) \quad (\text{by Eq. (11)}) \\ &= | \vee_{x, (1, \omega)} \begin{array}{c} y \\ \diagdown \quad \diagup \\ \end{array} \quad (\text{by Item (a) of Definition 3.7 and Eq. (13)}) \\ &= \begin{array}{c} y \\ \diagdown \quad \diagup \\ \omega \end{array}, \\ \begin{array}{c} x \\ \diagdown \quad \diagup \\ \end{array} \succ_\omega \begin{array}{c} y \\ \diagdown \quad \diagup \\ \end{array} &= \begin{array}{c} x \\ \diagdown \quad \diagup \\ \end{array} \succ_\omega (| \vee_{y, (1, 1)} |) \\ &= \left(\begin{array}{c} x \\ \diagdown \quad \diagup \\ \end{array} \prec_1 | + \begin{array}{c} x \\ \diagdown \quad \diagup \\ \end{array} \succ_\omega | \right) \vee_{y, (\omega, 1)} | \quad (\text{by Eq. (12)}) \\ &= \begin{array}{c} x \\ \diagdown \quad \diagup \\ \end{array} \vee_{y, (\omega, 1)} | \quad (\text{by Item (a) of Definition 3.7 and Eq. (13)}) \end{aligned}$$

$$= \begin{array}{c} x \\ \swarrow \\ \omega \\ \searrow \\ y \end{array}.$$

Example 3.9. Let $T = \begin{array}{c} z \\ \swarrow \\ \alpha \\ \searrow \\ x \\ \swarrow \\ \beta \\ \searrow \\ u \end{array}$ and $U = \begin{array}{c} y \\ \swarrow \\ \searrow \end{array}$ with $x, y, z, u \in X$ and $\alpha, \beta \in \Omega$. For $\omega \in \Omega$,

$$\begin{aligned} T <_{\omega} U &= \left(\begin{array}{c} z \\ \swarrow \\ \searrow \\ \vee_{x,(\alpha,\beta)} \end{array} \begin{array}{c} u \\ \swarrow \\ \searrow \end{array} \right) <_{\omega} \begin{array}{c} y \\ \swarrow \\ \searrow \end{array} \\ &= \begin{array}{c} z \\ \swarrow \\ \searrow \\ \vee_{x,(\alpha,\beta\omega)} \end{array} \left(\begin{array}{c} u \\ \swarrow \\ \searrow \end{array} <_{\omega} \begin{array}{c} y \\ \swarrow \\ \searrow \end{array} + \begin{array}{c} u \\ \swarrow \\ \searrow \\ >_{\beta} \end{array} \begin{array}{c} y \\ \swarrow \\ \searrow \end{array} \right) \\ &= \begin{array}{c} z \\ \swarrow \\ \searrow \\ \vee_{x,(\alpha,\beta\omega)} \end{array} \left(\begin{array}{c} u \\ \swarrow \\ \searrow \\ \omega \end{array} + \begin{array}{c} u \\ \swarrow \\ \searrow \\ \beta \end{array} \begin{array}{c} y \\ \swarrow \\ \searrow \end{array} \right) \quad (\text{by Example 3.8}) \\ &= \begin{array}{c} z \\ \swarrow \\ \alpha \\ \searrow \\ x \\ \swarrow \\ \beta\omega \\ \searrow \\ u \end{array} + \begin{array}{c} z \\ \swarrow \\ \alpha \\ \searrow \\ x \\ \swarrow \\ \beta\omega \\ \searrow \\ u \end{array}, \end{aligned}$$

$$\begin{aligned} T >_{\omega} U &= \begin{array}{c} z \\ \swarrow \\ \alpha \\ \searrow \\ x \\ \swarrow \\ \beta \end{array} >_{\omega} (| \vee_{y,(1,1)} |) = \left(\begin{array}{c} z \\ \swarrow \\ \alpha \\ \searrow \\ x \\ \swarrow \\ \beta \end{array} <_1 | + \begin{array}{c} z \\ \swarrow \\ \alpha \\ \searrow \\ x \\ \swarrow \\ \beta \end{array} >_{\omega} | \right) \vee_{y,(\omega,1)} | \\ &= \begin{array}{c} z \\ \swarrow \\ \alpha \\ \searrow \\ x \\ \swarrow \\ \beta \end{array} \vee_{y,(\omega,1)} | \quad (\text{by Item (a) of Definition 3.7 and Eq. (13)}) \\ &= \begin{array}{c} z \\ \swarrow \\ \alpha \\ \searrow \\ x \\ \swarrow \\ \beta \\ \searrow \\ \omega \\ \searrow \\ y \end{array}. \end{aligned}$$

Proposition 3.10. Let X be a set and let Ω be a semigroup. Then $(\text{DD}(X, \Omega), (<_{\omega}, >_{\omega})_{\omega \in \Omega})$ is a dendriform family algebra, generated by $\left\{ \begin{array}{c} x \\ \swarrow \\ \searrow \end{array} \mid x \in X \right\}$.

Proof. We proceed in two steps to prove the result.

Step 1: We first prove $(\text{DD}(X, \Omega), \{<_{\omega}, >_{\omega} \mid \omega \in \Omega\})$ is a dendriform family algebra. Let

$$T = T^l \vee_{x,(\alpha_1,\alpha_2)} T^r, U = U^l \vee_{y,(\beta_1,\beta_2)} U^r, W = W^l \vee_{z,(\gamma_1,\gamma_2)} W^r \in \text{DD}(X, \Omega).$$

Then we apply induction on $\text{dep}(T) + \text{dep}(U) + \text{dep}(W) \geq 3$. For the initial step $\text{dep}(T) + \text{dep}(U) + \text{dep}(W) = 3$, we have

$$T = \begin{array}{c} x \\ \swarrow \\ \searrow \end{array}, U = \begin{array}{c} y \\ \swarrow \\ \searrow \end{array} \quad \text{and} \quad W = \begin{array}{c} z \\ \swarrow \\ \searrow \end{array}$$

and so

$$\begin{aligned} &(T <_{\beta} U) <_{\gamma} W \\ &= \left(\begin{array}{c} x \\ \swarrow \\ \searrow \\ <_{\beta} \end{array} \begin{array}{c} y \\ \swarrow \\ \searrow \end{array} \right) <_{\gamma} \begin{array}{c} z \\ \swarrow \\ \searrow \end{array} = \left((| \vee_{x,(1,1)} |) <_{\beta} \begin{array}{c} y \\ \swarrow \\ \searrow \end{array} \right) <_{\gamma} \begin{array}{c} z \\ \swarrow \\ \searrow \end{array} \\ &= \left(| \vee_{x,(1,\beta)} \left(| <_{\beta} \begin{array}{c} y \\ \swarrow \\ \searrow \end{array} + | >_1 \begin{array}{c} y \\ \swarrow \\ \searrow \end{array} \right) \right) <_{\gamma} \begin{array}{c} z \\ \swarrow \\ \searrow \end{array} \quad (\text{by Eq. (11)}) \\ &= \left(| \vee_{x,(1,\beta)} \begin{array}{c} y \\ \swarrow \\ \searrow \end{array} \right) <_{\gamma} \begin{array}{c} z \\ \swarrow \\ \searrow \end{array} \quad (\text{by Item (a) in Definition 3.7 and Eq. (13)}) \\ &= | \vee_{x,(1,\beta\gamma)} \left(\begin{array}{c} y \\ \swarrow \\ \searrow \\ <_{\gamma} \end{array} \begin{array}{c} z \\ \swarrow \\ \searrow \end{array} + \begin{array}{c} y \\ \swarrow \\ \searrow \\ >_{\beta} \end{array} \begin{array}{c} z \\ \swarrow \\ \searrow \end{array} \right) \quad (\text{by Eq. (11)}) \\ &= | \vee_{x,(1,\beta\gamma)} \left(\begin{array}{c} y \\ \swarrow \\ \searrow \\ \gamma \end{array} + \begin{array}{c} y \\ \swarrow \\ \searrow \\ \beta \end{array} \begin{array}{c} z \\ \swarrow \\ \searrow \end{array} \right) \quad (\text{by Example 3.8}) \end{aligned}$$

$$\begin{aligned}
&= | \vee_{x,(1,\beta\gamma)} \left(| \prec_{\beta\gamma} \left(\begin{array}{c} y/z \\ \vee \\ \gamma \end{array} + \begin{array}{c} y/z \\ \vee \\ \beta \end{array} \right) + | \succ_1 \left(\begin{array}{c} y/z \\ \vee \\ \gamma \end{array} + \begin{array}{c} y/z \\ \vee \\ \beta \end{array} \right) \right) \\
&\quad \text{(by Item (a) in Definition 3.7 and Eq. (13))} \\
&= (| \vee_{x,(1,1)} |) \prec_{\beta\gamma} \left(\begin{array}{c} y/z \\ \vee \\ \gamma \end{array} + \begin{array}{c} y/z \\ \vee \\ \beta \end{array} \right) \quad \text{(by Eq. (11))} \\
&= \begin{array}{c} x \\ \vee \\ \prec_{\beta\gamma} \left(\begin{array}{c} y/z \\ \vee \\ \gamma \end{array} + \begin{array}{c} y/z \\ \vee \\ \beta \end{array} \right) \end{array} \\
&= \begin{array}{c} x \\ \vee \\ \prec_{\beta\gamma} \left(\begin{array}{c} y \\ \vee \\ \prec_{\gamma} \begin{array}{c} z \\ \vee \\ \end{array} + \begin{array}{c} y \\ \vee \\ \succ_{\beta} \begin{array}{c} z \\ \vee \\ \end{array} \end{array} \right) \\
&= T \prec_{\beta\gamma} (U \prec_{\gamma} W + U \succ_{\beta} W),
\end{aligned}$$

verifying Eq. (1). Next,

$$\begin{aligned}
&(T \succ_{\alpha} U) \prec_{\gamma} W \\
&= \left(\begin{array}{c} x \\ \vee \\ \succ_{\alpha} \begin{array}{c} y \\ \vee \\ \end{array} \right) \prec_{\gamma} \begin{array}{c} z \\ \vee \\ \end{array} = \left(\begin{array}{c} x \\ \vee \\ \succ_{\alpha} (| \vee_{y,(1,1)} |) \right) \prec_{\gamma} \begin{array}{c} z \\ \vee \\ \end{array} \\
&= \left(\left(\begin{array}{c} x \\ \vee \\ \succ_{\alpha} | + \begin{array}{c} x \\ \vee \\ \prec_1 | \end{array} \right) \vee_{y,(\alpha,1)} | \right) \prec_{\gamma} \begin{array}{c} z \\ \vee \\ \end{array} \quad \text{(by Eq. (12))} \\
&= \left(\begin{array}{c} x \\ \vee \\ \vee_{y,(\alpha,1)} | \end{array} \right) \prec_{\gamma} \begin{array}{c} z \\ \vee \\ \end{array} \quad \text{(by Item (a) in Definition 3.7 and Eq. (13))} \\
&= \begin{array}{c} x \\ \vee \\ \vee_{y,(\alpha,\gamma)} \left(| \prec_{\gamma} \begin{array}{c} z \\ \vee \\ \end{array} + | \succ_1 \begin{array}{c} z \\ \vee \\ \end{array} \right) \end{array} \quad \text{(by Eq. (11))} \\
&= \begin{array}{c} x \\ \vee \\ \vee_{y,(\alpha,\gamma)} \begin{array}{c} z \\ \vee \\ \end{array} \end{array} \quad \text{(by Item (a) in Definition 3.7 and Eq. (13))} \\
&= \left(\begin{array}{c} x \\ \vee \\ \succ_{\alpha} | + \begin{array}{c} x \\ \vee \\ \prec_1 | \end{array} \right) \vee_{y,(\alpha,\gamma)} \begin{array}{c} z \\ \vee \\ \end{array} \quad \text{(by Item (a) in Definition 3.7 and Eq. (13))} \\
&= \begin{array}{c} x \\ \vee \\ \succ_{\alpha} \left(| \vee_{y,(1,\gamma)} \begin{array}{c} z \\ \vee \\ \end{array} \right) \end{array} \quad \text{(by Eq. (12))} \\
&= \begin{array}{c} x \\ \vee \\ \succ_{\alpha} \left(| \vee_{y,(1,\gamma)} \left(| \prec_{\gamma} \begin{array}{c} z \\ \vee \\ \end{array} + | \succ_1 \begin{array}{c} z \\ \vee \\ \end{array} \right) \right) \end{array} \quad \text{(by Item (a) in Definition 3.7 and Eq. (13))} \\
&= \begin{array}{c} x \\ \vee \\ \succ_{\alpha} \left((| \vee_{y,(1,1)} |) \prec_{\gamma} \begin{array}{c} z \\ \vee \\ \end{array} \right) = \begin{array}{c} x \\ \vee \\ \succ_{\alpha} \left(\begin{array}{c} y \\ \vee \\ \prec_{\gamma} \begin{array}{c} z \\ \vee \\ \end{array} \end{array} \right) \\
&= T \succ_{\alpha} (U \prec_{\gamma} W),
\end{aligned}$$

verifying Eq. (2). Finally,

$$\begin{aligned}
&T \succ_{\alpha} (U \succ_{\beta} W) \\
&= \begin{array}{c} x \\ \vee \\ \succ_{\alpha} \left(\begin{array}{c} y \\ \vee \\ \succ_{\beta} \begin{array}{c} z \\ \vee \\ \end{array} \end{array} \right) = \begin{array}{c} x \\ \vee \\ \succ_{\alpha} \left(\begin{array}{c} y \\ \vee \\ \succ_{\beta} (| \vee_{z,(1,1)} |) \end{array} \right) \\
&= \begin{array}{c} x \\ \vee \\ \succ_{\alpha} \left(\left(\begin{array}{c} y \\ \vee \\ \succ_{\beta} | + \begin{array}{c} y \\ \vee \\ \prec_1 | \end{array} \right) \vee_{z,(\beta,1)} | \right) \end{array} \quad \text{(by Eq. (12))} \\
&= \begin{array}{c} x \\ \vee \\ \succ_{\alpha} \left(\begin{array}{c} y \\ \vee \\ \vee_{z,(\beta,1)} | \end{array} \right) \end{array} \quad \text{(by Item (a) in Definition 3.7 and Eq. (13))} \\
&= \left(\begin{array}{c} x \\ \vee \\ \succ_{\alpha} \begin{array}{c} y \\ \vee \\ \end{array} + \begin{array}{c} x \\ \vee \\ \prec_{\beta} \begin{array}{c} y \\ \vee \\ \end{array} \end{array} \right) \vee_{z,(\alpha\beta,1)} | \quad \text{(by Eq. (12))}
\end{aligned}$$

$$\begin{aligned}
&= \left(\begin{array}{c} x \quad y \\ \diagdown \quad / \\ \alpha \quad \beta \end{array} + \begin{array}{c} x \quad y \\ / \quad \diagdown \\ \alpha \quad \beta \end{array} \right) \vee_{z, (\alpha\beta, 1)} \mid \quad (\text{by Example 3.8}) \\
&= \left(\left(\begin{array}{c} x \quad y \\ \diagdown \quad / \\ \alpha \quad \beta \end{array} + \begin{array}{c} x \quad y \\ / \quad \diagdown \\ \alpha \quad \beta \end{array} \right) \succ_{\alpha\beta} \mid + \left(\begin{array}{c} x \quad y \\ \diagdown \quad / \\ \alpha \quad \beta \end{array} + \begin{array}{c} x \quad y \\ / \quad \diagdown \\ \alpha \quad \beta \end{array} \right) \prec_1 \mid \right) \vee_{z, (\alpha\beta, 1)} \mid \\
&\quad (\text{by Item (a) in Definition 3.7 and Eq. (13)}) \\
&= \left(\begin{array}{c} x \quad y \\ \diagdown \quad / \\ \alpha \quad \beta \end{array} + \begin{array}{c} x \quad y \\ / \quad \diagdown \\ \alpha \quad \beta \end{array} \right) \succ_{\alpha\beta} (\mid \vee_{z, (1, 1)} \mid) \quad (\text{by Eq. (12)}) \\
&= \left(\begin{array}{c} x \quad y \\ \diagdown \quad / \\ \alpha \quad \beta \end{array} \succ_{\alpha} \begin{array}{c} y \\ / \\ \beta \end{array} + \begin{array}{c} x \\ \diagdown \\ \alpha \end{array} \begin{array}{c} y \quad z \\ / \quad \diagdown \\ \beta \quad \gamma \end{array} \right) \succ_{\alpha\beta} \begin{array}{c} y \\ / \\ \beta \end{array} \quad (\text{by Example 3.8}) \\
&= (T \succ_{\alpha} U + T \prec_{\beta} U) \succ_{\alpha\beta} W.
\end{aligned}$$

This completes the proof of the initial step. Assume that the conclusion holds for $\text{dep}(T) + \text{dep}(U) + \text{dep}(W) = k$ and consider the induction step of $\text{dep}(T) + \text{dep}(U) + \text{dep}(W) = k + 1 \geq 4$. We have

$$\begin{aligned}
&(T \prec_{\beta} U) \prec_{\gamma} W \\
&= \left(T^l \vee_{x, (\alpha_1, \alpha_2\beta)} (T^r \prec_{\beta} U + T^r \succ_{\alpha_2} U) \right) \prec_{\gamma} W \quad (\text{by Eq. (11)}) \\
&= T^l \vee_{x, (\alpha_1, \alpha_2\beta\gamma)} \left((T^r \prec_{\beta} U + T^r \succ_{\alpha_2} U) \prec_{\gamma} W + (T^r \prec_{\beta} U + T^r \succ_{\alpha_2} U) \succ_{\alpha_2\beta} W \right) \\
&\quad (\text{by Eq. (11)}) \\
&= T^l \vee_{x, (\alpha_1, \alpha_2\beta\gamma)} \left((T^r \prec_{\beta} U) \prec_{\gamma} W + (T^r \succ_{\alpha_2} U) \prec_{\gamma} W + T^r \succ_{\alpha_2} (U \succ_{\beta} W) \right) \\
&\quad (\text{by the induction hypothesis and Eq. (3)}) \\
&= T^l \vee_{x, (\alpha_1, \alpha_2\beta\gamma)} \left(T^r \prec_{\beta\gamma} (U \prec_{\gamma} W + U \succ_{\beta} W) + T^r \succ_{\alpha_2} (U \prec_{\gamma} W) + T^r \succ_{\alpha_2} (U \succ_{\beta} W) \right) \\
&\quad (\text{by the induction hypothesis and Eq. (1)}) \\
&= T^l \vee_{x, (\alpha_1, \alpha_2\beta\gamma)} \left(T^r \prec_{\beta\gamma} (U \prec_{\gamma} W + U \succ_{\beta} W) + T^r \succ_{\alpha_2} (U \prec_{\gamma} W + U \succ_{\beta} W) \right) \\
&= (T^l \vee_{x, (\alpha_1, \alpha_2)} T^r) \prec_{\beta\gamma} (U \prec_{\gamma} W + U \succ_{\beta} W) \quad (\text{by Eq. (11)}) \\
&= T \prec_{\beta\gamma} (U \prec_{\gamma} W + U \succ_{\beta} W).
\end{aligned}$$

This proves Eq. (1) in Definition 2.6. Next, we prove Eq. (2):

$$\begin{aligned}
&(T \succ_{\alpha} U) \prec_{\gamma} W \\
&= \left((T \succ_{\alpha} U^l + T \prec_{\beta_1} U^l) \vee_{y, (\alpha\beta_1, \beta_2)} U^r \right) \prec_{\gamma} W \quad (\text{by Eq. (12)}) \\
&= (T \succ_{\alpha} U^l + T \prec_{\beta_1} U^l) \vee_{y, (\alpha\beta_1, \beta_2\gamma)} (U^r \prec_{\gamma} W + U^r \succ_{\beta_2} W) \quad (\text{by Eq. (11)}) \\
&= T \succ_{\alpha} \left(U^l \vee_{y, (\beta_1, \beta_2\gamma)} (U^r \prec_{\gamma} W + U^r \succ_{\beta_2} W) \right) \quad (\text{by Eq. (12)}) \\
&= T \succ_{\alpha} \left((U^l \vee_{y, (\beta_1, \beta_2)} U^r) \prec_{\gamma} W \right) \quad (\text{by Eq. (11)}) \\
&= T \succ_{\alpha} (U \prec_{\gamma} W).
\end{aligned}$$

Finally, we prove Eq. (3):

$$\begin{aligned}
&T \succ_{\alpha} (U \succ_{\beta} W) \\
&= T \succ_{\alpha} \left(U \succ_{\beta} (W^l \vee_{z, (\gamma_1, \gamma_2)} W^r) \right)
\end{aligned}$$

$$\begin{aligned}
&= T \succ_{\alpha} \left((U \succ_{\beta} W^l + U \prec_{\gamma_1} W^l) \vee_{z, (\beta\gamma_1, \gamma_2)} W^r \right) \quad (\text{by Eq. (12)}) \\
&= \left(T \succ_{\alpha} (U \succ_{\beta} W^l + U \prec_{\gamma_1} W^l) + T \prec_{\beta\gamma_1} (U \succ_{\beta} W^l + U \prec_{\gamma_1} W^l) \right) \vee_{z, (\alpha\beta\gamma_1, \gamma_2)} W^r \quad (\text{by Eq. (12)}) \\
&= \left(T \succ_{\alpha} (U \succ_{\beta} W^l) + T \succ_{\alpha} (U \prec_{\gamma_1} W^l) + T \prec_{\beta\gamma_1} (U \succ_{\beta} W^l + U \prec_{\gamma_1} W^l) \right) \vee_{z, (\alpha\beta\gamma_1, \gamma_2)} W^r \\
&= \left((T \succ_{\alpha} U + T \prec_{\beta} U) \succ_{\alpha\beta} W^l + (T \succ_{\alpha} U) \prec_{\gamma_1} W^l + (T \prec_{\beta} U) \prec_{\gamma_1} W^l \right) \vee_{z, (\alpha\beta\gamma_1, \gamma_2)} W^r \\
&\quad (\text{by the induction hypothesis and Eqs. (1)-(3)}) \\
&= \left((T \succ_{\alpha} U + T \prec_{\beta} U) \succ_{\alpha\beta} W^l + (T \succ_{\alpha} U + T \prec_{\beta} U) \prec_{\gamma_1} W^l \right) \vee_{z, (\alpha\beta\gamma_1, \gamma_2)} W^r \\
&= (T \succ_{\alpha} U + T \prec_{\beta} U) \succ_{\alpha\beta} (W^l \vee_{z, (\gamma_1, \gamma_2)} W^r) \quad (\text{by Eq. (12)}) \\
&= (T \succ_{\alpha} U + T \prec_{\beta} U) \succ_{\alpha\beta} W.
\end{aligned}$$

This completes the proof of step 1.

Step 2: We show that $\text{DD}(X, \Omega)$ is generated by $\left\{ \bigvee_x \mid x \in X \right\}$. Let $T = T^l \vee_{x, (\alpha_1, \alpha_2)} T^r$ with $x \in X$ and $\alpha_1, \alpha_2 \in \Omega^1$. We employ induction on $\text{dep}(T) \geq 1$. For the initial step of $\text{dep}(T) = 1$, we have

$$T = \mid \vee_{x, (1, 1)} \mid = \bigvee_x \in \left\{ \bigvee_x \mid x \in X \right\}.$$

Assume that the conclusion holds for T with $\text{dep}(T) = k$ and consider the induction step of $\text{dep}(T) = k + 1 \geq 2$. Since T^l and T^r can not be \mid simultaneously, there exist three cases to consider.

Case 1: $T^l = \mid$ and $T^r \neq \mid$. Then $\alpha_1 = 1$ and

$$\begin{aligned}
T &= T^l \vee_{x, (\alpha_1, \alpha_2)} T^r = \mid \vee_{x, (1, \alpha_2)} T^r \\
&= \mid \vee_{x, (1, \alpha_2)} (\mid \prec_{\alpha_2} T^r + \mid \succ_1 T^r) \quad (\text{by Item (a) in Definition 3.7 and Eq. (13)}) \\
&= (\mid \vee_{x, (1, 1)} \mid) \prec_{\alpha_2} T^r \quad (\text{by Eq. (11)}) \\
&= \bigvee_x \prec_{\alpha_2} T^r.
\end{aligned}$$

By the induction hypothesis, T^r is generated by $\left\{ \bigvee_x \mid x \in X \right\}$ and so T is generated by

$$\left\{ \bigvee_x \mid x \in X \right\}.$$

Case 2: $T^l \neq \mid$ and $T^r = \mid$. This case is similar to Case 1.

Case 3: $T^l \neq \mid$ and $T^r \neq \mid$. Then

$$\begin{aligned}
T &= T^l \vee_{x, (\alpha_1, \alpha_2)} T^r \\
&= T^l \vee_{x, (\alpha_1, \alpha_2)} (\mid \prec_{\alpha_2} T^r + \mid \succ_1 T^r) \quad (\text{by Item (a) in Definition 3.7 and Eq. (13)}) \\
&= (T^l \vee_{x, (\alpha_1, 1)} \mid) \prec_{\alpha_2} T^r \quad (\text{by Eq. (11)}) \\
&= \left((T^l \succ_{\alpha_1} \mid + T^l \prec_1 \mid) \vee_{x, (\alpha_1, 1)} \mid \right) \prec_{\alpha_2} T^r \quad (\text{by Item (a) in Definition 3.7 and Eq. (13)}) \\
&= \left(T^l \succ_{\alpha_1} (\mid \vee_{x, (1, 1)} \mid) \right) \prec_{\alpha_2} T^r \quad (\text{by Eq. (12)}) \\
&= \left(T^l \succ_{\alpha_1} \bigvee_x \right) \prec_{\alpha_2} T^r.
\end{aligned}$$

$$\begin{aligned}
&= \bar{f}\left(| \vee_{x,(1,\omega)} \left(| \prec_{\omega} \begin{array}{c} \diagup y \\ \diagdown \end{array} + | \succ_1 \begin{array}{c} \diagdown y \\ \diagup \end{array} \right) \right) \quad (\text{by Eq. (11)}) \\
&= \bar{f}\left(| \vee_{x,(1,\omega)} \begin{array}{c} \diagup y \\ \diagdown \end{array} \right) \quad (\text{by Item (a) in Definition 3.7 and Eq. (13)}) \\
&= f(x) \prec'_{\omega} \bar{f}\left(\begin{array}{c} \diagup y \\ \diagdown \end{array}\right) \quad (\text{by Eq. (15)}) \\
&= \bar{f}\left(\begin{array}{c} \diagdown x \\ \diagup \end{array}\right) \prec'_{\omega} \bar{f}\left(\begin{array}{c} \diagup y \\ \diagdown \end{array}\right) \quad (\text{by Eq. (14)}) \\
&= \bar{f}(T) \prec'_{\omega} \bar{f}(U).
\end{aligned}$$

Assume that the conclusion holds for T and U in $\text{DD}(X, \Omega)$ with $\text{dep}(T) + \text{dep}(U) = k$ and consider the case when $\text{dep}(T) + \text{dep}(U) = k + 1 \geq 3$, there are four cases to consider.

Case 1: $T^l = |$ and $T^r \neq |$. Then $\alpha_1 = 1$ and

$$\begin{aligned}
\bar{f}(T \prec_{\omega} U) &= \bar{f}(| \vee_{x,(1,\alpha_2)} T^r) \prec_{\omega} U \\
&= \bar{f}(| \vee_{x,(1,\alpha_2\omega)} (T^r \prec_{\omega} U + T^r \succ_{\alpha_2} U)) \quad (\text{by Eq. (11)}) \\
&= f(x) \prec'_{\alpha_2\omega} \bar{f}(T^r \prec_{\omega} U + T^r \succ_{\alpha_2} U) \quad (\text{by Eq. (15)}) \\
&= f(x) \prec'_{\alpha_2\omega} (\bar{f}(T^r) \prec'_{\omega} \bar{f}(U) + \bar{f}(T^r) \succ'_{\alpha_2} \bar{f}(U)) \\
&\quad (\text{by the induction hypothesis on } \text{dep}(T) + \text{dep}(U)) \\
&= (f(x) \prec'_{\alpha_2} \bar{f}(T^r)) \prec'_{\omega} \bar{f}(U) \quad (\text{by Eq. (1)}) \\
&= \bar{f}(| \vee_{x,(1,\alpha_2)} T^r) \prec'_{\omega} \bar{f}(U) \quad (\text{by Eq. (15)}) \\
&= \bar{f}(T) \prec'_{\omega} \bar{f}(U).
\end{aligned}$$

Case 2: $T^l \neq |$ and $T^r = |$. This case is similar to Case 1.

Case 3: $T^l \neq |$ and $T^r \neq |$. Then

$$\begin{aligned}
\bar{f}(T \prec_{\omega} U) &= \bar{f}((T^l \vee_{x,(\alpha_1,\alpha_2)} T^r) \prec_{\omega} U) \\
&= \bar{f}(T^l \vee_{x,(\alpha_1,\alpha_2\omega)} (T^r \succ_{\alpha_2} U + T^r \prec_{\omega} U)) \quad (\text{by Eq. (11)}) \\
&= (\bar{f}(T^l) \succ'_{\alpha_1} f(x)) \prec'_{\alpha_2\omega} \bar{f}(T^r \succ_{\alpha_2} U + T^r \prec_{\omega} U) \quad (\text{by Eq. (15)}) \\
&= (\bar{f}(T^l) \succ'_{\alpha_1} f(x)) \prec'_{\alpha_2\omega} (\bar{f}(T^r) \succ'_{\alpha_2} \bar{f}(U) + \bar{f}(T^r) \prec'_{\omega} \bar{f}(U)) \\
&\quad (\text{by the induction hypothesis on } \text{dep}(T) + \text{dep}(U)) \\
&= ((\bar{f}(T^l) \succ'_{\alpha_1} f(x)) \prec'_{\alpha_2} \bar{f}(T^r)) \prec'_{\omega} \bar{f}(U) \quad (\text{by Eq. (1)}) \\
&= \bar{f}(T^l \vee_{x,(\alpha_1,\alpha_2)} T^r) \prec'_{\omega} \bar{f}(U) \quad (\text{by Eq. (15)}) \\
&= \bar{f}(T) \prec'_{\omega} \bar{f}(U).
\end{aligned}$$

Case 4: $T^l = |$ and $T^r = |$. Then $T = \begin{array}{c} \diagdown x \\ \diagup \end{array}$ for some $x \in X$ and we have



$$\begin{aligned}
\bar{f}(T \prec_{\omega} U) &= \bar{f}(| \vee_{x,(1,\omega)} U) \\
&= f(x) \prec'_{\omega} \bar{f}(U) \\
&= \bar{f}(T) \prec'_{\omega} \bar{f}(U)
\end{aligned}$$

according to (15).

This completes the proof. \square

4. FREE TRIDENDRIFORM FAMILY ALGEBRAS

In this section, we construct free tridendriform family algebras in terms of typed valently decorated Schröder trees. For this, let us first recall the construction of free tridendriform algebras. See [21, 25] for more details.

4.1. Free tridendriform algebras. Let X be a set. For $n \geq 0$, let $T_{n,X}$ be the set of planar rooted trees with $n+1$ leaves and with vertices valently decorated by elements of X , in the sense that if a vertex has valence k , then the vertex is decorated by an element in X^{k-1} . For example, the vertex of  is decorated by $x \in X$ while the vertex of  is decorated by $(x, y) \in X^2$. Here are the first few of them.

$$T_{0,X} = \{\emptyset\}, \quad T_{1,X} = \left\{ \begin{array}{c} \diagup \\ x \\ \diagdown \end{array} \mid x \in X \right\}, \quad T_{2,X} = \left\{ \begin{array}{c} \diagup \quad \diagdown \\ y \quad x \\ \diagdown \end{array}, \begin{array}{c} \diagup \quad \diagdown \\ x \quad y \\ \diagdown \end{array}, \begin{array}{c} \diagup \quad \diagdown \\ x \quad y \\ \diagdown \end{array} \mid x, y \in X \right\},$$

$$T_{3,X} = \left\{ \begin{array}{c} \diagup \quad \diagdown \\ z \quad y \\ \diagdown \quad x \end{array}, \begin{array}{c} \diagup \quad \diagdown \\ y \quad z \\ \diagdown \quad x \end{array}, \begin{array}{c} \diagup \quad \diagdown \\ y \quad z \\ \diagdown \quad x \end{array}, \begin{array}{c} \diagup \quad \diagdown \\ x \quad y \quad z \\ \diagdown \end{array}, \begin{array}{c} \diagup \quad \diagdown \\ y \quad z \\ \diagdown \quad x \end{array}, \begin{array}{c} \diagup \quad \diagdown \\ x \quad y \\ \diagdown \quad z \end{array}, \dots \mid x, y, z \in X \right\}.$$

For $T^{(i)} \in T_{n_i, X}$, $0 \leq i \leq k$, and $x_1, \dots, x_k \in X$, the grafting \bigvee of $T^{(i)}$ over (x_1, \dots, x_k) is

$$\bigvee_{x_1, \dots, x_k}^{k+1} (T^{(0)}, \dots, T^{(k)}).$$

Any planar rooted tree T can be uniquely expressed as such a grafting of lower depth trees. For example,

$$\begin{array}{c} \diagup \quad \diagdown \\ x \quad y \\ \diagdown \end{array} = \bigvee_{x,y} (|, |, |).$$

Let $\text{DT}(X) := \bigoplus_{n \geq 1} \mathbf{k}T_{n,X}$. Define binary operations \langle, \rangle and \cdot on $\text{DT}(X)$ recursively on $\text{dep}(T) + \text{dep}(U)$ by

(a) $| \rangle T := T \langle | := T$, $| \langle T := T \rangle | := 0$ and $| \cdot T := T \cdot | := 0$ for $T \in T_{n,X}$ with $n \geq 1$.

(b) For $\bigvee_{x_1, \dots, x_m}^{m+1} (T^{(0)}, \dots, T^{(m)})$ and $\bigvee_{y_1, \dots, y_n}^{n+1} (U^{(0)}, \dots, U^{(n)})$, define

$$T \langle U := \bigvee_{x_1, \dots, x_{m-1}, x_m}^{m+1} (T^{(0)}, \dots, T^{(m-1)}, T^{(m)} \rangle U + T^{(m)} \langle U + T^{(m)} \cdot U),$$

$$T \rangle U := \bigvee_{y_1, y_2, \dots, y_n}^{n+1} (T \rangle U^{(0)} + T \langle U^{(0)} + T \cdot U^{(0)}, U^{(1)}, \dots, U^{(n)}),$$

$$T \cdot U := \bigvee_{x_1, \dots, x_{m-1}, x_m, y_1, \dots, y_n}^{m+n+1} (T^{(0)}, \dots, T^{(m-1)}, T^{(m)} \rangle U^{(0)} + T^{(m)} \langle U^{(0)} + T^{(m)} \cdot U^{(0)}, U^{(1)}, \dots, U^{(n)}).$$

Let $j : X \rightarrow \text{DT}(X)$ be the map defined by $j(x) = \begin{array}{c} \diagup \\ x \\ \diagdown \end{array}$ for any $x \in X$. The following result is well-known.

Theorem 4.1. [21, 25] *Let X be a set. Then $(\text{DT}(X), \langle, \rangle, \cdot)$, together with j , is the free tridendriform algebra on X .*

4.2. Free tridendriform family algebras. In this subsection, we introduce typed valently decorated Schröder trees to construct free tridendriform family algebras. Let us first propose

Definition 4.2. Let X and Ω be two sets. A X -valently decorated Ω -typed (abbreviated typed valently decorated) Schröder tree is a triple $T = (T, \text{dec}, \text{type})$, where

(a) T is a Schröder tree.

- (b) $\text{dec} : IV(T) \rightarrow \sqcup_{n \geq 1} X^n$ is a map sending any vertex of arity $n + 1$ to X^n .
(c) $\text{type} : IE(T) \rightarrow \Omega$ is a map.

Remark 4.3. The trees in $\text{DT}(X)$ employed to construct free tridendriform algebras are indeed X -valently decorated $\{\star\}$ -typed Schröder trees, where $\{\star\}$ is any set with one single element.

Let X be a set and let Ω be a semigroup. For $n \geq 0$, let $T_n := T_{n,X,\Omega}$ be the set of X -valently decorated Ω^1 -typed Schröder trees with $n + 1$ leaves, such that the leaves are decorated by the identity 1 in Ω^1 and internal edges are decorated by elements of Ω . Let us expose some examples. Note that the edge decoration 1 is omitted for convenience.

$$T_0 = \{\emptyset\}, \quad T_1 = \left\{ \begin{array}{c} x \\ \diagdown \quad \diagup \\ \text{---} \\ \diagup \quad \diagdown \\ y \end{array} \mid x \in X \right\}, \quad T_2 = \left\{ \begin{array}{c} y \\ \diagdown \quad \diagup \\ \alpha \quad x \\ \diagup \quad \diagdown \\ y \end{array}, \begin{array}{c} y \\ \diagdown \quad \diagup \\ x \quad \alpha \\ \diagup \quad \diagdown \\ y \end{array}, \begin{array}{c} x|y \\ \diagdown \quad \diagup \\ \text{---} \\ \diagup \quad \diagdown \\ y \end{array} \mid x, y \in X, \alpha \in \Omega \right\},$$

$$T_3 = \left\{ \begin{array}{c} z \\ \diagdown \quad \diagup \\ \beta \quad y \\ \diagdown \quad \diagup \\ \alpha \quad x \\ \diagup \quad \diagdown \\ y \end{array}, \begin{array}{c} y \\ \diagdown \quad \diagup \\ \alpha \quad \beta \\ \diagdown \quad \diagup \\ x \end{array}, \begin{array}{c} y \\ \diagdown \quad \diagup \\ z \\ \diagdown \quad \diagup \\ \alpha \quad x \\ \diagup \quad \diagdown \\ y \end{array}, \begin{array}{c} x|y|z \\ \diagdown \quad \diagup \\ \text{---} \\ \diagup \quad \diagdown \\ y \end{array}, \begin{array}{c} z \\ \diagdown \quad \diagup \\ x \quad \alpha \\ \diagdown \quad \diagup \\ y \end{array}, \dots \mid x, y, z \in X, \alpha, \beta \in \Omega \right\}.$$

Denote by

$$\text{DT}(X, \Omega) := \bigoplus_{n \geq 1} \mathbf{k}T_n.$$

For typed valently decorated Schröder trees $T^{(i)} \in T_{n_i}$, $0 \leq i \leq k$, and $x_1, \dots, x_k \in X, \alpha_0, \dots, \alpha_k \in \Omega^1$, the grafting \bigvee of $T^{(i)}$ over (x_1, \dots, x_k) and $(\alpha_0, \dots, \alpha_k)$ is

$$(16) \quad T = \bigvee_{x_1, \dots, x_k}^{k+1; \alpha_0, \dots, \alpha_k} (T^{(0)}, \dots, T^{(k)})$$

obtained by joining the $k + 1$ roots of $T^{(i)}$ to a new vertex valently decorated by (x_1, \dots, x_k) and decorating the new edges from left to right by $\alpha_0, \dots, \alpha_k$ respectively.

Example 4.4. Let $T^{(0)} = \begin{array}{c} x \\ \diagdown \quad \diagup \\ \text{---} \\ \diagup \quad \diagdown \\ y \end{array}$, $T^{(1)} = \begin{array}{c} y \\ \diagdown \quad \diagup \\ \text{---} \\ \diagup \quad \diagdown \\ z \end{array}$, $T^{(2)} = \begin{array}{c} z \\ \diagdown \quad \diagup \\ \text{---} \\ \diagup \quad \diagdown \\ y \end{array}$, $u, v \in X$ and $\alpha, \beta, \gamma \in \Omega$. Then

$$\bigvee_{u,v}^{3; \alpha, \beta, \gamma} \left(\begin{array}{c} x \\ \diagdown \quad \diagup \\ \text{---} \\ \diagup \quad \diagdown \\ y \end{array}, \begin{array}{c} y \\ \diagdown \quad \diagup \\ \text{---} \\ \diagup \quad \diagdown \\ z \end{array}, \begin{array}{c} z \\ \diagdown \quad \diagup \\ \text{---} \\ \diagup \quad \diagdown \\ y \end{array} \right) = \begin{array}{c} x \quad y \quad z \\ \diagdown \quad \diagup \quad \diagup \\ \alpha \quad u \quad \beta \quad v \quad \gamma \\ \diagup \quad \diagdown \\ y \end{array}.$$

Any typed valently decorated Schröder tree $T \in \text{DT}(X, \Omega)$ can be uniquely expressed as such a grafting of lower depth typed valently decorated Schröder trees in Eq. (16), and we define the breadth of T as the arity of the internal vertex adjacent to the root, i.e. $\text{bre}(T) := k + 1$. For example,

$$\begin{array}{c} x|y \\ \diagdown \quad \diagup \\ \text{---} \\ \diagup \quad \diagdown \\ y \end{array} = \bigvee_{x,y}^{3; 1,1,1} (|, |, |); \quad \text{bre} \left(\begin{array}{c} x|y \\ \diagdown \quad \diagup \\ \text{---} \\ \diagup \quad \diagdown \\ y \end{array} \right) = 3, \quad \text{and}$$

$$T = \begin{array}{c} y \\ \diagdown \quad \diagup \\ x \quad \alpha \quad u \\ \diagdown \quad \diagup \\ z \quad \beta \end{array} = \bigvee_{x,z}^{3; 1, \alpha, \beta} (|, \begin{array}{c} y \\ \diagdown \quad \diagup \\ \text{---} \\ \diagup \quad \diagdown \\ z \end{array}, \begin{array}{c} u \\ \diagdown \quad \diagup \\ \text{---} \\ \diagup \quad \diagdown \\ y \end{array}), \quad \text{bre}(T) = 3.$$

Definition 4.5. Let X be a set and let Ω be a semigroup. Define binary operations $(\prec_\omega, \succ_\omega)_{\omega \in \Omega}$ and \cdot on $\text{DT}(X, \Omega)$ recursively on $\text{dep}(T) + \text{dep}(U)$ by

- (a) $| \succ_\omega T := T \prec_\omega | := T$, $| \prec_\omega T := T \succ_\omega | := 0$ and $| \cdot T := T \cdot | := 0$ for $\omega \in \Omega$ and $T \in T_n$ with $n \geq 1$.

(b) Let

$$T = \bigvee_{x_1, \dots, x_m}^{m+1; \alpha_0, \dots, \alpha_m} (T^{(0)}, \dots, T^{(m)}) \in T_m,$$

$$U = \bigvee_{y_1, \dots, y_n}^{n+1; \beta_0, \dots, \beta_n} (U^{(0)}, \dots, U^{(n)}) \in T_n,$$

define

(17)

$$T <_{\omega} U := \bigvee_{x_1, \dots, x_{m-1}, x_m}^{m+1; \alpha_0, \dots, \alpha_{m-1}, \alpha_m \omega} (T^{(0)}, \dots, T^{(m-1)}, T^{(m)}) >_{\alpha_m} U + T^{(m)} <_{\omega} U + T^{(m)} \cdot U,$$

(18)

$$T >_{\omega} U := \bigvee_{y_1, y_2, \dots, y_n}^{n+1; \omega \beta_0, \beta_1, \dots, \beta_n} (T >_{\omega} U^{(0)} + T <_{\beta_0} U^{(0)} + T \cdot U^{(0)}, U^{(1)}, \dots, U^{(n)}),$$

(19)

$$T \cdot U := \bigvee_{x_1, \dots, x_{m-1}, x_m, y_1, \dots, y_n}^{m+n+1; \alpha_0, \dots, \alpha_{m-1}, \alpha_m \beta_0, \beta_1, \dots, \beta_n} (T^{(0)}, \dots, T^{(m-1)}, T^{(m)}) >_{\alpha_m} U^{(0)} + T^{(m)} <_{\beta_0} U^{(0)} + T^{(m)} \cdot U^{(0)},$$

$$U^{(1)}, \dots, U^{(n)}).$$

Here in Eq. (19) when $T^{(m)} = | = U^{(0)}$, we employ the convention that

$$(20) \quad | <_1 | + | >_1 | + | \cdot | := |,$$

and

$$(21) \quad | >_1 T := T <_1 | := T \text{ and } | <_1 T := T >_1 | := 0.$$

Let us expose some examples for better understanding.

Example 4.6. Let $T = \begin{array}{c} x \\ \diagdown \quad \diagup \\ | \end{array}$ and $U = \begin{array}{c} y \\ \diagdown \quad \diagup \\ | \end{array}$ with $x, y \in X$. For $\omega \in \Omega$,

$$\begin{aligned} \begin{array}{c} x \\ \diagdown \quad \diagup \\ | \end{array} <_{\omega} \begin{array}{c} y \\ \diagdown \quad \diagup \\ | \end{array} &= \bigvee_x^{2; 1, 1} (|, |) <_{\omega} \begin{array}{c} y \\ \diagdown \quad \diagup \\ | \end{array} \\ &= \bigvee_x^{2; 1, \omega} (|, | <_{\omega} \begin{array}{c} y \\ \diagdown \quad \diagup \\ | \end{array} + | >_1 \begin{array}{c} y \\ \diagdown \quad \diagup \\ | \end{array} + | \cdot \begin{array}{c} y \\ \diagdown \quad \diagup \\ | \end{array}) \quad (\text{by Eq. (17)}) \\ &= \bigvee_x^{2; 1, \omega} (|, \begin{array}{c} y \\ \diagdown \quad \diagup \\ | \end{array}) \quad (\text{by Item (a) in Definition 4.5 and Eq. (21)}) \\ &= \begin{array}{c} x \quad y \\ \diagdown \quad \diagup \\ \omega | \end{array}, \end{aligned}$$

$$\begin{aligned} \begin{array}{c} x \\ \diagdown \quad \diagup \\ | \end{array} >_{\omega} \begin{array}{c} y \\ \diagdown \quad \diagup \\ | \end{array} &= \begin{array}{c} x \\ \diagdown \quad \diagup \\ | \end{array} >_{\omega} \bigvee_y^{2; 1, 1} (|, |) \\ &= \bigvee_y^{2; \omega, 1} (\begin{array}{c} x \\ \diagdown \quad \diagup \\ | \end{array} <_1 | + \begin{array}{c} x \\ \diagdown \quad \diagup \\ | \end{array} >_{\omega} | + \begin{array}{c} x \\ \diagdown \quad \diagup \\ | \end{array} \cdot |, |) \quad (\text{by Eq. (18)}) \\ &= \bigvee_y^{2; \omega, 1} (\begin{array}{c} x \\ \diagdown \quad \diagup \\ | \end{array}, |) \quad (\text{by Item (a) in Definition 4.5 and Eq. (21)}) \end{aligned}$$

$$\begin{aligned} &= \begin{array}{c} x \quad y \\ \diagdown \quad \diagup \\ \omega | \end{array}, \\ \begin{array}{c} x \\ \diagdown \quad \diagup \\ | \end{array} \cdot \begin{array}{c} y \\ \diagdown \quad \diagup \\ | \end{array} &= \bigvee_x^{2; 1, 1} (|, |) \cdot \bigvee_y^{2; 1, 1} (|, |) \\ &= \bigvee_{x, y}^{3; 1, 1, 1} (|, | <_1 | + | >_1 | + | \cdot |, |) \quad (\text{by Eq. (19)}) \end{aligned}$$

$$\begin{aligned}
&= \bigvee_{x,y}^{3;1,1,1}(|, |, |) \quad (\text{by Eq. (20)}) \\
&= \bigvee_{\alpha} \begin{array}{c} y \\ \diagup \quad \diagdown \\ x \\ \alpha \\ \bigvee \\ y \end{array}.
\end{aligned}$$

Example 4.7. Let $T = \bigvee_{\alpha} \begin{array}{c} y \\ \diagup \quad \diagdown \\ x \\ \alpha \\ \bigvee \\ y \end{array}$, $U = \bigvee_{\beta} \begin{array}{c} z \\ \diagup \quad \diagdown \\ \beta \\ \bigvee \\ z \end{array}$ with $x, y, z \in X$ and $\alpha \in \Omega$. For $\beta \in \Omega$,

$$\begin{aligned}
T \succ_{\beta} U &= \bigvee_{\alpha} \begin{array}{c} y \\ \diagup \quad \diagdown \\ x \\ \alpha \\ \bigvee \\ y \end{array} \succ_{\beta} \bigvee_z^{2;1,1}(|, |) = \bigvee_z^{2;\beta,1} \left(\bigvee_{\alpha} \begin{array}{c} y \\ \diagup \quad \diagdown \\ x \\ \alpha \\ \bigvee \\ y \end{array} \succ_{\beta} | + \bigvee_{\alpha} \begin{array}{c} y \\ \diagup \quad \diagdown \\ x \\ \alpha \\ \bigvee \\ y \end{array} \prec_1 | + \bigvee_{\alpha} \begin{array}{c} y \\ \diagup \quad \diagdown \\ x \\ \alpha \\ \bigvee \\ y \end{array} \cdot |, | \right) \\
&= \bigvee_z^{2;\beta,1} \left(\bigvee_{\alpha} \begin{array}{c} y \\ \diagup \quad \diagdown \\ x \\ \alpha \\ \bigvee \\ y \end{array}, | \right) = \bigvee_{\beta} \begin{array}{c} y \\ \diagup \quad \diagdown \\ x \\ \alpha \\ \diagdown \quad \diagup \\ z \\ \beta \\ \bigvee \\ z \end{array}. \\
T \prec_{\beta} U &= \bigvee_x^{2;\alpha,1} \left(\bigvee_y \begin{array}{c} y \\ \diagup \quad \diagdown \\ \alpha \\ \bigvee \\ y \end{array}, | \right) \prec_{\beta} \bigvee_z \begin{array}{c} z \\ \diagup \quad \diagdown \\ \beta \\ \bigvee \\ z \end{array} = \bigvee_x^{2;\alpha,\beta} \left(\bigvee_y \begin{array}{c} y \\ \diagup \quad \diagdown \\ \alpha \\ \bigvee \\ y \end{array}, | \prec_{\beta} \bigvee_z \begin{array}{c} z \\ \diagup \quad \diagdown \\ \beta \\ \bigvee \\ z \end{array} + | \succ_1 \bigvee_z \begin{array}{c} z \\ \diagup \quad \diagdown \\ \beta \\ \bigvee \\ z \end{array} + | \cdot \bigvee_z \begin{array}{c} z \\ \diagup \quad \diagdown \\ \beta \\ \bigvee \\ z \end{array} \right) \\
&= \bigvee_x^{2;\alpha,\beta} \left(\bigvee_y \begin{array}{c} y \\ \diagup \quad \diagdown \\ \alpha \\ \bigvee \\ y \end{array}, \bigvee_z \begin{array}{c} z \\ \diagup \quad \diagdown \\ \beta \\ \bigvee \\ z \end{array} \right) = \bigvee_{\alpha} \begin{array}{c} y \\ \diagup \quad \diagdown \\ x \\ \alpha \\ \diagdown \quad \diagup \\ z \\ \beta \\ \bigvee \\ z \end{array}. \\
U \cdot T &= \bigvee_z^{2;1,1}(|, |) \cdot \bigvee_x^{2;\alpha,1} \left(\bigvee_y \begin{array}{c} y \\ \diagup \quad \diagdown \\ \alpha \\ \bigvee \\ y \end{array}, | \right) = \bigvee_{z,x}^{3;1,\alpha,1} \left(|, | \succ_1 \bigvee_y \begin{array}{c} y \\ \diagup \quad \diagdown \\ \alpha \\ \bigvee \\ y \end{array} + | \prec_{\alpha} \bigvee_y \begin{array}{c} y \\ \diagup \quad \diagdown \\ \alpha \\ \bigvee \\ y \end{array} + | \cdot \bigvee_y \begin{array}{c} y \\ \diagup \quad \diagdown \\ \alpha \\ \bigvee \\ y \end{array}, | \right) \\
&= \bigvee_{z,x}^{3;1,\alpha,1} \left(|, \bigvee_y \begin{array}{c} y \\ \diagup \quad \diagdown \\ \alpha \\ \bigvee \\ y \end{array}, | \right) = \bigvee_{\alpha} \begin{array}{c} y \\ \diagup \quad \diagdown \\ z \\ \alpha \\ \diagdown \quad \diagup \\ x \\ \beta \\ \bigvee \\ z \end{array}.
\end{aligned}$$

Proposition 4.8. Let X be a set and let Ω be a semigroup. Then $(\text{DT}(X, \Omega), (\prec_{\omega}, \succ_{\omega})_{\omega \in \Omega}, \cdot)$ is a tridendriform family algebra, generated by $\left\{ \bigvee_x \begin{array}{c} x \\ \diagup \quad \diagdown \\ \alpha \\ \bigvee \\ x \end{array} \mid x \in X \right\}$.

Proof. We divide into two steps to prove the result.

Step 1: we first prove $(\text{DT}(X, \Omega), \{\prec_{\omega}, \succ_{\omega} \mid \omega \in \Omega\}, \cdot)$ is a tridendriform family algebra. Let

$$\begin{aligned}
T &= \bigvee_{x_1, \dots, x_m}^{m+1; \alpha_0, \dots, \alpha_m} (T^{(0)}, \dots, T^{(m)}), \\
U &= \bigvee_{y_1, \dots, y_n}^{n+1; \beta_0, \dots, \beta_n} (U^{(0)}, \dots, U^{(n)}) \text{ and} \\
W &= \bigvee_{z_1, \dots, z_{\ell}}^{\ell+1; \gamma_0, \dots, \gamma_{\ell}} (W^{(0)}, \dots, W^{(\ell)})
\end{aligned}$$

be in $\text{DT}(X, \Omega)$. Then we apply induction on $\text{dep}(T) + \text{dep}(U) + \text{dep}(W) \geq 3$. For the initial step of $\text{dep}(T) + \text{dep}(U) + \text{dep}(W) = 3$, we have $T = \bigvee_{x_1 \dots x_m} \begin{array}{c} x_1 \dots x_m \\ \diagup \quad \dots \quad \diagdown \\ \alpha \\ \bigvee \\ x_1 \dots x_m \end{array}$, $U = \bigvee_{y_1 \dots y_n} \begin{array}{c} y_1 \dots y_n \\ \diagup \quad \dots \quad \diagdown \\ \beta \\ \bigvee \\ y_1 \dots y_n \end{array}$ and $W = \bigvee_{z_1 \dots z_{\ell}} \begin{array}{c} z_1 \dots z_{\ell} \\ \diagup \quad \dots \quad \diagdown \\ \gamma \\ \bigvee \\ z_1 \dots z_{\ell} \end{array}$ for some $x_1, \dots, x_m, y_1, \dots, y_n, z_1, \dots, z_{\ell} \in X$. Then

$$\begin{aligned}
&(T \prec_{\beta} U) \prec_{\gamma} W \\
&= \left(\bigvee_{x_1 \dots x_m} \begin{array}{c} x_1 \dots x_m \\ \diagup \quad \dots \quad \diagdown \\ \alpha \\ \bigvee \\ x_1 \dots x_m \end{array} \prec_{\beta} \bigvee_{y_1 \dots y_n} \begin{array}{c} y_1 \dots y_n \\ \diagup \quad \dots \quad \diagdown \\ \beta \\ \bigvee \\ y_1 \dots y_n \end{array} \right) \prec_{\gamma} \bigvee_{z_1 \dots z_{\ell}} \begin{array}{c} z_1 \dots z_{\ell} \\ \diagup \quad \dots \quad \diagdown \\ \gamma \\ \bigvee \\ z_1 \dots z_{\ell} \end{array} \\
&= \left(\bigvee_{x_1, \dots, x_m}^{m+1; 1, \dots, 1} (|, \dots, |) \prec_{\beta} \bigvee_{y_1 \dots y_n} \begin{array}{c} y_1 \dots y_n \\ \diagup \quad \dots \quad \diagdown \\ \beta \\ \bigvee \\ y_1 \dots y_n \end{array} \right) \prec_{\gamma} \bigvee_{z_1 \dots z_{\ell}} \begin{array}{c} z_1 \dots z_{\ell} \\ \diagup \quad \dots \quad \diagdown \\ \gamma \\ \bigvee \\ z_1 \dots z_{\ell} \end{array}
\end{aligned}$$

$$\begin{aligned}
&= \bigvee_{x_1, \dots, x_{m-1}, x_m}^{m+1; \alpha_0, \dots, \alpha_{m-1}, \alpha_m \beta \gamma} \left(T^{(0)}, \dots, T^{(m-1)}, T^{(m)} \prec_{\beta \gamma} (U \prec_{\gamma} W + U \succ_{\beta} W + U \cdot W) + T^{(m)} \succ_{\alpha_m} (U \prec_{\gamma} W + U \succ_{\beta} W + U \cdot W) + T^{(m)} \cdot (U \prec_{\gamma} W + U \succ_{\beta} W + U \cdot W) \right) \\
&= \bigvee_{x_1, \dots, x_m}^{m+1; \alpha_0, \dots, \alpha_m} (T^{(0)}, \dots, T^{(m)}) \prec_{\beta \gamma} (U \prec_{\gamma} W + U \succ_{\beta} W + U \cdot W) \quad (\text{by Eq. (17)}) \\
&= T \prec_{\beta \gamma} (U \prec_{\gamma} W + U \succ_{\beta} W + U \cdot W).
\end{aligned}$$

The proofs of Eqs. (5)-(10) are similar and are moved to Appendix.

Step 2: We show that $\text{DT}(X, \Omega)$ is generated by $\left\{ \bigvee_x \mid x \in X \right\}$. We employ induction on

$\text{dep}(T) \geq 1$. For the initial step $\text{dep}(T) = 1$, we have $T = \bigvee_{x_1 \cdot \dots \cdot x_m} = \bigvee_{x_1} \cdot \bigvee_{x_2} \cdot \dots \cdot \bigvee_{x_m}$

where $\bigvee_{x_i} \in \left\{ \bigvee_x \mid x \in X \right\}$ with $1 \leq i \leq m$. For any induction step $\text{dep}(T) \geq 2$, we apply a secondary induction on $\text{bre}(T) \geq 2$.

For the initial step $\text{bre}(T) = 2$, since $\text{dep}(T) \geq 2$, we have $T \neq \bigvee_x$ for any $x \in X$ and there are three cases to consider.

Case 1: $T^{(0)} = |$ and $T^{(1)} \neq |$. Then $\alpha_0 = 1$ and

$$\begin{aligned}
T &= \bigvee_x^{2; \alpha_0, \alpha_1} (T^{(0)}, T^{(1)}) = \bigvee_x^{2; 1, \alpha_1} (|, T^{(1)}) \\
&= \bigvee_x^{2; 1, \alpha_1} (|, | \prec_{\alpha_1} T^{(1)} + | \succ_1 T^{(1)} + | \cdot T^{(1)}) \quad (\text{by Item (a) in Definition 4.5 and Eq. (21)}) \\
&= \bigvee_x^{2; 1, 1} (|, |) \prec_{\alpha_1} T^{(1)} \quad (\text{by Eq. (17)}) \\
&= \bigvee_x \prec_{\alpha_1} T^{(1)}.
\end{aligned}$$

By the induction hypothesis, $T^{(1)}$ is generated by $\left\{ \bigvee_x \mid x \in X \right\}$ and so T is generated by

$$\left\{ \bigvee_x \mid x \in X \right\}.$$

Case 2: $T^{(0)} \neq |$ and $T^{(1)} = |$. This case is similar to Case 1.

Case 3: $T^{(0)} \neq |$ and $T^{(1)} \neq |$. Then

$$\begin{aligned}
T &= \bigvee_x^{2; \alpha_0, \alpha_1} (T^{(0)}, T^{(1)}) \\
&= \bigvee_x^{2; \alpha_0, \alpha_1} (T^{(0)}, | \prec_{\alpha_1} T^{(1)} + | \succ_1 T^{(1)} + | \cdot T^{(1)}) \quad (\text{by Item (a) in Definition 4.5 and Eq. (21)}) \\
&= \bigvee_x^{2; \alpha_0, 1} (T^{(0)}, |) \prec_{\alpha_1} T^{(1)} \quad (\text{by Eq. (17)}) \\
&= \left(\bigvee_x^{2; \alpha_0, 1} (T^{(0)} \succ_{\alpha_0} | + T^{(0)} \prec_1 | + T^{(0)} \cdot |, |) \right) \prec_{\alpha_1} T^{(1)} \quad (\text{by Item (a) in Definition 4.5 and Eq. (21)}) \\
&= \left(T^{(0)} \succ_{\alpha_0} \bigvee_x^{2; 1, 1} (|, |) \right) \prec_{\alpha_1} T^{(1)} \quad (\text{by Eq. (18)}) \\
&= \left(T^{(0)} \succ_{\alpha_0} \bigvee_x \right) \prec_{\alpha_1} T^{(1)}.
\end{aligned}$$

By the induction hypothesis, $T^{(0)}$ and $T^{(1)}$ are generated by $\left\{ \begin{array}{c} x \\ \diagdown \quad \diagup \\ | \end{array} \middle| x \in X \right\}$ and so T is generated by $\left\{ \begin{array}{c} x \\ \diagdown \quad \diagup \\ | \end{array} \middle| x \in X \right\}$.

Now let us consider the case when $\text{dep}(T) = k + 1 \geq 2$ and $\text{bre}(T) = m + 1 \geq 3$. Then we can write $T = \bigvee_{x_1, \dots, x_m}^{m+1; \alpha_0, \dots, \alpha_m} (T^{(0)}, \dots, T^{(m)})$, there exist four cases to consider.

Case 4: $T^{(0)} = |$ and $T^{(1)} \neq |$. Then $\alpha_0 = 1$ and

$$\begin{aligned} T &= \bigvee_{x_1, x_2, \dots, x_m}^{m+1; 1, \alpha_1, \dots, \alpha_m} (|, T^{(1)}, \dots, T^{(m)}) \\ &= \left(\bigvee_{x_1}^{2; 1, \alpha_1} (|, T^{(1)}) \right) \cdot \left(\bigvee_{x_2}^{2; 1, \alpha_2} (|, T^{(2)}) \right) \cdot \dots \cdot \left(\bigvee_{x_m}^{2; 1, \alpha_m} (|, T^{(m)}) \right) \\ &\quad \text{(by Item (a) in Definition 4.5 and Eq. (19))} \\ &= \left(\begin{array}{c} x_1 \\ \diagdown \quad \diagup \\ <_{\alpha_1} \end{array} T^{(1)} \right) \cdot \left(\begin{array}{c} x_2 \\ \diagdown \quad \diagup \\ <_{\alpha_2} \end{array} T^{(2)} \right) \cdot \dots \cdot \left(\begin{array}{c} x_m \\ \diagdown \quad \diagup \\ <_{\alpha_m} \end{array} T^{(m)} \right). \quad \text{(by Case 1)} \end{aligned}$$

By the induction hypothesis, $T^{(i)}$ with $1 \leq i \leq m$ are generated by $\left\{ \begin{array}{c} x \\ \diagdown \quad \diagup \\ | \end{array} \middle| x \in X \right\}$ and so T is generated by $\left\{ \begin{array}{c} x \\ \diagdown \quad \diagup \\ | \end{array} \middle| x \in X \right\}$.

Case 5: $T^{(0)} \neq |$ and $T^{(1)} = |$. This case is similar to Case 4.

Case 6: $T^{(0)} \neq |$ and $T^{(1)} \neq |$. Then

$$\begin{aligned} T &= \bigvee_{x_1, x_2, \dots, x_m}^{m+1; \alpha_0, \alpha_1, \dots, \alpha_m} (T^{(0)}, T^{(1)}, \dots, T^{(m)}) \\ &= \left(\bigvee_{x_1}^{2; \alpha_0, \alpha_1} (T^{(0)}, T^{(1)}) \right) \cdot \left(\bigvee_{x_2}^{2; 1, \alpha_2} (|, T^{(2)}) \right) \cdot \dots \cdot \left(\bigvee_{x_m}^{2; 1, \alpha_m} (|, T^{(m)}) \right) \\ &\quad \text{(by Item (a) in Definition 4.5 and Eq. (19))} \\ &= \left(\left(T^{(0)} \right) >_{\alpha_0} \begin{array}{c} x_1 \\ \diagdown \quad \diagup \\ <_{\alpha_1} \end{array} T^{(1)} \right) \cdot \left(\begin{array}{c} x_2 \\ \diagdown \quad \diagup \\ <_{\alpha_2} \end{array} T^{(2)} \right) \cdot \dots \cdot \left(\begin{array}{c} x_m \\ \diagdown \quad \diagup \\ <_{\alpha_m} \end{array} T^{(m)} \right). \quad \text{(by Case 1 and Case 3)} \end{aligned}$$

By the induction hypothesis, $T^{(i)}$ with $1 \leq i \leq m$ are generated by $\left\{ \begin{array}{c} x \\ \diagdown \quad \diagup \\ | \end{array} \middle| x \in X \right\}$ and so T is generated by $\left\{ \begin{array}{c} x \\ \diagdown \quad \diagup \\ | \end{array} \middle| x \in X \right\}$.

Case 7: $T^{(0)} = |$ and $T^{(1)} = |$. Then

$$\begin{aligned} T &= \bigvee_{x_1, x_2, x_3, \dots, x_m}^{m+1; 1, 1, \alpha_2, \dots, \alpha_m} (|, |, T^{(2)}, \dots, T^{(m)}) \\ &= \left(\bigvee_{x_1}^{2; 1, 1} (|, |) \right) \cdot \left(\bigvee_{x_2}^{2; 1, \alpha_2} (|, T^{(2)}) \right) \cdot \dots \cdot \left(\bigvee_{x_m}^{2; 1, \alpha_m} (|, T^{(m)}) \right) \\ &\quad \text{(by Item (a) in Definition 4.5 and Eq. (19))} \\ &= \begin{array}{c} x_1 \\ \diagdown \quad \diagup \\ <_{\alpha_1} \end{array} \cdot \left(\begin{array}{c} x_2 \\ \diagdown \quad \diagup \\ <_{\alpha_2} \end{array} T^{(2)} \right) \cdot \dots \cdot \left(\begin{array}{c} x_m \\ \diagdown \quad \diagup \\ <_{\alpha_m} \end{array} T^{(m)} \right). \quad \text{(by Case 1)} \end{aligned}$$

By the induction hypothesis, $T^{(i)}$ with $1 \leq i \leq m$ are generated by $\left\{ \begin{array}{c} x \\ \diagdown \quad \diagup \\ | \end{array} \middle| x \in X \right\}$ and so is T . This completes the proof. \square

The concept of free tridendriform family algebra is given as usual.

Definition 4.9. Let X be a set and let Ω be a semigroup. A free tridendriform family algebra on X is a tridendriform family algebra $(T, (<_{\omega}, >_{\omega})_{\omega \in \Omega}, \cdot)$ together with the map $j : X \rightarrow T$ that satisfies

the following universal property: for any tridendriform family algebra $(T', (\langle'_{\omega}, \succ'_{\omega})_{\omega \in \Omega}, \cdot')$ and map $f : X \rightarrow T'$, there is a unique tridendriform family algebra morphism $\bar{f} : T \rightarrow T'$ such that $f = \bar{f} \circ j$. The free tridendriform family algebra is unique up to isomorphism.

Let $j : X \rightarrow \text{DT}(X, \Omega)$ be the map defined by $j(x) = \bigvee_x$ for $x \in X$. Now we arrive at our main result in this section.

Theorem 4.10. *Let X be a set and let Ω be a semigroup. The tridendriform family algebra $(\text{DT}(X, \Omega), (\langle_{\omega}, \succ_{\omega})_{\omega \in \Omega}, \cdot)$, together with the map j , is the free tridendriform family algebra on X .*

Proof. By Proposition 4.8, we are left to prove $(\text{DT}(X, \Omega), (\langle_{\omega}, \succ_{\omega})_{\omega \in \Omega}, \cdot)$ satisfies the universal property. For this, let $(T', (\langle'_{\omega}, \succ'_{\omega})_{\omega \in \Omega}, \cdot')$ be a tridendriform family algebra. First of all, if there exists a tridendriform family algebra morphism $\bar{f} : \text{DT}(X, \Omega) \rightarrow T'$ extending $f : X \rightarrow T'$ in the sense that $\bar{f}(\bigvee_x) = f(x)$ for any $x \in X$, then such an \bar{f} is unique, due to the fact that the collection of trees $(\bigvee_x)_{x \in X}$ generate the tridendriform family algebra $\text{DT}(X, \Omega)$. Now we define a linear map

$$\bar{f} : \text{DT}(X, \Omega) \rightarrow T', \quad T \mapsto \bar{f}(T)$$

by induction on $\text{dep}(T) \geq 1$ and prove that it is a tridendriform family algebra morphism. We use the unique decomposition

$$(22) \quad \begin{aligned} T &= \bigvee_{x_1, \dots, x_m}^{m+1; \alpha_0, \dots, \alpha_m} (T^{(0)}, \dots, T^{(m)}) \\ &= \bigvee_{x_1}^{2; \alpha_0, \alpha_1} (T^{(0)}, T^{(1)}) \cdot \bigvee_{x_2}^{2; 1, \alpha_2} (|, T^{(2)}) \cdot \dots \cdot \bigvee_{x_m}^{2; 1, \alpha_m} (|, T^{(m)}), \end{aligned}$$

together with:

$$\begin{aligned} \bigvee_x^{2; \alpha_0, \alpha_1} (V, W) &= \bigvee_x \text{ if } V = W = |, \\ &= V \succ_{\alpha_0} \bigvee_x \text{ if } V \neq | \text{ and } W = |, \\ &= \bigvee_x \langle_{\alpha_1} W \text{ if } V = | \text{ and } W \neq |, \\ &= V \succ_{\alpha_0} \bigvee_x \langle_{\alpha_1} W \text{ if } V \neq | \text{ and } W \neq |, \end{aligned}$$

which uniquely defines $\bar{f}(T)$ in terms of $\bar{f}(T^{(0)}), \dots, \bar{f}(T^{(m)})$ and the operations $((\langle'_{\omega}, \succ'_{\omega})_{\omega \in \Omega}, \cdot')$. We are going to prove that \bar{f} is a morphism of tridendriform family algebras:

$$\bar{f}(T \langle_{\omega} U) = \bar{f}(T) \langle'_{\omega} \bar{f}(U), \quad \bar{f}(T \succ_{\omega} U) = \bar{f}(T) \succ'_{\omega} \bar{f}(U) \text{ and } \bar{f}(T \cdot U) = \bar{f}(T) \cdot' \bar{f}(U),$$

in which we only prove the first equation, as the proofs of the two other are similar. Starting from

$$T = \bigvee_{x_1, \dots, x_m}^{m+1; \alpha_0, \dots, \alpha_m} (T^{(0)}, \dots, T^{(m)}), \quad U = \bigvee_{y_1, \dots, y_n}^{n+1; \beta_0, \dots, \beta_n} (U^{(0)}, \dots, U^{(n)})$$

we compute:

$$\begin{aligned} &\bar{f}(T \langle_{\omega} U) \\ &= \bar{f} \left(\bigvee_{x_1, \dots, x_m}^{m+1; \alpha_0, \dots, \alpha_{m-1}, \alpha_m \omega} (T^{(0)}, \dots, T^{(m-1)}, T^{(m)} \succ_{\alpha_m} U + T^{(m)} \langle_{\omega} U + T^{(m)} \cdot U) \right) \\ &= \bar{f} \left(\bigvee_{x_1}^{2; \alpha_0, \alpha_1} (T^{(0)}, T^{(1)}) \cdot \bigvee_{x_2}^{2; 1, \alpha_2} (|, T^{(2)}) \cdot \dots \cdot \bigvee_{x_m}^{2; 1, \alpha_m \omega} (|, T^{(m)} \succ_{\alpha_m} U + T^{(m)} \langle_{\omega} U + T^{(m)} \cdot U) \right) \end{aligned}$$

$$\begin{aligned}
& \text{(from (22))} \\
& = \bar{f}\left(\bigvee_{x_1}^{2; \alpha_0, \alpha_1}(T^{(0)}, T^{(1)})\right) \cdot' \dots \cdot' \bar{f}\left(\bigvee_{x_m}^{x_m} \prec_{\alpha_m \omega} (T^{(m)} \succ_{\alpha_m} U + T^{(m)} \prec_{\omega} U + T^{(m)} \cdot U)\right) \\
& \text{(from the definition of } \bar{f}\text{)}
\end{aligned}$$

(23)

$$= \bar{f}\left(\bigvee_{x_1}^{2; \alpha_0, \alpha_1}(T^{(0)}, T^{(1)})\right) \cdot' \dots \cdot' \bar{f}\left(\left(\bigvee_{x_m}^{x_m} \prec_{\alpha_m} T^{(m)}\right) \prec_{\omega} U\right) \text{ (from Axiom (4)).}$$

Now we proceed by induction on $\text{dep}(T) + \text{dep}(U)$, together with a secondary induction on $\text{bre}(T)$. If $\text{dep}(T) + \text{dep}(U) = 2$ (main initial step), then $T^{(0)} = \dots = T^{(m)} = |$, and Equation (23) boils down to

$$\begin{aligned}
\bar{f}(T \prec_{\omega} U) &= \bar{f}\left(\bigvee_{x_1}^{x_1}\right) \cdot' \dots \cdot' \bar{f}\left(\bigvee_{x_{m-1}}^{x_{m-1}}\right) \cdot' \bar{f}\left(\bigvee_{x_m}^{x_m} \prec_{\omega} U\right) \\
&= \bar{f}\left(\bigvee_{x_1}^{x_1}\right) \cdot' \dots \cdot' \bar{f}\left(\bigvee_{x_{m-1}}^{x_{m-1}}\right) \cdot' \left(\bar{f}\left(\bigvee_{x_m}^{x_m}\right) \prec'_{\omega} \bar{f}(U)\right) \text{ (from the definition of } \bar{f}\text{)} \\
&= \left(\bar{f}\left(\bigvee_{x_1}^{x_1}\right) \cdot' \dots \cdot' \bar{f}\left(\bigvee_{x_{m-1}}^{x_{m-1}}\right) \cdot' \bar{f}\left(\bigvee_{x_m}^{x_m}\right)\right) \prec'_{\omega} \bar{f}(U) \text{ (from an iteration of Axiom (9))} \\
&= \bar{f}(T) \prec'_{\omega} \bar{f}(U).
\end{aligned}$$

If $\text{dep}(T) + \text{dep}(U) \geq 3$ and $\text{bre}(T) = 2$ (secondary initial step), then we have

$$\begin{aligned}
\bar{f}(T \prec_{\omega} U) &= \bar{f}\left(\bigvee_{x_1}^{2; \alpha_0, \alpha_1 \omega}(T^{(0)}, T^{(1)} \succ_{\alpha_1} U + T^{(1)} \prec_{\omega} U + T^{(1)} \cdot U)\right) \\
&= \bar{f}\left(T^{(0)} \succ_{\alpha_0} \bigvee_{x_1}^{x_1} \prec_{\alpha_1 \omega} (T^{(1)} \succ_{\alpha_1} U + T^{(1)} \prec_{\omega} U + T^{(1)} \cdot U)\right) \\
&= \bar{f}(T^{(0)}) \succ'_{\alpha_0} \bar{f}\left(\bigvee_{x_1}^{x_1}\right) \prec'_{\alpha_1 \omega} \bar{f}(T^{(1)} \succ_{\alpha_1} U + T^{(1)} \prec_{\omega} U + T^{(1)} \cdot U) \text{ (by definition of } \bar{f}\text{)} \\
&= \bar{f}(T^{(0)}) \succ'_{\alpha_0} \bar{f}\left(\bigvee_{x_1}^{x_1}\right) \prec'_{\alpha_1 \omega} \left(\bar{f}(T^{(1)}) \succ'_{\alpha_1} \bar{f}(U) + \bar{f}(T^{(1)}) \prec'_{\omega} \bar{f}(U) + \bar{f}(T^{(1)}) \cdot' \bar{f}(U)\right) \\
& \text{(by the main induction hypothesis)} \\
&= \bar{f}(T^{(0)}) \succ'_{\alpha_0} \left(\left(\bar{f}\left(\bigvee_{x_1}^{x_1}\right) \prec'_{\alpha_1} \bar{f}(T^{(1)})\right) \prec'_{\omega} \bar{f}(U)\right) \\
&= \left(\bar{f}(T^{(0)}) \succ'_{\alpha_0} \left(\bar{f}\left(\bigvee_{x_1}^{x_1}\right) \prec'_{\alpha_1} \bar{f}(T^{(1)})\right)\right) \prec'_{\omega} \bar{f}(U) \\
&= \bar{f}(T) \prec'_{\omega} \bar{f}(U) \text{ (again by definition of } \bar{f}\text{)}.
\end{aligned}$$

Finally, if $\text{dep}(T) + \text{dep}(U) \geq 3$ and $\text{bre}(T) \geq 3$, we can push further compilation (23) by using the secondary initial step for $\bigvee_{x_m}^{x_m} \prec_{\alpha_m} T^{(m)}$, which is of breadth two:

$$\begin{aligned}
\bar{f}(T \prec_{\omega} U) &= \bar{f}\left(\bigvee_{x_1}^{2; \alpha_0, \alpha_1}(T^{(0)}, T^{(1)})\right) \cdot' \dots \cdot' \left(\bar{f}\left(\bigvee_{x_m}^{x_m} \prec_{\alpha_m} T^{(m)}\right) \prec'_{\omega} \bar{f}(U)\right) \\
&= \left(\bar{f}\left(\bigvee_{x_1}^{2; \alpha_0, \alpha_1}(T^{(0)}, T^{(1)})\right) \cdot' \dots \cdot' \bar{f}\left(\bigvee_{x_m}^{x_m} \prec_{\alpha_m} T^{(m)}\right)\right) \prec'_{\omega} \bar{f}(U) \\
& \text{(by iterating Axiom (9))} \\
&= \bar{f}\left(\left(\bigvee_{x_1}^{2; \alpha_0, \alpha_1}(T^{(0)}, T^{(1)})\right) \cdot' \dots \cdot' \left(\bigvee_{x_m}^{x_m} \prec_{\alpha_m} T^{(m)}\right)\right) \prec'_{\omega} \bar{f}(U) \\
& \text{(by definition of } \bar{f}\text{)} \\
&= \bar{f}(T) \prec'_{\omega} \bar{f}(U).
\end{aligned}$$

The verification of the two identities $\bar{f}(T \succ_{\omega} U) = \bar{f}(T) \succ'_{\omega} \bar{f}(U)$ and $\bar{f}(T \cdot U) = \bar{f}(T) \cdot' \bar{f}(U)$ is similar and left to the reader. \square

Remark 4.11. We can deduce Theorem 3.12 from Theorem 4.10, as dendriform family algebras are tridendriform family algebras with a zero product \cdot . Indeed, the free dendriform family algebra generated by X is the quotient of $\text{DT}(X, \Omega)$ by the tridendriform family ideal I generated by $\text{DT}(X, \Omega) \cdot \text{DT}(X, \Omega)$. It is not difficult to show that I is linearly generated by all Ω -typed Schröder trees which are not binary, so the quotient is isomorphism to $\text{DD}(X, \Omega)$.

Acknowledgments: This work was supported by the National Natural Science Foundation of China (Grant No. 11771191 and 11861051). We thank the Referee for careful reading and useful suggestions which helped us to improve the presentation.

REFERENCES

- [1] G. Baxter, An analytic problem whose solution follows from a simple algebraic identity, *Pacific J. Math.* **10** (1960), 731-742. [3](#)
- [2] Y. Bruned, M. Hairer and L. Zambotti, Algebraic renormalisation of regularity structures, *Invent. math.* **215** (2019), 1039-1156. [2](#), [9](#)
- [3] F. Bergeron, G. Labelle and P. Leroux, Combinatorial species and tree-like structures, Encyclopedia of Mathematics and its Application, vol. 67, Cambridge University Press, Cambridge, 1998, Translated from the 1994 French original by Margaret Readdy, With a foreword by Gian-Carlo Rota. [2](#)
- [4] A. Connes and D. Kreimer, Hopf algebras, renormalization and non-commutative geometry, *Commun. Math. Phys.* **199** (1998), 203-242. [2](#)
- [5] A. Connes and D. Kreimer, Renormalization in quantum field theory and the Riemann-Hilbert problem. I. The Hopf algebra structure of graphs and the main theorem, *Commun. Math. Phys.* **210** (2000), 249-273. [2](#)
- [6] A. Connes and D. Kreimer, Renormalization in quantum field theory and the Riemann-Hilbert problem. II. The β -function, diffeomorphisms and the renormalization group, *Comm. Math. Phys.* **216** (2001), 215-241. [2](#)
- [7] K. Ebrahimi-Fard, J. Gracia-Bondia and F. Patras, A Lie theoretic approach to renormalization, *Comm. Math. Phys.* **276** (2007), 519-549. [1](#), [3](#)
- [8] K. Ebrahimi-Fard, D. Manchon and F. Patras, New identities in dendriform algebras, *J. Algebra* **320** (2008), 708-727. [2](#)
- [9] A. Frabetti, Leibniz homology of dialgebras of matrices, *J. Pure. Appl. Algebra* **129** (1998), 123-141. [1](#)
- [10] L. Foissy, Les algèbres de Hopf des arbres enracinés décorés II, *Bull. Sci. Math.* **126** (2002), 249-288. [2](#)
- [11] L. Foissy, Algebraic structures on typed decorated rooted trees, arXiv: 1811.07572v1 [math.RA] 19 Nov 2018. [9](#)
- [12] X. Gao, L. Guo and M. Rosenkranz, Free intergro-differential algebras and Gröbner-Shirshov basis, *J. Algebra* **442** (2015), 354-396. [2](#)
- [13] X. Gao, L. Guo and Y. Zhang, Matching Rota-Baxter algebras, matching dendriform algebras and matching pre-Lie algebras, arxiv:1909.10577 (2019). [2](#)
- [14] X. Gao, L. Guo and S. H. Zheng, Construction of free commutative integro-differential algebras by the method of Gröbner-Shirshov basis, *J. Algebra Appl.* **5** (2014), 1350160. [2](#)
- [15] L. Guo and W. Keigher, On free Baxter algebras: completions and the internal construction, *Adv. Math.* **151** (2000), 101-127. [2](#)
- [16] L. Guo and W. Keigher, On differential Rota-Baxter algebras, *J. Pure Appl. Algebra* **212** (2008), 522-540. [2](#)
- [17] L. Guo, Operated monoids, Motzkin paths and rooted trees, *J. Algebraic Combin.* **29** (2009), 35-62. [3](#)
- [18] L. Guo, An introduction to Rota-Baxter algebra, International Press, 2012. [1](#)
- [19] D. Kreimer and Erik Panzer, Hopf-algebraic renormalization of Kreimer's toy model, Master thesis, Handbook, <https://arxiv.org/abs/1202.3552>. [1](#), [3](#)
- [20] J. -L. Loday, Une version non commutative des algèbre de Lie: les algèbres de Leibniz, *Ens. Math.* **39** (1993), 269-293. [1](#), [2](#), [4](#), [8](#)
- [21] J. -L. Loday and M. R. Ronco, Hopf algebra of the planar binary trees, *Adv. Math.* **39** (1998), 293-309. [1](#), [2](#), [8](#), [17](#)

- [22] J. -L. Loday, Dialgebra, in Dialgebras and related operad, *Lecture Notes in Math.* **1763** (2002), 7-66. [8](#)
- [23] J. -L. Loday, Arithmetree, *J. Algebra* **258** (2002), 275-309. [1](#)
- [24] J. -L. Loday, Scindement d'associativité et algèbres de Hopf, the Proceedings of the Conference in honor of Jean Leray, Nantes, (2002), *Séminaire et Congrès(SMF)* **9** (2004), 155-172. [1](#)
- [25] J. -L. Loday and M. O. Ronco, Trialgebras and families of polytopes, in "Homotopy theory: relations with algebraic geometry, group cohomology, and algebraic K-theory", *Contemp. Math.* **346** (2004), 369-398. [1, 2, 4, 17](#)
- [26] M. R. Ronco, Eulerian idempotents and Milnor-Moore theorem for certain non-commutative Hopf algebras, *J. Algebra* **254** (2002), 152-172. [1, 8](#)
- [27] Richard J. Mathar, Topologically distinct sets of non-intersecting circles in the plane, arXiv:1603.00077,2016. [2](#)
- [28] Y. Y. Zhang and X. Gao, Free Rota-Baxter family algebras and (tri)dendriform family algebras, *Pacific J. Math.* **301** (2019), 741-766. [1, 2, 4, 5](#)

5. APPENDIX

The remaining proofs of Step 1 in Proposition [4.8](#).

$$\begin{aligned}
& (T \succ_{\alpha} U) \prec_{\gamma} W \\
&= \bigvee_{y_1, y_2, \dots, y_n}^{n+1; \alpha\beta_0, \beta_1, \dots, \beta_n} (T \succ_{\alpha} U^{(0)} + T \prec_{\beta_0} U^{(0)} + T \cdot U^{(0)}, U^{(1)}, \dots, U^{(n)}) \prec_{\gamma} W \\
&= \bigvee_{y_1, y_2, \dots, y_{n-1}, y_n}^{n+1; \alpha\beta_0, \beta_1, \dots, \beta_{n-1}, \beta_n \gamma} (T \succ_{\alpha} U^{(0)} + T \prec_{\beta_0} U^{(0)} + T \cdot U^{(0)}, U^{(1)}, \dots, U^{(n-1)}, \\
&\quad U^{(n)} \succ_{\beta_n} W + U^{(n)} \prec_{\gamma} W + U^{(n)} \cdot W) \\
&= T \succ_{\alpha} \left(\bigvee_{y_1, \dots, y_{n-1}, y_n}^{n+1; \beta_0, \dots, \beta_{n-1}, \beta_n \gamma} (U^{(0)}, \dots, U^{(n-1)}, U^{(n)} \succ_{\beta_n} W + U^{(n)} \prec_{\gamma} W + U^{(n)} \cdot W) \right) \\
&= T \succ_{\alpha} (U \prec_{\gamma} W).
\end{aligned}$$

$$\begin{aligned}
& T \succ_{\alpha} (U \succ_{\beta} W) \\
&= T \succ_{\alpha} \left(\bigvee_{z_1, z_2, \dots, z_{\ell}}^{\ell+1; \beta\gamma_0, \gamma_1, \dots, \gamma_{\ell}} (U \succ_{\beta} W^{(0)} + U \prec_{\gamma_0} W^{(0)} + U \cdot W^{(0)}, W^{(1)}, \dots, W^{(\ell)}) \right) \\
&= \bigvee_{z_1, z_2, \dots, z_{\ell}}^{\ell+1; \alpha\beta\gamma_0, \gamma_1, \dots, \gamma_{\ell}} \left(T \succ_{\alpha} (U \succ_{\beta} W^{(0)} + U \prec_{\gamma_0} W^{(0)} + U \cdot W^{(0)}) + T \prec_{\beta\gamma_0} (U \succ_{\beta} W^{(0)} + U \prec_{\gamma_0} W^{(0)} \right. \\
&\quad \left. + U \cdot W^{(0)}) + T \cdot (U \succ_{\beta} W^{(0)} + U \prec_{\gamma_0} W^{(0)} + U \cdot W^{(0)}), W^{(1)}, \dots, W^{(\ell)} \right) \\
&= \bigvee_{z_1, z_2, \dots, z_{\ell}}^{\ell+1; \alpha\beta\gamma_0, \gamma_1, \dots, \gamma_{\ell}} \left(T \succ_{\alpha} (U \succ_{\beta} W^{(0)}) + T \succ_{\alpha} (U \prec_{\gamma_0} W^{(0)}) + T \succ_{\alpha} (U \cdot W^{(0)}) + (T \prec_{\beta} U) \prec_{\gamma_0} W^{(0)} \right. \\
&\quad \left. + T \cdot (U \succ_{\beta} W^{(0)}) + T \cdot (U \prec_{\gamma_0} W^{(0)}) + T \cdot (U \cdot W^{(0)}), W^{(1)}, \dots, W^{(\ell)} \right) \\
&= \bigvee_{z_1, z_2, \dots, z_{\ell}}^{\ell+1; \alpha\beta\gamma_0, \gamma_1, \dots, \gamma_{\ell}} \left((T \succ_{\alpha} U + T \prec_{\beta} U + T \cdot U) \succ_{\alpha\beta} W^{(0)} + (T \succ_{\alpha} U + T \prec_{\beta} U + T \cdot U) \prec_{\gamma_0} W^{(0)} \right. \\
&\quad \left. + (T \succ_{\alpha} U + T \prec_{\beta} U + T \cdot U) \cdot W^{(0)}, W^{(1)}, \dots, W^{(\ell)} \right) \\
&= (T \succ_{\alpha} U + T \prec_{\beta} U + T \cdot U) \succ_{\alpha\beta} \left(\bigvee_{z_1, \dots, z_{\ell}}^{\ell+1; \gamma_0, \dots, \gamma_{\ell}} (W^{(0)}, \dots, W^{(\ell)}) \right) \\
&= (T \succ_{\alpha} U + T \prec_{\beta} U + T \cdot U) \succ_{\alpha\beta} W.
\end{aligned}$$

$$\begin{aligned}
& (T \succ_{\alpha} U) \cdot W \\
&= \bigvee_{y_1, y_2, \dots, y_n}^{n+1; \alpha\beta_0, \beta_1, \dots, \beta_n} (T \succ_{\alpha} U^{(0)} + T \prec_{\beta_0} U^{(0)} + T \cdot U^{(0)}, U^{(1)}, \dots, U^{(n)}) \cdot \bigvee_{z_1, \dots, z_{\ell}}^{\ell+1; \gamma_0, \dots, \gamma_{\ell}} (W^{(0)}, \dots, W^{(\ell)})
\end{aligned}$$

$$\begin{aligned}
&= \bigvee_{y_1, y_2, \dots, y_n, z_1, z_2, \dots, z_\ell}^{n+1+\ell; \alpha\beta_0, \beta_1, \dots, \beta_{n-1}, \beta_n \gamma_0, \gamma_1, \dots, \gamma_\ell} \left(T \succ_\alpha U^{(0)} + T \prec_{\beta_0} U^{(0)} + T \cdot U^{(0)}, U^{(1)}, \dots, U^{(n-1)}, \right. \\
&\quad \left. U^{(n)} \succ_{\beta_n} W^{(0)} + U^{(n)} \prec_{\gamma_0} W^{(0)} + U^{(n)} \cdot W^{(0)}, W^{(1)}, \dots, W^{(l)} \right) \\
&= T \succ_\alpha \bigvee_{y_1, \dots, y_{n-1}, y_n, z_1, \dots, z_\ell}^{n+1+\ell; \beta_0, \dots, \beta_{n-1}, \beta_n \gamma_0, z_1, \dots, z_\ell} \left(U^{(0)}, \dots, U^{(n-1)}, U^{(n)} \succ_{\beta_n} W^{(0)} + U^{(n)} \prec_{\gamma_0} W^{(0)} + U^{(n)} \cdot W^{(0)}, \right. \\
&\quad \left. W^{(1)}, \dots, W^{(l)} \right) \\
&= T \succ_\alpha (U \cdot W).
\end{aligned}$$

$$\begin{aligned}
&(T \prec_\beta U) \cdot W \\
&= \bigvee_{x_1, \dots, x_{m-1}, x_m}^{m+1; \alpha_0, \dots, \alpha_{m-1}, \alpha_m \beta} \left(T^{(0)}, \dots, T^{(m-1)}, T^{(m)} \succ_{\alpha_m} U + T^{(m)} \prec_\beta U + T^{(m)} \cdot U \right) \\
&\quad \cdot \bigvee_{z_1, \dots, z_\ell}^{\ell+1; \gamma_0, \dots, \gamma_\ell} (W^{(0)}, \dots, W^{(l)}) \\
&= \bigvee_{x_1, \dots, x_{m-1}, x_m, z_1, \dots, z_\ell}^{m+1+\ell; \alpha_0, \dots, \alpha_{m-1}, \alpha_m \beta \gamma_0, \gamma_1, \dots, \gamma_\ell} \left(T^{(0)}, \dots, T^{(m-1)}, T^{(m)} \succ_{\alpha_m} U + T^{(m)} \prec_\beta U + T^{(m)} \cdot U \right) \succ_{\alpha_m \beta} W^{(0)} \\
&\quad + (T^{(m)} \succ_{\alpha_m} U + T^{(m)} \prec_\beta U + T^{(m)} \cdot U) \prec_{\gamma_0} W^{(0)} + (T^{(m)} \succ_{\alpha_m} U + T^{(m)} \prec_\beta U \\
&\quad + T^{(m)} \cdot U) \cdot W^{(0)}, W^{(1)}, \dots, W^{(\ell)} \\
&= \bigvee_{x_1, \dots, x_{m-1}, x_m, z_1, \dots, z_\ell}^{m+1+\ell; \alpha_0, \dots, \alpha_{m-1}, \alpha_m \beta \gamma_0, \gamma_1, \dots, \gamma_\ell} \left(T^{(0)}, \dots, T^{(m-1)}, T^{(m)} \succ_{\alpha_m} (U \succ_\beta W^{(0)} + U \prec_{\gamma_0} W^{(0)} + U \cdot W^{(0)}) \right. \\
&\quad \left. + T^{(m)} \prec_{\beta \gamma_0} (U \succ_\beta W^{(0)} + U \prec_{\gamma_0} W^{(0)} + U \cdot W^{(0)}) + T^{(m)} \cdot (U \succ_\beta W^{(0)} + U \prec_{\gamma_0} W^{(0)} + U \cdot W^{(0)}), \right. \\
&\quad \left. W^{(1)}, \dots, W^{(\ell)} \right) \\
&= \bigvee_{x_1, \dots, x_m}^{m+1; \alpha_0, \dots, \alpha_m} (T^{(0)}, \dots, T^{(m)}) \cdot \bigvee_{z_1, z_2, \dots, z_\ell}^{\ell+1; \beta \gamma_0, \gamma_1, \dots, \gamma_\ell} (U \succ_\beta W^{(0)} + U \prec_{\gamma_0} W^{(0)} + U \cdot W^{(0)}, W^{(1)}, \dots, W^{(l)}) \\
&= T \cdot \bigvee_{z_1, z_2, \dots, z_\ell}^{\ell+1; \beta \gamma_0, \gamma_1, \dots, \gamma_\ell} (U \succ_\beta W^{(0)} + U \prec_{\gamma_0} W^{(0)} + U \cdot W^{(0)}, W^{(1)}, \dots, W^{(l)}) \\
&= T \cdot (U \succ_\beta W).
\end{aligned}$$

$$\begin{aligned}
&(T \cdot U) \prec_\gamma W \\
&= \bigvee_{x_1, \dots, x_m, y_1, y_2, \dots, y_n}^{m+1+n; \alpha_0, \dots, \alpha_{m-1}, \alpha_m \beta_0, \beta_1, \dots, \beta_n} \left(T^{(0)}, \dots, T^{(m-1)}, T^{(m)} \succ_{\alpha_m} U^{(0)} + T^{(m)} \prec_{\beta_0} U^{(0)} + T^{(m)} \cdot U^{(0)}, U^{(1)}, \dots, U^{(n)} \right) \\
&= \bigvee_{x_1, \dots, x_{m-1}, x_m, y_1, \dots, y_{n-1}, y_n}^{m+1+n; \alpha_0, \dots, \alpha_{m-1}, \alpha_m \beta_0, \beta_1, \dots, \beta_{n-1}, \beta_n \gamma} \left(T^{(0)}, \dots, T^{(m-1)}, T^{(m)} \succ_{\alpha_m} U^{(0)} + T^{(m)} \prec_{\beta_0} U^{(0)} + T^{(m)} \cdot U^{(0)}, \right. \\
&\quad \left. U^{(1)}, \dots, U^{(n-1)}, U^{(n)} \succ_{\beta_n} W + U^{(n)} \prec_\gamma W + U^{(n)} \cdot W \right) \\
&= \bigvee_{x_1, \dots, x_m}^{m+1; \alpha_0, \dots, \alpha_m} (T^{(0)}, \dots, T^{(m)}) \cdot \bigvee_{y_1, \dots, y_{n-1}, y_n}^{n+1; \beta_0, \dots, \beta_{n-1}, \beta_n \gamma} (U^{(0)}, \dots, U^{(n-1)}, U^{(n)} \succ_{\beta_n} W + U^{(n)} \prec_\gamma W + U^{(n)} \cdot W) \\
&= T \cdot (U \prec_\gamma W).
\end{aligned}$$

$$\begin{aligned}
&(T \cdot U) \cdot W \\
&= \bigvee_{x_1, \dots, x_{m-1}, x_m, y_1, y_2, \dots, y_n}^{m+1; \alpha_0, \dots, \alpha_{m-1}, \alpha_m \beta_0, \beta_1, \dots, \beta_n} \left(T^{(0)}, \dots, T^{(m-1)}, T^{(m)} \succ_{\alpha_m} U^{(0)} + T^{(m)} \prec_{\beta_0} U^{(0)} + T^{(m)} \cdot U^{(0)}, U^{(0)}, \dots, U^{(n)} \right) \\
&\quad \cdot \bigvee_{z_1, \dots, z_\ell}^{\ell+1; \gamma_0, \dots, \gamma_\ell} (W^{(0)}, \dots, W^{(l)}) \\
&= \bigvee_{x_1, \dots, x_{m-1}, x_m, y_1, \dots, y_{n-1}, y_n, z_1, \dots, z_\ell}^{m+1+n+\ell; \alpha_0, \dots, \alpha_{m-1}, \alpha_m \beta_0, \beta_1, \dots, \beta_{n-1}, \beta_n \gamma_0, \gamma_1, \dots, \gamma_\ell} \left(T^{(0)}, \dots, T^{(m-1)}, T^{(m)} \succ_{\alpha_m} U^{(0)} + T^{(m)} \prec_{\beta_0} U^{(0)} + T^{(m)} \cdot U^{(0)}, \right. \\
&\quad \left. U^{(0)}, \dots, U^{(n)} \right)
\end{aligned}$$

$$\begin{aligned}
& U^{(1)}, \dots, U^{(n-1)}, U^{(n)} \succ_{\beta_n} W^{(0)} + U^{(n)} \prec_{\gamma_0} W^{(0)} + U^{(n)} \cdot W^{(0)}, W^{(1)}, \dots, W^{(\ell)} \\
= & \bigvee_{x_1, \dots, x_m}^{m+1; \alpha_0, \dots, \alpha_m} (T^{(0)}, \dots, T^{(m)}) \cdot \bigvee_{y_1, \dots, y_{n-1}, y_n, z_1, \dots, z_\ell}^{n+1; \beta_0, \dots, \beta_{n-1}, \beta_n \gamma_0, \gamma_1, \dots, \gamma_\ell} (U^{(0)}, \dots, U^{(n-1)}, U^{(n)} \succ_{\beta_n} W^{(0)} \\
& + U^{(n)} \prec_{\gamma_0} W^{(0)} + U^{(n)} \cdot W^{(0)}, W^{(1)}, \dots, W^{(\ell)}) \\
= & T \cdot (U \cdot W).
\end{aligned}$$

SCHOOL OF MATHEMATICS AND STATISTICS, LANZHOU UNIVERSITY, LANZHOU, 730000, P. R. CHINA
Email address: zhangyy17@lzu.edu.cn

SCHOOL OF MATHEMATICS AND STATISTICS, KEY LABORATORY OF APPLIED MATHEMATICS AND COMPLEX SYSTEMS, LANZHOU UNIVERSITY, LANZHOU, 730000, P.R. CHINA
Email address: gaoling@lzu.edu.cn

LABORATOIRE DE MATHÉMATIQUES BLAISE PASCAL, CNRS–UNIVERSITÉ CLERMONT-AUVERGNE, 3 PLACE VASARÉLY, CS 60026, F63178 AUBIÈRE, FRANCE
Email address: Dominique.Manchon@uca.fr