



HAL
open science

Rapid and automatic discrimination between facial expressions in the human brain

Fanny Poncet, Jean Yves Baudouin, Milena Dzhelyova, Bruno Rossion, Arnaud Leleu

► **To cite this version:**

Fanny Poncet, Jean Yves Baudouin, Milena Dzhelyova, Bruno Rossion, Arnaud Leleu. Rapid and automatic discrimination between facial expressions in the human brain. *Neuropsychologia*, 2019, 129, pp.47-55. 10.1016/j.neuropsychologia.2019.03.006 . hal-02406196v2

HAL Id: hal-02406196

<https://hal.science/hal-02406196v2>

Submitted on 3 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Highlights

- We provide a neural marker for direct discrimination between facial expressions
- Specific brain signatures are isolated for each basic facial expression
- Fear and neutrality elicit larger and qualitatively distinct brain responses
- Neutral and fearful faces may have a particular status in the human brain

1
2
3
4
5 **Rapid and automatic discrimination between facial expressions in the human**
6
7 **brain**
8
9

10
11 Fanny Poncet¹, Jean-Yves Baudouin^{1,2}, Milena P Dzhelyova³, Bruno Rossion^{3,4,5}, Arnaud Leleu¹
12
13

14
15 ¹ Developmental Ethology and Cognitive Psychology group, Centre des Sciences du Goût et de
16 l'Alimentation, CNRS, Université Bourgogne Franche-Comté, Inra, AgroSup Dijon, F-21000 Dijon,
17 France

18 ² Laboratoire Développement, Individu, Processus, Handicap, Éducation (DIPHE), Département
19 Psychologie du Développement, de l'Éducation et des Vulnérabilités (PsyDÉV), Institut de
20 psychologie, Université de Lyon (Lumière Lyon 2), 69676 Bron cedex, France

21 ³ Psychological Sciences Research Institute, Institute of Neuroscience, University of Louvain (UCL),
22 1348 Louvain-la-Neuve, Belgium

23 ⁴ Université de Lorraine, CNRS, CRAN – UMR 7039, F-54000 Nancy, France

24 ⁵ Université de Lorraine, CHRU-Nancy, Service de Neurologie, F-54000, France
25
26
27
28
29

30 Corresponding authors: FP (Fanny.Poncet@u-bourgogne.fr) and AL (Arnaud.Leleu@u-bourgogne.fr)
31

32 Corresponding address : Équipe Éthologie développementale et Psychologie cognitive, Centre des
33 Sciences du Goût et de l'Alimentation, 9^E Boulevard Jeanne d'Arc, 21000 Dijon, France
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Abstract

Automatic responses to brief expression changes from a neutral face have been recently isolated in the human brain using fast periodic visual stimulation (FPVS) coupled with scalp electroencephalography (EEG). Based on these observations, here we isolate specific neural signatures for the rapid categorization of each of 5 basic expressions, i.e., when they are directly discriminated from all other facial expressions. Scalp EEG was recorded in 15 participants presented with pictures alternating at a rapid 6 Hz rate (i.e., one fixation/face, backward- and forward-masked). In different stimulation sequences, an expressive (angry, disgusted, happy, fearful, or sad) or a neutral face arose every 5 pictures (i.e., at $6/5 = 1.2$ Hz), among pictures of the same individual expressing the other emotions randomly. Frequency-domain analysis indicated a robust (i.e., recorded in every individual participant) and objective (i.e., at the predefined 1.2 Hz frequency and its harmonics) expression-specific brain response over occipito-temporal sites for each emotion and neutrality. In this context of variable expressions, while neural responses to the different expressions (Anger, Disgust, Happiness, Sadness) were dissimilar qualitatively, a much larger specific signature for neutral faces as compared to facial expressions was found. Interestingly, Fear also elicited a strong contrasted response with other facial expressions, associated with a specific neural signature over ventral occipito-temporal sites. Collectively, these findings reveal that specific EEG signatures for different facial expressions can be isolated in the human brain, pointing to partially different neural substrates. In addition, they provide support for a strong and highly selective neural response to fear at the system-level, in line with the importance of this emotional expression for biological survival.

Key-words: Fast Periodic Visual Stimulation, EEG, frequency-tagging, facial expression, discrimination, categorization.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1. Introduction

Quickly recognizing expressive facial cues about internal emotional states of others is a crucial ability in the human species, especially for adapted social interactions. Despite the wide variety of expressions, some of them have been described as universal basic emotions, being recognized by allocating distinct patterns of facial actions to discrete emotion categories (Darwin, 1872; Ekman, 1993; for meta-analyses, see Elfenbein & Ambady, 2002; Sauter & Eisner, 2013). More specifically, according to Ekman’s proposal (Ekman, 1992, 1994; Ekman & Friesen, 1971; Ekman, Friesen, & Ellsworth, 1972), facial expressions are perceived as belonging to qualitatively discrete emotional categories with sharp, clear-cut boundaries between each category (although see Barrett & Wager, 2006; Russell, 1980; Woodworth & Schlosberg, 1954).

At the neural level, isolating a specific signature for each facial expression has been rather challenging. A number of studies have shown that the processing of every facial expression elicits activity in face-selective regions of the inferior occipital gyrus (IOG), lateral fusiform gyrus (LatFG) and posterior superior temporal sulcus (pSTS) (Calder & Young, 2005; Haxby, Hoffman, Gobbini, 2000). The latter region is known to play an important role in processing dynamic changes in expression (Puce et al., 2003; Srinivasan et al., 2016), although significant responses to facial expressions have also been recorded in the LatFG and the ventral face processing system in general (e.g., Kawasaki et al., 2012). Outputs from the pSTS are thought to project to an extended face-processing network, including the amygdala, the insula, and the limbic system, for further analysis of facial expression. At this level, distinct and discrete brain regions are considered to be involved in the processing of the different basic facial emotion categories; namely happiness, anger, disgust, fear, and sadness (e.g., Vytal, & Hamann, 2010 for a meta-analysis). However, the specificity of these regions for processing various expressions has been questioned (Barrett & Wager, 2006; Barrett, Lindquist, Bliss-Moreau, Duncan, & Brennan, 2007). In addition, recent studies reported that emotional facial expressions also elicit separable patterns of brain activity in regions responding to all facial expressions, such as the anterior and posterior STS and the amygdala (Harris, Young, & Andrews, 2012, 2014; Said, Moore, Engell, Todorov, & Haxby, 2010; Whalen et al., 2013). In sum, neuroimaging studies have struggled to identify consistent specific patterns of neural activation for each major facial expression category.

With scalp electroencephalography (EEG), a technique that is not only advantageous for its high temporal resolution but also for being able to measure direct brain activity in various populations (e.g., neurotypical adults, children and infants, as well as neurological or psychiatric patients), specific neural signatures for some facial expressions have also been very difficult to isolate

1 when using standard event-related potentials (ERPs) approaches (for reviews, see Calvo &
2 Nummenmaa, 2016; Vuilleumier & Pourtois, 2007). The differential processing of emotional vs.
3 neutral faces is thought to primarily occur around 150-300ms post-stimulus onset over occipito-
4 temporal and occipito-parietal sites (e.g., Calvo, Marrero, & Beltrán, 2013; Leleu, Godard, Dollion,
5 Durand, Schaal, & Baudouin, 2015; Williams, Palmer, Liddell, Song, & Gordon, 2006). The
6 categorization of facial expressions in discrete categories is further considered from about 300ms
7 after stimulus-onset, at the level of several late components with distinct topographies across studies
8 (P3b: Luo, Feng, He, Wang, & Luo, 2010; late positive potential/LPP: Leppänen, Kauppinen, Peltola, &
9 Hietanen, 2007; slow positive waves/SPWs: Calvo & Beltrán, 2013; posterior negativity: Ashley,
10 Vuilleumier, & Swick, 2004). Conclusions about brain facial expressions categorization remain fragile
11 however, and no solid and systematic pattern has been reliably found for each emotion category (for
12 discussions, see Calvo & Nummenmaa, 2016; Dzhelyova et al., 2017). This is not surprising since ERPs
13 are generally obtained by contrasting the sudden onset of a facial expression stimulus to a uniform
14 visual field (i.e., background). The recorded brain response thus largely reflects the general visual
15 processing of the stimulus (including specific responses to faces or to all expressive faces), making it
16 difficult to isolate subtle differential neurophysiological activities across expressions.
17

18 To tackle this issue, recent studies have used Fast Periodic Visual Stimulation (FPVS)
19 associated with EEG recordings to show robust neural responses to brief expression *changes* directly
20 contrasted with neutral faces (Dzhelyova, Jacques & Rossion, 2017; Leleu, Dzhelyova, Rossion,
21 Brochard, Durand, Schaal & Baudouin, 2018). The FPVS-EEG approach relies on the brain's property
22 to synchronize its activity with stimuli displayed periodically (Adrian & Matthews, 1934), eliciting EEG
23 responses at the same frequency (Regan, 1989; Norcia, Appelbaum, Ales, Cottareau, & Rossion, 2015
24 for review). FPVS-EEG has high objectivity because responses are measured at a predetermined
25 frequency of stimulation. This response can have a high signal-to-noise ratio (SNR), even within a few
26 minutes of recording. By using a rapid base rate of visual stimulation and introducing a systematic
27 stimulus change at a lower periodic rate, FPVS-EEG designs allow dissociating two distinct brain
28 responses elicited at two different frequencies within a single stimulation sequence (Braddick,
29 Wattam-Bell & Atkinson, 1986; Liu-Shuang, Norcia & Rossion, 2014). The first response, tagged at the
30 base frequency, reflects the neural processing of all visual events occurring at each stimulus-onset, a
31 mixture of both low- and higher-level processes. Interestingly, the second response, tagged at the
32 lower frequency, is a direct (i.e., without post-hoc subtraction) marker of the *differential* neural
33 processes specifically elicited by the specific stimulus change in contrast to the base stimuli. In recent
34 years, this approach has been successfully used to investigate the categorization of face stimuli as
35 faces (against various nonface objects; e.g., Rossion et al., 2015), face individuation (Liu-Shuang et
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

al., 2014; 2016; Dzhelyova & Rossion, 2014a, 2014b), and facial expression processing (Dzhelyova et al., 2017; Leleu et al., 2018).

In the studies by Dzhelyova et al. (2017) and Leleu et al. (2018), specific expressions (either, happy, fear, or disgust, with anger and sadness in Leleu et al., 2018) were displayed regularly (at 1.2 or 1.33Hz) in different sequences with the neutral face of the same individual at the base frequency (6 or 12Hz), mimicking brief expression changes from neutrality. The face stimuli were displayed either upright or upside-down (Dzhelyova et al., 2017) or with linearly increasing intensities (Leleu et al., 2018). Discrimination of the expressions from the neutral face was indexed by the predefined frequency-tagged brain response to the target expression with: (1) a response to brief changes of facial expression over the right ventral and dorsal occipito-temporal sites, larger for upright than for upside-down faces, and increasing non-linearly with expression intensity; (2) variable amplitudes and topographies for the different expressions, with larger response for disgust than the other expressions, a more dorsal response for happiness, more restricted responses for fear and sadness, a more bilaterally distributed response to fear, and the lowest response for sadness. In both studies, significant responses were evidenced in every individual participant.

Here, we investigate further whether a brain response specifically indexing the categorization of facial expressions, i.e., their assignment into discrete categories from the differential recruitment of brain structures for each facial emotion, can be isolated using FPVS-EEG. Indeed, when contrasting a given facial expression with neutrality, the evoked response may elicit neural activities that are both specific for that expression and common to several if not all expressions. Some brain areas may respond similarly to all expressions, whereas some may respond to a subset or only to a single expression (e.g., Vytal & Hamann, 2010). Some regions may also respond to all expressions but with separable brain activity (Said et al., 2010). To dissociate the contributions of specific and common neural activities across expressions, we used a design in which variable facial expressions were randomly displayed at a 6 Hz base stimulation frequency (i.e., 6 stimuli per second, stimulus duration \approx 167 ms, single fixation limiting eye movements) and directly contrasted to another target expression periodically inserted every 5th stimulus, at a 1.2 Hz specific frequency (*Figure 1*). Hence, the brain response tagged at 6 Hz here reflects both low-level visual processes (e.g., contrast-change) elicited by the rapid stream of stimulation, and higher-level perceptual responses to the rapid change of facial expression. If common neural processes are involved for every facial expression (e.g., neural activities elicited by any facial movement), they are captured by this general visual response to the fast train of facial expressions. In contrast, the brain response tagged at 1.2 Hz is a direct signature of *expression categorization*, that is, the differential neural activities elicited for one specific expression against all other possible ones. To determine whether the neutral expression is discriminated from all of the other expressions, it was also included at the 1.2 Hz expression-specific frequency. We show

1 that the brain response to each facial expression can be quantified at the specific 1.2 Hz frequency
2 and its harmonics (i.e., integer multiples). The expression-specific response is significant in every
3 individual brain, characterized by different scalp topographies according to the emotion expressed,
4 and is larger for the categorization of neutrality and fear, thus reflecting the particular status of these
5 two facial expressions.
6
7
8
9

10 **2. Materials and methods**

11 **2.1. Participants**

12
13 Fifteen participants (10 females, mean age = 25.5 ± 4 (SD) years, range 20–33 years) were
14 included in the experiment and received financial compensation. All of them reported normal or
15 corrected-to-normal vision and none reported history of psychiatric or neurological disorders.
16 Participants provided written informed consent prior to the experiment and testing was conducted in
17 accordance with the Declaration of Helsinki. It is worth noting that we conducted an exploratory
18 study for which effect size was not pre-experimentally defined and adequate sample size difficult to
19 estimate. We further discuss this point in the Limitations section and provide power analysis
20 conducted on the observed effect size.
21
22
23
24
25
26
27
28
29
30

31 **2.2. Stimuli**

32
33 Images from 6 individual faces (01F, 02F, 05F, 07F, 09F, 10F) were taken from the NimStim
34 database (Tottenham et al., 2009). We selected for each face a neutral expression as well as an
35 expression of anger, disgust, fear, happiness and sadness (*Figure 1*). Surprise was not used due to its
36 ambiguity in terms of emotional valence (Kim et al., 2004), and the lack of evidence for categorical
37 boundaries between surprise and other expressions (Etcoff & Magee, 1992). Stimuli were cropped in
38 a medallion-shape window to remove information from the background and hairstyle and presented
39 on a mid-level gray color background (128/255 in grayscale). Images were set to a size of 6×4.8 cm
40 ($6 \times 4.8^\circ$ of visual angle at a viewing distance of 57 cm) and equated for mean luminance.
41
42
43
44
45
46
47
48

49 **2.3. Procedure**

50
51 The procedure was similar to the experiments of Dzhelyova et al. (2017) and Leleu et al. (2018). Fast
52 periodic visual stimulation (FPVS) was used with a base rate stimulation of 6 Hz (i.e. 6 images/s), and
53 an expression-specific rate of 1.2 Hz (same emotion every 5 pictures, see below). Using a 24-inch LED
54 monitor with a 60 Hz refresh rate and a resolution of 1920×1080 pixels, stimuli were displayed on a
55 mid-level gray background (128/255 in grayscale) through sinusoidal contrast modulation (0–100 %) at
56 the base frequency (6 Hz) with custom Java software (SinStim). At this rapid rate, each stimulus is
57
58
59
60
61
62
63
64
65

presented for about 167ms and reaches full contrast approximately 83ms after onset (half a cycle). This rapid mode of stimulation implies that perception occurs at a glance. To avoid low-level repetition effects, the size of the images was randomly varied between 95 % and 105 % (i.e., $5.7 \times 4.56^\circ$ and $6.3 \times 5.04^\circ$ of visual angle respectively) at every stimulation cycle. Sequences presenting one individual face and testing the perception of one specific emotional expression were organized so that one of the emotional expressions is periodically inserted at the expression-specific rate (i.e., $6/5 = 1.2$ Hz, ≈ 833 ms between two target expressive faces) among the other expressions randomly displayed at the base frequency (*Figure 1*). As a result, two brain responses reflecting two distinct processes elicited within a single stimulation sequence are projected to different frequencies in the EEG spectrum. EEG amplitude recorded at the 6 Hz base frequency and its harmonics (i.e., integer multiples of 6 Hz: 12 Hz, 18 Hz, etc.) reflects a general visual response to the rapid stream of face stimuli changing in contrast, size and expression. In contrast, the neural activity elicited at 1.2 Hz and harmonics is a direct marker of the automatic categorization of a specific facial expression with no contamination from other processes elicited by any change of facial expression. To determine whether the neutral expression is automatically categorized when directly contrasted against all the other expressions, it was also included at the expression-specific frequency. Hence, the design evaluated the specific response to the five emotional expressions (anger, disgust, fear, happiness and sadness) plus neutrality, for a total of 6 conditions. One advantage of this design is that it directly contrasts every expression category within a single stimulation sequence and thus isolates specific neural responses to one emotional expression against the others.

After EEG-cap placement, participants were seated in a light- and sound-isolated cabin and their head was held on a chinrest at 57 cm from the screen. Each participant was stimulated with 36 sequences: 6 conditions \times 6 individual faces. For each sequence, the stimulation lasted 35 s on average with a random pre-stimulation interval of 0.5 to 1 s presenting a blank screen, followed by a gradual fade-in of increasing contrast modulation depth (2.5s), and then by the stimulation (30s). A gradual fade-out lasted 1s afterward followed by a random blank post-stimulation interval (0.5 to 1s). The 36 sequences were divided in 6 blocks of 6 sequences, each block presenting one sequence per condition and alternating individual faces. The presentation orders of blocks and sequences within blocks were randomized across participants. The presentation order of the 5 irrelevant expressions at the base frequency followed cycles of the 5 expressions (e.g., anger, disgust, fear, sadness and neutrality when happiness was presented at the expression-specific frequency; *Figure 1*) whose order was randomized at each cycle. During each sequence, participants were instructed to look at a circle that was always present on the screen and located just below the eyes when the face stimuli were displayed. They had to press the space bar with both index fingers as fast as possible each time the circle changed to a square. These shape-changes briefly (200ms) intervened 5 random times with

1 a minimum 2 s interval between each change. This orthogonal task was used to ensure that
2 participants were attentive. When asked at the end of the experiment, all participants reported
3 having noticed expressive faces during the stimulation, but none detected their periodicity.
4
5

6 **2.4. EEG Acquisition**

7
8 Electroencephalographic (EEG) activity was continuously recorded from a BioSemi Active-
9 Two amplifier system (BioSemi, The Netherlands) with 64 Ag/AgCl electrodes located according to
10 the 10-10 classification system. During recording, the Common Mode Sense (CMS) active electrode
11 was used as reference and the Driven Right Leg (DRL) passive electrode was used as ground.
12
13 Electrode offset was reduced between $\pm 15 \mu\text{V}$ for each channel and EEG was digitalized at a 1024 Hz
14 sampling rate.
15
16
17
18
19
20

21 **2.5. EEG preprocessing**

22
23 EEG processing was conducted using Letswave 5 (<http://nocions.webnode.com/letswave>),
24 running on Matlab 2012 (Mathworks, USA), and largely followed analyses steps described in previous
25 studies (Dzhelyova et al., 2017; Leleu et al., 2018). The continuously recorded data were first
26 bandpass filtered at 0.1–100 Hz (butterworth filter, 4th order) and then downsampled to 256 Hz to
27 reduce file size and processing time. They were cropped into 35s-segments for each sequence (1s
28 before the fade-in and 0.5s after the fade-out). An Independent Component Analysis (ICA) was
29 computed (e.g., Makeig et al., 1996) and components corresponding to eye blinks (recorded over Fp
30 electrodes) and artifacts recorded over frontal and temporal electrodes were removed. Remaining
31 noisy or artifact-ridden electrodes (i.e., with amplitude exceeding $\pm 100 \mu\text{V}$) were rebuilt using linear
32 interpolation from the four nearest channels (mean number across participants: 0.4 channels, range:
33 0–2). All EEG epochs were finally re-referenced to a common average reference.
34
35
36
37
38
39
40
41
42
43
44

45 **2.6. EEG frequency-domain analysis**

46
47 Preprocessed data segments were cropped down into 30-sec epochs beginning just after the
48 fade-in and corresponding exactly to thirty-six 1.2 Hz cycles (for a total of 7680 time bins). The 6
49 segments corresponding to the 6 sequences for a specific emotion (i.e., 6 individual faces) were then
50 averaged in the time domain for all participants to reduce EEG activity non-phase-locked to the
51 stimuli. The resulting 6 averaged segments (i.e., all conditions) were finally transformed with fast
52 Fourier transform (FFT) and amplitude spectra were extracted for all channels with a high frequency
53 resolution of $1/30 = 0.033 \text{ Hz}$. Individual FFT data were grand-averaged across participants for group
54 analysis.
55
56
57
58
59
60
61
62
63
64
65

1 To determine the significant harmonics for both general visual (i.e., elicited by the rapid 6 Hz
2 base stimulation frequency) and expression-specific (i.e., elicited by the 1.2 Hz periodic presentation
3 of a given facial expression) responses, FFT grand-averaged data were pooled across electrodes and
4 conditions and Z-scores were calculated at each frequency bin by subtracting the mean surrounding
5 noise amplitude (i.e., estimated from the 20 surrounding frequency bins excluding the 2 immediately
6 neighboring and the 2 most extreme values, e.g. Dzhelyova et al., 2017; Leleu et al., 2018) and
7 dividing by its standard deviation. Harmonics were considered significant until Z-scores were no
8 longer above 1.64 ($p < .05$, one-tailed, signal > noise) for two consecutive harmonics. For the general
9 visual response, significant harmonics were found until the 8th harmonic (i.e., 48 Hz, harmonics were
10 not considered after the 50 Hz response elicited by AC power). For the expression-specific response,
11 significant harmonics were found until the 7th harmonic (i.e., 8.4 Hz).

12 Z-scores were then used as a data-driven whole-scalp approach to determine different
13 regions-of-interest (ROIs) to include in statistical analyses, separately for each brain response. FFT
14 data were first grand-averaged across conditions and significant harmonics for each response were
15 summed for each electrode (for the expression-specific response, the 5th harmonic corresponding to
16 the base frequency (i.e., 6 Hz) was excluded). Summed harmonics were used to quantify the overall
17 responses in single values (Retter & Rossion, 2016). Topographical differences were then normalized
18 according to the scalp-wide global power of each brain response (McCarthy & Wood, 1985).
19 Normalization allows identifying the electrodes over which the response is largest irrespective of its
20 global magnitude across the scalp (Leleu et al., 2018). Z-scores were calculated and the significant
21 electrodes were finally pooled together in three distinct regions of interest (ROIs) considered for
22 further statistical analyses (i.e., different ROIs and analyses for each brain response). For the
23 expression-specific response, ROIs comprised medial occipital (mO: channels Oz, Iz, O1, O2), and
24 lateral (right and left) occipito-temporal (rOT: channels P4, P8, P10, PO4, PO8, IOT: channels P3, P7,
25 P9, PO3, PO7) sites. Note that the left-hemispheric channels P3 and PO3 were not significant but
26 included in the IOT ROI to match the significant channels identified in the right hemisphere (i.e., P4
27 and PO4). For the general visual response, the mO region included 5 channels (Oz, Iz, O1, O2, POz),
28 and each lateral ROI included 3 channels (rOT: PO4, PO8, P8; IOT: PO3, PO7, P7).

29 Statistical analyses were separately carried on both responses (i.e., expression-specific and
30 general visual). To quantify the overall magnitude of each response in a single value expressed in
31 amplitude (μV), individual FFT amplitude spectra were first baseline corrected by subtracting the
32 mean surrounding noise (see above) and harmonics were summed until the 8th harmonic for the
33 general visual response, and until the 7th harmonic for the expression-specific response excluding the
34 one corresponding to the base frequency (i.e., 6 Hz). Repeated-measures ANOVAs (i.e., one for each
35 brain response) were conducted on these summed baseline-corrected amplitudes (BCA) with
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Expression (anger, disgust, fear, happiness, sadness and neutrality) and *ROI* (mO, IOT, rOT) as within-subject factors. Mauchly's test for sphericity violation was performed and Greenhouse-Geisser correction for degrees of freedom was applied whenever sphericity was violated. For significant effects, post-hoc comparisons were conducted using Tukey's HSD test. For visualization of both brain responses, signal-to-noise ratios (SNR) were calculated as summed uncorrected amplitudes divided by mean surrounding noise.

2.7. Scalp topography analysis

To assess whether the different facial expressions were associated with specific topographical patterns, we determined which electrodes across the whole scalp presented a significant expression-specific response on the grand-averaged uncorrected amplitudes summed across significant harmonics. Topographical differences were first normalized (McCarthy & Wood, 1985) to equalize the scalp-wide global magnitude of the response across emotions, and Z-scores were then calculated for each electrode. A repeated-measures ANOVA was also conducted on normalized summed BCA with *Expression* (anger, disgust, fear, happiness, sadness and neutrality) and *Electrode* (64 channels) as within-subject factors to determine whether the different emotions elicit distinguishable topographical patterns when amplitude is equalized across them (Dzhelyova et al., 2017). We focused on the interaction between *Expression* and *Electrode* that would indicate different topographical distributions between facial expressions.

2.8. Individual data analysis

To assess the significance of the expression-specific response at an individual level, we also calculated Z-scores (see section 2.6) in every participant for summed uncorrected amplitudes over the electrodes belonging to the defined ROIs, and then for all the other electrodes.

3. Results

3.1. Expression-specific response

Since only one specific facial expression periodically appeared at 1.2 Hz (i.e., every 5th stimulus) among all the other expressions randomly displayed at the 6 Hz base rate of stimulation, the brain response recorded at 1.2 Hz and its harmonics directly reflects single-glance categorization of a specific expression against the others regardless of expression-changes (captured in the general visual response elicited at the 6 Hz base rate). Visual inspection revealed that an expression-specific response was mainly recorded over lateral occipito-temporal and occipito-parietal regions, with

1 larger amplitudes for fear and neutrality (Fig. 2). These two expressions elicited specific responses of
2 high signal-to-noise ratio (SNR around 1.8, i.e. 80 % of signal increase) suggesting that they are
3 clearly discriminated when directly contrasted with all the other expressions. While the other facial
4 expressions elicited weaker responses, they were nonetheless all visible on the EEG spectra (SNR
5 around 1.4, i.e. 40 % of signal increase) except for happiness (but with a significant nevertheless, see
6 Z-scores below).
7
8
9

10 These observations from visual inspection were confirmed by the statistical analyses with a
11 significant main effect of *Expression* ($F(2.6, 36.6) = 12.35, \epsilon=0.52, \eta^2_p = 0.477, p<.001$), due to a larger
12 expression-specific response for fear (0.36 ± 0.05 (SEM) μV) and neutrality ($0.37 \pm 0.08 \mu\text{V}$) than for
13 the other 4 expressions (anger: $0.09 \pm 0.03 \mu\text{V}$, disgust: $0.14 \pm 0.03 \mu\text{V}$, happiness: $0.06 \pm 0.03 \mu\text{V}$,
14 sadness: $0.12 \pm 0.03 \mu\text{V}$, all $ps<.01$), and no other significant differences. This effect was qualified by
15 an interaction between *Expression* and *ROI* ($F(10, 140) = 2.48, \eta^2_p = 0.15, p=.009$) with a larger
16 response over mO ($0.45 \pm 0.11 \mu\text{V}$) than IOT ($0.25 \pm 0.06 \mu\text{V}$) only for neutrality ($p=.001$). However,
17 further inspection of the spatial distribution of the response for each facial expression suggested
18 variable topographies between expressions (Fig. 3).
19
20
21
22
23
24
25

26 To better identify the topographical patterns observed for each facial expression, we
27 eliminated the large amplitude differences between expressions by normalizing the response
28 amplitude across the whole scalp (McCarthy & Wood, 1985; see Dzhelyova et al., 2017). As can be
29 seen in the upper part of Figure 3, normalized summed baseline-corrected amplitudes (BCA) revealed
30 a variable topography of the expression-specific response across the different expressions.
31 Interestingly, while all facial expressions seem to elicit occipito-parietal activities, and present with a
32 right hemisphere advantage, fear is the only expression showing a more ventral topographical
33 distribution and a left hemisphere advantage. The repeated-measures ANOVA run on normalized
34 summed BCA and including the 64 recording channels revealed a significant interaction between
35 *Expression* and *Electrode* ($F(315,4410)=1.51, \eta^2_p = 0.10, p<.001$), confirming the variable spatial
36 distributions of the expression-specific response across facial expressions when amplitude
37 differences are controlled.
38
39
40
41
42
43
44
45
46

47 We also calculated Z-scores on normalized uncorrected amplitudes to determine which
48 electrodes presented a significant expression-specific response for each emotion (bottom part of
49 Figure 3). All facial expressions elicited a significant expression-specific response over at least 4
50 posterior electrodes. Responses to anger and sadness were centered over medial occipital sites while
51 responses to disgust and happiness were sparser over posterior regions. The categorization of fear
52 and neutrality elicited significant responses encompassing many posterior electrodes but with a
53 greater right hemispheric response only for neutrality. Significant electrodes also confirmed that an
54
55
56
57
58
59
60
61
62
63
64
65

1 expression-specific response was recorded over occipito-parietal channels for all facial expressions
2 except for fear.
3
4
5

6 **3.2. Individual data analysis**

7
8 Individual data analyses revealed the strength of the expression-specific response and its
9 reliability across participants. When combining all facial expressions, a significant expression-specific
10 response was recorded for every participant over at least 2 out of the 14 electrodes included in the
11 ROIs (*Figure 4*). When considering each facial expression separately, a significant expression-specific
12 response was recorded over at least 2 electrodes over the scalp for every participant and for every
13 facial expression. For fear in particular, the response was significant for every participant, with at
14 least 2 out of the 14 electrodes included in the ROIs. The same was observed for neutrality, with at
15 least 1 electrode included in the ROIs, except for one participant who nevertheless displayed a
16 significant response over another posterior electrode (P5). For disgust and happiness, 13 out of the
17 15 participants displayed a significant response over at least 1 electrode included in the ROIs, the 2
18 other participants presenting a significant response elsewhere over the scalp. Finally, for anger and
19 sadness, a significant response was observed over at least 1 electrode included in the ROIs for 12 out
20 of the 15 participants, the 3 other participants displaying a significant response elsewhere over the
21 scalp.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36

37 **3.3. General visual response and behavioral data**

38
39 The brain response recorded at the base frequency and harmonics reflects the processes
40 elicited by low- and high-level visual cues rapidly changing in the stream of stimulation (e.g., contrast,
41 expression-change). In line with previous studies (Dzhelyova et al., 2017; Leleu et al., 2018), we found
42 a clear medial occipital response of high amplitude (*Figure 2*). Accordingly, the main effect of *ROI*
43 ($F(2,28) = 21.14, \eta^2_p = 0.60, p < .001$) confirmed the larger response over mO sites ($3.68 \pm 0.42 \mu V$)
44 than over the other ROIs ($p < .01$, IOT: $1.56 \pm 0.17 \mu V$, rOT: $2.38 \pm 0.28 \mu V$), with 48.3 % of the overall
45 signal recorded over mO regions (respectively 20.5 % and 31.2 % for IOT and rOT), the response at
46 the rOT sites is also larger than over the IOT sites ($p < .05$). Importantly, neither the main effect of
47 *Expression* ($F < 1$) nor its interaction with *ROI* ($F(10,140) = 1.17, \eta^2_p = 0.07, p = .35$) were significant,
48 confirming that the facial expression displayed at the expression-specific frequency had no influence
49 on the general visual response, and that the global attentional level that typically modulates such
50 periodic brain activities (e.g., Kim et al., 2007) cannot account for the differences observed for the
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 expression-specific response. In addition, behavioral data for the orthogonal task showed that the
2 mean accuracy for detecting circle-to-square changes was 94.56% (SD=9.2), with a mean correct
3 response time of 457ms (SD=54). There were no differences between facial expressions for both
4 accuracy and response times (both $F_s < 1$).
5
6
7
8
9

10 **4. Discussion**

11
12 Here, we extend recent FPVS-EEG studies measuring facial expression discrimination
13 (Dzhelyova et al., 2017; Leleu et al., 2018) by showing that when directly contrasted against all the
14 other expressions, each facial expression elicits a specific signature over posterior brain areas in the
15 EEG spectrum, thus providing evidence for specific neural processes dedicated to each facial
16 expression. We further showed that rapid discrimination (i.e. at a glance) between facial expressions
17 and assignment of one expression into a specific emotion category are subtended by discriminable
18 activities at the brain level, with variable topographies between expressions. More specifically, fear
19 and neutrality elicit a larger response than anger, disgust, sadness, and happiness, revealing a
20 particular status for these two expressions among all the others. For fear, the response also differs
21 qualitatively, with a more ventral (i.e., occipito-temporal) and left-hemispheric response than for the
22 other expressions.
23
24
25
26
27
28
29
30
31

32 The main purpose of the study was to determine whether dissociable brain responses can be
33 observed for the assignment of facial expressions into discrete emotional categories. In other words,
34 our aim was to demonstrate that specific patterns of electrocortical activities can be recorded for
35 each facial expression when neural activities elicited for every facial expression (e.g., a neural process
36 that codes for facial movements in general) are eliminated. We controlled for those general neural
37 responses using a frequency-tagging approach in which an expression-specific response was
38 recorded at 1.2 Hz and harmonics by opposing a facial emotion to all the others, delineating the
39 aspects of the response that are specifically elicited by that given facial expression. Indeed, since
40 facial expressions were rapidly changing at the 6 Hz base rate of stimulation, neural activities
41 reflecting the processing of any change of facial expression were captured by the general visual
42 response recorded at 6 Hz and harmonics. To our knowledge, this is the first probing evidence of
43 unique EEG signatures for each basic facial expression. Indeed, in previous studies, brain responses
44 to variable facial expressions were generally confounded with other more general neural signals
45 (e.g., face-related responses, expression-change responses) when measuring a large and global EEG
46 response to the sudden onset of facial expression from a no-stimulus baseline as in typical ERP
47 studies (Calvo & Nummenmaa, 2016 for review), or a brain response to a change of expression from
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 a neutral face as in recent FPVS-EEG studies (Dzhelyova et al., 2017; Leleu et al., 2018). Importantly,
2 the different facial expressions did not modulate the general visual response elicited at 6 Hz and
3 harmonics, and the behavioral data obtained for the orthogonal shape-change detection task,
4 showing that the expression-specific responses are not accounted by attentional variations during
5 visual stimulation but truly reflect the differential processing of facial expressions.
6

7
8 With the present design, we were thus able to reveal specific neural markers of high SNR
9 (i.e., ≈ 1.4 in average, 40 % of signal increase compared with surrounding noise level) for each facial
10 emotion in a few minutes of recording (i.e., only 3 minutes of stimulation were analyzed for each
11 facial expression). These specific signatures were significant in every individual participant and
12 presented with clear distinct topographies and amplitudes for the five basic expressions of
13 happiness, anger, disgust, fear, and sadness, as well as for neutrality. While neutrality elicits a large
14 response over right occipito-temporal sites, anger and sadness elicit more focal responses over
15 medial occipital and parietal sites, predominantly over the right hemisphere for anger and more
16 medial for sadness. Responses to disgust and happiness are more laterally distributed and
17 encompass mostly central parietal regions, predominantly over the right hemisphere. In contrast,
18 fear elicits a large response encompassing all occipito-temporal sites, predominantly over the left
19 hemisphere. The EEG responses elicited with the present design may reflect the activity of brain
20 regions involved in parting facial expressions into discrete categories, such as the STS (Said et al.,
21 2010). The distinct topographies also suggest that neural activities originating from at least partially
22 distinct brain regions are recorded, such as those from the extended network involved in facial
23 emotion recognition (Calder & Young, 2005; Haxby et al., 2000; Vytal & Hamann, 2010). Indeed,
24 many electrodes responded only to a subset of expressions, and none responded to all expressions.
25 Thus, we propose that the present paradigm taps the activity of neural substrates specifically
26 involved in the coding of discrete facial emotions, as those isolated by brain imaging studies (e.g.,
27 Vytal & Hamann, 2010). These authors further underlined that the regions identified for each
28 emotion category respond reliably to facial expressions, but also to other kinds of materials, visual or
29 not. Thus, the expression-specific brain responses recorded here may also partly capture neural
30 activities from areas that code for the emotional content of stimuli, irrespectively of their physical
31 characteristics.
32

33
34 In this framework, both fear and neutrality appeared to elicit larger expression-specific
35 responses. For neutrality, a face without emotional expressive features was opposed to all
36 expressions of emotions. The observed large response for this non-emotional expression suggests
37 that some brain regions respond to the general dissociation between non-emotional vs. emotional
38 faces, that is, brain regions whose activity is related to the occurrence of (non-)expressive features,
39 whatever their emotional content. In that respect, we observed a response pattern that was more
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

typical of the right occipito-temporal activities usually reported for several face-related processes with the FPVS-EEG approach (e.g., Rossion, 2014a, 2014b). This suggests that the differential response to neutral faces originates from the core regions of the face-processing network (Calder & Young, 2005; Haxby et al., 2000). Considering the nature of the design – which contrasted expressive and non-expressive faces – the specific response to neutrality may be mainly driven by cortical regions involved in the processing of changeable aspects of faces, notably the STS (e.g., Srinivasan, Golomb, & Martinez, 2016). It may also reflect activity from the lateral fusiform gyrus, involved in the processing of invariant aspect of faces (Haxby et al., 2000), and also modulated by the emotional content of faces (Buchel & Dolan, 2000; George, Driver, & Dolan, 2001; Kawasaki et al., 2012; Morris et al., 1998; Vuilleumier, Armony, Driver, & Dolan, 2001).

For fear, the response displayed the larger amplitude compared with the other emotions, with a more occipito-temporal topography and a left hemisphere advantage. This suggests that fear holds a special position as a facial expression, clearly discriminated from the other expressions, possibly in line with a selective “fear network” in the brain centered on the amygdala (Öhman & Wiens, 2004). Accordingly, both neuroimaging (Vytal & Hamann, 2010) and lesion studies (Adolphs, Tranel, Damasio, & Damasio, 1994) have reliably associated fear with amygdala activity. It is unlikely that the response recorded in our study was elicited directly in the amygdala, considering the deep location of this structure. Rather, it could reflect the modulatory influence of this structure on the visual cortex. Such modulations were frequently suggested in brain imaging studies (Pessoa, McKenna, Gutierrez, & Ungerleider, 2002; Vuilleumier, Armony, Driver, & Dolan, 2001). Furthermore, such modulations were not observed in patients with amygdala damages (Vuilleumier, Richardson, Armony, Driver, & Dolan, 2004). Thus, the stronger response we recorded for fear at posterior occipito-temporal sites, in comparison to all other emotional expressions, may reflect the reentrance modulation of the amygdala to the visual cortex (Kawasaki, Tsuchiya, Kovach, Nourski, Oya, Howard, & Adolphs, 2012; Pourtois et al., 2010), or more generally the spreading of the response elicited by fear in the amygdala to occipito-temporal, STS, anterior temporal and orbitofrontal cortex (Krolak-Salmon, Henaff, Vighetto, Bertrand, & Mauguiere, 2004). Finally, it may reflect top-down influences from an amygdala-prefrontal system that increased monitoring of the environment to solve predictive uncertainty initiated by fearful expressions (see Whalen et al., 2013). In any event, thanks to FPVS-EEG, we were able to isolate a specific signature for fear compared with other facial expressions. It holds high promise for future research conducted on populations whose differential processing of fear can be impaired (e.g. individuals with autism spectrum disorders (ASD); Malaia, Cockerham, & Rublein, 2017), in line with the successful recent application of FPVS-EEG in clinical research for testing e.g. face individuation in individuals with ASD (Vettori et al., 2018) or expression-change perception in participants with 22q11.2 deletion syndrome (Leleu et al., 2019).

4.1 Limitations

One limitation of the present study is the relatively small sample size that was not pre-experimentally defined due to the exploratory nature of the study. However, thanks to the frequency-tagging approach which provides very high SNR, we were able to identify a significant expression-specific response in every participant. This suggests that this brain response to facial expression is reliable. Accordingly, power analysis conducted on BCA averaged across ROIs and facial expressions with a significance level $\alpha = .05$ (one-tailed, signal > noise = $BCA > 0$) revealed that with a mean amplitude of 0.19 ± 0.11 (SD) μV , we obtained a large effect size of 1.73 (Cohen's d) leading to maximal power $1-\beta = 1$. With this power, a similar effect could even be found with a smaller sample size $N = 8$. Power would still be high ($1-\beta = 0.83$) if we consider a smaller effect size of $d = 0.67$, corresponding to the lower limit of the 99 % confidence interval calculated with the observed effect size. Hence, despite a small sample size, the brain response identified here seems to reflect a robust neural marker of the visual discrimination of facial expressions.

5. Conclusions

Taken together, the present findings demonstrate that specific EEG signatures can be isolated and quantified for different facial expressions of basic emotions using FPVS-EEG and direct contrasts between a given facial expression and all the others. These specific neural markers are of high SNR, reliably elicited in every individual participant in a few minutes of recording, and present with variable topographies, suggesting that different neural substrates underlie the brain coding of each facial expression of emotion. In this context, fear and neutrality elicit quantitatively and qualitatively different brain responses, suggesting particular status for neutral (i.e., no emotion) and fearful (i.e., threatened) faces in the human brain. Finally, the robustness of the present approach offers a unique opportunity to evaluate single-glance facial expression processing with precise issues on specific emotions and/or for various populations.

Acknowledgments

The authors are grateful to Romain PatruX for help in data collection and to Renaud Brochard and the “Institut Universitaire de France” for assistance and financial support in performing EEG. This work received support from the “Conseil Régional Bourgogne Franche-Comté” (PARI grant to FP, JYB and AL, and FABER grant to AL), the FEDER (European Funding for Regional Economic Development) and the French “Investissements d’Avenir” program, project ISITE-BFC (contract ANR-15-IDEX-03). MD is supported by the Fonds National de la Recherche Scientifique (FNRS). The authors declare no competing interest.

References

- 1
2
3
4 Adolphs, R., Tranel, D., Damasio, H., Damasio, A. R., 1994. Impaired recognition of emotion in facial
5 expressions following bilateral damage to the human amygdala. *Nature* 372 (6507), 669–672.
6
7 Adrian, E. D., Matthews, B. H. C., 1934. The Berger rhythm: potential changes from the occipital lobes
8 in man. *Brain* 57 (4), 355-385.
9
10 Ashley, V., Vuilleumier, P., Swick, D., 2004. Time course and specificity of event-related potentials to
11 emotional expressions. *Neuroreport* 15 (1), 211-216.
12
13 Barrett, L. F., Lindquist, K. A., Bliss-Moreau, E., Duncan, S., Gendron, M., Mize, J., Brennan, L., 2007.
14 Of mice and men: natural kinds of emotions in the mammalian brain? A response to
15 Panksepp and Izard. *Perspectives on Psychological Science*, 2 (3), 297–312.
16
17 Barrett, L. F., Wager, T., 2006. The structure of emotion: Evidence from the neuroimaging studies.
18 *Current Directions in Psychological Science*, 15 (2), 79–83.
19
20 Braddick, O. J., Wattam-Bell, J., Atkinson, J., 1986. Orientation-specific cortical responses develop in
21 early infancy. *Nature* 320, 617–619.
22
23 Buchel, C., Dolan, R. J., 2000. Classical fear conditioning in functional neuroimaging. *Current Opinion*
24 *in Neurobiology*, 10 (2), 219–223.
25
26 Calder, A. J., Young, A. W., 2005. Understanding the recognition of facial identity and facial
27 expression. *Nature Reviews Neuroscience* 6, 641–651.
28
29 Calvo, M. G., Nummenmaa, L., 2016. Perceptual and affective mechanisms in facial expression
30 recognition: an integrative review. *Cognition and Emotion* 30 (6), 1081–1106.
31
32 Calvo, M. G., Beltrán, D., 2013. Recognition advantage of happy faces: Tracing the neurocognitive
33 processes. *Neuropsychologia* 51 (11), 2051–2060.
34
35 Calvo, M. G., Marrero, H., Beltrán, D., 2013. When does the brain distinguish between genuine and
36 ambiguous smiles? An ERP study. *Brain and Cognition* 81 (2), 237–246.
37
38 Darwin, C. 1872. *The expression of the emotions in man and animals*. London, UK: John Murray.
39 Retrieved from <http://darwin-online.org.uk>.
40
41 Dzhelyova, M., Rossion, B., 2014a. Supra-additive contribution of shape and surface information to
42 individual face discrimination as revealed by fast periodic visual stimulation. *Journal of Vision*
43 14 (14, 15), 1-14.
44
45 Dzhelyova, M., Rossion, B., 2014b. The effect of parametric stimulus size variation on individual face
46 discrimination indexed by fast period visual stimulation. *BMC Neuroscience* 15 (87), 1-12.
47
48 Dzhelyova, M., Jacques, C., Rossion, B., 2017. At a Single Glance: Fast Periodic Visual Stimulation
49 Uncovers the Spatio-Temporal Dynamics of Brief Facial Expression Changes in the Human
50 Brain. *Cerebral Cortex* 27 (8), 4106-4123.
51
52 Ekman, P., 1992. An argument for basic emotions. *Cognition and Emotion* 6 (3-4), 169-200.
53
54 Ekman, P., 1993. Facial expression and emotion. *American Psychologist* 48 (4), 384-392.
55
56 Ekman, P., 1994. Strong evidence for universals in facial expressions: a reply to Russell's mistaken
57 critique. *Psychological Bulletin* 115 (2), 268–287.
58
59 Ekman, P., Friesen, W. V., 1971. Constants across cultures in the face and emotion. *Journal of*
60 *Personality and Social Psychology* 17 (2), 124-129.
61
62
63
64
65

- 1 Ekman, P., Friesen, W. V., Ellsworth, P., 1972. *Emotion in the Human Face: Guidelines for Research*
2 *and an Integration of Findings*. New York: Pergamon Press.
- 3 Elfenbein, H. A., Ambady, N., 2002. On the universality and cultural specificity of emotion
4 recognition: a meta-analysis. *Psychological Bulletin* 128 (2), 203–235.
- 5 Etcoff, N. L., Magee, J. J., 1992. Categorical perception of facial expressions. *Cognition* 44 (3), 227-
6 240.
- 7
8 George, N., Driver, J., Dolan, R. J., 2001. Seen gaze-direction modulates fusiform activity and its
9 coupling with other brain areas during face processing. *NeuroImage* 13 (6), 1102–1112.
- 10
11 Harris, R. J., Young, A. W., Andrews, T. J., 2012. Morphing between expressions dissociates
12 continuous from categorical representations of facial expression in the human brain.
13 *Proceedings of the National Academy of Sciences* 109 (51), 21164–21169.
- 14
15 Harris, R. J., Young, A. W., Andrews, T. J., 2014. Dynamic stimuli demonstrate a categorical
16 representation of facial expression in the amygdala. *Neuropsychologia* 56, 47-52.
- 17
18 Haxby, J. V., Hoffman, E. A., Gobbini, M. I., 2000. The distributed human neural system for face
19 perception. *Trends in Cognitive Sciences* 4 (6), 223–233.
- 20
21 Kawasaki, H., Tsuchiya, N., Kovach, C. K., Nourski, K. V., Oya, H., Howard, M. A., Adolphs, R., 2012.
22 Processing of facial emotion in the human fusiform gyrus. *Journal of Cognitive Neuroscience*
23 24 (6), 1358–1370.
- 24
25 Kim, Y. J., Grabowecy, M., Paller, K. A., Muthu, K., Suzuki, S., 2007. Attention induces
26 synchronization-based response gain in steady-state visual evoked potentials. *Nature*
27 *Neuroscience* 10 (1), 117–125.
- 28
29 Kim, H., Somerville, L., Johnstone, T., Polis, S., Alexander, A., Shin, L., Whalen, P., 2004. Contextual
30 modulation of amygdala responsivity to surprised faces. *Journal of Cognitive Neuroscience* 16
31 (10), 1730–1745.
- 32
33 Krolak-Salmon, P., Henaff, M. A., Vighetto, A., Bertrand, O., Mauguiere, F., 2004. Early amygdala
34 reaction to fear spreading in occipital, temporal, and frontal cortex: A depth electrode ERP
35 study in human. *Neuron* 42 (4), 665–676.
- 36
37 Leleu, A., Dzhelyova, M., Rossion, B., Brochard, R., Durand, K., Schaal, B., Baudouin J.-Y., 2018. Tuning
38 functions for automatic detection of brief changes of facial expression in the human
39 brain. *NeuroImage* 179, 235-251.
- 40
41 Leleu, A., Favre, E., Yailian, A., Fumat, H., Klamm, J., Amado, I., Baudouin, J.-Y., Franck, N., Demily, C.,
42 2019. An implicit and reliable neural measure quantifying impaired visual coding of facial
43 expression: evidence from the 22q11.2 deletion syndrome. *Translational Psychiatry* 9, 67.
- 44
45 Leleu, A., Godard, O., Dollion, N., Durand, K., Schaal, B., Baudouin, J.-Y., 2015. Contextual odors
46 modulate the visual processing of emotional facial expressions: An ERP study.
47 *Neuropsychologia* 77, 366-379.
- 48
49 Leppänen, J. M., Kauppinen, P., Peltola, M. J., Hietanen, J. K., 2007. Differential electrocortical
50 responses to increasing intensities of fearful and happy emotional expressions. *Brain*
51 *Research* 1166, 103–109.
- 52
53 Liu-Shuang, J., Norcia, A. M., Rossion, B., 2014. An objective index of individual face discrimination in
54 the right occipito-temporal cortex by means of fast periodic oddball stimulation.
55 *Neuropsychologia* 52, 57-72.
- 56
57
58
59
60
61
62
63
64
65

- 1 Liu-Shuang, J., Torfs, K., Rossion, B., 2016. An objective electrophysiological marker of face
2 individualisation impairment in acquired prosopagnosia with fast periodic visual stimulation.
3 *Neuropsychologia* 83, 100-113.
- 4 Luo, W., Feng, W., He, W., Wang, N.-Y., Luo, Y.-J., 2010. Three stages of facial expression processing:
5 ERP study with rapid serial visual presentation. *NeuroImage* 49 (2), 1857–1867.
- 6
7 Makeig, S., Bell, A. J., Jung, T.-P., Sejnowski, T.J., 1996. Independent component analysis of
8 electroencephalographic data. In: D. Touretzky, M. Mozer & M. Hasselmo (Eds.) *Advances in*
9 *Neural Information Processing Systems* 8, 145–151. Cambridge, MA: MIT Press.
- 10
11 Malaia, E., Cockerham, D., Rublein, K., 2017, Visual integration of fear and anger emotional cues by
12 children on the autism spectrum and neurotypical peers: An EEG study. *Neuropsychologia* (in
13 press).
- 14
15 McCarthy, G., Wood, C. C., 1985. Scalp distributions of event-related potentials: An ambiguity
16 associated with analysis of variance models. *Electroencephalography and Clinical*
17 *Neurophysiology* 62 (3), 203–208.
- 18
19 Morris, J. S., Friston, K. J., Buchel, C., Frith, C. D., Young, A. W., Calder, A. J., et al., 1998. A
20 neuromodulatory role for the human amygdala in processing emotional facial expressions.
21 *Brain* 121, 47–57.
- 22
23 Norcia, A. M., Appelbaum, L. G., Ales, J. M., Cottareau, B. R., Rossion, B., 2015. The steady-state
24 visual evoked potential in vision research: A review. *Journal of Vision* 15 (6): 4, 1-46.
- 25
26 Öhman, A., Wiens, S., 2004. The concept of an evolved fear module and cognitive theories of
27 anxiety. In A. S. R. Manstead, N. Frijda & A. Fischer (Eds.) *Feelings and Emotions: The*
28 *Amsterdam Symposium*. Cambridge: Cambridge University Press.
- 29
30 Pessoa, L., McKenna, M., Gutierrez, E., Ungerleider, L. G., 2002. Neural processing of emotional faces
31 requires attention. *Proceedings of the National Academy of Sciences of the United States of*
32 *America* 99 (17), 11458–11463.
- 33
34 Pourtois, G., Spinelli, L., Seeck, M., Vuilleumier, P., 2010. Modulation of face processing by emotional
35 expression and gaze direction during intracranial recordings in right fusiform cortex. *Journal*
36 *of cognitive neuroscience* 22 (9), 2086-2107.
- 37
38 Puce, A., Syngienotis, A., Thompson, J. C., Abbott, D. F., Wheaton, K. J., Castiello, U., 2003. The
39 human temporal lobe integrates facial form and motion: evidence from fMRI and ERP
40 studies. *NeuroImage* 19 (3), 861–869.
- 41
42 Regan, D., 1989. *Human Brain Electrophysiology: Evoked Potentials and Evoked Magnetic Fields in*
43 *Science and Medicine*. Elsevier, New York.
- 44
45 Retter, T. L., Rossion, B., 2016. Uncovering the neural magnitude and spatio-temporal dynamics of
46 natural image categorization in a fast visual stream. *Neuropsychologia* 91, 9–28.
- 47
48 Rossion, B., Torfs, K., Jacques, C., Liu-Shuang, J., 2015. Fast periodic presentation of natural images
49 reveals a robust face-selective electrophysiological response in the human brain. *Journal of*
50 *Vision* 15 (1): 18, 1-18.
- 51
52 Rossion, B., 2014a. Understanding face perception by means of human electrophysiology. *Trends in*
53 *Cognitive Sciences* 18 (6), 310-318.
- 54
55 Rossion, B., 2014b. Understanding individual face discrimination by means of fast periodic visual
56 stimulation. *Experimental Brain Research* 232 (6), 1599-1621.
- 57
58 Russell, J. A., 1980. A circumplex model of affect. *Journal of Personality and Social Psychology* 39,
59 1161-1178.
- 60
61
62
63
64
65

- 1 Said, C. P., Moore, C. D., Engell, A. D., Todorov, A., Haxby, J. V., 2010. Distributed representations of
2 dynamic facial expressions in the superior temporal sulcus. *Journal of Vision* 10 (5): 11, 1-12.
- 3 Sauter, D. A., Eisner, F., 2013. Commonalities outweigh differences in the communication of
4 emotions across human cultures. *Proceedings of the National Academy of Sciences of the*
5 *United States of America* 110 (3), E180.
- 6 Srinivasan, R., Golomb, J. D., Martinez, A. M., 2016. A neural basis of facial action recognition in
7 humans. *The Journal of Neuroscience*. 36 (16), 4434–4442.
- 8 Tottenham, N., Tanaka, J. W., Leon, A. C., McCarry, T., Nurse, M., Hare, T. A., Nelson, C. A., 2009. The
9 NimStim set of facial expressions: judgments from untrained research participants. *Psychiatry*
10 *Research* 168 (3), 242–249.
- 11 Vettori, S., Dzhelyova, M., Van der Donck, S., Jacques, C., Steyaert, J., Rossion, B., Boets, B., 2018.
12 Reduced neural sensitivity to rapid individual face discrimination in autism spectrum
13 disorder. *NeuroImage: Clinical* (in press).
- 14 Vuilleumier, P., Pourtois, G., 2007. Distributed and interactive brain mechanisms during emotion face
15 perception: evidence from functional neuroimaging. *Neuropsychologia* 45 (1), 174–194.
- 16 Vuilleumier, P., Armony, J. L., Driver, J., Dolan, R. J., 2001. Effects of attention and emotion on face
17 processing in the human brain: An event-related fMRI study. *Neuron* 30 (3), 829–841.
- 18 Vuilleumier, P., Richardson, M. P., Armony, J. L., Driver, J., Dolan, R. J., 2004. Distant influences of
19 amygdala lesion on visual cortical activation during emotional face processing. *Nature*
20 *Neuroscience* 7 (11), 1271–1281.
- 21 Vytal, K., Hamann, S., 2010. Neuroimaging support for discrete neural correlates of basic emotions: A
22 voxel-based meta-analysis. *Journal of Cognitive Neuroscience* 22 (12), 2864–2885.
- 23 Whalen, P. J., Raila, H., Bennett, R., Mattek, A., Brown, A., Taylor, J., et al., 2013. Neuroscience and
24 facial expressions of emotion: The role of amygdala-prefrontal interactions. *Emotion Review*
25 *5* (1), 78–83.
- 26 Williams, L. M., Palmer, D., Liddell, B. J., Song, L., Gordon, E., 2006. The ‘when’ and ‘where’ of
27 perceiving signals of threat versus non-threat. *NeuroImage* 31 (1), 458–467.
- 28 Woodworth, R. S., Schlosberg, H., 1954. *Experimental Psychology: Revised Edition*. Henry Holt, New
29 York.
- 30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure captions

Figure 1. Stimuli and experimental design. Example of one individual female face with a neutral expression or expressing 5 basic emotions (anger, disgust, fear, happiness and sadness). Faces were presented through sinusoidal contrast variation and 10 % randomized size variation with all emotions randomly displayed at a 6 Hz base rate (1 cycle \approx 167ms) except for one which was periodically inserted at a lower rate of 1.2 Hz (every 5th cycle \approx 833ms between each expression). This design thus isolates the specific response to a given emotion at the 1.2 Hz frequency regardless of expression-changes occurring at the 6 Hz frequency.

Figure 2. Expression-specific and general visual response for each facial expression. A. Expression-specific response for each facial expression. Left part: FFT signal-to-noise ratio (SNR) spectra for channels included in the regions-of-interest (ROIs) and for the 6 facial expressions. SNR for the expression-specific response are visible at the 1.2 Hz frequency and its harmonics (i.e., integer multiples, e.g., 2.4 Hz, 3.6 Hz). Middle part: FFT signal-to-noise ratio (SNR) spectra calculated on the sum of significant harmonics of the 1.2 Hz expression-specific rate (F: until the 7th harmonic, i.e. 8.4 Hz, excluding the 6 Hz base frequency) for channels included in the ROIs and for the 6 facial expressions. Note the high SNR for each facial expression except happiness, with large responses for fear and neutrality (SNR \approx 1.8, 80 % of signal increase) and weaker but still identifiable responses for the other expressions (SNR \approx 1.4, 40 % of signal increase). Right part: 3D-topographical maps (posterior view) of summed baseline-corrected amplitudes (BCA) for the expression-specific response. B. General visual response elicited at the base rate. Left part: FFT signal-to-noise ratio (SNR) spectra calculated on the sum of significant harmonics of the 6 Hz base rate (F: until the 8th harmonic, i.e. 48 Hz) for channels included in the regions-of-interest (ROIs) and for the 6 facial expressions. Note that SNR is high and does not differ between the 6 facial expressions (SNR \approx 7, 700 % of single increase). Right part: 3D-topographical maps (posterior view) of summed baseline-corrected amplitudes (BCA) for the general visual response.

Figure 3. Scalp topography of the expression-specific response. Upper part: 3D-topographical maps (posterior view) of normalized summed baseline-corrected amplitudes (BCA) of the expression-specific response for each facial expression. The scaling of the response equalizes global amplitude across expressions and reveals specific topographical patterns for each expression category. Bottom

part: posterior electrodes with a significant ($Z > 1.64$, $p < .05$, one-tailed, signal > noise) response. For each expression, at least 4 electrodes were significant, and no electrode significantly responded to all expression categories. Note that neutrality and fear presented numerous significant electrodes. The response to fear was restricted to occipito-temporal channels compared with all other expressions.

Figure 4. Individual data: expression-specific response recorded for each participant. Left part: 3D-topographical maps (posterior view) of the expression-specific response averaged across the 6 facial expressions (amplitude is scaled for each participant and individual maximum amplitudes are indicated above each map). Right part: 3D-topographical maps (posterior view) of the electrodes presenting a significant (i.e., $Z > 1.64$, $p < .05$, one-tailed, signal > noise) expression-specific response averaged across facial expressions for each individual participant. Electrodes in black are those included in the ROIs, those in gray are other posterior electrodes.

Figure 1
[Click here to download high resolution image](#)




Figure 2
[Click here to download high resolution image](#)




Figure 3
[Click here to download high resolution image](#)





Figure 4
[Click here to download high resolution image](#)



Author contributions

FP, JYB, MD, BR and AL designed research; FP, JYB and AL collected data; FP, JYB and AL analyzed data; FP, JYB and AL wrote the paper; FP, JYB, MD, BR and AL revised the paper.