



HAL
open science

Congruence between fine-scale genetic breaks and dispersal potential in an estuarine seaweed across multiple transition zones

Katy R Nicastro, Jorge Assis, Ester Álvares Serrão, Gareth A. Pearson, João Neiva, Myriam Valero, Rita Jacinto, Gerardo I Zardi

► **To cite this version:**

Katy R Nicastro, Jorge Assis, Ester Álvares Serrão, Gareth A. Pearson, João Neiva, et al.. Congruence between fine-scale genetic breaks and dispersal potential in an estuarine seaweed across multiple transition zones. *ICES Journal of Marine Science*, 2020, 77 (1), pp.371-378. 10.1093/icesjms/fsz179 . hal-02406672

HAL Id: hal-02406672

<https://hal.science/hal-02406672>


Submitted on 12 Dec 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Congruence between fine-scale genetic breaks and dispersal potential in an estuarine seaweed across multiple transition zones.

Journal:	<i>ICES Journal of Marine Science</i>
Manuscript ID	ICESJMS-2019-167.R2
Manuscript Types:	Original Article
Date Submitted by the Author:	04-Sep-2019
Complete List of Authors:	Nicastro, Katy; CCMAR Assis, Jorge; CCMAR Serrão, Ester; University of Algarve, CCMAR- Centre of Marine Sciences Pearson, Gareth; CCMAR Neiva, Joao; CCMAR Jacinto, Rita; CCMAR Valero, Myriam; CNRS Zardi, Gerardo; Rhodes University, Dept Zoology and Entomology
Keyword:	Biogeography, physical modelling, gene flow, <i>Fucus</i> spp.

SCHOLARONE™
Manuscripts

1
2
3 1 **Congruence between fine-scale genetic breaks and dispersal potential in an**
4
5 2 **estuarine seaweed across multiple transition zones.**
6
7
8 3

9
10 4 Katy R. Nicastró¹, Jorge Assis¹, Ester A. Serrão¹, Gareth A. Pearson¹, João Neiva¹, Myriam
11
12 5 Valero², Rita Jacinto¹, Gerardo I. Zardi^{3*}
13
14

15 6 ¹CCMAR-CIMAR – Associated Laboratory, University of Algarve, Campus de Gambelas,
16 7 Faro, 8005–139, Portugal

18 8 ²UMI 3614 Evolutionary Biology and Ecology of Algae, CNRS, Sciences Sorbonne
19 9 Université, UC, UACH, Station Biologique de Roscoff, Roscoff Cedex, France

21
22 10 ³Department of Zoology and Entomology, Rhodes University, Grahamstown, 6140, South
23 11 Africa

24 12 *corresponding author: zardi73@yahoo.it, +351 927738340
25
26
27 13

28
29
30 14 **Additional emails:**

31
32 15 KRN: katynicastro@gmail.com
33
34

35 16 JA: jorgemfa@me.com
36
37

38 17 EAS: eserrao@ualg.pt
39
40

41 18 GAP: gpearson@ualg.pt
42
43

44 19 JN: jmneiva@ualg.pt
45
46

47 20 MV: myriam.valero@sb-roscoff.fr
48
49

50 21 RJ: ritajacinto85@gmail.com
51
52 22

53
54 23 **Running head:** Congruence between fine-scale genetic breaks and dispersal potential.
55
56

57 24 **Keywords:** Biogeography, physical modelling, gene flow, *Fucus* spp.
58
59
60

25 **ABSTRACT**

26 Genetic structure in biogeographical transition zones can be shaped by several factors
27 including limited dispersal across barriers, admixture following secondary contact, differential
28 selection and mating incompatibility. A striking example is found in Northwest France and
29 Northwest Spain , where the estuarine seaweed *Fucus ceranoides* L. exhibits sharp, regional
30 genetic clustering. This pattern has been related to historical population fragmentation and
31 divergence into distinct glacial refugia, followed by post-glacial expansion and secondary
32 contact. The contemporary persistence of sharp ancient genetic breaks between nearby
33 estuaries has been attributed to prior colonization effects (density barriers) but the effect of
34 oceanographic barriers has not been tested. Here, through a combination of mesoscale
35 sampling (15 consecutive populations) and population genetic data (mtlGS) in NW France,
36 we define regional genetic disjunctions similar to those described in NW Iberia. Most
37 importantly, using high resolution dispersal simulations for Brittany and Iberian populations,
38 we provide evidence for a central role of contemporary hydrodynamics in maintaining genetic
39 breaks across these two major biogeographic transition zones. Our findings further show the
40 importance of a comprehensive understanding of oceanographic regimes in hydrodynamically
41 complex coastal regions to explain the maintenance of sharp genetic breaks along
42 continuously populated coastlines.

44 **INTRODUCTION**

45 Fueled by current concerns about the impacts of global change on biodiversity, there is a
46 renewed interest in understanding the processes affecting a species' range dynamics as well
47 as the factors shaping its genetic diversity. Generally, both historical and contemporary
48 processes are invoked to explain the distribution and phylogeography of a species. Tectonic
49 events and past climate fluctuations created topographic, environmental and hydrologic
50 barriers that were key in shaping phylogeographic structure of many plant and animal species.
51 Of these, the Last Glacial Maximum is probably the most significant and recent historical event

1
2
3 52 (e.g., Hewitt, 2000; Barnes et al., 2002; Liang et al., 2017; Neiva et al., 2018). During this time,
4
5 53 ice sheets covered much of current cold and temperate zones of the Northern Hemisphere,
6
7 54 triggering range fragmentation and contractions of entire ecosystems to more southern
8
9 55 latitudes, while during the present interglacial, widespread range shifts and poleward
10
11 56 expansions have occurred.

12
13 57 It is predicted that genetic diversity is lowest in recently colonized areas and highest in refugial
14
15 58 areas where long-term persistence was possible (e.g., Hewitt, 1996; Comps et al., 2001;
16
17 59 Hewitt, 2004; Kennedy et al., 2017). However, contemporary demographic processes can
18
19 60 either mask or even erase genetic signatures of population expansions or retreats (Smith et
20
21 61 al., 2011). High levels of gene flow may homogenize genetic diversity among previously
22
23 62 structured populations. On the other hand, nonrandom mating with individuals in close
24
25 63 geographic proximity can generate genetic structuring within a continuous population (Slatkin,
26
27 64 1993). Intrinsic characteristics of the species such as dispersal ability, migration patterns and
28
29 65 changes in population densities can further confound any historical signature (e.g., Mims et al.,
30
31 66 2015; Chust et al., 2016; Assis et al., 2018)

32
33 67 Biogeographical transition regions provide a good framework for exploring and understanding
34
35 68 genetic structuring of species through space and time. These are areas of overlap and
36
37 69 segregation between different biotic components, and geographically separated clades often
38
39 70 coincide with these transition zones (Ferro and Morrone, 2014). In the marine realm,
40
41 71 phylogeographic breaks in biogeographical transition regions are generally attributed to
42
43 72 historical processes or contemporary dispersal barriers, such as upwelling phenomena and
44
45 73 currents that may limit along-shore dispersal, environmental differences boosting local
46
47 74 adaptation and/or reproductive strategies maintaining self-recruitment (Gilg and Hilbish, 2003;
48
49 75 Zardi et al., 2007; Selkoe et al., 2010).

50
51 76 There are well known biogeographical boundaries in all oceans, some of which are associated
52
53 77 with oceanographic features that generate contemporary barriers to dispersal. For example,
54
55 78 the strong southward-flowing Agulhas Current, which runs about 10 km offshore along most
56
57 79 of South Africa's east and southeast coast, impedes larval dispersal and thus promotes local

1
2
3 80 adaptation (Teske et al., 2011; Zardi et al., 2011). Dispersal and gene flow around Point
4
5 81 Conception in southern California are also strongly affected by extensive upwelling of colder
6
7 82 sub-surface waters and by the southward California Current (Wares et al., 2001; Hohenlohe,
8
9 83 2004; Johansson et al., 2015).

10
11 84 Here, we investigated the contribution of contemporary oceanographic connectivity in
12
13 85 maintaining geographically separated genetic clades within a continuously distributed
14
15 86 seaweed species along the biogeographic transition areas of Brittany (northwest France,
16
17 87 Europe) and northwest Iberia (northern Spain, Europe; Fig. 1). These two areas are highly
18
19 88 relevant for studying this question and marine connectivity in general. Both areas delimit the
20
21 89 boundaries between cold-temperate and warm-temperate regions (Spalding et al., 2007), they
22
23 90 are refugial zones for numerous species (Provan, 2013) and, over the past decades, they
24
25 91 have undergone significant changes in macroalgae assemblages due to climate change
26
27 92 (Nicastro et al., 2013; Gallon et al., 2014; Assis et al., 2017).

28
29
30 93 In this study, we use phylogeographic analyses and Lagrangian Particle Simulations
31
32 94 (LPS) coupled with network analyses to evaluate the levels of congruence between dispersal
33
34 95 potential and the patterns of present-day genetic differentiation and diversity in the estuarine
35
36 96 seaweed *Fucus ceranoides* along these two biogeographic transition zones. This species is
37
38 97 perennial, dioecious and restricted to estuarine intertidal areas. Furoid seaweeds have no
39
40 98 planktonic dispersal stage and have restricted gamete dispersal (Serrão et al., 1997).
41
42 99 However, adult individuals can achieve long distance gene flow via the rafting of whole or
43
44 100 partially detached thalli with reproductive structures (Thiel and Haye, 2006; McKenzie and
45
46 101 Bellgrove, 2008), a form of population connectivity strongly influenced by hydrodynamic forces
47
48 102 and coastal topography. Currently, *F. ceranoides* is distributed from northern Portugal to
49
50 103 northern Norway (Lein, 1984) and Iceland (Munda, 1999), covering both past non-glaciated
51
52 104 and glaciated regions of Europe. Previous studies have shown the occurrence of two divergent
53
54 105 genetic lineages in this species between the south and the north of Europe, the
55
56 106 phylogeographic break being localized in Northwest France near the English Channel (Neiva
57
58 107 et al., 2012a; Neiva et al., 2012b). Specifically, the dominant Northwest Iberian haplotypes of

1
2
3 108 this species was found in southern Brittany but not after the genetic break in northern Brittany
4
5 109 (Neiva et al., 2010). We discuss three scenarios for the concordance between present-day
6
7 110 oceanographic dispersal barriers and the observed genetic differentiation among *F.*
8
9 111 *ceranoides* estuarine populations: 1) Contemporary oceanographic barriers to dispersal are
10
11 112 responsible for the origin, the position and the maintenance (i.e. delaying genetic
12
13 113 homogenization) of the genetic differentiation; 2) Contemporary oceanographic circulation
14
15 114 patterns explain the position, the maintenance but not the origin of the genetic differentiation;
16
17 115 3) Reproductive isolation, in addition to contemporary oceanographic patterns, limits genetic
18
19 116 homogenisation thus contributing to the position and the maintenance of the genetic
20
21 117 differentiation.
22
23
24 118

27 119 **MATERIALS**

30 120 **Genetic data**

31
32 121 The two study areas (NW France and NW Spain) are peninsulas characterized by complex
33
34 122 and variable circulation patterns (Puillat et al., 2004; Varela et al., 2005; Ruiz-Villarreal et al.,
35
36 123 2006; Ayata et al., 2011). Because of that, sampling of only a limited number of localities would
37
38 124 be inappropriate. We therefore decided to sample a small number of individuals at as many
39
40 125 sites along the two coastlines as possible, rather than obtaining large numbers of individuals
41
42 126 from a limited number of sites (e.g., Sotka et al., 2004; Teske et al., 2007). This decision was
43
44 127 also supported by the very low genetic diversity within localities for this species (Neiva et al.,
45
46 128 2010; 2012b; Neiva et al., 2012a). As a result, two sets of sequences of *Fucus ceranoides*
47
48 129 were prepared and analyzed separately. The first data set (Data set 1) comprised sequences
49
50 130 of individuals collected in the estuaries of all major rivers between Hennebont (HB, southern
51
52 131 Brittany) and Camarel (CM, northern Brittany) in northwest France (n = 2 from each site;
53
54 132 Table SI1). The second dataset (Data set 2) consisted of sequences previously analysed in
55
56 133 Neiva et al. (Neiva et al., 2010; 2012b; Neiva et al., 2012a) and sampled across northern Iberia
57
58 134 between Viana do Castelo (VIA, northern Portugal) and Porcia (POR, northern Spain). To
59
60

1
2
3 135 allow a comparison between data sets, a random subsample of 2 individuals from each site
4
5 136 was used for Data set 2. The random subsampling in NW Spain was repeated to ensure
6
7 137 consistency in the results obtained.
8

9 138 Sampling was performed in 2014-2015 and conducted with similar *criteria* for individuals used
10
11 139 in both data sets. All collection sites were characterized by monospecific belts of *F.*
12
13 140 *ceranoides* attached to hard substrata and were exposed to steep salinity fluctuations
14
15 141 throughout the tidal cycle. At each site, 5-10 cm tips of apical vegetative tissue were collected
16
17 142 from individuals sampled in the mid distributional range of the species. Neighboring sites were
18
19 143 at an average proximity of about 50 (± 15) and 33 (± 17) km for Data sets 1 and 2 respectively.
20
21 144 All samples were individually stored dehydrated in silica-gel crystals until DNA extraction.
22

23 145 *DNA isolation and sequencing of data set 1*

24
25
26 146 To compile Data set 1, genomic DNA was extracted from approximately 10 mg of dried tissue
27
28 147 using the Nucleospin® Multi-96 plant kit (Macherey-Nagel Duren, Germany), according to the
29
30 148 manufacturer's protocol. Individuals were sequenced for the mitochondrial 23 S/trnK
31
32 149 intergenic spacer (mtIGS, Neiva et al., 2010). Primer sequences and amplification details were
33
34 150 the same as in Neiva et al. (Neiva et al., 2010; 2012b; Neiva et al., 2012a). Amplified fragments
35
36 151 were run in an ABI PRISM 3130xl automated capillary sequencer (Applied Biosystems,
37
38 152 CCMAR Portugal). MtDNA sequences were aligned, proofread and edited in GENEIOUS 3.8
39
40 153 (Drummond et al., 2011).
41
42

43 154 *Data analyses*

44
45 155 For both data sets, haplotype frequencies were estimated using DnaSP 5.0 (Librado and
46
47 156 Rozas, 2009). The relationships among the MtIGS haplotypes were inferred using statistical
48
49 157 parsimony with Tcs v. 1.13 (Clement et al., 2000). Because additional subsampled dataset for
50
51 158 the Iberia provided similar results (Table SI2 and Fig. SI1), only one was used for the
52
53 159 simulations.
54

55 160 **Dispersal simulations**

56
57 161 The main oceanographic regions in northwest France and northwest Iberia (~550km and
58
59 162 ~600km of coastlines, respectively) were identified by coupling Lagrangian Particle
60

1
2
3 163 Simulations (LPS) with network analyses (least cost distance and community algorithm, e.g.,
4
5 164 Assis et al., 2015; Klein et al., 2016; Assis et al., 2018). The simulations used daily data of
6
7 165 ocean currents assembled from the Hybrid Coordinate Ocean Model (HYCOM), a resolution
8
9 166 product with a spatial resolution of 0.08° (approx. 6–9 km), forced by wind speed, wind stress,
10
11 167 precipitation and heat flux. This model can resolve oceanic fronts, meandering currents,
12
13 168 filaments and eddies (Chassignet et al., 2007), important mesoscale processes to properly
14
15 169 simulate ocean dispersal (Assis et al., 2015; Klein et al., 2016).

16
17
18 170 Both regions of simulation were gridded to a common spatial resolution of 0.01° (approx. 1km).
19
20 171 A polygon representing global coastlines - OpenStreetMap geographic information (Haklay
21
22 172 and Weber, 2008)- was used to define intertidal source and sink cells. Passive particles
23
24 173 simulating rafts of *F. ceranoides* adult individuals were released from each gridded cell every
25
26 174 12 hours and allowed to drift for 60 full days; an extreme period for long-lived rafts of brown
27
28 175 macroalgae (Monteiro et al., 2016; Assis et al., 2018). This approach aimed to capture the
29
30 176 rare, long-distance dispersal events, allowing gene flow at the scales of both regions (Monteiro
31
32 177 et al., 2016; Assis et al., 2018). After the 60 days period, or when ending up on shore, the
33
34 178 particles were removed from the simulation. The geographic position of all particles was
35
36 179 calculated every hour with bilinear interpolation on the ocean velocity fields (with a spatial
37
38 180 resolution of 0.08°), while combining a 4th Order Runge-Kutta adaptive time-step on the path
39
40 181 equations (e.g., Lett et al., 2008; Klein et al., 2016).

41
42
43 182 The degree of connectivity between all pairs of gridded cells was determined by dividing the
44
45 183 number of unique particles released from cell *i* that ended up in cell *j*, by the total number of
46
47 184 particles released from cell *i*. To account for the inter-annual variability in the ocean data, the
48
49 185 simulations ran independently per year (from January to December), for the most recent 10-
50
51 186 year period of data, available in HYCOM (i.e., 2003 to 2012). Asymmetrical connectivity
52
53 187 matrices were determined by averaging the outcomes of the annual simulations.

54
55
56 188 The connectivity matrices were used in network analyses (i.e., graph theory) to infer the major
57
58 189 oceanographic regions of northwest Iberia and northwest France. To this end, network
59
60 190 percolation removed weak probabilities to a threshold maintaining all cells (nodes) connected

1
2
3 191 into a single network (Rozenfeld et al. 2008; Cunha et al., 2017), while maximizing modularity,
4
5 192 which quantified the strength of the backbone structure (or goodness of fit) of the networks
6
7 193 (Newman, 2006). This allowed the removal of surplus connections with unimportant
8
9 194 information. The leading eigenvector algorithm (Newman, 2006) was applied to the percolated
10
11 195 networks to assign a unique membership to the nodes. This approach allowed the detection
12
13 196 of communities in the networks (e.g., Munwes et al., 2010), which in practice translated into a
14
15 197 delineation of oceanographic regions in northwest Iberia and northwest France structured by
16
17 198 connectivity of ocean currents (Assis et al., 2018). The statistical significance of the
18
19 199 membership assignment to the nodes was inferred by testing the proportion of 9999
20
21 200 membership randomizations that retrieved a higher modularity than that observed.
22
23

24 201 Lagrangian Particle Simulations and network analyses were performed in R (R Development
25
26 202 Core Team, 2016) using the packages: data.table (Dowle et al., 2019), dismo (Hijmans et al.,
27
28 203 2017), igraph (Kamvar et al., 2014), parallel (Team, 2018), raster (Lamigueiro et al., 2018)
29
30 204 and vegan (Oksanen et al., 2018).
31
32
33
34

35 206 **RESULTS**

36
37 207 In Data set 1 (NW France), six mtIGS haplotypes were identified in 30 individuals of *Fucus*
38
39 208 *ceranoides* in 15 sampled sites. In NW France, the network analyses showed two dominant
40
41 209 haplotypes (A and D) plus three derived ones (B, C and E), each private to one population
42
43 210 (Fig 1a). Haplotype frequency distribution revealed a geographical segregation of the two main
44
45 211 haplotypes. Haplotype 1 was present in individuals from regions R1 and R2 while Haplotype
46
47 212 D was restricted to more northern sites within region R3.

48
49 213 Out of 415 sequences retrieved from the GenBank, 52 sequences were randomly selected for
50
51 214 Data set 2 (northwest Iberia). The network showed seven main haplotypes of which two were
52
53 215 shared among six to ten populations and three were shared by two to three populations. The
54
55 216 remaining two haplotypes were private to one single population. In the frequency distribution,
56
57 217 the main haplotype was restricted to region R5 and haplotype F was present only in region
58
59
60

1
2
3 218 R4. Haplotype A was present in regions R3 and R1 while haplotype B was confined to region

4
5 219 **R2. Dispersal simulations**

6
7 220 The LPS using HYCOM ocean currents over the 10-year period released 7300 particles per
8
9 221 cell (7.80×10^6 and 7.88×10^6 particles in total in northwest France and northwest Iberia,
10
11 222 respectively). Particles drifted for longer distances in northwest Iberia than in northwest France
12
13 223 (up to 431.2 km; Table 1; Fig. SI2). The maximum period of drifting time was also higher in
14
15 224 northwest Iberia (26.7 days), but on average, particles drifted for longer periods in northwest
16
17 225 France (3.61 ± 2.63 days). The maximum probability of connectivity between the pairs of cells
18
19 226 was observed in northwest France (western Brittany; 0.721). However, the average cell
20
21 227 probabilities within regions did not vary considerably (Table 1; Fig. SI2).

22
23
24 228 The assignment of oceanographic regions performed by the leading eigenvector algorithm
25
26 229 (network analysis) for northwest France and northwest Iberia showed significant modularity
27
28 230 values of 0.41 and 0.57 (p-values < 0.001), respectively. The algorithm identified three regions
29
30 231 in northwest France (Fig. 1a), with breaks in Penmarch and Porspoder, and five regions in
31
32 232 northwest Iberia (Fig. 1b), with breaks in Ria de Arousa, Corrubedo, Camelle (northern Costa
33
34 233 da Morte) and Cabo Ortegal (Cariño). The average probability of connectivity within the
35
36 234 oceanographic regions of northwest France (diagonal of Fig. 1d) was 10-fold higher than
37
38 235 between regions, with the exception of those between R1 and R2 (Fig. 1d), which were of the
39
40 236 same order of magnitude. Following a similar pattern, the probabilities of connectivity in
41
42 237 northwest Iberia were 10 to 100-fold higher within regions than between regions (diagonal of
43
44 238 Fig. 1e), with a marked increase as the distance between groups increased.

45
46
47
48 239

49
50
51 240 **DISCUSSION**

52
53
54 241 Our results show an overall strong match between oceanographic regions identified
55
56 242 by dispersal simulations and fine-scale genetic discontinuities in the estuarine seaweed *Fucus*
57
58 243 *ceranoides* inhabiting north Atlantic shores (Fig. 1).

1
2
3 244 Predictions of oceanographic transport made with Lagrangian particle simulations are
4
5 245 highly sensitive to the inner spatial and temporal resolution of circulation models (Putman and
6
7 246 He, 2013). While we preserved the raw circulation processes modelled by Hycom at the scales
8
9 247 of days and tens of kilometers, a main limitation may arise if additional oceanographic
10
11 248 processes occurring at smaller scales are important for realistic connectivity events shaping
12
13 249 the genetic structure of *F. ceranoides*. For instance, circulation models with coarse spatial
14
15 250 resolutions may underestimate drifting times up to a factor of ~2. In the same way, weekly or
16
17 251 even daily temporal time steps may not be a realistic representation of what organisms
18
19 252 continuously experience (Fossette et al., 2012). Regardless of such potential limitations, our
20
21 253 results compared with independent genetic data, as well as additional studies comparing
22
23 254 connectivity estimates with satellite-tracked organisms (e.g., Fossette et al., 2012) suggest
24
25 255 that, overall, particles advected by Hycom data provide a reliable estimate of the main
26
27 256 processes shaping current flow.

28
29
30 257 Previous studies have shown that mtIGS differentiation and contemporary *F.*
31
32 258 *ceranoides* genetic structuring in NW Iberia is most likely the result of complex, past range
33
34 259 dynamics (Neiva et al., 2012a). High levels of endemism and diversity highlighted by genetic
35
36 260 analyses of *F. ceranoides* populations inhabiting this region indicate long-term persistence in
37
38 261 glacial refugia. Despite the refugial role played by NW Iberia, sea level changes associated
39
40 262 with glacial/inter-glacial expansion and melting of ice-sheets had significant effects on near-
41
42 263 shore habitats (Chao et al., 2002; Roucoux et al., 2005). The current fine-scale genetic breaks
43
44 264 in Iberian *F. ceranoides* are the result of past fragmentation and divergence of populations
45
46 265 into distinct refugia (estuarine refugia within regional refugia), followed by expansion and
47
48 266 secondary contact of vicariant phylogroups (Neiva et al., 2012b).

49
50
51 267 In addition to southern European refugia, the ice-free paleo-shores of northwest
52
53 268 France (western Brittany) together with south western Ireland and the English Channel have
54
55 269 been recognized as northern periglacial refugia for several species (*Palmaria palmata* (Provan
56
57 270 et al., 2005); *Celleporella hyalina* (Gomez et al., 2007); *Fucus serratus* (Hoarau et al., 2007);
58
59 271 *Neomysis integer* (Remerie et al., 2009); *Ascophyllum nodosum* (Olsen et al., 2010)

1
2
3 272 Potentially, periodic sea level changes associated to Pleistocene glacial/interglacial cycles
4
5 273 could have caused continuous rearrangements of estuaries triggering *F. ceranoides*
6
7 274 population contraction/expansion dynamics similar to those described for Iberian shores;
8
9 275 these rearrangements could have then eventually led to the sharp genetic differentiation in
10
11 276 Brittany reported here. In the light of these earlier findings and observations the scenario of
12
13 277 contemporary oceanographic barriers being responsible for the origin the genetic
14
15 278 differentiation among *F. ceranoides* estuarine populations is rejected. Our results point to the
16
17 279 preponderant role of oceanography in determining the position and possibly maintaining the
18
19 280 break between vicariant lineages. This is further supported by previous studies showing
20
21 281 shared haplotypes between Brittany (southern) and Iberia (Neiva et al., 2010; Neiva et al.,
22
23 282 2012a; Neiva et al., 2012b).

24
25
26 283 The persistent integrity of the sharp, fine-scale genetic discontinuities at secondary
27
28 284 contact zones can have a number of non-exclusive explanations. Previous studies assessing
29
30 285 this fine-scale phylogeographic structure within *F. ceranoides* in NW Iberia have highlighted
31
32 286 that contemporary dispersal between established populations is effectively too low to erase
33
34 287 historical divergence stemming from past fragmentation processes (Neiva et al., 2012b). In
35
36 288 this instance, sporadic inter-estuarine dispersal and density barriers have been invoked as the
37
38 289 main driver of limited connectivity.

39
40
41 290 Although post-glacial range expansion shows that *F. ceranoides* can effectively drift
42
43 291 across large spatial scales (Neiva et al., 2012a), dispersal among colonized estuaries is
44
45 292 expected to be limited. *F. ceranoides* is dioecious, and therefore prerequisites for successful
46
47 293 long-distance, inter-estuarine colonization are dispersal of fertile male and female fronds and
48
49 294 synchronous gamete release to produce zygotes at the new location. In addition, coastal
50
51 295 topography and estuarine morphology can curb circulation patterns, retention times and,
52
53 296 consequently, connectivity through drifting (Muhlin et al., 2008; Nicastro et al., 2008; Pardo et
54
55 297 al., 2019). The sheltered nature of estuarine habitats can significantly limit gamete dispersal
56
57 298 and the intrinsic features of geomorphology of each estuary can modulate circulation velocity
58
59 299 and intensity (Day et al., 1989). It has been suggested that pronounced meander curvatures

1
2
3 300 contribute to reduced dispersal efficiency and the variable patterns of gene flow between
4
5 301 coastal and estuarine habitats (Zardi et al., 2013).
6

7 302 Density barrier effects are usually particularly marked in species such as *F. ceranoides*
8
9 303 characterized by rapid population growth and consequent habitat saturation. The dense,
10
11 304 monospecific *F. ceranoides* canopies typical of European northern Atlantic estuaries act as a
12
13 305 demographic buffer against numerically rare inter-estuarine immigrants favoring the
14
15 306 conservation of pre-existing genetic structure.
16
17

18 307 Clearly, the abundance of estuaries and their proximity strongly suggest that distance
19
20 308 can be excluded from the list of determinants maintaining *F. ceranoides* genetic breaks along
21
22 309 these shores. In NW Iberia, the mean distance between populations inhabiting neighboring
23
24 310 estuaries is not significantly different than that between bordering populations across
25
26 311 phylogeographic breaks (Neiva et al., 2012b). Similar distributional patterns can be found in
27
28 312 Brittany where several edge populations are spatially closer to populations across regional
29
30 313 genetic disjunctions than they are to their adjacent population within the same genetic clade.
31
32 314 While some studies carried out between cold- and warm-temperate marine ecosystems along
33
34 315 the northwest coast of France have invoked distinct mesoscale hydrographic features as
35
36 316 drivers of genetic patterns (Goldson et al., 2001; Jolly et al., 2005; Couceiro et al., 2013;
37
38 317 Almeida et al., 2017), others have highlighted lack of evidence for the role of hydrodynamics
39
40 318 as dispersal determinants for the observed genetic discontinuities along NW Iberian shores
41
42 319 (Neiva et al., 2012b). The latter works have also stressed the difficulties to track drifters'
43
44 320 movements at a scale relevant for the organism to estimate migration rates among estuaries
45
46 321 and phylogroups. In NW Iberia, circulation dynamics are complex with high seasonal variability
47
48 322 and lack of persistent oceanographic patterns (Ruiz-Villarreal et al., 2006; Alvarez et al.,
49
50 323 2009). Here, through the use of large scale, dispersal simulations we reveal several
51
52 324 oceanographic regions matching haplotype segregation. Our findings add important evidence
53
54 325 to previous conclusions and support the scenario of key, mesoscale oceanographic processes
55
56 326 having a determinant role in explaining the position of the observed high levels of regional
57
58 327 genetic divergence. We also hypothesized that mesoscale oceanographic dispersal barriers
59
60

1
2
3 328 are key to the maintenance of inter-estaurine genetic differentiation. In a neutral model of
4
5 329 secondary contact following allopatric differentiation, signs of secondary intergradation are
6
7 330 generally observed around oceanographic barriers (Woodruff, 1973; Barton and Hewitt, 1985;
8
9 331 Bierne et al., 2011). Admixed nuclear background has only been reported between two
10
11 332 neighboring *F. ceranoides* Iberian populations (Neiva et al., 2012b), indicating that individuals
12
13 333 belonging to distinct phylogroups can interbreed. However, no signs of hybridization have
14
15 334 been observed in other Iberian or French populations. The geographically restricted and
16
17 335 limited lineage admixture suggests that other factors may be at play in maintaining fine-scale
18
19 336 genetic differentiation. In particular, incipient reproductive isolation (pre- or post-zygotic) can
20
21 337 depress gene flow between divergent phylogroups (e.g., Tellier et al., 2011). Under this
22
23 338 scenario, limited dispersal across oceanographic barriers explains the position of the genetic
24
25 339 discontinuity while the delay in homogenisation is mainly explained by endogenous
26
27 340 components of reproductive isolation. Future studies assessing spatial and temporal
28
29 341 reproductive dynamics of distinct *F. ceranoides* lineages will be crucial to provide a direct
30
31 342 testing of this hypothesis.
32
33

343 **Conclusions**

344 Our study highlights the need to combine evidence from multiple sources for a comprehensive
35
36 345 understanding of ecological and evolutionary mechanisms linked to phylogeographic breaks.
37
38 346 These conclusions are of great significance for other organisms with sporadic and spatially
39
40 347 limited dispersal, helping clarify the apparent inconsistency of extensive and sharp genetic
41
42 348 differentiation in geographically restricted regions. In addition to theoretical evolutionary
43
44 349 relevance, the identification of contemporary dynamics contributing to the maintenance of
45
46 350 significant units of intraspecific biodiversity is critical for efficient approaches of management
47
48 351 and conservation efforts.
49
50
51
52
53

54 352

55

56 353 **ACKNOWLEDGMENTS**

57
58
59
60

1
2
3 354 This research was funded through projects BIODIVERSA/004/2015,
4
5 355 IF/01413/2014/CP1217/CT0004 and UID/Multi/04326/2019 by the Foundation for Science
6
7 356 and Technology (FCT - MEC, Portugal) and further supported by the National Research
8
9 357 Foundation of South Africa (Grant number 64801) and through the fellowship grant
10
11 358 SFRH/BPD/88935/2012 funded by FCT and SFRH/BPD/111003/2015 jointly funded by a
12
13 359 Pew Marine Fellowship (USA) and FCT. We thank two anonymous reviewers for their
14
15
16 360 suggestions and comments.
17
18 361

362 REFERENCES

- 363 Almeida, S. C., Nicastro, K. R., Zardi, G. I., Pearson, G. A., Valero, M., and Serrão, E. A.
364 2017. Reproductive strategies and population genetic structure of *Fucus* spp. across a
365 northeast Atlantic biogeographic transition. *Aquatic Living Resources*, 30: 16.
- 366 Alvarez, I., Ospina-Alvarez, N., Pazos, Y., Bernardez, P., Campos, M., Gomez-Gesteira, J.,
367 Alvarez-Ossorio, M., et al. 2009. A winter upwelling event in the Northern Galician Rias:
368 frequency and oceanographic implications. *Estuarine, Coastal and Shelf Science*, 82: 573-
369 582.
- 370 Assis, J., Bercibar, E., Claro, B., Alberto, F., Reed, D., Raimondi, P., and Serrão, E. 2017.
371 Major shifts at the range edge of marine forests: the combined effects of climate changes
372 and limited dispersal. *Scientific Reports*, 7: 44348.
- 373 Assis, J., Serrão, E. A., Coelho, N. C., Tempera, F., Valero, M., Claro, B., and Alberto, F.
374 2018. Past climate changes and strong oceanographic barriers structured low latitude
375 genetic relics for the golden kelp *Laminaria ochroleuca*. *Journal of Biogeography*.
- 376 Assis, J., Zupan, M., Nicastro, K. R., Zardi, G. I., McQuaid, C. D., and Serrão, E. A. 2015.
377 Oceanographic conditions limit the spread of a marine invader along southern African
378 shores. *PLoS ONE*, 10(6):e0128124.

- 1
2
3 379 Ayata, S.-D., Stolba, R., Comtet, T., and Thiébaud, É. 2011. Meroplankton distribution and its
4
5 380 relationship to coastal mesoscale hydrological structure in the northern Bay of Biscay (NE
6
7 381 Atlantic). *Journal of Plankton Research*, 33: 1193-1211.
8
9 382 Barnes, I., Matheus, P., Shapiro, B., Jensen, D., and Cooper, A. 2002. Dynamics of
10
11 383 Pleistocene Population Extinctions in Beringian Brown Bears. *Science*, 295: 2267-2270.
12
13 384 Barton, N. H., and Hewitt, G. M. 1985. Analysis of hybrid zones. *Annual Review of Ecology
14
15 385 and Systematics*, 16: 113-148.
16
17 386 Bierne, N., Welch, J., Loire, E., Bonhomme, F., and David, P. 2011. The coupling
18
19 387 hypothesis: why genome scans may fail to map local adaptation genes. *Molecular Ecology*,
20
21 388 20: 2044-2072.
22
23 389 Chao, R. B., Casais, M. C., Cortizas, A. M. n., Alberti, A. P., and Paz, M. V. 2002. Holocene
24
25 390 evolution on Galician coast (NW Spain): an example of paraglacial dynamics. *Quaternary
26
27 391 International*, 93: 149-159.
28
29 392 Chassignet, E. P., Hurlburt, H. E., Smedstad, O. M., Halliwell, G. R., Hogan, P. J., Wallcraft,
30
31 393 A. J., Baraille, R., et al. 2007. The HYCOM (hybrid coordinate ocean model) data
32
33 394 assimilative system. *Journal of Marine Systems*, 65: 60-83.
34
35 395 Chust, G., Villarino, E., Chenuil, A., Irigoien, X., Bizsel, N., Bode, A., Broms, C., et al. 2016.
36
37 396 Dispersal similarly shapes both population genetics and community patterns in the marine
38
39 397 realm. *Scientific reports*, 6: 28730.
40
41 398 Clement, M., Posada, D., and Crandall, K. A. 2000. TCS: a computer program to estimate
42
43 399 gene genealogies. *Molecular Ecology*, 9: 1657-1659.
44
45 400 Comps, B., Gömöry, D., Letouzey, J., Thiébaud, B., and Petit, R. 2001. Diverging trends
46
47 401 between heterozygosity and allelic richness during postglacial colonization in the European
48
49 402 beech. *Genetics*, 157: 389-397.
50
51 403 Couceiro, L., Robuchon, M., Destombe, C., and Valero, M. 2013. Management and
52
53 404 conservation of the kelp species *Laminaria digitata*: using genetic tools to explore the
54
55 405 potential exporting role of the MPA "Parc naturel marin d'Iroise". *Aquatic Living Resources*,
56
57 406 26: 197-205.
58
59
60

- 1
2
3 407 Day, J. W., Hall, C. A. S., Kemp, W. M., and A., Y.-A. 1989. Estuarine Ecology, John Wiley
4 and Sons, New York.
5
6 408
7 409 Dowle, M., Srinivasan, A., Gorecki, J., Chirico, M., Stetsenko, P., Short, T., Lianoglou, S., et
8
9 410 al. 2019. Package 'data.table'. Extension of 'data.frame'.
10
11 411 Drummond, A., Ashton, B., Buxton, S., Cheung, M., Cooper, A., Duran, C., Field, M., et al.
12
13 412 2011. Geneious v5. 4.
14
15 413 Ferro, I., and Morrone, J. J. 2014. Biogeographical transition zones: a search for conceptual
16
17 414 synthesis. *Biological Journal of the Linnean Society*, 113: 1-12.
18
19 415 Fossette, S., Putman, N. F., Lohmann, K. J., Marsh, R., and Hays, G. C. 2012. A biologist's
20
21 416 guide to assessing ocean currents: a review. *Marine Ecology Progress Series*, 457: 285-301.
22
23 417 Gallon, R. K., Robuchon, M., Leroy, B., Le Gall, L., Valero, M., and Feunteun, E. 2014.
24
25 418 Twenty years of observed and predicted changes in subtidal red seaweed assemblages
26
27 419 along a biogeographical transition zone: inferring potential causes from environmental data.
28
29 420 *Journal of Biogeography*, 41: 2293-2306.
30
31 421 Gilg, M. R., and Hilbish, T. J. 2003. The geography of marine larval dispersal: coupling
32
33 422 genetics with fine-scale physical oceanography. *Ecology*, 84: 2989-2998.
34
35 423 Goldson, A., Hughes, R., and Gliddon, C. 2001. Population genetic consequences of larval
36
37 424 dispersal mode and hydrography: a case study with bryozoans. *Marine Biology*, 138: 1037-
38
39 425 1042.
40
41 426 Gomez, A., Hughes, R. N., Wright, P. J., Carvalho, G. R., and Lunt, D. H. 2007.
42
43 427 Mitochondrial DNA phylogeography and mating compatibility reveal marked genetic
44
45 428 structuring and speciation in the NE Atlantic bryozoan *Celleporella hyalina*. *Molecular*
46
47 429 *Ecology*, 16: 2173-2188.
48
49 430 Haklay, M., and Weber, P. 2008. Openstreetmap: User-generated street maps. *IEEE*
50
51 431 *Pervasive Computing*, 7: 12-18.
52
53 432 Hewitt, G. 2000. The genetic legacy of the Quaternary ice ages. *Nature*, 405: 907-913.
54
55 433 Hewitt, G. M. 1996. Some genetic consequences of ice ages, and their role in divergence
56
57 434 and speciation. *Biological Journal of the Linnean Society*, 58: 247-276.
58
59
60

- 1
2
3 435 Hewitt, G. M. 2004. Genetic consequences of climatic oscillations in the Quaternary.
4
5 436 Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences,
6
7 437 359: 183-195.
8
9 438 Hijmans, R. J., Phillips, S., Leathwick, J., Elith, J., and Hijmans, M. R. J. 2017. Package
10
11 439 'dismo'. Circles, 9: 1-68.
12
13 440 Hoarau, G., Coyer, J. A., Veldsink, J. H., Stam, W. T., and Olsen, J. L. 2007. Glacial refugia
14
15 441 and recolonization pathways in the brown seaweed *Fucus serratus*. Molecular Ecology, 16:
16
17 442 3606-3616.
18
19 443 Hohenlohe, P. A. 2004. Limits to gene flow in marine animals with planktonic larvae: models
20
21 444 of *Littorina* species around Point Conception, California. Biological Journal of the Linnean
22
23 445 Society, 82: 169-187.
24
25 446 Johansson, M. L., Alberto, F., Reed, D. C., Raimondi, P. T., Coelho, N. C., Young, M. A.,
26
27 447 Drake, P. T., et al. 2015. Seascape drivers of *Macrocystis pyrifera* population genetic
28
29 448 structure in the northeast Pacific. Molecular Ecology, 24: 4866-4885.
30
31 449 Jolly, M. T., Jollivet, D., Gentil, F., Thiébaud, É., and Viard, F. 2005. Sharp genetic break
32
33 450 between Atlantic and English Channel populations of the polychaete *Pectinaria koreni*, along
34
35 451 the North coast of France. Heredity, 94: 23.
36
37 452 Kamvar, Z. N., Tabima, J. F., and Grünwald, N. J. 2014. Poppr: an R package for genetic
38
39 453 analysis of populations with clonal, partially clonal, and/or sexual reproduction. PeerJ, 2:
40
41 454 e281.
42
43 455 Kennedy, J. P., Garavelli, L., Truelove, N. K., Devlin, D. J., Box, S. J., Chérubin, L. M., and
44
45 456 Feller, I. C. 2017. Contrasting genetic effects of red mangrove (*Rhizophora mangle* L.) range
46
47 457 expansion along West and East Florida. Journal of Biogeography, 44: 335-347.
48
49 458 Klein, M., Teixeira, S., Assis, J., Serrão, E. A., Gonçalves, E. J., and Borges, R. 2016. High
50
51 459 interannual variability in connectivity and genetic pool of a temperate clingfish matches
52
53 460 oceanographic transport predictions. PloS one, 11: e0165881.
54
55 461 Lamigueiro, O. P., Hijmans, R., and Lamigueiro, M. O. P. 2018. Package 'rasterVis'.
56
57
58
59
60

- 1
2
3 462 Lein, T. E. 1984. Distribution, reproduction, and ecology of *Fucus ceranoides*
4
5 463 L.(Phaeophyceae) in Norway. *Sarsia*, 69: 75-81.
6
7 464 Lett, C., Verley, P., Mullon, C., Parada, C., Brochier, T., Penven, P., and Blanke, B. 2008. A
8
9 465 Lagrangian tool for modelling ichthyoplankton dynamics. *Environmental Modelling &*
10
11 466 *Software*, 23: 1210-1214.
12
13 467 Liang, Y., He, D., Jia, Y., Sun, H., and Chen, Y. 2017. Phylogeographic studies of
14
15 468 schizothoracine fishes on the central Qinghai-Tibet Plateau reveal the highest known glacial
16
17 469 microrefugia. *Scientific Reports*, 7: 10983.
18
19 470 Librado, P., and Rozas, J. 2009. DnaSP v5: a software for comprehensive analysis of DNA
20
21 471 polymorphism data. *Bioinformatics*, 25: 1451-1452.
22
23 472 McKenzie, P. F., and Bellgrove, A. 2008. Dispersal of *Hormosira banksii* (Phaeophyceae) via
24
25 473 detached fragments: reproductive viability and longevity. *Journal of Phycology*, 44: 1108-
26
27 474 1115.
28
29 475 Mims, M. C., Phillipsen, I. C., Lytle, D. A., Kirk, E. E. H., and Olden, J. D. 2015. Ecological
30
31 476 strategies predict associations between aquatic and genetic connectivity for dryland
32
33 477 amphibians. *Ecology*, 96: 1371-1382.
34
35 478 Monteiro, C. A., Paulino, C., Jacinto, R., Serrão, E. A., and Pearson, G. A. 2016. Temporal
36
37 479 windows of reproductive opportunity reinforce species barriers in a marine broadcast
38
39 480 spawning assemblage. *Scientific reports*, 6: 29198.
40
41 481 Muhlin, J. F., Engel, C. R., Stessel, R., Weatherbee, R. A., and Brawley, S. H. 2008. The
42
43 482 influence of coastal topography, circulation patterns, and rafting in structuring populations of
44
45 483 an intertidal alga. *Molecular Ecology*, 17: 1198-1210.
46
47 484 Munda, I. M. 1999. The benthic algal vegetation of land-locked fjords in southeastern
48
49 485 Iceland. *Hydrobiologia*, 393: 169.
50
51 486 Munwes, I., Geffen, E., Roll, U., Friedmann, A., Daya, A., Tikochinski, Y., and Gafny, S.
52
53 487 2010. The change in genetic diversity down the core-edge gradient in the eastern spadefoot
54
55 488 toad (*Pelobates syriacus*). *Molecular Ecology*, 19: 2675-2689.
56
57
58
59
60

- 1
2
3 489 Neiva, J., Paulino, C., Nielsen, M. M., Krause-Jensen, D., Saunders, G. W., Assis, J.,
4
5 490 Bárbara, I., et al. 2018. Glacial vicariance drives phylogeographic diversification in the
6
7 491 amphiboreal kelp *Saccharina latissima*. *Scientific reports*, 8: 1112.
8
9 492 Neiva, J., Pearson, G. A., Valero, M., and Serrão, E. A. 2010. Surfing the wave on a
10
11 493 borrowed board: range expansion and spread of introgressed organellar genomes in the
12
13 494 seaweed *Fucus ceranoides* L. *Molecular Ecology*, 19: 4812-4822.
14
15 495 Neiva, J., Pearson, G. A., Valero, M., and Serrão, E. A. 2012a. Drifting fronds and drifting
16
17 496 alleles: range dynamics, local dispersal and habitat isolation shape the population structure
18
19 497 of the estuarine seaweed *Fucus ceranoides*. *Journal of Biogeography*, 39: 1167-1178.
20
21 498 Neiva, J., Pearson, G. A., Valero, M., and Serrão, E. A. 2012b. Fine-scale genetic breaks
22
23 499 driven by historical range dynamics and ongoing density-barrier effects in the estuarine
24
25 500 seaweed *Fucus ceranoides* L. *BMC Evolutionary Biology*, 12: 78.
26
27 501 Newman, M. E. 2006. Modularity and community structure in networks. *Proceedings of the*
28
29 502 *national academy of sciences*, 103: 8577-8582.
30
31 503 Nicastro, K. R., Zardi, G. I., McQuaid, C. D., Teske, P. R., and Barker, N. P. 2008. Coastal
32
33 504 topography drives genetic structure in marine mussels. *Marine Ecology Progress Series*,
34
35 505 368: 189-195.
36
37 506 Nicastro, K. R., Zardi, G. I., Teixeira, S., Neiva, J., Serrao, E. A., and Pearson, G. A. 2013.
38
39 507 Shift happens: trailing edge contraction associated with recent warming trends threatens a
40
41 508 distinct genetic lineage in the marine macroalga *Fucus vesiculosus*. *BMC Biology*, 11: 6.
42
43 509 Oksanen, J., Blanchet, F., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P., et
44
45 510 al. 2018. *vegan: a community ecology package*. R package verions 2.4-6.
46
47 511 Olsen, J. L., Zechman, F. W., Hoarau, G., Coyer, J. A., Stam, W. T., Valero, M., and Åberg,
48
49 512 P. 2010. The phylogeographic architecture of the fucoid seaweed *Ascophyllum nodosum*: an
50
51 513 intertidal 'marine tree' and survivor of more than one glacial–interglacial cycle. *Journal of*
52
53 514 *Biogeography*, 37: 842-856.
54
55 515 Pardo, C., Guillemain, M.-L., Pena, V., Barbara, I., Valero, M., and Barreiro, R. 2019. Local
56
57 516 coastal configuration rather than latitudinal gradient shape clonal diversity and genetic

- 1
2
3 517 structure of *Phymatolithon calcareum* maerl beds in North European Atlantic. *Frontiers in*
4
5 518 *Marine Science*, 6: 149.
6
7 519 Provan, J. 2013. The effects of past, present and future climate change on range-wide
8
9 520 genetic diversity in northern North Atlantic marine species. *Frontiers of Biogeography*, 5(1).
10
11 521 Provan, J., Wattier, R. A., and Maggs, C. A. 2005. Phylogeographic analysis of the red
12
13 522 seaweed *Palmaria palmata* reveals a Pleistocene marine glacial refugium in the English
14
15 523 Channel. *Molecular Ecology*, 14: 793-803.
16
17 524 Puillat, I., Lazure, P., Jégou, A., Lampert, L., and Miller, P. 2004. Hydrographical variability
18
19 525 on the French continental shelf in the Bay of Biscay, during the 1990s. *Continental Shelf*
20
21 526 *Research*, 24: 1143-1163.
22
23 527 Putman, N. F., and He, R. 2013. Tracking the long-distance dispersal of marine organisms:
24
25 528 sensitivity to ocean model resolution. *Journal of the Royal Society Interface*, 10: 20120979.
26
27 529 Remerie, T., Vierstraete, A., Weekers, P. H., Vanfleteren, J. R., and Vanreusel, A. 2009.
28
29 530 Phylogeography of an estuarine mysid, *Neomysis integer* (Crustacea, Mysida), along the
30
31 531 north-east Atlantic coasts. *Journal of Biogeography*, 36: 39-54.
32
33 532 Roucoux, K., De Abreu, L., Shackleton, N., and Tzedakis, P. 2005. The response of NW
34
35 533 Iberian vegetation to North Atlantic climate oscillations during the last 65kyr. *Quaternary*
36
37 534 *Science Reviews*, 24: 1637-1653.
38
39 535 Ruiz-Villarreal, M., González-Pola, C., del Río, G. D., Lavin, A., Otero, P., Piedracoba, S.,
40
41 536 and Cabanas, J. 2006. Oceanographic conditions in North and Northwest Iberia and their
42
43 537 influence on the Prestige oil spill. *Marine Pollution Bulletin*, 53: 220-238.
44
45 538 Selkoe, K. A., Watson, J. R., HORIN, T. B., IACCHEI, M., MITARAI, S., SIEGEL, D. A.,
46
47 539 GAINES, S. D., et al. 2010. Taking the chaos out of genetic patchiness: seascape genetics
48
49 540 reveals ecological and oceanographic drivers of genetic patterns in three temperate reef
50
51 541 species. *Molecular Ecology*, 19: 3708-3726.
52
53 542 Serrão, E. A., Kautsky, L., Lifvergren, T., and Brawley, S. H. 1997. Gamete dispersal and
54
55 543 pre-recruitment mortality in Baltic *Fucus vesiculosus*. *Phycologia*, 36 101-102.
56
57
58
59
60

- 1
2
3 544 Slatkin, M. 1993. Isolation by distance in equilibrium and non-equilibrium populations.
4
5 545 Evolution, 47: 264-279.
6
7 546 Smith, B. T., Escalante, P., Baños, B. E. H., Navarro-Sigüenza, A. G., Rohwer, S., and
8
9 547 Klicka, J. 2011. The role of historical and contemporary processes on phylogeographic
10
11 548 structure and genetic diversity in the Northern Cardinal, *Cardinalis cardinalis*. BMC
12
13 549 Evolutionary Biology, 11: 136.
14
15 550 Sotka, E. E., Wares, J. P., Barth, J. A., Grosberg, R. K., and Palumbi, S. R. 2004. Strong
16
17 551 genetic clines and geographical variation in gene flow in the rocky intertidal barnacle
18
19 552 *Balanus glandula*. Molecular Ecology, 13: 2143-2156.
20
21 553 Spalding, M. D., Fox, H. E., Allen, G. R., Davidson, N., Ferdana, Z. A., Finlayson, M.,
22
23 554 Halpern, B. S., et al. 2007. Marine ecoregions of the world: a bioregionalization of coastal
24
25 555 and shelf areas. Bioscience, 57: 573-583.
26
27 556 Team, R. C. 2018. R: A language and environment for statistical computing; 2015.
28
29 557 Tellier, F., Tapia, J., Faugeron, S., Destombe, C., and Valero, M. 2011. The *lessonia*
30
31 558 *nigrescens* species complex (laminariales, phaeophyceae) shows strict parapatry and
32
33 559 complete reproductive isolation in a secondary contact zone 1. Journal of Phycology, 47:
34
35 560 894-903.
36
37 561 Teske, P., Papadopoulos, I., Zardi, G., McQuaid, C., Edkins, M., Griffiths, C., and Barker, N.
38
39 562 2007. Implications of life history for genetic structure and migration rates of southern African
40
41 563 coastal invertebrates: planktonic, abbreviated and direct development. Marine Biology, 152:
42
43 564 697-711.
44
45 565 Teske, P. R., Von Der Heyden, S., McQuaid, C. D., and Barker, N. P. 2011. A review of
46
47 566 marine phylogeography in southern Africa. South African Journal of Science, 107: 43-53.
48
49 567 Thiel, M., and Haye, P. 2006. The ecology of rafting in the marine environment-III.
50
51 568 Biogeographical and evolutionary consequences (in Oceanography and marine biology: An
52
53 569 annual review-Volume 48, pp. 323-429).
54
55
56
57
58
59
60

- 1
2
3 570 Varela, R. A., Rosón, G., Herrera, J. L., Torres-López, S., and Fernández-Romero, A. 2005.
4
5 571 A general view of the hydrographic and dynamical patterns of the Rías Baixas adjacent sea
6
7 572 area. *Journal of Marine Systems*, 54: 97-113.
8
9 573 Wares, J. P., Gaines, S., and Cunningham, C. W. 2001. A comparative study of asymmetric
10
11 574 migration events across a marine biogeographic boundary. *Evolution*, 55: 295-306.
12
13 575 Woodruff, D. S. 1973. Natural hybridization and hybrid zones. *Systematic Biology*, 22: 213-
14
15 576 218.
16
17 577 Zardi, G., McQuaid, C., Teske, P., and Barker, N. 2007. Unexpected genetic structure of
18
19 578 mussel populations in South Africa: indigenous *Perna perna* and invasive *Mytilus*
20
21 579 *galloprovincialis*. *Marine Ecology Progress Series*, 337: 135-144.
22
23 580 Zardi, G., Nicastro, K., McQuaid, C., Hancke, L., and Helmuth, B. 2011. The combination of
24
25 581 selection and dispersal helps explain genetic structure in intertidal mussels. *Oecologia*, 165:
26
27 582 947-958.
28
29 583 Zardi, G. I., Nicastro, K. R., Ferreira Costa, J., Serrão, E. A., and Pearson, G. A. 2013.
30
31 584 Broad scale agreement between intertidal habitats and adaptive traits on a basis of
32
33 585 contrasting population genetic structure. *Estuarine, Coastal and Shelf Science*, 131: 140-
34
35 586 148.
36
37 587
38
39
40
41
42 588
43
44
45 589
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **590 FIGURE LEGEND**
4
5


6 **591 Fig. 1.** Haplotype distribution of *Fucus ceranoides* in the coastlines of (a) Northwester France
7
8 **592** and (panel b) Northwest Iberia. The colors along coastlines (a, b) depict the different
9
10 **593** oceanographic regions (R) identified in network analysis. A pairwise matrix of directional
11
12 **594** connectivity between oceanographic regions identified in (d) western Brittany and (e)
13
14 **595** northwest Iberia. Matrix header colors according to the oceanographic regions depicted in the
15
16 **596** top panels a and b. Haplotype frequency and differentiation degree in (f) western Brittany and
17
18 **597** (g) northwest Iberia. Note that there is no correspondence of haplotypes between the two
19
20 **598** panels (i.e., each dataset was treated separately).
21
22
23
24 **599**

25
26 **600 TABLES**
27

28 **601 Table 1.** Maximum and average distances (km), drifting time (days) and probabilities produced
29
30 **602** by the particles connecting different cells for the lagrangian particle simulations running in
31
32 **603** Northwest France and Northwest Spain..
33
34
35 **604**

Region	Distance (km)		Time (days)		Probability	
	Maximum	Mean (\pm SD)	Maximum	Mean (\pm SD)	Maximum	Mean (\pm SD)
W Brittany	382.6	62.6 \pm 54.6	24.6	3.61 \pm 2.63	0.721	0.004 \pm 0.020
NW Iberia	431.2	105.9 \pm 90.8	26.7	2.09 \pm 1.81	0.607	0.004 \pm 0.021

36
37
38
39
40
41
42
43
44
45 **605**
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



1
2
3 1 **Congruence between fine-scale genetic breaks and dispersal potential in an**
4
5 2 **estuarine seaweed across multiple transition zones.**
6
7

8 3

9
10 4 Katy R. Nicastr¹, Jorge Assis¹, Ester A. Serrão¹, Gareth A. Pearson¹, João Neiva¹, Myriam
11
12 5 Valero², Rita Jacinto¹, Gerardo I. Zardi^{3*}
13
14

15 6 ¹CCMAR-CIMAR – Associated Laboratory, University of Algarve, Campus de Gambelas,
16 7 Faro, 8005–139, Portugal

18 8 ²UMI 3614 Evolutionary Biology and Ecology of Algae, CNRS, Sciences Sorbonne
19 9 Université, UC, UACH, Station Biologique de Roscoff, Roscoff Cedex, France

21
22 10 ³Department of Zoology and Entomology, Rhodes University, Grahamstown, 6140, South
23 11 Africa

24 12 *corresponding author: zardi73@yahoo.it, +351 927738340
25
26

27 13

28
29
30 14 **Additional emails:**

31
32
33 15 KRN: katynicastro@gmail.com

34
35 16 JA: jorgemfa@me.com

36
37
38 17 EAS: eserrao@ualg.pt

39
40
41 18 GAP: gpearson@ualg.pt

42
43
44 19 JN: jmneiva@ualg.pt

45
46
47 20 MV: myriam.valero@sb-roscoff.fr

48
49
50 21 RJ: ritajacinto85@gmail.com
51
52

53 22

54
55 23 **Running head:** Congruence between fine-scale genetic breaks and dispersal potential.

56
57 24 **Keywords:** Biogeography, physical modelling, gene flow, *Fucus* spp.
58
59
60

25 **ABSTRACT**

26 Genetic structure in biogeographical transition zones can be shaped by several factors
27 including limited dispersal across barriers, admixture following secondary contact, differential
28 selection and mating incompatibility. A striking example is found in Northwest France and
29 Northwest Spain , where the estuarine seaweed *Fucus ceranoides* L. exhibits sharp, regional
30 genetic clustering. This pattern has been related to historical population fragmentation and
31 divergence into distinct glacial refugia, followed by post-glacial expansion and secondary
32 contact. The contemporary persistence of sharp ancient genetic breaks between nearby
33 estuaries has been attributed to prior colonization effects (density barriers) but the effect of
34 oceanographic barriers has not been tested. Here, through a combination of mesoscale
35 sampling (15 consecutive populations) and population genetic data (mtlGS) in NW France,
36 we define regional genetic disjunctions similar to those described in NW Iberia. Most
37 importantly, using high resolution dispersal simulations for Brittany and Iberian populations,
38 we provide evidence for a central role of contemporary hydrodynamics in maintaining genetic
39 breaks across these two major biogeographic transition zones. Our findings further show the
40 importance of a comprehensive understanding of oceanographic regimes in hydrodynamically
41 complex coastal regions to explain the maintenance of sharp genetic breaks along
42 continuously populated coastlines.

44 **INTRODUCTION**

45 Fueled by current concerns about the impacts of global change on biodiversity, there is a
46 renewed interest in understanding the processes affecting a species' range dynamics as well
47 as the factors shaping its genetic diversity. Generally, both historical and contemporary
48 processes are invoked to explain the distribution and phylogeography of a species. Tectonic
49 events and past climate fluctuations created topographic, environmental and hydrologic
50 barriers that were key in shaping phylogeographic structure of many plant and animal species.
51 Of these, the Last Glacial Maximum is probably the most significant and recent historical event

1
2
3 52 (e.g., Hewitt, 2000; Barnes et al., 2002; Liang et al., 2017; Neiva et al., 2018). During this time,
4
5 53 ice sheets covered much of current cold and temperate zones of the Northern Hemisphere,
6
7 54 triggering range fragmentation and contractions of entire ecosystems to more southern
8
9 55 latitudes, while during the present interglacial, widespread range shifts and poleward
10
11 56 expansions have occurred.

12
13 57 It is predicted that genetic diversity is lowest in recently colonized areas and highest in refugial
14
15 58 areas where long-term persistence was possible (e.g., Hewitt, 1996; Comps et al., 2001;
16
17 59 Hewitt, 2004; Kennedy et al., 2017). However, contemporary demographic processes can
18
19 60 either mask or even erase genetic signatures of population expansions or retreats (Smith et
20
21 61 al., 2011). High levels of gene flow may homogenize genetic diversity among previously
22
23 62 structured populations. On the other hand, nonrandom mating with individuals in close
24
25 63 geographic proximity can generate genetic structuring within a continuous population (Slatkin,
26
27 64 1993). Intrinsic characteristics of the species such as dispersal ability, migration patterns and
28
29 65 changes in population densities can further confound any historical signature (e.g., Mims et al.,
30
31 66 2015; Chust et al., 2016; Assis et al., 2018)

32
33 67 Biogeographical transition regions provide a good framework for exploring and understanding
34
35 68 genetic structuring of species through space and time. These are areas of overlap and
36
37 69 segregation between different biotic components, and geographically separated clades often
38
39 70 coincide with these transition zones (Ferro and Morrone, 2014). In the marine realm,
40
41 71 phylogeographic breaks in biogeographical transition regions are generally attributed to
42
43 72 historical processes or contemporary dispersal barriers, such as upwelling phenomena and
44
45 73 currents that may limit along-shore dispersal, environmental differences boosting local
46
47 74 adaptation and/or reproductive strategies maintaining self-recruitment (Gilg and Hilbish, 2003;
48
49 75 Zardi et al., 2007; Selkoe et al., 2010).

50
51 76 There are well known biogeographical boundaries in all oceans, some of which are associated
52
53 77 with oceanographic features that generate contemporary barriers to dispersal. For example,
54
55 78 the strong southward-flowing Agulhas Current, which runs about 10 km offshore along most
56
57 79 of South Africa's east and southeast coast, impedes larval dispersal and thus promotes local

1
2
3 80 adaptation (Teske et al., 2011; Zardi et al., 2011). Dispersal and gene flow around Point
4
5 81 Conception in southern California are also strongly affected by extensive upwelling of colder
6
7 82 sub-surface waters and by the southward California Current (Wares et al., 2001; Hohenlohe,
8
9 83 2004; Johansson et al., 2015).

10
11 84 Here, we investigated the contribution of contemporary oceanographic connectivity in
12
13 85 maintaining geographically separated genetic clades within a continuously distributed
14
15 86 seaweed species along the biogeographic transition areas of Brittany (northwest France,
16
17 87 Europe) and northwest Iberia (northern Spain, Europe; Fig. 1). These two areas are highly
18
19 88 relevant for studying this question and marine connectivity in general. Both areas delimit the
20
21 89 boundaries between cold-temperate and warm-temperate regions (Spalding et al., 2007), they
22
23 90 are refugial zones for numerous species (Provan, 2013) and, over the past decades, they
24
25 91 have undergone significant changes in macroalgae assemblages due to climate change
26
27 92 (Nicastro et al., 2013; Gallon et al., 2014; Assis et al., 2017).

28
29
30 93 In this study, we use phylogeographic analyses and Lagrangian Particle Simulations
31
32 94 (LPS) coupled with network analyses to evaluate the levels of congruence between dispersal
33
34 95 potential and the patterns of present-day genetic differentiation and diversity in the estuarine
35
36 96 seaweed *Fucus ceranoides* along these two biogeographic transition zones. This species is
37
38 97 perennial, dioecious and restricted to estuarine intertidal areas. Furoid seaweeds have no
39
40 98 planktonic dispersal stage and have restricted gamete dispersal (Serrão et al., 1997).
41
42 99 However, adult individuals can achieve long distance gene flow via the rafting of whole or
43
44 100 partially detached thalli with reproductive structures (Thiel and Haye, 2006; McKenzie and
45
46 101 Bellgrove, 2008), a form of population connectivity strongly influenced by hydrodynamic forces
47
48 102 and coastal topography. Currently, *F. ceranoides* is distributed from northern Portugal to
49
50 103 northern Norway (Lein, 1984) and Iceland (Munda, 1999), covering both past non-glaciated
51
52 104 and glaciated regions of Europe. Previous studies have shown the occurrence of two divergent
53
54 105 genetic lineages in this species between the south and the ~~North-north~~ of Europe, the
55
56 106 phylogeographic break being localized in Northwest France near the English Channel (Neiva
57
58 107 et al., 2012a; Neiva et al., 2012b). Specifically, the dominant Northwest Iberian haplotypes of

1
2
3 108 this species was found in southern Brittany but not after the genetic break in northern Brittany
4
5 109 (Neiva et al., 2010). We discuss three scenarios for the concordance between present-day
6
7 110 oceanographic dispersal barriers and the observed genetic differentiation among *F.*
8
9 111 *ceranoides* estuarine populations: 1) Contemporary oceanographic barriers to dispersal are
10
11 112 responsible for the origin, the position and the maintenance (i.e. delaying genetic
12
13 113 homogenization) of the genetic differentiation; 2) Contemporary oceanographic circulation
14
15 114 patterns explain the position, the maintenance but not the origin of the genetic differentiation;
16
17 115 3) Reproductive isolation, in addition to contemporary oceanographic patterns, limits genetic
18
19 116 homogenisation thus contributing to the position and the maintenance of the genetic
20
21 117 differentiation.
22
23
24 118

27 119 **MATERIALS**

30 120 **Genetic data**

31
32 121 The two study areas (NW France and NW Spain) are peninsulas characterized by complex
33
34 122 and variable circulation patterns (Puillat et al., 2004; Varela et al., 2005; Ruiz-Villarreal et al.,
35
36 123 2006; Ayata et al., 2011). Because of that, sampling of only a limited number of localities would
37
38 124 be inappropriate. We therefore decided to sample a small number of individuals at as many
39
40 125 sites along the two coastlines as possible, rather than obtaining large numbers of individuals
41
42 126 from a limited number of sites (e.g., Sotka et al., 2004; Teske et al., 2007). This decision was
43
44 127 also supported by the very low genetic diversity within localities for this species (Neiva et al.,
45
46 128 2010; 2012b; Neiva et al., 2012a). As a result, two sets of sequences of *Fucus ceranoides*
47
48 129 were prepared and analyzed separately. The first data set (Data set 1) comprised sequences
49
50 130 of individuals collected in the estuaries of all major rivers between Hennebont (HB, southern
51
52 131 Brittany) and Camarel (CM, northern Brittany) in northwest France (n = 2 from each site;
53
54 132 Table SI1). The second dataset (Data set 2) consisted of sequences previously analysed in
55
56 133 Neiva et al. (Neiva et al., 2010; 2012b; Neiva et al., 2012a) and sampled across northern Iberia
57
58 134 between Viana do Castelo (VIA, northern Portugal) and Porcia (POR, northern Spain). To
59
60

1
2
3 135 allow a comparison between data sets, a random subsample of 2 individuals from each site
4
5 136 was used for Data set 2. The random subsampling in NW Spain was repeated to ensure
6
7 137 consistency in the results obtained.
8

9 138 Sampling was performed in 2014-2015 and conducted with similar *criteria* for individuals used
10
11 139 in both data sets. All collection sites were characterized by monospecific belts of *F.*
12
13 140 *ceranoides* attached to hard substrata and were exposed to steep salinity fluctuations
14
15 141 throughout the tidal cycle. At each site, 5-10 cm tips of apical vegetative tissue were collected
16
17 142 from individuals sampled in the mid distributional range of the species. Neighboring sites were
18
19 143 at an average proximity of about 50 (± 15) and 33 (± 17) km for Data sets 1 and 2 respectively.
20
21 144 All samples were individually stored dehydrated in silica-gel crystals until DNA extraction.
22

23 145 *DNA isolation and sequencing of data set 1*

24
25
26 146 To compile Data set 1, genomic DNA was extracted from approximately 10 mg of dried tissue
27
28 147 using the Nucleospin® Multi-96 plant kit (Macherey-Nagel Duren, Germany), according to the
29
30 148 manufacturer's protocol. Individuals were sequenced for the mitochondrial 23 S/trnK
31
32 149 intergenic spacer (mtIGS, Neiva et al., 2010). Primer sequences and amplification details were
33
34 150 the same as in Neiva et al. (Neiva et al., 2010; 2012b; Neiva et al., 2012a). Amplified fragments
35
36 151 were run in an ABI PRISM 3130xl automated capillary sequencer (Applied Biosystems,
37
38 152 CCMAR Portugal). MtDNA sequences were aligned, proofread and edited in GENEIOUS 3.8
39
40 153 (Drummond et al., 2011).
41
42

43 154 *Data analyses*

44
45 155 For both data sets, haplotype frequencies were estimated using DnaSP 5.0 (Librado and
46
47 156 Rozas, 2009). The relationships among the MtIGS haplotypes were inferred using statistical
48
49 157 parsimony with Tcs v. 1.13 (Clement et al., 2000). Because additional subsampled dataset for
50
51 158 the Iberia provided similar results (Table SI2 and Fig. SI1), only one was used for the
52
53 159 simulations.
54

55 160 **Dispersal simulations**

56
57 161 The main oceanographic regions in northwest France and northwest Iberia (~550km and
58
59 162 ~600km of coastlines, respectively) were identified by coupling Lagrangian Particle
60

1
2
3 163 Simulations (LPS) with network analyses ~~–least cost and community algorithms–~~(least cost
4
5 164 distance and community algorithm, e.g., Assis et al., 2015; Klein et al., 2016; Assis et al.,
6
7 165 2018). The simulations used daily data of ocean currents assembled from the Hybrid
8
9 166 Coordinate Ocean Model (HYCOM), a ~~high~~-resolution product with a spatial resolution of 0.08°
10
11 167 (approx. 6–9 km), ~~forced~~ by wind speed, wind stress, precipitation and heat flux. This model
12
13
14 168 can resolve oceanic fronts, meandering currents, filaments and eddies (Chassignet et al.,
15
16 169 2007), important mesoscale processes to properly simulate ocean dispersal (Assis et al.,
17
18 170 2015; Klein et al., 2016).

19
20 171 Both regions of simulation were gridded to a common spatial resolution of 0.01° (approx. 1km).
21
22 172 A polygon representing global coastlines - OpenStreetMap geographic information (Haklay
23
24 173 and Weber, 2008)- was used to define intertidal source and sink cells. Passive particles
25
26 174 simulating rafts of *F. ceranoides* adult individuals were released from each gridded cell every
27
28 175 12 hours and allowed to drift for 60 full days; an extreme period for long-lived rafts of brown
29
30 176 macroalgae (Monteiro et al., 2016; Assis et al., 2018). This approach aimed to capture the
31
32 177 rare, long-distance dispersal events, allowing gene flow at the scales of both regions (Monteiro
33
34 178 et al., 2016; Assis et al., 2018). After the 60 days period, or when ending up on shore, the
35
36 179 particles were removed from the simulation. The geographic position of all particles was
37
38 180 calculated every hour with bilinear interpolation on the ocean velocity fields (with a spatial
39
40 181 resolution of 0.08°), while combining a 4th Order Runge-Kutta adaptive time-step on the path
41
42 182 equations (e.g., Lett et al., 2008; Klein et al., 2016).

43
44
45 183 The degree of connectivity between all pairs of gridded cells was determined by dividing the
46
47 184 number of unique particles released from cell *i* that ended up in cell *j*, by the total number of
48
49 185 particles released from cell *i*. To account for the inter-annual variability in the ocean data, the
50
51 186 simulations ran independently per year (from January to December), for the most recent 10-
52
53 187 year period of data, available in HYCOM (i.e., 2003 to 2012). Asymmetrical connectivity
54
55 188 matrices were determined by averaging the outcomes of the annual simulations.

56
57
58 189 The connectivity matrices were used in network analyses (i.e., graph theory) to infer the major
59
60 190 oceanographic regions of northwest Iberia and northwest France. To this end, network

1
2
3 191 percolation removed weak probabilities to a threshold maintaining all cells (nodes) connected
4
5 192 into a single network (Rozenfeld et al. 2008; Cunha et al., 2017), while maximizing modularity,
6
7 193 which quantified the strength of the backbone structure (or goodness of fit) of the networks
8
9 194 (Newman, 2006). This allowed the removal of surplus connections with unimportant
10
11 195 information. The leading eigenvector algorithm (Newman, 2006) was applied to the percolated
12
13 196 networks to assign a unique membership to the nodes. This approach allowed the detection
14
15 197 of communities in the networks (e.g., Munwes et al., 2010), which in practice translated into a
16
17 198 delineation of oceanographic regions in northwest Iberia and northwest France structured by
18
19 199 connectivity of ocean currents (Assis et al., 2018). The statistical significance of the
20
21 200 membership assignment to the nodes was inferred by testing the proportion of 9999
22
23 201 membership randomizations that retrieved a higher modularity than that observed.
24
25

26 202 Lagrangian Particle Simulations and network analyses were performed in R (R Development
27
28 203 Core Team, 2016) using the packages: data.table (Dowle et al., 2019), dismo (Hijmans et al.,
29
30 204 2017), igraph (Kamvar et al., 2014), parallel (Team, 2018), raster (Lamigueiro et al., 2018)
31
32 205 and vegan (Oksanen et al., 2018).
33
34
35 206

36 207 **RESULTS**

37
38 208 In Data set 1 (NW France), six mtIGS haplotypes were identified in 30 individuals of *Fucus*
39
40 209 *ceranoides* in 15 sampled sites. In NW France, the network analyses showed two dominant
41
42 210 haplotypes (A and D) plus three derived ones (B, C and E), each private to one population
43
44 211 (Fig 1a). Haplotype frequency distribution revealed a geographical segregation of the two main
45
46 212 haplotypes. Haplotype 1 was present in individuals from regions R1 and R2 while Haplotype
47
48 213 D was restricted to more northern sites within region R3.
49

50
51 214 Out of 415 sequences retrieved from the GenBank, 52 sequences were randomly selected for
52
53 215 Data set 2 (northwest Iberia). The network showed seven main haplotypes of which two were
54
55 216 shared among six to ten populations and three were shared by two to three populations. The
56
57 217 remaining two haplotypes were private to one single population. In the frequency distribution,
58
59 218 the main haplotype was restricted to region R5 and haplotype F was present only in region
60

1
2
3 219 R4. Haplotype A was present in regions R3 and R1 while haplotype B was confined to region

4
5 220 **R2. Dispersal simulations**

6
7 221 The LPS using HYCOM ocean currents over the 10-year period released 7300 particles per
8
9 222 cell (7.80×10^6 and 7.88×10^6 particles in total in northwest France and northwest Iberia,
10
11 223 respectively). Particles drifted for longer distances in northwest Iberia than in northwest France
12
13 224 (up to 431.2 km; Table 1; Fig. SI2). The maximum period of drifting time was also higher in
14
15 225 northwest Iberia (26.7 days), but on average, particles drifted for longer periods in northwest
16
17 226 France (3.61 ± 2.63 days). The maximum probability of connectivity between the pairs of cells
18
19 227 was observed in northwest France (western Brittany; 0.721). However, the average cell
20
21 228 probabilities within regions did not vary considerably (Table 1; Fig. SI2).

22
23
24 229 The assignment of oceanographic regions performed by the leading eigenvector algorithm
25
26 230 (network analysis) for northwest France and northwest Iberia showed significant modularity
27
28 231 values of 0.41 and 0.57 (p-values < 0.001), respectively. The algorithm identified three regions
29
30 232 in northwest France (Fig. 1a), with breaks in Penmarch and Porspoder, and five regions in
31
32 233 northwest Iberia (Fig. 1b), with breaks in Ria de Arousa, Corrubedo, Camelle (northern Costa
33
34 234 da Morte) and Cabo Ortegal (Cariño). The average probability of connectivity within the
35
36 235 oceanographic regions of northwest France (diagonal of Fig. 1d) was 10-fold higher than
37
38 236 between regions, with the exception of those between R1 and R2 (Fig. 1d), which were of the
39
40 237 same order of magnitude. Following a similar pattern, the probabilities of connectivity in
41
42 238 northwest Iberia were 10 to 100-fold higher within regions than between regions (diagonal of
43
44 239 Fig. 1e), with a marked increase as the distance between groups increased.

45
46
47
48 240

49
50
51 241 **DISCUSSION**

52
53
54 242 Our results show an overall strong match between oceanographic regions identified
55
56 243 by dispersal simulations and fine-scale genetic discontinuities in the estuarine seaweed *Fucus*
57
58 244 *ceranoides* inhabiting north Atlantic shores (Fig. 1).

1
2
3 245 Predictions of oceanographic transport made with Lagrangian particle simulations are
4
5 246 highly sensitive to the inner spatial and temporal resolution of circulation models (Putman and
6
7 247 He, 2013). While we preserved the raw circulation processes modelled by Hycom at the scales
8
9 248 of days and tens of kilometers, a main limitation may arise if additional oceanographic
10
11 249 processes occurring at smaller scales are important for realistic connectivity events shaping
12
13 250 the genetic structure of *F. ceranoides*. For instance, circulation models with coarse spatial
14
15 251 resolutions may underestimate drifting times up to a factor of ~2. In the same way, weekly or
16
17 252 even daily temporal time steps may not be a realistic representation of what organisms
18
19 253 continuously experience (Fossette et al., 2012). Regardless of such potential limitations, our
20
21 254 results compared with independent genetic data, as well as additional studies comparing
22
23 255 connectivity estimates with satellite-tracked organisms (e.g., Fossette et al., 2012) suggest
24
25 256 that, overall, particles advected by Hycom data provide a reliable estimate of the main
26
27 257 processes shaping current flow.

28
29
30 258 Previous studies have shown that mtIGS differentiation and contemporary *F.*
31
32 259 *ceranoides* genetic structuring in NW Iberia is most likely the result of complex, past range
33
34 260 dynamics (Neiva et al., 2012a). High levels of endemism and diversity highlighted by genetic
35
36 261 analyses of *F. ceranoides* populations inhabiting this region indicate long-term persistence in
37
38 262 glacial refugia. Despite the refugial role played by NW Iberia, sea level changes associated
39
40 263 with glacial/inter-glacial expansion and melting of ice-sheets had significant effects on near-
41
42 264 shore habitats (Chao et al., 2002; Roucoux et al., 2005). The current fine-scale genetic breaks
43
44 265 in Iberian *F. ceranoides* are the result of past fragmentation and divergence of populations
45
46 266 into distinct refugia (estuarine refugia within regional refugia), followed by expansion and
47
48 267 secondary contact of vicariant phylogroups (Neiva et al., 2012b).

49
50
51 268 In addition to southern European refugia, the ice-free paleo-shores of northwest
52
53 269 France (western Brittany) together with south western Ireland and the English Channel have
54
55 270 been recognized as northern periglacial refugia for several species (*Palmaria palmata* (Provan
56
57 271 et al., 2005); *Celleporella hyalina* (Gomez et al., 2007); *Fucus serratus* (Hoarau et al., 2007);
58
59 272 *Neomysis integer* (Remerie et al., 2009); *Ascophyllum nodosum* (Olsen et al., 2010)

1
2
3 273 Potentially, periodic sea level changes associated to Pleistocene glacial/interglacial cycles
4
5 274 could have caused continuous rearrangements of estuaries triggering *F. ceranoides*
6
7 275 population contraction/expansion dynamics similar to those described for Iberian shores;
8
9 276 these rearrangements could have then eventually led to the sharp genetic differentiation in
10
11 277 Brittany reported here. In the light of these earlier findings and observations the scenario of
12
13 278 contemporary oceanographic barriers being responsible for the origin the genetic
14
15 279 differentiation among *F. ceranoides* estuarine populations is rejected. Our results point to the
16
17 280 preponderant role of oceanography in determining the position and possibly maintaining the
18
19 281 break between vicariant lineages. This is further supported by previous studies showing
20
21 282 shared haplotypes between Brittany (southern) and Iberia (Neiva et al., 2010; Neiva et al.,
22
23 283 2012a; Neiva et al., 2012b).

24
25
26 284 The persistent integrity of the sharp, fine-scale genetic discontinuities at secondary
27
28 285 contact zones can have a number of non-exclusive explanations. Previous studies assessing
29
30 286 this fine-scale phylogeographic structure within *F. ceranoides* in NW Iberia have highlighted
31
32 287 that contemporary dispersal between established populations is effectively too low to erase
33
34 288 historical divergence stemming from past fragmentation processes (Neiva et al., 2012b). In
35
36 289 this instance, sporadic inter-estuarine dispersal and density barriers have been invoked as the
37
38 290 main driver of limited connectivity.

39
40
41 291 Although post-glacial range expansion shows that *F. ceranoides* can effectively drift
42
43 292 across large spatial scales (Neiva et al., 2012a), dispersal among colonized estuaries is
44
45 293 expected to be limited. *F. ceranoides* is dioecious, and therefore prerequisites for successful
46
47 294 long-distance, inter-estuarine colonization are dispersal of fertile male and female fronds and
48
49 295 synchronous gamete release to produce zygotes at the new location. In addition, coastal
50
51 296 topography and estuarine morphology can curb circulation patterns, retention times and,
52
53 297 consequently, connectivity through drifting (Muhlin et al., 2008; Nicastro et al., 2008; Pardo et
54
55 298 al., 2019). The sheltered nature of estuarine habitats can significantly limit gamete dispersal
56
57 299 and the intrinsic features of geomorphology of each estuary can modulate circulation velocity
58
59 300 and intensity (Day et al., 1989). It has been suggested that pronounced meander curvatures

1
2
3 301 contribute to reduced dispersal efficiency and the variable patterns of gene flow between
4
5 302 coastal and estuarine habitats (Zardi et al., 2013).
6

7 303 Density barrier effects are usually particularly marked in species such as *F. ceranoides*
8
9 304 characterized by rapid population growth and consequent habitat saturation. The dense,
10
11 305 monospecific *F. ceranoides* canopies typical of European northern Atlantic estuaries act as a
12
13 306 demographic buffer against numerically rare inter-estuarine immigrants favoring the
14
15 307 conservation of pre-existing genetic structure.
16
17

18 308 Clearly, the abundance of estuaries and their proximity strongly suggest that distance
19
20 309 can be excluded from the list of determinants maintaining *F. ceranoides* genetic breaks along
21
22 310 these shores. In NW Iberia, the mean distance between populations inhabiting neighboring
23
24 311 estuaries is not significantly different than that between bordering populations across
25
26 312 phylogeographic breaks (Neiva et al., 2012b). Similar distributional patterns can be found in
27
28 313 Brittany where several edge populations are spatially closer to populations across regional
29
30 314 genetic disjunctions than they are to their adjacent population within the same genetic clade.
31
32 315 While some studies carried out between cold- and warm-temperate marine ecosystems along
33
34 316 the northwest coast of France have invoked distinct mesoscale hydrographic features as
35
36 317 drivers of genetic patterns (Goldson et al., 2001; Jolly et al., 2005; Couceiro et al., 2013;
37
38 318 Almeida et al., 2017), others have highlighted lack of evidence for the role of hydrodynamics
39
40 319 as dispersal determinants for the observed genetic discontinuities along NW Iberian shores
41
42 320 (Neiva et al., 2012b). The latter works have also stressed the difficulties to track drifters'
43
44 321 movements at a scale relevant for the organism to estimate migration rates among estuaries
45
46 322 and phylogroups. In NW Iberia, circulation dynamics are complex with high seasonal variability
47
48 323 and lack of persistent oceanographic patterns (Ruiz-Villarreal et al., 2006; Alvarez et al.,
49
50 324 2009). Here, through the use of large scale, dispersal simulations we reveal several
51
52 325 oceanographic regions matching haplotype segregation. Our findings add important evidence
53
54 326 to previous conclusions and support the scenario of key, mesoscale oceanographic processes
55
56 327 having a determinant role in explaining the position of the observed high levels of regional
57
58 328 genetic divergence. We also hypothesized that mesoscale oceanographic dispersal barriers
59
60

1
2
3 329 are key to the maintenance of inter-estaurine genetic differentiation. In a neutral model of
4
5 330 secondary contact following allopatric differentiation, signs of secondary intergradation are
6
7 331 generally observed around oceanographic barriers (Woodruff, 1973; Barton and Hewitt, 1985;
8
9 332 Bierne et al., 2011). Admixed nuclear background has only been reported between two
10
11 333 neighboring *F. ceranoides* Iberian populations (Neiva et al., 2012b), indicating that individuals
12
13 334 belonging to distinct phylogroups can interbreed. However, no signs of hybridization have
14
15 335 been observed in other Iberian or French populations. The geographically restricted and
16
17 336 limited lineage admixture suggests that other factors may be at play in maintaining fine-scale
18
19 337 genetic differentiation. In particular, incipient reproductive isolation (pre- or post-zygotic) can
20
21 338 depress gene flow between divergent phylogroups (e.g., Tellier et al., 2011). Under this
22
23 339 scenario, limited dispersal across oceanographic barriers explains the position of the genetic
24
25 340 discontinuity while the delay in homogenisation is mainly explained by endogenous
26
27 341 components of reproductive isolation. Future studies assessing spatial and temporal
28
29 342 reproductive dynamics of distinct *F. ceranoides* lineages will be crucial to provide a direct
30
31 343 testing of this hypothesis.

344 **Conclusions**

345 Our study highlights the need to combine evidence from multiple sources for a comprehensive
346 understanding of ecological and evolutionary mechanisms linked to phylogeographic breaks.
347 These conclusions are of great significance for other organisms with sporadic and spatially
348 limited dispersal, helping clarify the apparent inconsistency of extensive and sharp genetic
349 elines differentiation in geographically restricted regions. In addition to theoretical evolutionary
350 relevance, the identification of contemporary dynamics contributing to the maintenance of
351 significant units of intraspecific biodiversity is critical for efficient approaches of management
352 and conservation efforts.

353

354 **ACKNOWLEDGMENTS**

1
2
3 355 This research was funded through projects BIODIVERSA/004/2015,
4
5 356 IF/01413/2014/CP1217/CT0004 and UID/Multi/04326/2019 by the Foundation for Science
6
7 357 and Technology (FCT - MEC, Portugal) and further supported by the National Research
8
9 358 Foundation of South Africa (Grant number 64801) and through the fellowship grant
10
11 359 SFRH/BPD/88935/2012 funded by FCT and SFRH/BPD/111003/2015 jointly funded by a
12
13 360 Pew Marine Fellowship (USA) and FCT. We thank two anonymous reviewers for their
14
15
16 361 suggestions and comments.
17
18 362

363 REFERENCES

- 364 Almeida, S. C., Nicastro, K. R., Zardi, G. I., Pearson, G. A., Valero, M., and Serrão, E. A.
365 2017. Reproductive strategies and population genetic structure of *Fucus* spp. across a
366 northeast Atlantic biogeographic transition. *Aquatic Living Resources*, 30: 16.
- 367 Alvarez, I., Ospina-Alvarez, N., Pazos, Y., Bernardez, P., Campos, M., Gomez-Gesteira, J.,
368 Alvarez-Ossorio, M., et al. 2009. A winter upwelling event in the Northern Galician Rias:
369 frequency and oceanographic implications. *Estuarine, Coastal and Shelf Science*, 82: 573-
370 582.
- 371 Assis, J., Bercibar, E., Claro, B., Alberto, F., Reed, D., Raimondi, P., and Serrão, E. 2017.
372 Major shifts at the range edge of marine forests: the combined effects of climate changes
373 and limited dispersal. *Scientific Reports*, 7: 44348.
- 374 Assis, J., Serrão, E. A., Coelho, N. C., Tempera, F., Valero, M., Claro, B., and Alberto, F.
375 2018. Past climate changes and strong oceanographic barriers structured low latitude
376 genetic relics for the golden kelp *Laminaria ochroleuca*. *Journal of Biogeography*.
- 377 Assis, J., Zupan, M., Nicastro, K. R., Zardi, G. I., McQuaid, C. D., and Serrão, E. A. 2015.
378 Oceanographic conditions limit the spread of a marine invader along southern African
379 shores. *PLoS ONE*, 10(6):e0128124.

- 1
2
3 380 Ayata, S.-D., Stolba, R., Comtet, T., and Thiébaud, É. 2011. Meroplankton distribution and its
4
5 381 relationship to coastal mesoscale hydrological structure in the northern Bay of Biscay (NE
6
7 382 Atlantic). *Journal of Plankton Research*, 33: 1193-1211.
8
9 383 Barnes, I., Matheus, P., Shapiro, B., Jensen, D., and Cooper, A. 2002. Dynamics of
10
11 384 Pleistocene Population Extinctions in Beringian Brown Bears. *Science*, 295: 2267-2270.
12
13 385 Barton, N. H., and Hewitt, G. M. 1985. Analysis of hybrid zones. *Annual Review of Ecology
14
15 386 and Systematics*, 16: 113-148.
16
17 387 Bierne, N., Welch, J., Loire, E., Bonhomme, F., and David, P. 2011. The coupling
18
19 388 hypothesis: why genome scans may fail to map local adaptation genes. *Molecular Ecology*,
20
21 389 20: 2044-2072.
22
23 390 Chao, R. B., Casais, M. C., Cortizas, A. M. n., Alberti, A. P., and Paz, M. V. 2002. Holocene
24
25 391 evolution on Galician coast (NW Spain): an example of paraglacial dynamics. *Quaternary
26
27 392 International*, 93: 149-159.
28
29 393 Chassignet, E. P., Hurlburt, H. E., Smedstad, O. M., Halliwell, G. R., Hogan, P. J., Wallcraft,
30
31 394 A. J., Baraille, R., et al. 2007. The HYCOM (hybrid coordinate ocean model) data
32
33 395 assimilative system. *Journal of Marine Systems*, 65: 60-83.
34
35 396 Chust, G., Villarino, E., Chenuil, A., Irigoien, X., Bizsel, N., Bode, A., Broms, C., et al. 2016.
36
37 397 Dispersal similarly shapes both population genetics and community patterns in the marine
38
39 398 realm. *Scientific reports*, 6: 28730.
40
41 399 Clement, M., Posada, D., and Crandall, K. A. 2000. TCS: a computer program to estimate
42
43 400 gene genealogies. *Molecular Ecology*, 9: 1657-1659.
44
45 401 Comps, B., Gömöry, D., Letouzey, J., Thiébaud, B., and Petit, R. 2001. Diverging trends
46
47 402 between heterozygosity and allelic richness during postglacial colonization in the European
48
49 403 beech. *Genetics*, 157: 389-397.
50
51 404 Couceiro, L., Robuchon, M., Destombe, C., and Valero, M. 2013. Management and
52
53 405 conservation of the kelp species *Laminaria digitata*: using genetic tools to explore the
54
55 406 potential exporting role of the MPA "Parc naturel marin d'Iroise". *Aquatic Living Resources*,
56
57 407 26: 197-205.
58
59
60

- 1
2
3 408 Day, J. W., Hall, C. A. S., Kemp, W. M., and A., Y.-A. 1989. Estuarine Ecology, John Wiley
4 and Sons, New York.
5
6 409
7 410 Dowle, M., Srinivasan, A., Gorecki, J., Chirico, M., Stetsenko, P., Short, T., Lianoglou, S., et
8
9 411 al. 2019. Package 'data.table'. Extension of 'data.frame'.
10
11 412 Drummond, A., Ashton, B., Buxton, S., Cheung, M., Cooper, A., Duran, C., Field, M., et al.
12
13 413 2011. Geneious v5. 4.
14
15 414 Ferro, I., and Morrone, J. J. 2014. Biogeographical transition zones: a search for conceptual
16
17 415 synthesis. *Biological Journal of the Linnean Society*, 113: 1-12.
18
19 416 Fossette, S., Putman, N. F., Lohmann, K. J., Marsh, R., and Hays, G. C. 2012. A biologist's
20
21 417 guide to assessing ocean currents: a review. *Marine Ecology Progress Series*, 457: 285-301.
22
23 418 Gallon, R. K., Robuchon, M., Leroy, B., Le Gall, L., Valero, M., and Feunteun, E. 2014.
24
25 419 Twenty years of observed and predicted changes in subtidal red seaweed assemblages
26
27 420 along a biogeographical transition zone: inferring potential causes from environmental data.
28
29 421 *Journal of Biogeography*, 41: 2293-2306.
30
31 422 Gilg, M. R., and Hilbish, T. J. 2003. The geography of marine larval dispersal: coupling
32
33 423 genetics with fine-scale physical oceanography. *Ecology*, 84: 2989-2998.
34
35 424 Goldson, A., Hughes, R., and Gliddon, C. 2001. Population genetic consequences of larval
36
37 425 dispersal mode and hydrography: a case study with bryozoans. *Marine Biology*, 138: 1037-
38
39 426 1042.
40
41 427 Gomez, A., Hughes, R. N., Wright, P. J., Carvalho, G. R., and Lunt, D. H. 2007.
42
43 428 Mitochondrial DNA phylogeography and mating compatibility reveal marked genetic
44
45 429 structuring and speciation in the NE Atlantic bryozoan *Celleporella hyalina*. *Molecular*
46
47 430 *Ecology*, 16: 2173-2188.
48
49 431 Haklay, M., and Weber, P. 2008. Openstreetmap: User-generated street maps. *IEEE*
50
51 432 *Pervasive Computing*, 7: 12-18.
52
53 433 Hewitt, G. 2000. The genetic legacy of the Quaternary ice ages. *Nature*, 405: 907-913.
54
55 434 Hewitt, G. M. 1996. Some genetic consequences of ice ages, and their role in divergence
56
57 435 and speciation. *Biological Journal of the Linnean Society*, 58: 247-276.
58
59
60

- 1
2
3 436 Hewitt, G. M. 2004. Genetic consequences of climatic oscillations in the Quaternary.
4
5 437 Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences,
6
7 438 359: 183-195.
8
9 439 Hijmans, R. J., Phillips, S., Leathwick, J., Elith, J., and Hijmans, M. R. J. 2017. Package
10
11 440 'dismo'. *Circles*, 9: 1-68.
12
13 441 Hoarau, G., Coyer, J. A., Veldsink, J. H., Stam, W. T., and Olsen, J. L. 2007. Glacial refugia
14
15 442 and recolonization pathways in the brown seaweed *Fucus serratus*. *Molecular Ecology*, 16:
16
17 443 3606-3616.
18
19 444 Hohenlohe, P. A. 2004. Limits to gene flow in marine animals with planktonic larvae: models
20
21 445 of *Littorina* species around Point Conception, California. *Biological Journal of the Linnean*
22
23 446 *Society*, 82: 169-187.
24
25 447 Johansson, M. L., Alberto, F., Reed, D. C., Raimondi, P. T., Coelho, N. C., Young, M. A.,
26
27 448 Drake, P. T., et al. 2015. Seascape drivers of *Macrocystis pyrifera* population genetic
28
29 449 structure in the northeast Pacific. *Molecular Ecology*, 24: 4866-4885.
30
31 450 Jolly, M. T., Jollivet, D., Gentil, F., Thiébaud, É., and Viard, F. 2005. Sharp genetic break
32
33 451 between Atlantic and English Channel populations of the polychaete *Pectinaria koreni*, along
34
35 452 the North coast of France. *Heredity*, 94: 23.
36
37 453 Kamvar, Z. N., Tabima, J. F., and Grünwald, N. J. 2014. Poppr: an R package for genetic
38
39 454 analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ*, 2:
40
41 455 e281.
42
43 456 Kennedy, J. P., Garavelli, L., Truelove, N. K., Devlin, D. J., Box, S. J., Chérubin, L. M., and
44
45 457 Feller, I. C. 2017. Contrasting genetic effects of red mangrove (*Rhizophora mangle* L.) range
46
47 458 expansion along West and East Florida. *Journal of Biogeography*, 44: 335-347.
48
49 459 Klein, M., Teixeira, S., Assis, J., Serrão, E. A., Gonçalves, E. J., and Borges, R. 2016. High
50
51 460 interannual variability in connectivity and genetic pool of a temperate clingfish matches
52
53 461 oceanographic transport predictions. *PloS one*, 11: e0165881.
54
55 462 Lamigueiro, O. P., Hijmans, R., and Lamigueiro, M. O. P. 2018. Package 'rasterVis'.
56
57
58
59
60

- 1
2
3 463 Lein, T. E. 1984. Distribution, reproduction, and ecology of *Fucus ceranoides*
4
5 464 L.(Phaeophyceae) in Norway. *Sarsia*, 69: 75-81.
6
7 465 Lett, C., Verley, P., Mullon, C., Parada, C., Brochier, T., Penven, P., and Blanke, B. 2008. A
8
9 466 Lagrangian tool for modelling ichthyoplankton dynamics. *Environmental Modelling &*
10
11 467 *Software*, 23: 1210-1214.
12
13 468 Liang, Y., He, D., Jia, Y., Sun, H., and Chen, Y. 2017. Phylogeographic studies of
14
15 469 schizothoracine fishes on the central Qinghai-Tibet Plateau reveal the highest known glacial
16
17 470 microrefugia. *Scientific Reports*, 7: 10983.
18
19 471 Librado, P., and Rozas, J. 2009. DnaSP v5: a software for comprehensive analysis of DNA
20
21 472 polymorphism data. *Bioinformatics*, 25: 1451-1452.
22
23 473 McKenzie, P. F., and Bellgrove, A. 2008. Dispersal of *Hormosira banksii* (Phaeophyceae) via
24
25 474 detached fragments: reproductive viability and longevity. *Journal of Phycology*, 44: 1108-
26
27 475 1115.
28
29 476 Mims, M. C., Phillipsen, I. C., Lytle, D. A., Kirk, E. E. H., and Olden, J. D. 2015. Ecological
30
31 477 strategies predict associations between aquatic and genetic connectivity for dryland
32
33 478 amphibians. *Ecology*, 96: 1371-1382.
34
35 479 Monteiro, C. A., Paulino, C., Jacinto, R., Serrão, E. A., and Pearson, G. A. 2016. Temporal
36
37 480 windows of reproductive opportunity reinforce species barriers in a marine broadcast
38
39 481 spawning assemblage. *Scientific reports*, 6: 29198.
40
41 482 Muhlin, J. F., Engel, C. R., Stessel, R., Weatherbee, R. A., and Brawley, S. H. 2008. The
42
43 483 influence of coastal topography, circulation patterns, and rafting in structuring populations of
44
45 484 an intertidal alga. *Molecular Ecology*, 17: 1198-1210.
46
47 485 Munda, I. M. 1999. The benthic algal vegetation of land-locked fjords in southeastern
48
49 486 Iceland. *Hydrobiologia*, 393: 169.
50
51 487 Munwes, I., Geffen, E., Roll, U., Friedmann, A., Daya, A., Tikochinski, Y., and Gafny, S.
52
53 488 2010. The change in genetic diversity down the core-edge gradient in the eastern spadefoot
54
55 489 toad (*Pelobates syriacus*). *Molecular Ecology*, 19: 2675-2689.
56
57
58
59
60

- 1
2
3 490 Neiva, J., Paulino, C., Nielsen, M. M., Krause-Jensen, D., Saunders, G. W., Assis, J.,
4
5 491 Bárbara, I., et al. 2018. Glacial vicariance drives phylogeographic diversification in the
6
7 492 amphiboreal kelp *Saccharina latissima*. *Scientific reports*, 8: 1112.
8
9 493 Neiva, J., Pearson, G. A., Valero, M., and Serrão, E. A. 2010. Surfing the wave on a
10
11 494 borrowed board: range expansion and spread of introgressed organellar genomes in the
12
13 495 seaweed *Fucus ceranoides* L. *Molecular Ecology*, 19: 4812-4822.
14
15 496 Neiva, J., Pearson, G. A., Valero, M., and Serrão, E. A. 2012a. Drifting fronds and drifting
16
17 497 alleles: range dynamics, local dispersal and habitat isolation shape the population structure
18
19 498 of the estuarine seaweed *Fucus ceranoides*. *Journal of Biogeography*, 39: 1167-1178.
20
21 499 Neiva, J., Pearson, G. A., Valero, M., and Serrão, E. A. 2012b. Fine-scale genetic breaks
22
23 500 driven by historical range dynamics and ongoing density-barrier effects in the estuarine
24
25 501 seaweed *Fucus ceranoides* L. *BMC Evolutionary Biology*, 12: 78.
26
27 502 Newman, M. E. 2006. Modularity and community structure in networks. *Proceedings of the*
28
29 503 *national academy of sciences*, 103: 8577-8582.
30
31 504 Nicastro, K. R., Zardi, G. I., McQuaid, C. D., Teske, P. R., and Barker, N. P. 2008. Coastal
32
33 505 topography drives genetic structure in marine mussels. *Marine Ecology Progress Series*,
34
35 506 368: 189-195.
36
37 507 Nicastro, K. R., Zardi, G. I., Teixeira, S., Neiva, J., Serrao, E. A., and Pearson, G. A. 2013.
38
39 508 Shift happens: trailing edge contraction associated with recent warming trends threatens a
40
41 509 distinct genetic lineage in the marine macroalga *Fucus vesiculosus*. *BMC Biology*, 11: 6.
42
43 510 Oksanen, J., Blanchet, F., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P., et
44
45 511 al. 2018. *vegan: a community ecology package*. R package versions 2.4-6.
46
47 512 Olsen, J. L., Zechman, F. W., Hoarau, G., Coyer, J. A., Stam, W. T., Valero, M., and Åberg,
48
49 513 P. 2010. The phylogeographic architecture of the fucoid seaweed *Ascophyllum nodosum*: an
50
51 514 intertidal 'marine tree' and survivor of more than one glacial-interglacial cycle. *Journal of*
52
53 515 *Biogeography*, 37: 842-856.
54
55 516 Pardo, C., Guillemain, M.-L., Pena, V., Barbara, I., Valero, M., and Barreiro, R. 2019. Local
56
57 517 coastal configuration rather than latitudinal gradient shape clonal diversity and genetic

- 1
2
3 518 structure of *Phymatolithon calcareum* maerl beds in North European Atlantic. *Frontiers in*
4
5 519 *Marine Science*, 6: 149.
6
7 520 Provan, J. 2013. The effects of past, present and future climate change on range-wide
8
9 521 genetic diversity in northern North Atlantic marine species. *Frontiers of Biogeography*, 5(1).
10
11 522 Provan, J., Wattier, R. A., and Maggs, C. A. 2005. Phylogeographic analysis of the red
12
13 523 seaweed *Palmaria palmata* reveals a Pleistocene marine glacial refugium in the English
14
15 524 Channel. *Molecular Ecology*, 14: 793-803.
16
17 525 Puillat, I., Lazure, P., Jégou, A., Lampert, L., and Miller, P. 2004. Hydrographical variability
18
19 526 on the French continental shelf in the Bay of Biscay, during the 1990s. *Continental Shelf*
20
21 527 *Research*, 24: 1143-1163.
22
23 528 Putman, N. F., and He, R. 2013. Tracking the long-distance dispersal of marine organisms:
24
25 529 sensitivity to ocean model resolution. *Journal of the Royal Society Interface*, 10: 20120979.
26
27 530 Remerie, T., Vierstraete, A., Weekers, P. H., Vanfleteren, J. R., and Vanreusel, A. 2009.
28
29 531 Phylogeography of an estuarine mysid, *Neomysis integer* (Crustacea, Mysida), along the
30
31 532 north-east Atlantic coasts. *Journal of Biogeography*, 36: 39-54.
32
33 533 Roucoux, K., De Abreu, L., Shackleton, N., and Tzedakis, P. 2005. The response of NW
34
35 534 Iberian vegetation to North Atlantic climate oscillations during the last 65kyr. *Quaternary*
36
37 535 *Science Reviews*, 24: 1637-1653.
38
39 536 Ruiz-Villarreal, M., González-Pola, C., del Río, G. D., Lavin, A., Otero, P., Piedracoba, S.,
40
41 537 and Cabanas, J. 2006. Oceanographic conditions in North and Northwest Iberia and their
42
43 538 influence on the Prestige oil spill. *Marine Pollution Bulletin*, 53: 220-238.
44
45 539 Selkoe, K. A., Watson, J. R., HORIN, T. B., IACCHEI, M., MITARAI, S., SIEGEL, D. A.,
46
47 540 GAINES, S. D., et al. 2010. Taking the chaos out of genetic patchiness: seascape genetics
48
49 541 reveals ecological and oceanographic drivers of genetic patterns in three temperate reef
50
51 542 species. *Molecular Ecology*, 19: 3708-3726.
52
53 543 Serrão, E. A., Kautsky, L., Lifvergren, T., and Brawley, S. H. 1997. Gamete dispersal and
54
55 544 pre-recruitment mortality in Baltic *Fucus vesiculosus*. *Phycologia*, 36 101-102.
56
57
58
59
60

- 1
2
3 545 Slatkin, M. 1993. Isolation by distance in equilibrium and non-equilibrium populations.
4
5 546 Evolution, 47: 264-279.
6
7 547 Smith, B. T., Escalante, P., Baños, B. E. H., Navarro-Sigüenza, A. G., Rohwer, S., and
8
9 548 Klicka, J. 2011. The role of historical and contemporary processes on phylogeographic
10
11 549 structure and genetic diversity in the Northern Cardinal, *Cardinalis cardinalis*. BMC
12
13 550 Evolutionary Biology, 11: 136.
14
15 551 Sotka, E. E., Wares, J. P., Barth, J. A., Grosberg, R. K., and Palumbi, S. R. 2004. Strong
16
17 552 genetic clines and geographical variation in gene flow in the rocky intertidal barnacle
18
19 553 *Balanus glandula*. Molecular Ecology, 13: 2143-2156.
20
21 554 Spalding, M. D., Fox, H. E., Allen, G. R., Davidson, N., Ferdana, Z. A., Finlayson, M.,
22
23 555 Halpern, B. S., et al. 2007. Marine ecoregions of the world: a bioregionalization of coastal
24
25 556 and shelf areas. Bioscience, 57: 573-583.
26
27 557 Team, R. C. 2018. R: A language and environment for statistical computing; 2015.
28
29 558 Tellier, F., Tapia, J., Faugeton, S., Destombe, C., and Valero, M. 2011. The *lessonia*
30
31 559 *nigrescens* species complex (laminariales, phaeophyceae) shows strict parapatry and
32
33 560 complete reproductive isolation in a secondary contact zone 1. Journal of Phycology, 47:
34
35 561 894-903.
36
37 562 Teske, P., Papadopoulos, I., Zardi, G., McQuaid, C., Edkins, M., Griffiths, C., and Barker, N.
38
39 563 2007. Implications of life history for genetic structure and migration rates of southern African
40
41 564 coastal invertebrates: planktonic, abbreviated and direct development. Marine Biology, 152:
42
43 565 697-711.
44
45 566 Teske, P. R., Von Der Heyden, S., McQuaid, C. D., and Barker, N. P. 2011. A review of
46
47 567 marine phylogeography in southern Africa. South African Journal of Science, 107: 43-53.
48
49 568 Thiel, M., and Haye, P. 2006. The ecology of rafting in the marine environment-III.
50
51 569 Biogeographical and evolutionary consequences (in Oceanography and marine biology: An
52
53 570 annual review-Volume 48, pp. 323-429).
54
55
56
57
58
59
60

- 1
2
3 571 Varela, R. A., Rosón, G., Herrera, J. L., Torres-López, S., and Fernández-Romero, A. 2005.
4
5 572 A general view of the hydrographic and dynamical patterns of the Rías Baixas adjacent sea
6
7 573 area. *Journal of Marine Systems*, 54: 97-113.
8
9 574 Wares, J. P., Gaines, S., and Cunningham, C. W. 2001. A comparative study of asymmetric
10
11 575 migration events across a marine biogeographic boundary. *Evolution*, 55: 295-306.
12
13 576 Woodruff, D. S. 1973. Natural hybridization and hybrid zones. *Systematic Biology*, 22: 213-
14
15 577 218.
16
17 578 Zardi, G., McQuaid, C., Teske, P., and Barker, N. 2007. Unexpected genetic structure of
18
19 579 mussel populations in South Africa: indigenous *Perna perna* and invasive *Mytilus*
20
21 580 *galloprovincialis*. *Marine Ecology Progress Series*, 337: 135-144.
22
23 581 Zardi, G., Nicastro, K., McQuaid, C., Hancke, L., and Helmuth, B. 2011. The combination of
24
25 582 selection and dispersal helps explain genetic structure in intertidal mussels. *Oecologia*, 165:
26
27 583 947-958.
28
29 584 Zardi, G. I., Nicastro, K. R., Ferreira Costa, J., Serrão, E. A., and Pearson, G. A. 2013.
30
31 585 Broad scale agreement between intertidal habitats and adaptive traits on a basis of
32
33 586 contrasting population genetic structure. *Estuarine, Coastal and Shelf Science*, 131: 140-
34
35 587 148.
36
37 588
38
39
40
41
42 589
43
44
45 590
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 591 **FIGURE LEGEND**
4
5

6 592 **Fig. 1.** Haplotype distribution of *Fucus ceranoides* in the coastlines of (a) Northwester France
7
8 593 and (panel b) Northwest Iberia. The colors along coastlines (a, b) depict the different
9
10 594 oceanographic regions (R) identified in network analysis. A pairwise matrix of directional
11
12 595 connectivity between oceanographic regions identified in (d) western Brittany and (e)
13
14 596 northwest Iberia. Matrix header colors according to the oceanographic regions depicted in the
15
16 597 top panels a and b. Haplotype frequency and differentiation degree in (f) western Brittany and
17
18 598 (g) northwest Iberia. Note that there is no correspondence of haplotypes between the two
19
20 599 panels (i.e., each dataset was treated separately).
21
22
23

24 600

25
26 601 **TABLES**
27

28 602 **Table 1.** Maximum and average distances (km), drifting time (days) and probabilities produced
29
30 603 by the particles connecting different cells for the lagrangian particle simulations running in
31
32 604 Northwest France and Northwest Spain..
33
34

35 605

Region	Distance (km)		Time (days)		Probability	
	Maximum	Mean (\pm SD)	Maximum	Mean (\pm SD)	Maximum	Mean (\pm SD)
W Brittany	382.6	62.6 \pm 54.6	24.6	3.61 \pm 2.63	0.721	0.004 \pm 0.020
NW Iberia	431.2	105.9 \pm 90.8	26.7	2.09 \pm 1.81	0.607	0.004 \pm 0.021

36
37
38
39
40
41
42
43
44
45 606
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60