



# ON THE BETTI MAP ASSOCIATED TO ABELIAN LOGARITHMS

Yves Andre, Pietro Xxx Corvaja, Umberto Zannier

## ► To cite this version:

Yves Andre, Pietro Xxx Corvaja, Umberto Zannier. ON THE BETTI MAP ASSOCIATED TO ABELIAN LOGARITHMS. 2019. hal-02408097

**HAL Id: hal-02408097**

**<https://hal.science/hal-02408097>**

Preprint submitted on 12 Dec 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ON THE BETTI MAP ASSOCIATED TO ABELIAN LOGARITHMS

Y. ANDRÉ, P. CORVAJA, U. ZANNIER

ABSTRACT. (draft 5)

## 1. PROBLEM AND MAIN RESULTS.

**1.1.** Let  $A$  be a complex abelian variety of dimension  $g$ . We write  $\Lambda$  for the its *period lattice*  $H_1(A(\mathbb{C}), \mathbb{Z})$ , identified with the kernel of the exponential map  $\exp_A : \text{Lie } A \rightarrow A$ .

We fix a complex point  $\sigma$  of  $A$ , and let  $\Lambda_\sigma$  be the lattice formed by the pairs  $(\ell \in \text{Lie } A, m \in \mathbb{Z})$  such that  $\exp_A(\ell) = m\sigma$ . It sits in an extension

$$(1.1) \quad 0 \rightarrow \Lambda \rightarrow \Lambda_\sigma \xrightarrow{(\ell, m) \mapsto m} \mathbb{Z}\sigma \rightarrow 0,$$

and we fix a splitting  $\lambda : \mathbb{Z}\sigma \rightarrow \Lambda_\sigma$  (*i.e.* a branch of the “abelian logarithm”). On the other hand, let us consider the first projection  $\Lambda_\sigma \xrightarrow{(\ell, m) \mapsto \ell} \text{Lie } A$  and its  $\mathbb{R}$ -linear extension  $\Lambda_\sigma \otimes_{\mathbb{Z}} \mathbb{R} \rightarrow \text{Lie } A$ . Since the composed map

$$(1.2) \quad \Lambda \otimes_{\mathbb{Z}} \mathbb{R} \rightarrow \Lambda_\sigma \otimes_{\mathbb{Z}} \mathbb{R} \rightarrow \text{Lie } A$$

is an isomorphism, one has a canonical projection  $\Lambda_\sigma \otimes_{\mathbb{Z}} \mathbb{R} \rightarrow \Lambda \otimes_{\mathbb{Z}} \mathbb{R}$ , and the image of  $\lambda(\sigma)$  in  $\Lambda \otimes_{\mathbb{Z}} \mathbb{R}$  can be expressed, in terms of a basis  $\underline{\ell}$  of  $\Lambda$ , by  $2g$  real coordinates: the *Betti coordinates*<sup>1</sup> of  $\sigma$ .

**1.2.** We are actually interested in the relative setting. Let  $S$  be a smooth connected complex algebraic variety.

Let  $A \xrightarrow{f} S$  be an abelian scheme of relative dimension  $g$ . Its Lie algebra  $\text{Lie } A$  is a rank  $g$  vector bundle on  $S$ . For our purpose, there will be no

---

*Date:* June 16, 2016.

*1991 Mathematics Subject Classification.*

*Key words and phrases.*

<sup>1</sup>the terminology seems due to D. Bertrand and refers to the fact that  $\Lambda_\sigma$  is nothing but the *Betti realization of the 1-motive*  $[\mathbb{Z} \xrightarrow{1 \mapsto \sigma} A]$  in the sense of [D2, X]. Note that the Betti coordinates depend not only on  $\sigma$  but also on the auxiliary choice of a splitting  $\lambda$  and of a basis of  $\Lambda$ .

loss of generality<sup>2</sup> in assuming, as we shall do, that  $S$  is affine and that this vector bundle is trivial, i.e.  $\Gamma \text{Lie } A$  is a free  $\Gamma \mathcal{O}_S$ -module of rank  $g$ .

Let  $\sigma : S \rightarrow A$  be a section of  $f$ . The above constructions extend to this relative setting as follows. Let  $\tilde{S}$  be a universal covering of  $S(\mathbb{C})$ , with its canonical structure of complex analytic manifold. The kernel of  $\exp_A$  is a locally constant sheaf on  $S(\mathbb{C})$ , which can be viewed as a constant lattice  $\Lambda$  on  $\tilde{S}$ . Similarly, one constructs an exact sequence of lattices on  $\tilde{S}$

$$(1.3) \quad 0 \rightarrow \Lambda \rightarrow \Lambda_\sigma \xrightarrow{(\ell, m) \mapsto m} \mathbb{Z}\sigma \rightarrow 0$$

and fixes a splitting  $\lambda : \mathbb{Z}\sigma \rightarrow \Lambda_\sigma$ .

Let  $\tilde{S}^{\text{real}}$  be  $\tilde{S}$  viewed as a real-analytic manifold and let  $\mathcal{O}^{\mathbb{R}}(\tilde{S}^{\text{real}})$  (*resp.*  $\mathcal{O}(\tilde{S}^{\text{real}})$ ) be the ring of real-valued (*resp.* complex valued) real-analytic functions on  $\tilde{S}$ . The composed map

$$(1.4) \quad \Lambda \otimes_{\mathbb{Z}} \mathcal{O}^{\mathbb{R}}(\tilde{S}^{\text{real}}) \rightarrow \Lambda_\sigma \otimes_{\mathbb{Z}} \mathcal{O}^{\mathbb{R}}(\tilde{S}^{\text{real}}) \rightarrow \Gamma \text{Lie } A \otimes_{\Gamma \mathcal{O}_S} \mathcal{O}(\tilde{S}^{\text{real}})$$

is an isomorphism of free  $\mathcal{O}^{\mathbb{R}}(\tilde{S}^{\text{real}})$ -modules. One thus has a canonical projection  $\Lambda_\sigma \otimes_{\mathbb{Z}} \mathcal{O}^{\mathbb{R}}(\tilde{S}^{\text{real}}) \rightarrow \Lambda \otimes_{\mathbb{Z}} \mathcal{O}^{\mathbb{R}}(\tilde{S}^{\text{real}})$ , and the image of  $\lambda(\sigma)$  in  $\Lambda \otimes_{\mathbb{Z}} \mathcal{O}^{\mathbb{R}}(\tilde{S}^{\text{real}})$  can be expressed, in terms of a basis  $\underline{\ell}$  of the lattice  $\Lambda$ , by a real-analytic map (the *Betti map*)

$$(1.5) \quad \beta = \beta(\sigma, \lambda, \underline{\ell}) : \tilde{S} \rightarrow \mathbb{R}^{2g}.$$

**1.3.** We assume for convenience that  $A \rightarrow S$  is *principally polarized*, which allows to consider to the *moduli map*  $\mu : S \rightarrow \mathcal{A}_g$ . We denote by  $A_{\bar{\eta}}$  the geometric generic fiber, and we set

$$(1.6) \quad d = \dim \mu(S).$$

We are interested in the range of  $\beta$ . It is easy to see that the its dimension is  $\leq \min(2d, 2g)$ . *Under what condition is the range of  $\beta$  maximal, i.e. of dimension  $\min(2d, 2g)$ ?*

This may not be the case for various trivial reasons (*e.g.* when the  $\sigma$  is a torsion section, or when  $A \xrightarrow{f} S$  is the pull-back of an abelian scheme defined over a base of dimension  $< d \leq g$ ).

**1.4.** Here is a sample of concrete results [not our final results, hopefully!]:

**1.4.1. Theorem.** *Assume that  $g \leq \min(3, d)$  and that the section  $\sigma$  is not contained in a proper subgroup scheme of  $A$ . Then the range of  $\beta$  has real dimension  $2g$ . More precisely, the restriction of  $\beta$  to the inverse image of a dense Zariski-open subset of  $S$  is a submersion.*

---

<sup>2</sup>since  $S$  can be covered by such affine open subsets.

The proof relies on an analysis of the Kodaira-Spencer map.

**1.4.2. Theorem.** *Assume that  $\sigma$  is not contained in a proper subgroup scheme of  $A$ . Assume moreover that  $d \leq \min(2, g)$ . Then  $\beta$  has real image  $2d$ . More precisely, the restriction of  $\beta$  to the inverse image of a dense Zariski-open subset of  $S$  is an immersion.*

**1.5.** *[Motivation for the problem, and application of the results (if any)... General conjecture ....]*

## 2. AN INFINITESIMAL APPROACH, AND ITS HOLOMORPHIC TRANSLATION.

**2.1.** Our approach is infinitesimal. In fact, the problem we address is the following:

*under what condition is  $\beta$  almost everywhere an immersion, resp. a submersion (according to whether  $d \leq g$  or  $g \leq d$ ) ?*

**2.1.1. Proposition.** *For every  $s \in \tilde{S}$ ,  $\text{Ker } \beta(s)$  is a complex vector subspace of  $\text{Lie } A_s \cong \mathbb{R}^{2g}$ . Moreover the fibers of  $\beta$  are complex-analytic subspaces of  $\tilde{S}$ , and all non-empty fibers have dimension  $\geq 2d - r$ , where  $r$  is the generic rank of  $\beta$ .*

*Proof.* - [cf. your prop 2.1 + 2.2]. □

**2.2.** We may first replace  $S$  by a finite etale covering and assume that for some  $n \geq 3$ , the  $n$ -torsion sections are defined over  $S$ , then replace it by its image in the moduli space  $\mathcal{A}_{g,n}$  of principally polarized abelian varieties of dimension  $g$  with level  $n$ -structure. We thus consider the rank of a jacobian matrix of size  $(2d, 2g)$ , and for this purpose, we introduce the following notation:

$(\omega_1, \dots, \omega_g, \eta_1, \dots, \eta_g)$  is a symplectic basis of  $\mathcal{H}_{dR}^1(A/S)$  with  $\omega_1, \dots, \omega_g \in \Omega_A$ ,

$(\gamma_1, \dots, \gamma_g)$  is a symplectic basis of  $\Lambda$ ,

$\Omega_1 := \left( \int_{\gamma_i} \omega_j \right)_{i,j=1,\dots,g}$ ,  $\Omega_2 := \left( \int_{\gamma_{i+g}} \omega_j \right)_{i,j=1,\dots,g}$ ,  $Z := \Omega_1 \cdot \Omega_2^{-1} = {}^t Z$

(which takes values in  $\mathfrak{H}_g$ ),  $L := \left( \int^\sigma \omega_j \right)_{j=1,\dots,g} \cdot \Omega_2^{-1}$ .

Let  $r$  be  $\min(d, g)$ . In the next step, we transform the vanishing of a  $2r$ - $2r$ -minor involving real-analytic functions by the vanishing of a  $r$ - $r$ -minor of a related matrix involving only complex-analytic functions on  $\tilde{S}$ . In order to do this, we take advantage of the monodromy, using a theorem of linear independence of abelian logarithms with respect to periods [A1, th. 2] (cf. also [A2, §1]).

At the end, we come along with the following translation of the problem (under extra assumptions):

**2.2.1. Proposition.** (with  $S \subset \mathcal{A}_{g,n}$ ). Assume that  $\sigma$  is not contained in a proper sub-group scheme. Then for any  $s \in \tilde{S}$ ,  $\beta$  is an immersion, resp. a submersion (according to whether  $d \leq g$  or  $g \leq d$ ) at  $s$  if and only if for some  $(\lambda_1, \dots, \lambda_g) \in \mathbb{C}^g$ , the matrix

$$(2.1) \quad ((\partial_i L)(s) + \sum_{k=1, \dots, g} \lambda_k (\partial_i Z_{kj})(s))_{i=1, \dots, d, j=1, \dots, g}$$

has maximal rank (where the  $\partial_i$  form a basis of the tangent space  $T_{\tilde{S}}(s)$ ).

This is already the case if for some  $(\lambda_1, \dots, \lambda_g) \in \mathbb{C}^g$ , the matrix

$$(2.2) \quad (\sum_{j=1, \dots, g} \lambda_k (\partial_i Z_{kj})(s))_{i=1, \dots, d, j=1, \dots, g}$$

has maximal rank.

*Proof.* ... [your §3.5].

The second assertion follows from the first, putting an arbitrary coefficient  $\lambda_0$  in front of  $(\partial_i L)(s)$  and letting  $\lambda_0$  tend to 0.  $\square$

*Remark.* For  $d = g$ , the negation of the conclusion leads to a special case of “Dieudonné’s problem”, which is notoriously difficult: describe vector spaces of singular matrices  $Z$ , resp. singular symmetric matrices, cf. e.g. [L]<sup>3</sup>.

### 3. ROLE OF THE KODAIRA-SPENCER MAP.

**3.1.** We now consider the following condition:

(3.1)

For every  $s$  in an dense open affine subset of  $S$ , the matrix (2.2) has maximal rank.

We shall translate it in terms of the Kodaira-Spencer map

$$\theta : T_S \otimes \Omega_A \rightarrow \Omega_A^\vee = \text{Lie } A,$$

where  $\Omega_A$  denotes the locally free  $\mathcal{O}_S$ -module of invariant 1-forms on  $A$ . We refer to [A3, 1.4, 2.1] for background about the Kodaira-Spencer map of a polarized abelian scheme  $A$  of relative dimension  $g$  over a smooth  $\mathbb{C}$ -scheme  $S$ . For any  $\partial \in \Gamma T_S$ , we denote by  $\theta_\partial$  the corresponding map  $\Omega_A \rightarrow \Omega_A^\vee$ , which belongs to  $\text{Hom}_{\text{sym}}(\Omega_A, \Omega_A^\vee)$ . Let us recall that in terms of the symplectic basis  $\omega_1, \dots, \omega_g, \eta_1, \dots, \eta_g$ , the matrix of  $\theta_\partial$  is given by the lower left quadrant  $T_\partial$  of the Gauss-Manin connection written in the form

$$\partial \begin{pmatrix} \Omega_1 & N_1 \\ \Omega_2 & N_2 \end{pmatrix} = \begin{pmatrix} \Omega_1 & N_1 \\ \Omega_2 & N_2 \end{pmatrix} \begin{pmatrix} R_\partial & S_\partial \\ T_\partial & U_\partial \end{pmatrix},$$

where  $R_\partial, S_\partial, T_\partial, U_\partial \in M_g(\mathcal{O}(S))$ . Moreover  $T_\partial$  is a symmetric matrix.

<sup>3</sup>we are grateful to M. Brion for this reference.

**3.1.1. Lemma.** *Assume that  $d = g$ . Then condition (3.1) is equivalent to:*

$$(3.2) \quad \exists \omega \in \Gamma \Omega_A, \forall \partial \in \Gamma T_S \setminus 0, \theta_{\partial} \cdot \omega \neq 0.$$

*Proof.* One has

$$\begin{aligned} \partial Z &= (\partial \Omega_1 - \Omega_1 \Omega_2^{-1} \partial \Omega_2) \Omega_2^{-1} = ((\Omega_1 R_{\partial} + N_1 T_{\partial}) - \Omega_1 \Omega_2^{-1} (\Omega_2 R_{\partial} + N_2 T_{\partial})) \Omega_2^{-1} \\ &= (N_1 - \Omega_1 \Omega_2^{-1} N_2) T_{\partial} \Omega_2^{-1} = -\Omega_2^{-1} ({}^t \Omega_1 N_2 - {}^t \Omega_2 N_1) T_{\partial} \Omega_2^{-1} = -2\pi i \Omega_2^{-1} T_{\partial} \Omega_2^{-1}. \end{aligned}$$

If  $\vec{\mu} = -2\pi i \vec{\lambda} \Omega_2^{-1}$ , one has

$$\det\left(\sum_{k=1, \dots, g} \lambda_k (\partial_i Z_{kj})(s)\right) = \det\left(\sum_{k=1, \dots, g} \mu_k (T_{\partial_i})_{kj}(s)\right) \cdot (\det \Omega_2)^{-1}.$$

Setting  $\omega = \sum \mu_k \omega_k$ , one has  $\sum \mu_k (T_{\partial_i})_{kj} = \theta_{\partial_i}(\omega) \cdot \eta_j$ . Thus (3.1) is equivalent to: there is a dense open affine subset  $S' \subset S$  such that

$$\forall s \in S'(\mathbb{C}), \exists \omega_s \in \Omega^1(A_s), \forall \partial_s \in T_{S,s} \setminus 0, (\theta(s)_{\partial_s})(\omega_s) \neq 0.$$

It is easy to see that this condition is equivalent in turn to (3.2) (for the direct implication, any  $\omega$  lifting  $\omega_s$  for some  $s$  will do).  $\square$

**3.2.** In [A3], a condition similar to (3.2) is studied (cf. [A3, lemma 1.4.5]), where the quantifiers are interchanged:

$$(3.3) \quad \forall \partial \in \Gamma T_S \setminus 0, \exists \omega \in \Gamma \Omega_A, \theta_{\partial} \cdot \omega \neq 0.$$

**3.2.1. Theorem.** [A3, def. 4.2.1, th. 4.2.2] *In the restricted PE case, condition (3.3) is satisfied.*  $\square$

**3.2.2. Lemma.** *If  $d = g$ , (3.2)  $\Rightarrow$  (3.3).*

*Proof.* Let  $\theta^{\omega} : T_S \rightarrow \Omega_A^{\vee}$  the contraction of  $\theta$  by  $\omega \in \Omega_A$ . Condition (3.2) is equivalent to the injectivity of  $\theta^{\omega}$  for some  $\omega$ . Since  $d = g$ , this is equivalent to the generic surjectivity of  $\theta^{\omega}$  for some  $\omega$ , which implies the generic surjectivity of  $\theta$ , which is in turn equivalent to (3.3).  $\square$

**3.2.3. Question.** Does the converse of the lemma holds?

**3.2.4. Remark.** Let  $\eta$  be the generic point of  $S$  and  $P(\Omega_{A_{\eta}})$  (resp.  $P(T_{\eta})$ ) be the projective space of lines on  $\Omega_{A_{\eta}}$  (resp.  $T_S$ ). We denote by  $\bar{\omega}$  (resp.  $\bar{\partial}$ ) the class of  $\omega$  in  $P(\Omega_{A_{\eta}})$  (resp.  $\partial$  in  $P(T_{\eta})$ ).

One could try to answer the question in general by explaining the “logical seesaw” between quantifiers from (3.3) to (3.2) through a “geometric seesaw” provided by the incidence variety  $\Sigma \subset P(\Omega_{A_{\eta}}) \times P(T_{\eta})$  formed by pairs  $(\bar{\omega}, \bar{\partial})$  such that  $\theta_{\partial} \cdot \omega = 0$ , and let  $p_1, p_2$  be the projections on the

factors. Let further  $\check{P}_h \subset P(T_\eta)$  be the locally closed subvariety formed by elements  $\bar{\partial}$  such that  $\theta_{\bar{\partial}, \eta}$  has rank  $h$ . One has

$$p_2(\Sigma) \subset \prod_{h=1}^{h=g-1} \check{P}_h \subset P(T_\eta) = \prod_{h=1}^{h=g} \check{P}_h,$$

and

$$\dim p_2^{-1}\{\partial\} = g - 1 - h \text{ if } \partial \in \check{P}_h.$$

By (3.3),  $\check{P}_g$  is non empty and in fact dense (the condition is open in  $\partial$ ), hence  $\dim p_2(\Sigma) \leq g - 2$ .

The *key problem* is whether one has

$$(3.4) \quad \dim \check{P}_{h-1} < \dim \check{P}_h$$

(which is equivalent to  $\dim \check{P}_h \neq \emptyset$  since  $\overline{\check{P}_h} \supset \check{P}_{h-1}$ ) for  $e < h < g$ , where  $e$  is the first index such that  $\check{P}_e \neq \emptyset$ .

If so, then  $\dim \check{P}_h \leq h - 1$  and  $\dim \Sigma \leq g - 2$ , so  $p_1$  is not dominant, and any  $\bar{\omega}$  outside the image of  $p_1$  will do.

We have only partial answers to 3.2.3.

**3.2.5. Proposition.** *If  $d = g \leq 3$ , (3.3)  $\Rightarrow$  (3.2).*

*Proof.* We use the previous remark: the case  $g \leq 2$  is trivial ((3.4) is satisfied). For  $g = 3$ ,  $\dim \overline{\check{P}_2} = 1$ , and it suffices to show that  $\overline{\check{P}_1}$  is of dimension 0 or empty. Let us identify  $P(T_\eta)$  with a plane in the projective space of  $S^2 \text{Lie } A_\eta$ . Let  $\Delta_h \subset P(S^2 \text{Lie } A_\eta)$  be the locus of matrices of rank  $h$ . Then  $\Delta_1$  may be identified with the Veronese surface  $V \subset \mathbb{P}^5$ , since

its elements are the symmetric matrices of the form 
$$\begin{pmatrix} x_0^2 & x_0x_1 & x_0x_2 \\ x_0x_1 & x_1^2 & x_1x_2 \\ x_0x_2 & x_1x_2 & x_2^2 \end{pmatrix},$$

and  $P(T_\eta) \cap V$  may be identified with  $\overline{\check{P}_1}$ . It is enough to show that there any plane  $P \subset \mathbb{P}^5$  such that  $P \cap V$  is a curve is contained in  $\overline{\Delta_2}$ . Such a plane would be cut by three hyperplanes, which would correspond to three non collinear quadratic forms  $q_0, q_1, q_2$  in  $x_0, x_1, x_2$ . If  $P \cap V$  is a curve, then one could choose the  $q_i$ 's of the form  $\ell \cdot x_i$ , where  $\ell$  is a linear form in  $x_0, x_1, x_2$ ; by change of coordinates,  $\ell$  may be taken to be  $x_0$ , and it is then clear that  $P \subset \overline{\Delta_2}$ .  $\square$

**3.2.6. Proposition.** *If  $\mu(S)$  is a sufficiently general subvariety of dimension  $g$  of a special subvariety  $S_1$  of  $\mathcal{A}_g$  of restricted PEL type (cf. [A3, 4.2.1]), then (3.3)  $\Rightarrow$  (3.2).*

*Proof.* Let  $D := \text{End } A_\eta$  (assumed to be a division algebra),  $E$  a maximal subfield of  $D$  containing  $F$  (and chosen to be CM except if  $D$  is a totally real field) and  $E^+$  the maximal totally real subfield of  $E$ . One has a decomposition  $\Omega_A = \bigoplus_{\nu: E \rightarrow \mathbb{C}} \Omega_A \otimes_{E \otimes \mathbb{C}, \nu} \mathbb{C}$ . Fix  $\omega \in \Omega_A$  with non-zero components  $\omega_\nu$ . Fix  $s \in S(\mathbb{C})$  such that  $\omega_{\nu, s}$  remains non-zero. With the notation of [A3, rem. 4.2.3 i)], the tangent space  $T_{S_1, s}$  decomposes as  $\bigoplus_{\nu: E \rightarrow \mathbb{C}} T_{S_1, s, \nu}$ , and the universal Kodaira-Spencer map on the family of PEL type induces an isomorphism  $T_{S_1, s, \nu} \cong S^2 \Omega_{A, s, \nu}^\vee$  or  $T_{S_1, s, \nu} \cong \bigotimes^2 \Omega_{A, s, \nu}^\vee$  (according to whether  $D$  is of type I or II, or of type III or IV). In particular the contraction  $(\theta_1^{\omega_\nu})_s : T_{S_1, s, \nu} \rightarrow \Omega_{A, s, \nu}^\vee$  is surjective, hence also  $(\theta_1^\omega)_s : T_{S_1, s} \rightarrow \Omega_{A, s}^\vee$ . Since  $S$  is sufficiently general in  $S_1$  of dimension  $g$ , then  $T_S$  is transverse to the kernel of  $\theta_1^\omega$  on a suitable dense open subset of  $S$  [details should be checked].  $\square$

**3.3. A counterexample.** We indicate here what seems to be an example with smallest  $g > 0$  of a subvariety (*resp.* a special subvariety) of dimension  $d$  of  $\mathcal{A}_{g, n}$  with  $d = g$ , such that the restriction of the universal abelian scheme does not satisfy condition (3.3) (hence not (3.1); in this example the Betti map is nowhere a submersion). In fact  $d = g = 5$  (*resp.* 6).

Let us first treat the modular example. Let  $E^+$  be a real quadratic field and  $E$  a totally imaginary quadratic extension of  $E^+$ . Let  $A \rightarrow S$  be a principally polarized abelian scheme of relative dimension 6, with level  $n \geq 3$  structure, such that  $\text{End}_S A = \mathcal{O}_E$ , and with Shimura type  $(r_\nu, s_\nu) = (1, 2), (1, 2)$  (for the two embeddings  $\nu_1, \nu_2$  of  $E^+$  in  $\mathbb{C}$ ). According to [S], there exists such a family, with base  $S$  a special subvariety of dimension  $\sum r_\nu \cdot s_\nu = 6$  of  $\mathcal{A}_{6, n}$ . By functoriality,  $\theta$  commutes with the  $\mathcal{O}_L$ -action, hence respects the decomposition  $\mathcal{H}_{dR}^1(A/S) = \bigoplus_{\rho: E \rightarrow \mathbb{C}} \mathcal{H}_\rho$ . For every  $\rho$ , and general  $\partial$ , the restriction of the Kodaira-Spencer map  $\theta_\partial$  to  $(\Omega_A)_\rho$  has rank 1 (either the source or the target is of rank 1), hence  $\theta_\partial$  has rank  $4 < 6$ .

The non modular example consists in taking a general hypersurface section in a modular example with  $d = 6, g = 5$ . Here  $E$  is a quadratic imaginary field, and  $\text{End}_S A = \mathcal{O}_E$  with Shimura type  $(r, s) = (2, 3)$ . For each complex embedding  $\rho$  of  $E$ , the restriction of the Kodaira-Spencer map  $\theta_\partial$  to  $(\Omega_A)_\rho$  has rank 2 (either the source or the target is of rank 2), hence  $\theta_\partial$  has rank  $4 < 5$ .

**3.3.1. Remark.** It would be interesting to go beyond, and settle (3.2) for instance in the case of maximal monodromy  $Sp_{2g}$ . Let  $A \rightarrow S$  be the jacobian of the universal hyperelliptic curve of genus  $g > 0$ . Here  $S$  is the affine space  $\mathcal{M}_{0, 2g+2} \cong (\mathbb{P}^1 \setminus \{0, 1, \infty\})^{2g-1}$ . By Torelli's theorem, one has  $\dim \mu(S) = 2g - 1$ . On the other hand, according to [M2, th. 5.6],  $\text{End } A_{\bar{\eta}} = \mathbb{Z}$  (in fact, it is known that the monodromy representation,



which is a specialization of the Burau representation of the braid group, has Zariski dense image in  $Sp_{2g}$ , cf. [aC]). Hence (3.2) would imply that, after replacing  $(\mathbb{P}^1 \setminus \{0, 1, \infty\})^{2g-1}$  by a suitable dense Zariski-open affine subset,  $\beta$  is a submersion. [There are explicit calculations (without detail) by Bryant of  $\theta$  in this case on MathOverflow (tag: hyperelliptic) which should settle the matter]

**3.4.** Let us turn to the proof of th. 1.4.1. [sketch] One may replace  $S$  by a finite étale covering and assume that  $A/S$  has a level  $n$  structure for some  $n \geq 3$ , then replace  $S$  by a smooth dense affine open subset of its image in  $\mathcal{A}_{g,n}$ . By assumption,  $d = \dim \mu(S) \geq g$ . Taking a sufficiently general locally closed affine subvariety  $S' \subset S$  of dimension  $g$ , the restriction to  $S'$  of the abelian scheme and of the section satisfy the same assumption. Moreover, one checks that the condition  $d \geq g$  implies that we are in the restricted PE case<sup>4</sup>. Replacing  $S$  by  $S'$ , one is in the situation  $d = g$ . One can then combine prop. 2.2.1, 3.2.5 and th. 3.2.1. [A similar deduction holds with prop. 3.2.6 in place of 3.2.5].  $\square$

#### 4. PROOF OF THEOREM 1.5.1.

[your §4 and 5].

#### REFERENCES

- [aC] N. A'CAMPO – *Tresses, monodromie et le groupe symplectique*, Comment. Math. Helv. **54** (2) (1979), 318-327.
- [A1] Y. ANDRÉ – *Mumford-Tate groups of mixed Hodge structures and the theorem of the fixed part*, Compos. Math. **82** (1992), 1-24.
- [A2] Y. ANDRÉ – *Groupes de Galois motiviques et périodes*, Séminaire Bourbaki 7/11/2015.
- [A3] Y. ANDRÉ – *On the Kodaira-Spencer map of abelian schemes*, companion paper.
- [D2] P. DELIGNE – *Théorie de Hodge III*, Publ. Math. I.H.E.S. **44** (1974), 5-77.
- [L] L. LOVÁSZ – *Singular spaces of matrices and their application in combinatorics*, Bol. Soc. Braz. Mat. **20** (1989), 87-99.
- [M2] S. MORI – *The endomorphism rings of some abelian varieties*, Japan J. Math **2**, 1 (1976), 109-130.
- [S] G. SHIMURA – *On an analytic family of polarized abelian varieties and automorphic functions*, Ann. of Maths. **78**, 1 (1963), 149-192.

---

<sup>4</sup>here one needs a generalization of the notion of restricted PE to the case of an abelian scheme whose generic fiber is not necessarily simple. This does not pose problem.