



HAL
open science

Progression and projection for hip surgery in France, 2008-2070: Epidemiologic study with trend and projection analysis

Roger Erivan, Guillaume Villatte, Julien Dartus, Nicolas Reina, Stéphane Descamps, Stéphane Boisgard

► To cite this version:

Roger Erivan, Guillaume Villatte, Julien Dartus, Nicolas Reina, Stéphane Descamps, et al.. Progression and projection for hip surgery in France, 2008-2070: Epidemiologic study with trend and projection analysis. *Orthopaedics & Traumatology: Surgery & Research*, 2019, 105 (7), pp.1227-1235. 10.1016/j.otsr.2019.07.021 . hal-02409392

HAL Id: hal-02409392

<https://hal.science/hal-02409392v1>

Submitted on 20 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Original article

Progression and projection for hip surgery in France, 2008-2070: epidemiologic study with trend and projection analysis

Roger **ERIVAN**^{a*}, Guillaume **VILLATTE**^a, Julien **DARTUS**^{b,c}, Nicolas **REINA**^d, Stéphane **DESCAMPS**^a,
Stéphane **BOISGARD**^a

a: Université Clermont Auvergne, CHU Clermont-Ferrand, CNRS, SIGMA Clermont, ICCF, F-63000
Clermont-Ferrand, France

b: Université de Lille Nord de France, 59000 Lille, France

c: Service d'Orthopédie, Hôpital Roger Salengro, Centre Hospitalier et Universitaire de Lille, Place de
Verdun 59037 Lille, France

d: Hôpital Pierre Paul Riquet, CHU Toulouse, Toulouse, France

* Corresponding author: Orthopedic and Trauma Surgery Department, Hôpital Gabriel Montpied,
CHU de Clermont Ferrand BP 69, 63003 Clermont Ferrand, France

Phone: +33 4 73 751 535

Email: rerivan@chu-clermontferrand.fr

Abstract

Introduction

Hip replacement was declared “operation of the century” in tribute to the functional improvement it provides. Frequency is increasing, but it is difficult to estimate the actual number of procedures performed and the expected progression, because of changes in indications and lengthening life-expectancy, and also, in France, because there is no registry. As data are lacking in France, we conducted an investigation 1) to update the number of hip surgeries in France, and 2) to forecast progression over the coming decades, considering extreme scenarios.

Hypothesis

The number of hip procedures can be expected to increase considerably over the coming 50 years.

Material and method

A study was conducted to analyze national coding data for the number of hip surgeries performed in France. Two scenarios were defined: one taking account of population progression and age structure, the other also extrapolating trends observed over recent years. Current hip surgery activity in France was measured, and progression estimated according to population changes.

Results

In 2018 in France, 183,139 procedures were coded as principally concerning the hip. There was a clear predominance of reconstruction procedures, with 148,965 primary hip replacements, 124,251 of which were total. There were 19,304 hip replacement revision procedures. There were strong regional differences in revision according to the type of center performing surgery ($p < 0.0001$). Between 2018 and 2050, primary hip replacement could be expected to increase by 41.9% or 114.3% and hip surgery overall by 42.0% or 98.3%, depending on the scenario.

Discussion

The present results are subject to future technological breakthroughs and medical discoveries, but forecast a major increase in hip surgery requirements. These results extend the present state of medical knowledge.

Level of evidence: IV, descriptive epidemiological study

Keywords: epidemiology; hip; hip replacement; progression; prediction; national data

1. Introduction

The number of orthopedic surgery procedures has been constantly increasing worldwide over recent decades [1–3], for multiple reasons: improvement in biomaterials and in surgical techniques, contribution of imaging, optimization of perioperative care, and patients' demand to be able to age in good health with minimal functional disability [3,4]. One of the most important developments has been in total hip arthroplasty (THA), which has now been consecrated “operation of the century” by some authors [5] because of the clear functional improvement achieved, and especially definitive abolition of pain, although risks remain [6–8]. Increasing numbers of patients are having resort to orthopedic surgery, as the population in France ages and constantly grows [9].

Estimating the number of procedures and their progression is made difficult by changes in indications and by the extension of life-expectancy. Scandinavian countries have registries [10–12], while other countries, such as France, uses codes for surgical procedures that enable the total number of operations related to hip surgery to be quantified. Numerous parameters, such as technical innovations [13], population changes, and health-system funding changes, hinder attempts to forecast and plan for future rates of hip surgery [5,11].

To the best of our knowledge, such data are not available for France, and we therefore conducted a study 1) to update the number of hip surgeries in France, and 2) to forecast progression over the coming decades, considering the extreme scenarios. The study hypothesis was that the number of hip surgeries can be expected to increase over the next 50 years.

2. Material and method

2.1 Material

Data for numbers and types of hip surgery were obtained from the DIAMANT data-base of the French PMSI medical information program on hospital admissions in medical, surgical, obstetric and odontologic (MCO) departments according to principal intervention as coded on the CCAM medical acts classification system. Data were collected by collating the standardized discharge reports, containing administrative, demographic, medical and therapeutic information, with the CCAM codes. Registration numbers for legal entities on the FINESS national health-care and social establishments file were taken from the DRESS research, studies, assessment and statistics directorate. The PMSI data-base provided patient gender and age-group. The anonymized discharge reports provided the dates (years) of the procedure in question. On the basis of these data, patients were age-grouped as 0-14, 15-29, 30-44, 45-59, 60-74, 75-89 and ≥ 90 years of age. Legal entity and FINESS data were used to classify health-care structures as public, private non-profit, or for-profit.

The study analyzed French national data. Statistical data were obtained from the National Institute of Statistics and Economic Studies (INSEE), for population demographics and progression over the coming years based on the present French birth-rate of 1.96. This kind of study uses low, medium and high estimates according to various parameters: life-expectancy, fecundity and migration. The present study used the medium scenario: i.e., mean values for the various parameters. It also took account of INSEE's predicted variations per age-group to calculate a progression factor according to patient-age per procedure [9].

The coding information search was started in March 2019, so as to have complete data for 2018. The DIAMANT data-base uses the following procedure: each center sent its full 2018 data set at the beginning of February 2019; then the hospital admissions information agency (ATIH) extracts all information in the data-bank on the 20th of each month and makes it available on a secure server on the 21st, and the data is then integrated into the DIAMANT base within 5 working days.

2.2 Methods

Data were categorized for simplified analysis: surgical or non-surgical hip dysplasia, primary hip arthroplasty (total or partial), hip implant dislocation, hip implant removal, hip implant revision, hip arthroscopy, and tumor surgery. Table 1 shows codes per category.

Two scenarios were applied for each category, based on the DRESS method [14]:

- Scenario 1: Constant characteristics. This scenario simply carries forward the parameters as measured on the PMSI data-base for the last full year (2018), treating them as unchanging over time. Thus, it takes account only of changes in population and age structure. It serves as a kind of baseline, with 2018 values adjusted for expected population change. The model was adjusted for each procedure according to changes in age-groups predicted by the INSEE data. For example, numbers for procedures mainly performed on younger patients increase less than for procedures performed on more elderly patients, as the population tends to age year on year. Thus, for each procedure and each decade, we calculated a procedure-weighting factor based on age distribution per procedure and INSEE data. Table 2 shows a few examples.
- Scenario 2: Extrapolated progression. This scenario carries forward trends observed over the period 2012-2018, adjusted for population change as in scenario 1. For each procedure, we estimated the progression in the number of procedures, using a linear regression model based on the data for recent years, extrapolated into the future.

Geographic descriptions were based on raw data, averaging the number of procedures over the period 2012-2018, with regionally based calculation according to the type of health-care structure. We also calculated the total number of procedures and number per head of the population.

2.3 Statistics

Statistical calculations were made on Excel™ (Microsoft, Redmond, WA) and XLSTAT™ software (Addinsoft, New York, NY). The global projections per procedure were made by summing the

individual projections for each age-group, weighted according to the INSEE data. Independent models were thus constructed for each procedure according to its target population and relevant age-groups. Results for quantitative variables were reported as numbers or mean \pm standard deviation and range, with 95% confidence intervals for estimates and first-order risk set at 5%. Variations in number of procedures were reported as numbers, without statistical calculations as these were real numbers rather than estimates, and the observed differences were by definition exact. There were no missing data; all data were analyzed.

3. Results

3.1 Data for 2018

In 2018, which was the last analyzable year, 183,139 procedures in France were coded as principally involving hip surgery. Distribution per category is shown in Table 3.

There was clear predominance of reconstruction surgery, with 150,060 primary hip arthroplasties, including 125,227 THAs. There were 19,457 revision procedures, amounting to 11.5% of implant surgeries. There were in all 180,180 procedures for primary hip arthroplasty, revision and associated complications, amounting to 98.3% of hip surgeries.

3.2 Progression 2012-2018

For the period 2012-2018, a mean $177,355 \pm 5,075$ procedures [range, 169,741-183,139] were performed per year, with an increase of 7.9% over the period (fig. 1).

At national level, taking all procedures together, 76,349 (43.0%) were performed in a public-sector hospital, 12,767 (7.2%) in a non-profit private structure, and 88,239 (49.8%) in a for-profit structure. Between 2012 and 2018, there was an increase of 10.4%, from 71,899 to 79,387, in the public sector, 14.2%, from 11,927 to 13,616, in the private non-profit sector, and 4.9%, from 85,915 to 90,136 in the for-profit sector. Over the period as a whole, 59.6% of procedures concerned female patients

(105,614 procedures), with an increase of 7.6%, from 99,208 to 106,776, and an increase of 8.7% in males, from 66,509 to 72,325 procedures.

Figure 2 maps the geographic distribution of primary THA according to the 2012-2018 averages, and Figure 3 maps implant revision. There were marked regional differences for primary, revision and complications surgery. Primary THA ranged from a mean 16.0 procedures per 1,000 persons per year between 2012 and 2018 in Corsica to 26.4 in Brittany; revision ranged from 1.5 procedures per 1,000 persons per year in Corsica to 3.7 in Bourgogne Franche-Comté. Primary THA rates in public, non-profit and profit structures varied non-significantly between regions ($p=0,965$), while revision rates varied significantly ($p<0.0001$), with numerous disparities, as shown in Table 4.

3.3 Foreseeable progression over coming decades according to data for recent years

Foreseeable progression over coming years was estimated on the 2 scenarios described above (see Methods): scenario 1, taking account only of age, with constant activity compared to 2018; and scenario 2, further taking account progression per procedure over recent years to forecast coming progression.

3.3.1 Dysplasia

The number of procedures for hip dysplasia decreased by 4.3% between 2012 and 2018, from 1,636 to 1,565 (fig. 4). Procedures concerned female patients in a mean 60.9% of cases (976/1,603). For the period 2018-2050, scenario 1 predicted a 5.5% increase, and scenario 2 a 12.6% increase ($p=0.201$).

3.3.2 Arthroscopy

Hip arthroscopy procedures increased by 58.6% between 2012 and 2018, from 789 to 1,251. Figure 5 shows the projections. Procedures concerned female patients in a mean 44.7% of cases (425/950).

For the period 2018-2050, scenario 1 predicted a 5.6% increase, and scenario 2 a 189.9% increase ($p=0.003$).

3.3.3 Primary THA

Primary THA is the most frequent hip procedure, and increased by 10.5% between 2012 and 2018, from 135,746 to 150,060 (fig. 6). Procedures concerned female patients in a mean 60% of cases (86,254/143,679). For the period 2018-2050, scenario 1 predicted a 41.9% increase, and scenario 2 a 114.3% increase ($p=0.0001$).

3.3.4 Revision

Hip implant revision decreased slightly between 2012 and 2018, by 0.4%, from 19,533 to 19,457 procedures (fig. 7). Procedures concerned female patients in a mean 56.1% of cases (10,936/19,499). For the period 2018-2050, scenario 1 predicted a 43.8% increase, and scenario 2 a 43.5% increase ($p=0.642$). In this instance, the two scenarios were virtually equivalent, as levels were more or less constant between 2012 and 2018, and the only relevant factor was numbers per age-group.

3.3.5 Hip dislocation

Hip dislocation rates decreased over the years in the elderly, while increasing in the overall French population. The decrease between 2012 and 2018 was 10.7%, from 11,947 to 10,663 procedures. The 2 scenarios diverged (fig. 8). Procedures concerned female patients in a mean 61.1% of cases (6,915/11,312). For the period 2018-2050, scenario 1 predicted a 51.0% increase, and scenario 2 a 39.2% decrease ($p<0.0001$).

3.3.6 Overall activity

Overall, the number of hip procedures increased yearly between 2012 and 2018, by 7.9%, from 169,741 to 183,139 (fig. 9). For the period 2018-2050, scenario 1 predicted a 42.0% increase, and scenario 2 a 98.3% increase ($p=0.001$).

4. Discussion

The present study provided precise figures for hip surgery in France and for foreseeable progression over coming years. At national level, there were disparities between procedures and between public/private sectors, notably for revision surgeries. Hip surgery was principally arthroplastic. Overall progression in hip surgery up to 2050 can be expected to be either 42.0% or 98.3% depending on the scenario; the study hypothesis was thus confirmed: overall hip surgery activity can be expected to increase in coming decades.

The study showed hip surgery to be an important part of orthopedic surgery in France, with numerous procedures performed. These data are comparable to those for other countries and to previous analyses of French data on both primary arthroplasty [3,10,15] and revision [4]. Arthroplasty and the management of arthroplasty complications account for most hip surgery procedures. In 2007 in the USA, Kurtz et al. [1] reported results similar to the present, but on a single scenario using a Poisson model with non-linear regression, more appropriate to knee arthroplasty. Ferguson et al. [16], in 2018, reported that rates of THA in developed countries were increasing by 30-40% every 10 years, comparable to the present estimates and lying between the 2 present scenarios.

In regard to our secondary objective, hip surgery activity can be expected to grow in the coming years and decades. Scenario 1 is to be seen as a baseline, with no change in the frequency of procedures, whereas scenario 2 takes additional account of trends in recent years. The data are obviously only estimates, and it seems reasonable to imagine that the real figures for each procedure will generally lie somewhere between the two scenarios. The regional distributions show considerable disparities, due to the local health-care offer (centers specializing in certain types of surgery, to patient demographics), and to medical demographics. The probable increase in activity may be slowed by national-level capacity to undertake these procedures, as a constant mean increase, although possible, is in reality liable to be curbed [17]. Regarding hip dislocation, our data

did not allow us to take account of the impact of dual mobility designs [13]. It would seem reasonable to expect an overall increase of 10% per decade in France, although with some variations: primary THA and hip arthroscopy are likely to increase more than revision and dysplasia surgeries due to improved prevention and the ever younger age of the patients in question.

The study involved several limitations. 1) There may be a bias related to lack of coding and non-exhaustive inventorying of procedures. There may also be classification bias related to the coding system: we simply trusted the system, without checking each file – which would not have been feasible. Nevertheless, in the French health system surgery is budgeted in terms of coded procedures, and most procedures are therefore doubtless coded, especially since the “activity-based tariffication” system was introduced. 2) Although we sought to be exhaustive, some procedures, and particularly the less frequent ones, may have been overlooked. We took account only of principal codings, and may therefore have underestimated certain procedures: the procedure coded as “principal” is the one bringing in the largest budget, but is not necessarily the one that actually motivated the surgery. However, this applies mainly in traumatology. 3) Another limitation is that population growth is not homogeneous but shows geographic variations. We did not take this into account, as the analysis would have been too complicated. As stated in the Methods section, we used the medium scenario for the INSEE data, which introduces uncertainty and may falsify the population growth estimates. Likewise, for progression under scenario 2, we applied linear regression, whereas exponential or logarithmic progression would be equally conceivable. 4) We did not take account of change in medical demographics, and notably of changes in the number of orthopedic surgeons or the increasing proportion of female practitioners, both of which are liable to vary over time and can impact the relation between supply and demand. 5) The use of linear regression may be particularly questionable in the case of hip arthroscopy, which is a new field, with strong growth potential; a comparable study in a few years’ time could show whether the expected slow-down has occurred. 6) Finally, estimates were based on the present state of medicine: if a cure were to be found for osteoarthritis, that would indeed be a game-changer; however, nothing of the

sort presently exists. Likewise, some new technology or developments in materials and implants might revolutionize the scenarios – but the present results can only be founded on the present state of medical knowledge.

5. Conclusion

Hip surgery is one of the oldest branches of orthopedics, and hip arthroplasty has, over the years, become a major procedure in France, due to the functional benefit and improved quality of life it ensures. The number of procedures has been incessantly increasing for more than 10 years and, according to forecasts, can be expected to continue to increase over the coming decades. Projections for 2050, taking account of population aging, suggest increases of 41.9-114.3% for THA and 42.0-98.3% for hip surgery as a whole. These estimates are subject to technological breakthroughs and medical discoveries, but describe a perspective of major increase in orthopedic surgery requirements.

Disclosures of interest: RE, GV, JD, NR and SD have no conflicts of interest to disclose in relation to the present article. Elsewhere, SB is a consultant for Zimmer-Biomet.

Funding: none

Author contributions: RE: project design, data study, article writing and correction. GV: project design and article writing. JD and NR: re-editing and correction. SD: project supervision and article writing. SB: project supervision.

References

- [1] Kurtz S, Ong K, Lau E, Mowat F, Halpern M. Projections of Primary and Revision Hip and Knee Arthroplasty in the United States from 2005 to 2030. *J Bone Joint Surg Am* 2007;89:780-5.
- [2] Putman S, Girier N, Girard J, Pasquier G, Migaud H, Chazard E. Épidémiologie des prothèses de hanche en France : analyse de la base nationale du PMSI de 2008 à 2014. *Rev Chir Orthop Traumatol* 2017;103:S90.
- [3] Maradit Kremers H, Larson DR, Crowson CS, Kremers WK, Washington RE, et al. Prevalence of Total Hip and Knee Replacement in the United States. *J Bone Joint Surg Am* 2015;97:1386–97.
- [4] Gwam CU, Mistry JB, Mohamed NS, Thomas M, Bigart KC, et al. Current epidemiology of revision Total Hip Arthroplasty in the United States: National inpatient sample 2009 to 2013. *J Arthroplasty* 2017;32:2088–92.
- [5] Learmonth ID, Young C, Rorabeck C. The operation of the century: total hip replacement. *The Lancet* 2007;370:1508–19.
- [6] Reina N, Delaunay C, Chiron P, Ramdane N, Hamadouche M, Société française de chirurgie orthopédique et traumatologique. Infection as a cause of primary total hip arthroplasty revision and its predictive factors. *Orthop Traumatol Surg Res* 2013;99:555–61.
- [7] Ehlinger M, Delaunay C, Karoubi M, Bonnomet F, Ramdane N, et al. Revision of primary total hip arthroplasty for peri-prosthetic fracture: A prospective epidemiological study of 249 consecutive cases in France. *Orthop Traumatol Surg Res* 2014;100:657–62.
- [8] Fessy MH, Putman S, Viste A, Isida R, Ramdane N, et al. What are the risk factors for dislocation in primary total hip arthroplasty? A multicenter case-control study of 128 unstable and 438 stable hips. *Orthop Traumatol Surg Res* 2017;103:663–8.
- [9] Nathalie Blanpain GB. Projections de population à l’horizon 2070 - Insee Première - 1619 n.d. <https://www.insee.fr/fr/statistiques/2496228> Accessed May 1, 2019
- [10] Malchau H, Herberts P, Eisler T, Garellick G, Söderman P. The Swedish Total Hip Replacement Register. *J Bone Joint Surg Am* 2002;84-A Suppl 2:2–20.

- [11] Rolfson O, Bohm E, Franklin P, Lyman S, Denissen G, et al. Patient-reported outcome measures in arthroplasty registries. *Acta Orthop* 2016;87:9–23.
- [12] Nemes S, Rolfson O, W-Dahl A, Garellick G, Sundberg M, et al. Historical view and future demand for knee arthroplasty in Sweden. *Acta Orthop* 2015;86:426–31.
- [13] Epinette JA, Lafuma A, Robert J, Doz M. Cost-effectiveness model comparing dual-mobility to fixed-bearing designs for total hip replacement in France. *Orthop Traumatol Surg Res* 2016;102:143–8.
- [14] France. Dossiers solidarité et santé. Paris: Direction de la recherche, des études, de l'évaluation et des statistiques; 2007. <https://drees.solidarites-sante.gouv.fr/etudes-et-statistiques/> Accessed May 1, 2019
- [15] Bozic KJ, Kurtz SM, Lau E, Ong K, Vail TP, Berry DJ. The epidemiology of revision Total Hip Arthroplasty in the United States. *J Bone Joint Surg Am* 2009;91:128-133.
- [16] Ferguson RJ, Palmer AJ, Taylor A, Porter ML, Malchau H, Glyn-Jones S. Hip replacement. *The Lancet* 2018;392:1662–71.
- [17] Erivan R, Chaput T, Villatte G, Ollivier M, Descamps S, Boisgard S. Ten-year epidemiological study in an orthopaedic and trauma surgery centre: Are there risks involved in increasing scheduled arthroplasty volume without increasing resources? *Orthop Traumatol Surg Res* 2018;104:1283-1289.

Figure legends

Figure 1: Progression in annual number of procedures related to the hip, 2012-2018; $p= 2.0E-18$

Figure 2: Map of French national distribution of primary THA: means for 2012-2018 (THA: total hip arthroplasty).

Figure 3: Map of French national distribution of THA revision and complications surgery: means for 2012-2018 (THA: total hip arthroplasty)

Figure 4: Progression in number of procedures related to hip dysplasia for 2012-2018 and trends up to 2070 according to scenarios 1 and 2 ($p=0.201$)

Figure 5: Progression in number of procedures with hip arthroscopy for 2012-2018 and trends up to 2070 according to scenarios 1 and 2 ($p=0.003$)

Figure 6: Progression in number of primary arthroplasties for 2012-2018 and trends up to 2070 according to scenarios 1 and 2 ($p=0.0001$)

Figure 7: Progression in number of hip arthroplasty revision procedures for 2012-2018 and trends up to 2070 according to scenarios 1 and 2 ($p=0.642$)

Figure 8: Progression in number of procedures for hip dislocation for 2012-2018 and trends up to 2070 according to scenarios 1 and 2 ($p<0.0001$)

Figure 9: Overall progression in number of hip surgeries for 2012-2018 and trends up to 2070 according to scenarios 1 and 2 ($p=0.001$)

Table 1: Procedure codes.

| Class | Sub-class | PMSI - CCAM Procedure |
|--|--|---|
| Dysplasia | Non-surgical | NEEP001 – Scheduled traction reduction of non-traumatic unstable hip, age >12 months |
| | | NEEP003 - Scheduled traction reduction of non-traumatic unstable hip, age <12 months |
| | | NEEP006 - Scheduled harness traction reduction of non-traumatic unstable hip, age >12 months |
| | Surgical | NEEA004 – Arthrotomy reduction of non-traumatic unstable hip |
| | | NEMA003 – Supra-acetabular hip bone block + extra-articular proximal femoral osteotomy |
| | | NEMA017 - Supra-acetabular hip bone block osteoplasty + greater trochanter osteotomy |
| | | NEMA018 – Hip arthroplasty femoral cup |
| | | NEMA020 - Acetabuloplasty + femoral osteotomy |
| | | NEMA021 - Supra-acetabular hip bone block osteoplasty/acetabuloplasty |
| Primary hip arthroplasty | THA | NEKA010 - THA + metallic acetabular reinforcement + femoral graft reconstruction |
| | | NEKA012 - THA + acetabular reconstruction and/or femoral graft reconstruction |
| | | NEKA013 - THA after hip fusion |
| | | NEKA014 - THA + metallic acetabular reinforcement |
| | | NEKA015 - THA after internal fixation osteotomy and/or femoral cervicocephalic implant |
| | | NEKA016 - THA + femoral shaft osteotomy |
| | | NEKA017 - THA + femoral head lowering in acetabulum |
| | | NEKA019 - THA after hip fusion + metallic acetabular reinforcement |
| | | NEKA020 - THA |
| | NEKA021 - THA + femoral head lowering in acetabulum +realignment osteotomy / femoral alignment | |
| PHA | NEKA018 - Cervicocephalic implant | |
| NEKA011 - Cervicocephalic implant + mobile cup | | |
| THA dislocation | Non-surgical | NEEP002 - Reduction of hip implant dislocation |
| | | NEEP004 - Reduction of hip dislocation + rigid full-leg splint |
| | | NEEP005 - Reduction of hip dislocation + femoral head fracture |
| | | NEEP007 - Reduction of hip dislocation + continuous traction |
| | Surgical | NEEA001 - Reduction of hip dislocation by arthrotomy |
| | | NEEA002 - Reduction of hip implant dislocation by arthrotomy |
| | | NEEA003 - Reduction of hip dislocation + femoral head fracture fixation with arthrotomy |
| | | NEDA002 – Inert block + lowering of greater trochanter with or without THA head/neck exchange |
| Implant revision | | NEKA001 – THA acetabular + femoral exchange + compacted graft without internal fixation |
| | | NEKA002 - THA acetabular and/or femoral exchange + acetabular and/or femoral bone reconstruction |
| | | NEKA003 - THA acetabular + femoral exchange without bone reconstruction |
| | | NEKA004 – THA acetabular insert exchange |
| | | NEKA005 - THA acetabular and/or femoral exchange + acetabular and/or femoral internal fixation |
| | | NEKA006 - THA acetabular + femoral exchange + reconstruction + acetabular and/or femoral internal fixation |
| | | NEKA007 - THA acetabular and/or femoral exchange + compacted graft reconstruction without internal fixation |
| | | NEKA008 - THA acetabular + femoral exchange + reconstruction and/or acetabular and/or femoral internal fixation |
| | | NEKA009 - THA acetabular and/or femoral exchange without bone reconstruction |
| | | NEKA022 – THA femoral cervicocephalic implant exchange |
| | | NELA001 – THA reimplantation + bone reconstruction |

| | |
|------------------|---|
| Implant ablation | NELA002 - THA reimplantation without bone reconstruction NELA003 – Acetabular component implantation in ipsilateral femoral cervicocephalic implant bearer NEGA001 – THA ablation + acetabular and/or femoral bone reconstruction NEGA002 - THA ablation NEGA003 - THA ablation + trochanteroilic coaptation NEGA004 - Ablation of PTA insert + acetabular component implantation NEGA005 - Ablation of femoral cervicocephalic implant |
| Tumoral | NEMA011 – Massive and/or made-top-measure hip implant for segmental and/or hip bone defect NEMA019 – Sacroiliac arthroplasty with segmental bone defect + direct fusion NEMA013 – Hip arthroplasty after en-bloc resection + femur/hip fixation |
| Arthroscopy | NEJC001 – Arthroscopic hip debridement NEFC001 - Arthroscopic hip synovectomy NEQC001 - Arthroscopic hip exploration |

THA: total hip arthroplasty; PHA: partial hip arthroplasty; PMSI: *Programme de médicalisation du système d'informations* medical information system; CCAM: *Classification Commune des Actes médicaux* medical acts classification.

Table 2: Examples of weighting factors per procedure according to population change per relevant age-group.

| Procedure – principal only | 2020 | 2025 | 2030 | 2035 | 2040 | 2050 | 2060 | 2070 |
|--|-------|-------|-------|-------|-------|-------|-------|-------|
| NEEA004 – Non-traumatic unstable hip reduction by arthrotomy | 1.004 | 1.009 | 1.010 | 1.017 | 1.028 | 1.045 | 1.054 | 1.067 |
| NEEP001 – Scheduled non-traumatic unstable hip reduction by traction age > 12 months | 1.010 | 1.034 | 1.056 | 1.077 | 1.093 | 1.124 | 1.150 | 1.173 |
| NEGA002 - Ablation of THA | 1.031 | 1.119 | 1.216 | 1.284 | 1.317 | 1.393 | 1.477 | 1.532 |
| NEKA004 – THA acetabular Insert exchange | 1.034 | 1.132 | 1.242 | 1.318 | 1.353 | 1.443 | 1.544 | 1.607 |
| NEKA005 – THA acetabular and/or femoral component exchange + acetabular and/or femoral internal fixation | 1.038 | 1.149 | 1.272 | 1.357 | 1.396 | 1.501 | 1.622 | 1.697 |
| NEKA011 – Cervicocephalic implant + mobile cup | 1.056 | 1.226 | 1.418 | 1.547 | 1.601 | 1.827 | 2.106 | 2.263 |
| NEKA020 - THA | 1.027 | 1.106 | 1.191 | 1.252 | 1.282 | 1.339 | 1.397 | 1.438 |
| NEMA017 – Supra-acetabular hip-bone block osteoplasty + greater trochanter osteotomy | 1.002 | 0.999 | 0.991 | 0.992 | 1.002 | 1.008 | 1.004 | 1.011 |

THA: total hip arthroplasty

Table 3: Number of hip surgeries in 2018

| class | sub-class | number in 2018 |
|------------------------|------------------|-----------------------|
| Dysplasia | total | 1,565 |
| | non-surgical | 382 |
| | surgical | 1,183 |
| Arthroscopy | | 1,251 |
| Primary implant | total | 150,060 |
| | THA | 125,227 |
| | PHA | 24,833 |
| THA dislocation | total | 10,663 |
| | non-surgical | 10,199 |
| | surgical | 464 |
| Revision | | 19,457 |
| Tumoral | | 143 |

THA: total hip arthroplasty; PHA: partial hip arthroplasty.

Table 4: Disparities in hip implant revision surgeries according to region and type of health-care structure. Values in **bold** are significant at p=0.05; the further from zero, the greater the significance; positive values show a greater number of procedures in this type of center, negative values a smaller number.

| Residues (Adjusted) | | | |
|-------------------------------------|---------------|--------------------|--------------------|
| | public sector | private non-profit | private for-profit |
| Alsace-Champagne-Ardenne-Lorraine | 4.297 | 4.579 | -6.652 |
| Aquitaine-Limousin-Poitou-Charentes | -2.041 | -1.025 | 2.566 |
| Auvergne-Rhône-Alpes | -0.061 | 4.091 | -2.054 |
| Bourgogne-Franche-Comté | 3.096 | -4.442 | -0.791 |
| Bretagne | 4.184 | -2.652 | -2.801 |
| Centre | -3.411 | -7.331 | 7.191 |
| Corse | -3.497 | -2.010 | 4.526 |
| Guadeloupe | 2.069 | -1.253 | -1.415 |
| Guyane | -1.801 | 8.131 | -2.407 |
| Île-de-France | -3.169 | 15.169 | -4.681 |
| Languedoc-Roussillon-Midi-Pyrénées | 0.015 | -1.255 | 0.634 |
| Martinique | 1.742 | -1.506 | -0.959 |
| Mayotte | 1.656 | -0.420 | -1.435 |
| Nord-Pas-de-Calais-Picardie | 3.913 | 2.507 | -5.198 |
| Normandie | -2.524 | -8.504 | 6.912 |
| Pays de la Loire | -3.177 | -10.313 | 8.499 |
| Provence-Alpes-Côte d'Azur | -1.084 | -2.120 | 2.177 |
| Réunion | 0.383 | -2.094 | 0.701 |






Figure 1:

Breakdown of activity sector



1: represents frequency per 10,000 inhabitants for each type of institution





average annual number of THA between 2012 and 2018 per 10,000

total = average annual number of THA between 2012 and 2018

ranking of regions

• University Hospital






Figure 4:




Figure 5:




Figure 6:




Figure 7:




Figure 8:




Figure 9: