



**HAL**  
open science

# Reliability of the functional measures of the corticospinal pathways to dorsiflexor muscles during maximal voluntary contractions

Robin Souron, Adrien Farabet, Guillaume Y. Millet, Thomas Lapole

## ► To cite this version:

Robin Souron, Adrien Farabet, Guillaume Y. Millet, Thomas Lapole. Reliability of the functional measures of the corticospinal pathways to dorsiflexor muscles during maximal voluntary contractions. *Journal of the Neurological Sciences*, 2016, 369 (11), pp.368-374. 10.1016/j.jns.2016.09.003. hal-02409556

**HAL Id: hal-02409556**

**<https://hal.science/hal-02409556v1>**

Submitted on 4 Sep 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Reliability of the functional measures of the corticospinal pathways to**  
2 **dorsiflexor muscles during maximal voluntary contractions**

3 Robin Souron<sup>1</sup>, Adrien Farabet<sup>1</sup>, Guillaume Y Millet<sup>2</sup> & Thomas Lapole<sup>1</sup>

4 <sup>1</sup>Univ Lyon, UJM-Saint-Etienne, LIBM, F-42023, SAINT-ETIENNE, France

5 <sup>2</sup>Human Performance Laboratory, Faculty of Kinesiology, University of Calgary,  
6 Canada

7  
8 Running title: Reliability of tibialis anterior corticospinal functional properties

9  
10 Corresponding author:

11 Thomas LAPOLE

12 Laboratoire Interuniversitaire de Biologie de la Motricité

13 Bâtiment IRMIS

14 10 rue de la Marandière

15 42270 Saint Priest en Jarez

16 04 77 42 18 91

17 [thomas.lapole@univ-st-etienne.fr](mailto:thomas.lapole@univ-st-etienne.fr)

18 **ABSTRACT**

19  
20 This study aimed to evaluate the intra- and inter-day reliability of transcranial  
21 magnetic stimulation (TMS)-related measurements recorded from the tibialis anterior  
22 (TA) muscle. Thirteen healthy young men and women ( $23 \pm 4$  years) performed 3  
23 testing sessions to assess intra- (i.e., two sessions performed the same day) and  
24 inter-day (i.e. two sessions performed one week apart) reliability of (i) dorsiflexion  
25 cortical maximal voluntary activation level ( $VA_{TMS}$ ), (ii) TA corticospinal excitability  
26 assessed through the amplitude of the motor evoked potentials (MEP) recorded  
27 during 100, 75 and 50% maximal voluntary contractions (MVC), and (iii) intracortical  
28 inhibition investigated via the cortical silent period (CSP) recorded at the same  
29 %MVC. Absolute (i.e., coefficient of variation (CV) and standard error of the mean  
30 (SEM)), and relative (i.e., intraclass correlation coefficients (ICC)) reliability  
31 parameters were calculated.  $VA_{TMS}$  demonstrated excellent intra- and inter-day  
32 reliabilities (ICC: 0.80 and 0.99; CV: 1.7 and 0.8%, respectively). MEPs and CSPs  
33 presented moderate to excellent intra- and inter-day reliabilities, while input-output  
34 curves extracted parameters presented highly variable outcomes. These results  
35 suggest that most TA corticospinal measurements during voluntary contraction can  
36 be used to quantify corticospinal adaptations after acute (e.g. fatigue) or long term  
37 (e.g. training) interventions.

38  
39  
40 Keywords: transcranial magnetic stimulation; tibialis anterior; cortical voluntary  
41 activation; corticospinal excitability, Boltzmann modelling.

43 **ABBREVIATIONS**

1	44	CSP	Cortical silent period
2			
3	45	CSP <sub>50</sub>	Cortical silent period during a 50% MVC submaximal contraction
4			
5	46	CSP <sub>75</sub>	Cortical silent period during a 75% MVC submaximal contraction
6			
7	47	CSP <sub>MVC</sub>	Cortical silent period during a MVC
8			
9	48	CV	Coefficient of variation
10			
11	49	EMG	Electromyographic
12			
13	50	ERT	Estimated resting twitch amplitude
14			
15	51	I <sub>50</sub>	Stimulus intensity required to obtain a MEP of half the size of the
16	52		plateau
17			
18	53	ICC	Intraclass correlation coefficient
19			
20	54	k	The slope parameter of the Boltzmann modelling
21			
22	55	M <sub>MVC</sub>	M-wave amplitude during a MVC
23			
24	56	MEP	Motor evoked potential
25			
26	57	MEP <sub>50</sub>	Motor evoked potential amplitude during a 50% MVC submaximal
27	58		contraction
28			
29	59	MEP <sub>75</sub>	Motor evoked potential amplitude during a 75% MVC submaximal
30	60		contraction
31			
32	61	MEP <sub>MVC</sub>	Motor evoked potential amplitude during a MVC
33			
34	62	MVC	Maximal voluntary contraction
35			
36	63	P	The function plateau of Boltzmann modelling
37			
38	64	RMS	Root mean square
39			
40	65	SEM	Standard error of the mean
41			
42	66	SIT	Superimposed twitch amplitude
43			
44	67	SIT <sub>MVC</sub>	Superimposed twitch amplitude during a maximal voluntary contraction
45			
46	68	TA	Tibialis anterior
47			
48	69	TMS	Transcranial magnetic stimulation
49	70	VA <sub>TMS</sub>	Voluntary activation assessed with TMS

## 1. Introduction

It is well established that training, and especially strength training, induces several adaptations to the central nervous system leading, *in fine*, to an augmentation of maximal voluntary contraction (MVC) force. The investigation of maximal voluntary activation is critical to assess neural adaptations. Voluntary activation refers to the neural drive delivered to a muscle during a voluntary contraction. The most common way to estimate maximal voluntary activation is to use the interpolated twitch technique [1], where a supramaximal stimulus is delivered over a motor nerve during an MVC to measure the extra force evoked by the stimulation. Using this method, changes in neural drive can be identified as at or above the site of stimulation of the motor axons, without distinction between spinal and supraspinal mechanisms [2]. Since training may induce both spinal [3] and supraspinal adaptations [4], transcranial magnetic stimulation (TMS) has been proposed as an alternative technique to ascertain whether adaptations occur throughout the corticospinal pathway [5]. Indeed, the evocation of a superimposed twitch elicited by TMS during MVC has been suggested to highlight a suboptimal drive from the motor cortex [5-7], although the cause of  $VA_{TMS}$  reduction can be due to modulation at any level of the corticospinal network (i.e. through reduced spinal and/or motoneuronal excitability).

The assessment of cortical voluntary activation using TMS (i.e.,  $VA_{TMS}$ ) cannot be performed for all muscles since some criteria must be met, especially having strong corticomotoneuronal projections [8].  $VA_{TMS}$  has been shown to be reliable for the elbow flexors [9] or the knee extensors [8], but not in dorsiflexor muscles thus far, despite the functional role of these muscles in particular during locomotion [10, 11]. The tibialis anterior (TA), contributing to almost half of the voluntary torque produced in ankle dorsiflexion (the remainder presumably being provided by the long extensors of the toes) [12], is known to have the strongest corticospinal projections among lower limb muscles [13]. Even though dorsiflexor muscles  $VA_{TMS}$  has already been assessed in fatigue studies [14, 15], no study has yet investigated  $VA_{TMS}$  reliability for this muscle group. Such an approach is necessary since the constant oscillation in the elements of the central nervous system and methodological factors (e.g. subject population, muscle considered, position for magnetic stimulation) are two main sources that may affect the stability of TMS measurements [16]. Indeed, motor-

105 evoked potential (MEP) elicited by TMS, which is known to represent the excitability  
106 of cortical neurons and/or motoneuron pools of the stimulated muscles [17], are  
107 inherently variable. For instance, MEP amplitude may be influenced by circadian  
108 variation [18]. To reduce variability, MEP amplitude is commonly expressed as ratio  
109 to M-wave to rise above the membrane excitability and conductivity influence. MEPs  
110 should also be measured during a stable voluntary contraction to further reduce  
111 variability [16, 19]. Among the studies that investigated TA MEP reliability [14, 16,  
112 20], only one met these last two criteria and reported high inter-day reliability for  
113 MEPs measured during submaximal lengthening and shortening contractions [16].  
114 During maximal isometric contraction, absolute MEP was also reported to be reliable  
115 [14], as it was at rest [14, 20]. Hence, intra-day and inter-day reliability of normalized  
116 TA MEPs during submaximal and maximal isometric contractions has yet to be  
117 determined.. Moreover, although input-output curves (i.e. the relation between TMS  
118 intensity and MEP amplitude) [19, 21] are recognized as a more sensitive measure of  
119 corticospinal excitability [22, 23], its reliability for the TA muscle has only been proven  
120 during a relaxed condition [20].

121  
122 Although several studies have investigated the reliability of TA TMS-related  
123 parameters [14, 16, 20], this topic needs further investigation. Specifically, dorsiflexor  
124  $VA_{TMS}$  must be examined to better understand short- and long-term adaptations  
125 within the central nervous system during acute and long term interventions. This  
126 would provide complementary knowledge with regard to previous TMS studies  
127 demonstrating changes in TA corticospinal excitability after strength training [24, 25].  
128 Therefore, the present study is intended to evaluate the intra- and inter-day reliability  
129 of  $VA_{TMS}$  and associated corticospinal functional properties (e.g. MEP amplitude) for  
130 dorsiflexor muscles.

131 **2. Material and methods**

132

133 **2.1. Participants.** Thirteen young, healthy young subjects (9 men and 4 women,  $23 \pm$   
134 4 years; height:  $177 \pm 8$  cm; body mass:  $69 \pm 10$  kg) participated in this study. The  
135 subjects' level of weekly physical activity was reported to be between 0 and 5 hours  
136 per week. Written, informed consent was obtained from all subjects prior to their  
137 participation and this study conformed to the standards from latest revision of the  
138 Helsinki Declaration and was approved by the local ethics committee. All subjects  
139 were free of lower-limb injury during the previous three months, contraindications to  
140 TMS, acute and chronic neurological disorders, and trauma. Before testing, each  
141 subject was informed about the nature and the aim of this study, as well as risks and  
142 discomfort associated with electrical and magnetic stimulation. They were instructed  
143 to abstain from caffeine for a minimum of 12 h before each session.

144

145 **2.2. Design of the study.** Subjects came for three visits to the laboratory at the  
146 same time of the day with a 1-week interval between visits. The first visit was a  
147 familiarization session of the entire experimental protocol. In order to assess the  
148 inter-day reliability, the same protocol was performed during the second and third  
149 visits, i.e., session 1 (S1) and session 3 (S3), respectively. Except for the TMS  
150 recruitment curve and the peripheral nerve stimulation setup (see below), the whole  
151 experimental protocol was performed twice with a 30-min resting period between  
152 measurement sessions during one of these visits (randomly designed), allowing the  
153 assessment of intra-day reliability (S1 and session 2 (S2)) (Fig. 1). The 30 minutes  
154 resting period began at the end of S1 and finished at the beginning of S2. Hence,  
155 subjects were invited to stay in a sitting position on the custom-built chair during the  
156 30-min period.

157

158 **2.3. Torque and electromyographic recordings.** Dorsiflexion torque was measured  
159 during voluntary contractions by a calibrated, instrumented pedal (CS1060 300 Nm,  
160 FGP Sensors, Les Clayes Sous Bois, France). Subjects were seated upright in a  
161 custom-built chair with a hip, knee, and ankle angle of 90, 120, and 90°, respectively.  
162 The foot was securely attached to the pedal with a custom designed hook and loop  
163 fastener. Electromyographic (EMG) activity of the right TA and soleus (SOL) was  
164 recorded with pairs of self-adhesive surface electrodes (Meditrace 100, Covidien,

165 Mansfield, USA) in bipolar configuration with a 30-mm interelectrode distance and the  
166 reference on the medial malleolus. SOL electrodes were placed 2 cm inferior to the  
167 insertion of the gastrocnemii on the Achilles tendon. TA electrodes were placed on  
168 the muscle belly parallel to the longitudinal axis of the muscle at one-third of the  
169 distance between the head of the fibula and the tip of the medial malleolus, according  
170 to SENIAM recommendations [26]. Skin impedance was measured using a classic  
171 impedance analyser (Fluke 87V Ex, Fluke, Wallisellen, Switzerland) and low  
172 impedance (<5 k $\Omega$ ) between electrodes was obtained by shaving, gently abrading the  
173 skin and then cleaning it with isopropyl alcohol. Signals were amplified with an octal  
174 bio-amplifier (ML138, ADInstruments; common mode rejection ratio = 85 dB, gain =  
175 500), bandpass filtered (5-500 Hz), and analogue-to-digital converted at a sampling  
176 rate of 2000 Hz by PowerLab system (16/30—ML880/P, ADInstruments, Bella Vista,  
177 Australia). All data was analyzed offline using Labchart 7 software (ADInstruments).

178  
179 **2.4. Stimulation.** Two types of stimulation were used. Specifically, electrical  
180 stimulation of the tibialis anterior motor nerve (i.e., peroneal nerve) and transcranial  
181 magnetic stimulation (TMS) over the motor hotspot for the TA muscle. For stimulation  
182 of the tibialis anterior motor nerve, a bipolar bar stimulating electrode with 30-mm  
183 anode-cathode spacing (Bipolar Felt Pad Stimulating Electrode Part number  
184 E.SB020/4mm, Digitimer) was positioned next to the fibular head. Single rectangular  
185 pulses with 0.5ms duration and 400 V maximal output voltage were delivered to the  
186 right peroneal nerve via a constant-current stimulator (DS7A, Digitimer, Welwyn  
187 Garden City, Hertfordshire, UK). The optimal site of stimulation was determined as  
188 the location eliciting the greatest TA M-wave for a given stimulus intensity. This site  
189 was marked with semi-permanent ink. To determine the optimal intensity of electrical  
190 stimulation, the intensity was increased by 5 mA until the magnitude of resting TA M-  
191 wave plateaued. The number of stimulations to find the optimal stimulating intensity  
192 varied among subjects (i.e., between 3 and 10). This optimal stimulating intensity was  
193 then increased by 50% to ensure supramaximal intensity of stimulation during  
194 contraction. Electrical stimulation configuration was determined at the beginning of  
195 the S1 and S3 to avoid any potentiation effects. The stimulation intensity used to  
196 evoke M-wave during MVC ranged between 30 and 97.5 mA. The mean stimulation  
197 intensities were  $47.6 \pm 20.2$  mA in S1 (the same intensity was used for S2) and 48.2



198 ± 17.0 mA in S3. The whole procedure of determining stimulation site and intensity  
199 was performed in approximately 3 min.

200  
201 For the TMS technique, motor evoked potentials (MEPs) and superimposed twitches  
202 (SITs) were evoked during voluntary isometric dorsiflexion via a Magstim 200<sup>2</sup>  
203 magnetic stimulator (Magstim 200<sup>2</sup>, The Magstim Company Ltd, Whitland, UK) with a  
204 110-mm double cone coil. The coil was positioned over the leg area of the left motor  
205 cortex along the nasal-inion axis to induce a postero-anterior current. The coil  
206 position to obtain an optimal activation of the TA is known to be 0.5-1 cm posterior  
207 and along the anteroposterior plane of the vertex [21]. This location was drawn on  
208 the swim cap every centimeter from 1 cm anterior to 3 cm posterior to the vertex  
209 along the nasal-inion line and also to 2 cm over the left motor cortex area (i.e., 8  
210 marks on the swim cap). Thus, a single stimulation was delivered at each of the  
211 specified anatomical positions (i.e. 8 stimulations in total) and the optimal coil position  
212 was chosen as the site eliciting the largest SIT and TA MEP amplitude with minimal  
213 SOL MEP amplitude in response to stimulation at 50% maximal stimulator output  
214 during brief voluntary dorsiflexion at 10% MVC. The whole procedure was performed  
215 in approximately 2 min.

216  
217 Subjects were provided with real-time feedback of their target force, which was  
218 displayed on a screen. This position was marked on a swim cap that was secured to  
219 the scalp. During voluntary contractions, TMS was delivered once the subject had  
220 stabilized the force on the target level. Subjects were instructed to re-contract as  
221 strongly and as fast as possible immediately after TMS delivery to permit accurate  
222 measurement of the cortical silent period (CSP). Electrodes for peripheral nerve  
223 stimulation as well as the TMS coil were placed and hand-held by the same  
224 investigator for all subjects and all visits.

## 2.5. Neuromuscular measurements.

227 **2.5.1. MVC.** After a standardized warm-up consisting of 10 submaximal voluntary  
228 contractions performed between 30-80% of the MVC obtained during the  
229 familiarization session, subjects performed three maximal dorsiflexion contractions  
230 lasting approximately 3 s, with a 1-min resting period between trials to avoid fatigue.  
231 Electrical stimulation was delivered during MVCs to obtain the maximal M-wave

232 peak-to-peak amplitude recorded during MVC ( $M_{MVC}$ ) (Fig. 1.B). MVC was  
233 determined as the highest torque value achieved before the electrical stimulation. TA  
234 EMG root mean square (RMS) was calculated as the mean over a 500-ms period  
235 after maximal force had reached a plateau and was then normalized to  $M_{MVC}$ .

236  
237 **2.5.2. Input-output curves.** TA TMS input-output curves were only assessed once  
238 during each visit to the laboratory to determine the optimal stimulation intensity.  
239 Hence, only inter-day reliability was investigated for this curve. TMS was delivered  
240 over the hotspot. Since corticospinal excitability is increased with muscle activity [27],  
241 input-output curves were determined during brief voluntary dorsiflexion at 20% MVC.  
242 Four consecutive 2-3 s contractions, with a 10 second rest between contractions,  
243 were performed at 20, 30, 40, 50, 60, 70 and 80% maximal stimulator output  
244 (randomly ordered), and a TMS pulse was delivered during each of these  
245 contractions (i.e., 28 stimulations were initially delivered). If a TA MEP plateau was  
246 not confirmed from these intensities, higher TMS intensities were investigated. The  
247 optimal stimulation intensity obtained from the input-output curves was chosen when  
248 TA MEP and SIT plateaued. The MEP evoked in the antagonist SOL was also  
249 measured to ensure it stayed at a low value (Fig 3). The whole procedure was  
250 performed in approximately 10 min.

251  
252 The aforementioned method has been recently reported to be appropriate for  
253 determining TMS intensity in the quadriceps [28]. The size of the TA MEPs was  
254 measured as the peak-to-peak amplitude of the non-rectified EMG response and  
255 expressed relative to  $M_{MVC}$ , and then averaged and plotted against the stimulus  
256 intensity. Then Boltzmann modelling was performed on input-output curves in order  
257 to apply a nonlinear regression analysis (Fig. 2.A) [21, 24, 29]:

$$\text{MEPampl} = \frac{P}{1 + \exp\left[\frac{150 - I}{k}\right]}$$

258 This function allows the determination of the relation between MEP amplitude  
259 (MEPampl) and the TMS intensity (I, in percentage of the stimulator output). Three  
260 parameters can be extracted from this relationship: the maximum MEP defined by the  
261 function (P, in %  $M_{MVC}$ ), the stimulus intensity at which the MEP size is 50% of the  
262 maximal MEP ( $I_{50}$ , in percentage of the stimulator output), and the slope parameter (k)

263 that is inversely proportional to the maximal function steepness [21]. The peak slope  
264 of the function was also calculated to provide an indication of the maximal rate of  
265 increase in MEP amplitude with stimulus intensity. The peak slope of the function is  
266 its tangent at  $I_{50}$  [19] and was further calculated by the following equation, taking into  
267 account the slope parameter (i.e.,  $k$ ) and the maximum MEP defined by the previous  
268 equation (i.e.,  $P$ ) [19, 24]:

$$\text{Peak slope} = \frac{k \times P}{4}$$

**2.5.3. Cortical voluntary activation.** Cortical voluntary activation ( $VA_{TMS}$ ) during  
maximal effort was calculated with the modified twitch interpolation technique as  
initially developed by Todd et al. [7]. Stimulus intensity used for the investigation of  
cortical voluntary activation was defined as the lowest stimulus intensity tested  
eliciting maximal SIT and TA MEP amplitude from the input-output curve test  
performed for corticospinal excitability measurement. The same intensity was used  
for intra-session reliability, but was not necessarily the same between S1 and S3  
(i.e., inter-day). The stimulus intensities used to assess  $VA_{TMS}$  were  $65.3 \pm 7.7\%$  and  
 $65.3 \pm 8.7\%$  of the maximal stimulator output in S1 and S3, respectively.  
Corticospinal excitability increases substantially during the transition from relaxed to  
contracted muscle states [30], thus underestimating TMS evoked twitch in the  
relaxed muscle. Hence, estimated resting twitch (ERT) was determined by linear  
regression of the relation between SIT amplitude evoked when TMS was delivered at  
100, 75 and 50% MVC and voluntary force (Fig. 2.B) [7]. This relation was  
extrapolated and the y-intercept was interpreted as the ERT amplitude (Fig. 4).  
 $VA_{TMS}$  was assessed with the equation:

$$VA_{TMS} = (1 - SIT_{MVC} / ERT) \times 100$$

where  $SIT_{MVC}$  is the superimposed twitch evoked during MVC. Three series were  
performed and averaged with contractions separated by 30 s and series by 2 min.  
Thus, the linear regression was run out with the 3 values from each series, and  
 $VA_{TMS}$  was then averaged. ERT was excluded and  $VA_{TMS}$  was not calculated when  
the linear regression was not linear within a series ( $r^2 < 0.9$ ) [31]. One series was  
excluded from the data analysis for 5 subjects (out of 39 series in total) and the  
average was done with 2 series rather than 3 in this case. ERT and  $SIT_{MVC}$  were  
determined on each series separately before being averaged. Mean  $R^2$  of the linear

294 regression was  $0.97 \pm 0.04$ ,  $0.97 \pm 0.05$  and  $0.95 \pm 0.03$  for S1, S2 and S3,  
295 respectively. MEPs and CSPs elicited by TMS during 100, 75 and 50% MVC  
296 contractions were also calculated and averaged over the 3 sets. Peak-to-peak  
297 amplitude of MEPs were measured and normalized to  $M_{MVC}$  ( $MEP_{MVC}$ ,  $MEP_{75}$  and  
298  $MEP_{50}$ , respectively). Since it was difficult to precisely determine the beginning and  
299 the end of the MEP, due to its polyphasic form, MEP area was not calculated. The  
300 duration of the CSPs were determined visually and defined as the duration from the  
301 TMS stimulus to the return of continuous voluntary EMG ( $CSP_{MVC}$ ,  $CSP_{75}$  and  $CSP_{50}$ ,  
302 respectively) (Fig. 2.C) Two investigators performed the CSP analysis, and the two  
303 values were averaged. The treatment was redone by a 3<sup>rd</sup> investigator when the  
304 difference between the two was higher than 5%. This large error occurred only for  
305 one subject (in that case, the average was performed on the values that were these  
306 closest to each other), showing that analysing CSP duration visually was not  
307 investigator-dependant and was thus an appropriate method.

**2.6. Statistical analysis.** As recommended by Hopkins [32], intra-session and inter-  
308 day reliabilities were assessed using the intra-class correlation coefficient (ICC),  
309 standard error of the mean (SEM) and coefficient of variation (CV). The ICC  
310 represents a measure of relative reliability showing the degree to which individuals  
311 maintain their position in a sample with repeated measurements [33]. According to  
312 Fleiss [34], ICC greater than 0.75 represents an excellent reliability; an ICC ranging  
313 between 0.74 and 0.40 represents a moderate to good reliability and an ICC smaller  
314 than 0.40 represents poor reliability. The SEM scores were also calculated to quantify  
315 absolute reliability and give an indication of the precision of a recorded parameter.  
316 Thus, a low SEM value means a strong reliability and vice versa [35]. The CV  
317 ((standard deviation (SD) / mean) was calculated to represent the degree to which  
318 repeated measurements vary for individuals [33]. Data are presented as means  $\pm$   
319 SD.  
320

321 **3. Results**

1  
2 322

3 323 **3.1. MVC and cortical voluntary activation ( $VA_{TMS}$ )**

4  
5 324 The results of intra- and inter-day reliabilities of MVC,  $M_{MVC}$ ,  $RMS/M_{MVC}$  and  $VA_{TMS}$   
6  
7 325 are presented in Table 1. Intra-day ICCs were 0.96 and 0.80 while inter-day ICCs  
8  
9 326 were 0.95 and 0.99, for MVC and  $VA_{TMS}$ , respectively. For these two parameters, all  
10  
11 327 CVs for intra- and inter-day conditions were lower than 4.1%, and were associated to  
12  
13 328 low SEM values.  $VA_{TMS}$  across the three testing sessions was  $96.0 \pm 4.9\%$ . Mean  
14  
15 329  $SIT_{MVC}$  and ERT values as well as their associated ICC, CV and SEM values are  
16  
17 330 reported in Table 1.

18 331

19  
20 332 **3.2. Corticospinal excitability and intracortical inhibition**

21  
22 333 Fig 2.C represents typical torque and EMG recordings (i.e., MEP amplitude and silent  
23  
24 334 period duration) in response to TMS during 100, 75 and 50% MVC contractions.  
25  
26 335 Mean results for CSP and MEP recorded at 100, 75 and 50% MVC are presented in  
27  
28 336 Table 1 as well as their reliability outcomes. For these two parameters, ICCs ranged  
29  
30 337 from 0.51 to 0.99 and CVs from 3.5% to 14.9%. The parameters of the input-output  
31  
32 338 curves are presented in Table 2.  $R^2$  of the Boltzmann sigmoid function were  $0.83 \pm$   
33  
34 339  $0.12$  and  $0.88 \pm 0.10$  for S1 and S3, respectively. For the four parameters extracted,  
35  
36 340 ICC values ranged from 0.71 to 0.87 and CVs from 8.0% to 43.9%.

37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

#### 341 4. Discussion

1 342 This study aimed to assess the intra- and inter-day reliability of several TMS-related  
2  
3 343 measures ( $VA_{TMS}$ , MEP amplitude, CSP, input-output curve parameters) related to  
4  
5 344 corticospinal output to a voluntarily activated muscle responsible for dorsiflexion in  
6  
7 345 healthy subjects. The main findings of the present study are that i) dorsiflexor  
8  
9 346 muscles  $VA_{TMS}$  was reported for the first time to be highly reliable, ii) MEPs and  
10  
11 347 CSPs showed reliable intra- and inter-day outcomes and iii) highly variable results  
12  
13 348 are reported for Boltzman modelling extracted parameters.

14 349  
15  
16 350 **4.1. MVC and cortical voluntary activation.** As expected, MVC presented good  
17  
18 351 intra-and inter-day reliability. To the best of our knowledge, cortical voluntary  
19  
20 352 activation level assessment using TMS during MVCs has only been demonstrated to  
21  
22 353 be reliable for elbow flexors [9] or knee extensors muscles [6, 8]. Although previous  
23  
24 354 fatigue studies investigated dorsiflexor  $VA_{TMS}$  [14, 15], the present study is the first  
25  
26 355 one to investigate its intra- and inter-day reliability.  $VA_{TMS}$ , as well as ERT,  
27  
28 356 demonstrated good to excellent intra-and inter-day reliabilities. An important result of  
29  
30 357 the present study is that  $VA_{TMS}$  recorded between days (i.e., inter-day reliability)  
31  
32 358 showed excellent reliability (i.e., 0.99 ICC, 0.8% CV and 0.4 SEM). Since neural  
33  
34 359 adaptations have been reported after strength training through CSP [24] and MEP  
35  
36 360 [25] increases, our results suggest that  $VA_{TMS}$  may be a relevant parameter to inform  
37  
38 361 on possible changes in the maximal neural drive from the motor cortex in longitudinal  
39  
40 362 studies although this may also reflect on a modification of spinal excitability.

41 363  
42 364  $VA_{TMS}$  was found to be  $96.0 \pm 4.9\%$  across the three testing sessions, highlighting  
43  
44 365 the inability of the nervous drive originating from the motor cortex to fully activate  
45  
46 366 motoneurons [5]. Voluntary activation measured in our study appears to be higher  
47  
48 367 compared to cortical voluntary activation recorded for elbow flexors (91-95%) [9, 31,  
49  
50 368 36-38] or knee extensors muscles (89-93%) [6, 8, 39, 40], and seem to be in  
51  
52 369 agreement with others showing that TA activation, quantified by the classical  
53  
54 370 superimposed stimulation method, was reported to be maximal or near maximal in a  
55  
56 371 healthy young population [41]. This may be explained by the fact that the TA has the  
57  
58 372 strongest corticospinal projections among all lower limb muscles [13].

59 373

374 Our values also appear to be greater than those reported by Ross et al., (2007) and  
375 Mileva et al., (2012) who found  $VA_{TMS}$  to be  $75.5 \pm 16.0\%$  and  $89.5 \pm 2.9\%$ ,  
376 respectively [14, 15]. These discrepancies may be due to differences in the TMS  
377 intensity and/or the type of TMS coil used to evaluate  $VA_{TMS}$ . Ross et al., (2007)  
378 evaluated the  $VA_{TMS}$  at 100% of maximal stimulator output using a 70 mm figure-of-  
379 eight coil [15], while a 110-mm double-cone coil was used in the present study. This  
380 difference in coil size in the present study may have caused an increased activation  
381 of the antagonist muscles leading to misinterpretation of the evoked superimposed  
382 twitches. In the study of Mileva et al., (2012), TMS intensity was set at 120% of the  
383 resting motor threshold [14]. According to a recent study, this may not have been  
384 optimal to fully activate the motor cortical neurons since such intensity selection does  
385 not precisely identify the minimum stimulus intensity needed to elicit SIT and MEPs of  
386 maximal amplitude [28]. The authors of the present study suggest that determining  
387 TMS intensity from an input-output curve (i.e., using optimal intensity identified as the  
388 minimum stimulus intensity needed to elicit MEPs of maximal amplitude with minimal  
389 co-activation) is more appropriate to assess corticospinal excitability and  $VA_{TMS}$ .

**4.2. Corticospinal excitability.** Input-output curves performed in the present study  
392 to determine TMS intensity for  $VA_{TMS}$  assessment were further used to characterize  
393 the excitability of the corticospinal pathway, through the Boltzmann modeling [21, 42].  
394 According to the ICC and SEM, the extracted parameters (i.e., the function plateau,  
395 the stimulus intensity required to obtain a MEP amplitude of half the size of the  
396 plateau, the slope parameter, and the peak slope) demonstrated good reliability and  
397 these values are in agreement with previous studies performed on first dorsal  
398 interosseous [19] or TA [20]. However, the present study demonstrated high CVs for  
399 the slope parameter and the peak slope (43.9 and 40.5%, respectively). The high  
400 reported CVs may question the reliability of these two parameters in the current  
401 study. Although Carroll et al., (2001) suggested that the slope parameter (i.e., the k  
402 parameter in our study) might be a more reliable index of corticospinal excitability  
403 than the peak slope for the first dorsal interosseous, similar reliability was reported for  
404 these two parameters in the present study [19]. Nevertheless, regarding the high CV  
405 described above, it seems difficult to use these two parameters to investigate the  
406 chronic and/or acute adaptations of TA corticospinal excitability.

408 While input-output curves were used in the present study to investigate TA  
409 corticospinal excitability during a low level of contraction, corticospinal excitability was  
410 also assessed at higher levels of contraction (100, 75 and 50% MVC) through MEP  
411 recordings. The current results present higher MEP values during submaximal (i.e.,  
412 MEP<sub>50</sub> and MEP<sub>75</sub>) than during maximal voluntary contractions (i.e., MEP<sub>MVC</sub>), which  
413 is in agreement with other results obtained on the TA [16]. It has been suggested that  
414 the probability for the TMS pulse to be evoked during the refractory period is higher  
415 during a maximal (higher discharger rate) rather than a submaximal voluntary  
416 contraction, leading to a lower MEP amplitude [8]. MEP<sub>MVC</sub> presented good intra- and  
417 inter-day reliabilities. These results are consistent with a recent study that found that  
418 the TA has a good inter-day reliability (ICC: 0.80) for MEPs recorded during MVC  
419 [14]. Even if ICC values are lower in our experiment (ICC: 0.51 and 0.52, for intra-  
420 and inter-day condition respectively), we can conclude that there was a moderate to  
421 good reliability regarding the low associate CV and SEM values, informing about  
422 small intra-subject variation. One aim of the present study was to investigate the  
423 intra-day reliability of MEP amplitude, which is known to represent corticospinal  
424 pathway excitability. Our results have shown that MEPs recorded at maximal or  
425 submaximal levels of voluntary contraction may be used as reliable parameters to  
426 investigate acute phenomenon such as fatigue. The good reliability observed in the  
427 present study is also in agreement with several studies performed on knee extensors  
428 [8] or elbow flexors [9]. Yet, unlike the studies where the reliability was independent  
429 on force level, in the present study, MEP reliability was higher at low level of  
430 voluntary contraction (i.e., MEP<sub>50</sub> and MEP<sub>75</sub>) than during MVC (i.e., MEP<sub>MVC</sub>). The  
431 good reliability observed for MEPs recorded during submaximal contraction is  
432 consistent with previous studies investigating the knee extensors [8], first dorsal  
433 interosseous [43], elbow flexor [9] or TA [16] muscles. The lower reliability observed  
434 for MEPs during MVC is consistent with previous studies, suggesting that high  
435 contraction intensities may cause a larger desynchronization of the action potential at  
436 the membrane level [16, 19]. Hence, the excitability of corticospinal pathway of the  
437 TA muscle can be reliably tested through the recording of MEP amplitude in  
438 response to TMS during maximal and submaximal contractions for acute  
439 interventions (e.g., fatigue) or longitudinal studies (e.g. training).



441 **4.3. Intracortical inhibition.** Cortical silent periods (CSP) associated with MEPs  
1 442 were also measured at 100, 75 and 50% MVC (i.e.,  $CSP_{MVC}$ ,  $CSP_{75}$  and  $CSP_{50}$ ,  
2 443 respectively). When applied during a voluntary contraction, TMS may provide  
3 444 information about GABA mediated intracortical inhibition through the assessment of  
4 445 the CSP [44]. Although the exact mechanisms of the CSP remain unclear, it is  
5 446 accepted that the initial part is mainly due to reduced spinal excitability while the final  
6 447 part of the CSP is cortical in origin [45]. CSP results from the activation of inhibitory  
7 448 cortical neurons projecting onto the pyramidal cells in the motor cortex [46], and is  
8 449 thought to be mainly influenced by GABA (gamma aminobutyric acid)  
9 450 neurotransmission [47]. In the present study,  $CSP_{MVC}$  was highly reliable, as shown  
10 451 by high ICC and low CV-SEM values for both intra- and inter-day comparisons. To  
11 452 date, only one study recorded CSP in TA during MVC [14] and similar ICC (0.93)  
12 453 were observed. In the present study,  $CSP_{75}$  and  $CSP_{50}$  were also found to be highly  
13 454 reliable, both between S1 and S2 (intra-day) and between S1 and S3 (inter-day), in  
14 455 agreement with previous studies that assessed CSP during submaximal voluntary  
15 456 contractions in TA [16, 20], first dorsal interosseous [43] and the extensor digitorum  
16 457 communis [48] muscles. Hence, CSP can be reliably used during maximal and  
17 458 submaximal dorsiflexion to investigate TA intracortical inhibition.  
18 459

34 460 As most reliability studies, the principal limitation of the present work is that TMS-  
35 461 related parameters were investigated for a healthy young population, and that current  
36 462 outcomes may not be applicable to other populations (i.e., clinical, elderly).  
37 463

41 464 In conclusion, the present study reports for the first time the excellent intra- and inter-  
42 465 day reliability for  $VA_{TMS}$  of dorsiflexor muscles. The outcomes were found to be highly  
43 466 reliable within and between days. A good reliability in the assessment of corticospinal  
44 467 excitability and intra-cortical inhibition directed towards the TA was also observed.  
45 468 TMS can thus be used to better understand the effects of training interventions on TA  
46 469 neural adaptations (e.g., neural drive from the cortex, corticospinal pathway  
47 470 excitability) in a healthy young population. This has important practical consequences  
48 471 because, besides the fact that TA has the most corticospinal projections among all  
49 472 lower limb muscles, it also plays an important functional role in daily life abilities such  
50 473 as lifting the foot in the swing phase during locomotion.  
51 474

475 **Acknowledgments**

1  
2 476 The authors sincerely thank Régis Bonnefoy for technical assistance as well as Aude  
3  
4 477 Carlet-Mallet and Michael James Asmussen for revision. The present study was  
5  
6 478 supported by a doctoral research grant from the University Jean Monnet of Saint-  
7 479 Etienne (France) assigned to Robin Souron.  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

480 **References**

- 1  
2 481  
3 482 [1] Merton PA. Voluntary strength and fatigue. *J Physiol.* 1954;123:553-64.  
4  
5 483 [2] Gandevia SC. Spinal and supraspinal factors in human muscle fatigue. *Physiol*  
6  
7 484 *Rev.* 2001;81:1725-89.  
8  
9 485 [3] Giesebrecht S, van Duinen H, Todd G, Gandevia SC, Taylor JL. Training in a  
10 486 ballistic task but not a visuomotor task increases responses to stimulation of human  
11  
12 487 corticospinal axons. *J Neurophysiol.* 2012;107:2485-92.  
13  
14 488 [4] Aagaard P, Simonsen EB, Andersen JL, Magnusson P, Dyhre-Poulsen P. Neural  
15  
16 489 adaptation to resistance training: changes in evoked V-wave and H-reflex responses.  
17  
18 490 *J Appl Physiol.* 2002;92:2309-18.  
19  
20 491 [5] Goodall S, Howatson G, Romer L, Ross E. Transcranial magnetic stimulation in  
21 492 sport science: a commentary. *Eur J Sport Sci.* 2014;14 Suppl 1:S332-40.  
22  
23 493 [6] Sidhu SK, Bentley DJ, Carroll TJ. Cortical voluntary activation of the human knee  
24  
25 494 extensors can be reliably estimated using transcranial magnetic stimulation. *Muscle*  
26  
27 495 *Nerve.* 2009;39:186-96.  
28  
29 496 [7] Todd G, Taylor JL, Gandevia SC. Measurement of voluntary activation of fresh  
30 497 and fatigued human muscles using transcranial magnetic stimulation. *J Physiol.*  
31  
32 498 2003;551:661-71.  
33  
34 499 [8] Goodall S, Romer LM, Ross EZ. Voluntary activation of human knee extensors  
35  
36 500 measured using transcranial magnetic stimulation. *Exp Physiol.* 2009;94:995-1004.  
37  
38 501 [9] Todd G, Taylor JL, Gandevia SC. Reproducible measurement of voluntary  
39 502 activation of human elbow flexors with motor cortical stimulation. *J Appl Physiol.*  
40  
41 503 2004;97:236-42.  
42  
43 504 [10] Billot M, Simoneau E, Van Hoecke J, Martin A. Coactivation at the ankle joint is  
44  
45 505 not sufficient to estimate agonist and antagonist mechanical contribution. *Muscle*  
46  
47 506 *Nerve.* 2010;41:511-8.  
48  
49 507 [11] Hase K, Stein RB. Analysis of rapid stopping during human walking. *J*  
50 508 *Neurophysiol.* 1998;80:255-61.  
51  
52 509 [12] Marsh E, Sale D, McComas AJ, Quinlan J. Influence of joint position on ankle  
53  
54 510 dorsiflexion in humans. *J Appl Physiol Respir Environ Exerc Physiol.* 1981;51:160-7.  
55  
56 511 [13] Perez MA, Lungholt BK, Nyborg K, Nielsen JB. Motor skill training induces  
57  
58 512 changes in the excitability of the leg cortical area in healthy humans. *Exp Brain Res.*  
59  
60 513 2004;159:197-205.  
61  
62  
63  
64  
65

- 514 [14] Mileva KN, Sumners DP, Bowtell JL. Decline in voluntary activation contributes  
1 515 to reduced maximal performance of fatigued human lower limb muscles. *Eur J Appl*  
2 516 *Physiol.* 2012;112:3959-70.
- 5 517 [15] Ross EZ, Middleton N, Shave R, George K, Nowicky A. Corticomotor excitability  
6 518 contributes to neuromuscular fatigue following marathon running in man. *Exp Physiol.*  
7 519 2007;92:417-26.
- 10 520 [16] Tallent J, Goodall S, Hortobagyi T, St Clair Gibson A, French DN, Howatson G.  
11 521 Repeatability of corticospinal and spinal measures during lengthening and shortening  
12 522 contractions in the human tibialis anterior muscle. *PLoS One.* 2012;7:e35930.
- 16 523 [17] Mills K. *Magnetic stimulation of the human nervous system: Oxford University*  
17 524 *Press; 1999.*
- 19 525 [18] Strutton PH, Catley M, Davey NJ. Stability of corticospinal excitability and grip  
20 526 force in intrinsic hand muscles in man over a 24-h period. *Physiol Behav.*  
21 527 2003;79:679-82.
- 25 528 [19] Carroll TJ, Riek S, Carson RG. Reliability of the input-output properties of the  
26 529 cortico-spinal pathway obtained from transcranial magnetic and electrical stimulation.  
27 530 *J Neurosci Methods.* 2001;112:193-202.
- 31 531 [20] Cacchio A, Cimini N, Alosi P, Santilli V, Marrelli A. Reliability of transcranial  
32 532 magnetic stimulation-related measurements of tibialis anterior muscle in healthy  
33 533 subjects. *Clin Neurophysiol.* 2009;120:414-9.
- 36 534 [21] Devanne H, Lavoie BA, Capaday C. Input-output properties and gain changes in  
37 535 the human corticospinal pathway. *Exp Brain Res.* 1997;114:329-38.
- 40 536 [22] Boroojerdi B, Battaglia F, Muellbacher W, Cohen LG. Mechanisms influencing  
41 537 stimulus-response properties of the human corticospinal system. *Clin Neurophysiol.*  
42 538 2001;112:931-7.
- 45 539 [23] Ridding MC, Rothwell JC. Stimulus/response curves as a method of measuring  
46 540 motor cortical excitability in man. *Electroencephalogr Clin Neurophysiol.*  
47 541 1997;105:340-4.
- 50 542 [24] Christie A, Kamen G. Cortical inhibition is reduced following short-term training in  
51 543 young and older adults. *Age.* 2014;36:749-58.
- 54 544 [25] Griffin L, Cafarelli E. Transcranial magnetic stimulation during resistance training  
55 545 of the tibialis anterior muscle. *J Electromyogr Kinesiol.* 2007;17:446-52.

- 546 [26] Hermens HJ, Freriks B, Disselhorst-Klug C, Rau G. Development of  
1 547 recommendations for SEMG sensors and sensor placement procedures. J  
2  
3 548 Electromyogr Kinesiol. 2000;10:361-74.
- 5 549 [27] Lazzaro VD, Restuccia D, Oliviero A, Profice P, Ferrara L, Insola A, et al. Effects  
6  
7 550 of voluntary contraction on descending volleys evoked by transcranial stimulation in  
8  
9 551 conscious humans. J Physiol. 1998;508:625-33.
- 10 552 [28] Temesi J, Gruet M, Rupp T, Verges S, Millet GY. Resting and active motor  
11  
12 553 thresholds versus stimulus-response curves to determine transcranial magnetic  
13  
14 554 stimulation intensity in quadriceps femoris. J Neuroeng Rehabil. 2014;11:40.
- 15  
16 555 [29] Kaelin-Lang A, Luft AR, Sawaki L, Burstein AH, Sohn YH, Cohen LG. Modulation  
17  
18 556 of human corticomotor excitability by somatosensory input. J Physiol. 2002;540:623-  
19  
20 557 33.
- 21  
22 558 [30] Ugawa Y, Terao Y, Hanajima R, Sakai K, Kanazawa I. Facilitatory effect of tonic  
23  
24 559 voluntary contraction on responses to motor cortex stimulation. Electroencephalogr  
25  
26 560 Clin Neurophysiol. 1995;97:451-4.
- 27 561 [31] Hunter SK, Butler JE, Todd G, Gandevia SC, Taylor JL. Supraspinal fatigue does  
28  
29 562 not explain the sex difference in muscle fatigue of maximal contractions. J Appl  
30  
31 563 Physiol. 2006;101:1036-44.
- 32  
33 564 [32] Hopkins WG. Measures of reliability in sports medicine and science. Sports Med.  
34  
35 565 2000;30:1-15.
- 36 566 [33] Suzuki S, Hayami A, Suzuki M, Watanabe S, Hutton RS. Reductions in  
37  
38 567 recruitment force thresholds in human single motor units by successive voluntary  
39  
40 568 contractions. Exp Brain Res. 1990;82:227-30.
- 41  
42 569 [34] Fleiss JL. The measurement of interrater agreement. Statistical methods for  
43  
44 570 rates and proportions. 1981;2:212-36.
- 45 571 [35] Larsson B, Karlsson S, Eriksson M, Gerdle B. Test-retest reliability of EMG and  
46  
47 572 peak torque during repetitive maximum concentric knee extensions. J Electromyogr  
48  
49 573 Kinesiol. 2003;13:281-7.
- 50  
51 574 [36] Hunter SK, Todd G, Butler JE, Gandevia SC, Taylor JL. Recovery from  
52  
53 575 supraspinal fatigue is slowed in old adults after fatiguing maximal isometric  
54  
55 576 contractions. J Appl Physiol. 2008;105:1199-209.
- 56 577 [37] Keller-Ross ML, Pereira HM, Pruse J, Yoon T, Schlinder-DeLap B, Nielson KA,  
57  
58 578 et al. Stressor-induced increase in muscle fatigability of young men and women is  
59  
60 579 predicted by strength but not voluntary activation. J Appl Physiol. 2014;116:767-78.

- 580 [38] Kennedy DS, McNeil CJ, Gandevia SC, Taylor JL. Fatigue-related firing of distal  
1 581 muscle nociceptors reduces voluntary activation of proximal muscles of the same  
2  
3 582 limb. *J Appl Physiol.* 2014;116:385-94.
- 5 583 [39] Temesi J, Rupp T, Martin V, Arnal PJ, Feasson L, Verges S, et al. Central  
6 584 fatigue assessed by transcranial magnetic stimulation in ultratrail running. *Med Sci*  
7 585 *Sports Exerc.* 2014;46:1166-75.
- 10 586 [40] Zghal F, Martin V, Thorkani A, Arnal PJ, Tabka Z, Cottin F. Effects of endurance  
11 587 training on the maximal voluntary activation level of the knee extensor muscles. *Eur J*  
12 588 *Appl Physiol.* 2013.
- 16 589 [41] Klass M, Baudry S, Duchateau J. Aging does not affect voluntary activation of  
17 590 the ankle dorsiflexors during isometric, concentric, and eccentric contractions. *J Appl*  
18 591 *Physiol.* 2005;99:31-8.
- 21 592 [42] Capaday C. Neurophysiological methods for studies of the motor system in freely  
22 593 moving human subjects. *J Neurosci Methods.* 1997;74:201-18.
- 25 594 [43] Orth M, Rothwell JC. The cortical silent period: intrinsic variability and relation to  
26 595 the waveform of the transcranial magnetic stimulation pulse. *Clin Neurophysiol.*  
27 596 2004;115:1076-82.
- 31 597 [44] Terao Y, Ugawa Y, Hanajima R, Machii K, Furubayashi T, Mochizuki H, et al.  
32 598 Predominant activation of I1-waves from the leg motor area by transcranial magnetic  
33 599 stimulation. *Brain Res.* 2000;859:137-46.
- 36 600 [45] Chen R, Lozano AM, Ashby P. Mechanism of the silent period following  
37 601 transcranial magnetic stimulation. Evidence from epidural recordings. *Exp Brain Res.*  
38 602 1999;128:539-42.
- 41 603 [46] Inghilleri M, Berardelli A, Cruccu G, Manfredi M. Silent period evoked by  
42 604 transcranial stimulation of the human cortex and cervicomedullary junction. *J Physiol.*  
43 605 1993;466:521-34.
- 47 606 [47] Werhahn KJ, Kunesch E, Noachtar S, Benecke R, Classen J. Differential effects  
48 607 on motorcortical inhibition induced by blockade of GABA uptake in humans. *J*  
49 608 *Physiol.* 1999;517 ( Pt 2):591-7.
- 52 609 [48] Wolf SL, Butler AJ, Campana GI, Parris TA, Struys DM, Weinstein SR, et al.  
53 610 Intra-subject reliability of parameters contributing to maps generated by transcranial  
54 611 magnetic stimulation in able-bodied adults. *Clin Neurophysiol.* 2004;115:1740-7.

614 **Figures captions**

1 615

2

3 616

4

5 617

6

7 618

8

9 619

10

11 620

12

13 621

14

15 622

16

17 623

18

19 624

20

21 625

22 626

23

24 627

25

26 628

27

28 629

29

30 630

31 631

32

33 632

34

35 633

36

37 634

38

39 635

40

41 636

42

43 637

44

45 638

46

47 639

48

49 640

50

51 641

52

53 642

54

55 643

56

57

58

59

60

61

62

63

64

65

**Fig.1.** Overview of the experimental design of the study (A) and the neuromuscular testing protocol (B). The measurements were performed during session 1 (S1), session 2 (S2) and session 3 (S3) except for peripheral nerve stimulation (PNS) and transcranial magnetic stimulation (TMS) setup which were performed only for S1 and S3.

**Fig. 2.** (A) Typical TA MEP input-output curve. Mean TA MEP amplitudes (white squares) are presented as a function of TMS intensity from 20 to 80% of maximal stimulator output. The *solid line* represents the Boltzman sigmoid function fit to that set of data. *Dashed lines* indicate  $I_{50}$  (i.e. stimulus intensity required to obtain a MEP amplitude of half the size of the plateau,  $\frac{1}{2} P$ ) and P (i.e. the maximum MEP defined by the function). (B) Superimposed twitch force evoked with TMS during maximal (i.e. 100% MVC) and submaximal (i.e. 75% and 50% MVC) voluntary contractions. (C) Example of electromyographic recording in response to transcranial magnetic stimulation (TMS) during maximal (100% MVC) and submaximal voluntary contractions (75% and 50% MVC). The time interval between black and grey arrows represents the way to assess cortical silent period (CSP) duration. Black arrows represent the time of TMS stimulation. Grey arrows represent the resumption of continuous voluntary EMG. Dotted arrows represent the MEP peak-to-peak amplitude. Data are from a representative subject.

**Fig. 3.** Input-output curve for a representative subject. The optimal stimulation intensity was selected when MEP TA plateaued and when MEP SOL was small. For this subject, 60% of the maximal stimulator output was chosen.

**Fig.4.** Amplitude of superimposed twitches (SIT) produced by TMS during contraction at 50% (triangles), 75% (circles) and 100% MVC (squares). The linear regression was extrapolated and the y-intercept (dashed line) was interpreted as the ERT amplitude. Data from a representative subject.

1  
2  
3  
4  
5  
6 644  
7 645  
8 646  
9

**Table 1.** Intra- and inter-day reliabilities of neuromuscular parameters recorded during maximal (MVC) and submaximal (75% and 50% MVC) voluntary contractions

	Mean ± SD			Intra-day			Inter-day		
	S1	S2	S3	ICC	CV (%)	SEM	ICC	CV (%)	SEM
<b>MVC (N.m)</b>	<b>52.4 ± 12.6</b>	<b>52.6 ± 12.9</b>	<b>51.4 ± 12.1</b>	<b>0.96</b>	<b>3.2 ± 2.4</b>	<b>2.7</b>	<b>0.95</b>	<b>4.1 ± 3.0</b>	<b>2.7</b>
<b>RMS / M<sub>MVC</sub> (%)</b>	<b>9.9 ± 2.3</b>	<b>10.8 ± 2.2</b>	<b>10.5 ± 2.6</b>	<b>0.87</b>	<b>7.7 ± 5.2</b>	<b>0.82</b>	<b>0.68</b>	<b>7.9 ± 6.1</b>	<b>1.4</b>
<b>VA<sub>TMS</sub> (%)</b>	<b>95.7 ± 5.1</b>	<b>95.9 ± 4.7</b>	<b>96.4 ± 4.9</b>	<b>0.80</b>	<b>1.7 ± 1.5</b>	<b>2.3</b>	<b>0.99</b>	<b>0.8 ± 0.9</b>	<b>0.4</b>
<b>SIT<sub>MVC</sub> (N.m)</b>	<b>0.76 ± 0.97</b>	<b>0.67 ± 0.88</b>	<b>0.60 ± 0.40</b>	<b>0.73</b>	<b>46.4 ± 40.5</b>	<b>0.5</b>	<b>0.95</b>	<b>38.3 ± 41.7</b>	<b>0.2</b>
<b>ERT (N.m)</b>	<b>16.4 ± 8.1</b>	<b>16.2 ± 4.9</b>	<b>17.4 ± 7.3</b>	<b>0.47</b>	<b>13.2 ± 13.2</b>	<b>4.9</b>	<b>0.94</b>	<b>13.9 ± 14.5</b>	<b>1.8</b>
<b>M<sub>MVC</sub> (mV)</b>	<b>5.5 ± 1.1</b>	<b>5.7 ± 1.1</b>	<b>6.2 ± 1.7</b>	<b>0.94</b>	<b>6.5 ± 3.7</b>	<b>0.3</b>	<b>0.36</b>	<b>14.1 ± 8.7</b>	<b>1.1</b>
<b>MEP<sub>50</sub> (mV)</b>	<b>3.1 ± 0.6</b>	<b>3.1 ± 0.8</b>	<b>3.3 ± 1.2</b>	<b>0.83</b>	<b>8.6 ± 6.3</b>	<b>0.28</b>	<b>0.55</b>	<b>13.1 ± 9.3</b>	<b>0.42</b>
<b>MEP<sub>75</sub> (mV)</b>	<b>3.5 ± 0.8</b>	<b>3.5 ± 0.8</b>	<b>3.6 ± 0.9</b>	<b>0.92</b>	<b>5.6 ± 4.6</b>	<b>0.23</b>	<b>0.92</b>	<b>6.7 ± 6.5</b>	<b>0.26</b>
<b>MEP<sub>MVC</sub> (mV)</b>	<b>3.1 ± 0.6</b>	<b>3.1 ± 0.8</b>	<b>3.3 ± 1.2</b>	<b>0.80</b>	<b>8.9 ± 6.9</b>	<b>0.31</b>	<b>0.47</b>	<b>11.2 ± 10.7</b>	<b>0.67</b>
<b>MEP<sub>50</sub> (% M<sub>MVC</sub>)</b>	<b>65.2 ± 14.7</b>	<b>65.3 ± 11.4</b>	<b>61.9 ± 14.9</b>	<b>0.75</b>	<b>10.3 ± 7.5</b>	<b>6.6</b>	<b>0.70</b>	<b>13.4 ± 8.1</b>	<b>8.1</b>
<b>MEP<sub>75</sub> (% M<sub>MVC</sub>)</b>	<b>75.8 ± 19.1</b>	<b>74.7 ± 13.7</b>	<b>74.4 ± 18.9</b>	<b>0.86</b>	<b>7.1 ± 5.0</b>	<b>6.1</b>	<b>0.88</b>	<b>7.5 ± 7.4</b>	<b>6.4</b>
<b>MEP<sub>MVC</sub> (% M<sub>MVC</sub>)</b>	<b>57.0 ± 14.0</b>	<b>50.1 ± 12.3</b>	<b>57.0 ± 14.4</b>	<b>0.51</b>	<b>14.9 ± 8.1</b>	<b>8.7</b>	<b>0.52</b>	<b>9.9 ± 10.4</b>	<b>9.2</b>
<b>CSP<sub>50</sub> (ms)</b>	<b>246 ± 58</b>	<b>266 ± 67</b>	<b>230 ± 58</b>	<b>0.84</b>	<b>6.4 ± 7.4</b>	<b>25.2</b>	<b>0.79</b>	<b>7.8 ± 9.2</b>	<b>26.2</b>
<b>CSP<sub>75</sub> (ms)</b>	<b>261 ± 64</b>	<b>271 ± 63</b>	<b>246 ± 59</b>	<b>0.99</b>	<b>3.5 ± 3.9</b>	<b>6.4</b>	<b>0.86</b>	<b>7.6 ± 8.3</b>	<b>23.3</b>
<b>CSP<sub>MVC</sub>(ms)</b>	<b>250 ± 51</b>	<b>267 ± 54</b>	<b>229 ± 62</b>	<b>0.98</b>	<b>7.3 ± 10.9</b>	<b>7.4</b>	<b>0.95</b>	<b>8.6 ± 11.6</b>	<b>13.2</b>

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49



647 Absolute data and CV are expressed as mean  $\pm$  standard deviation  
 1 648 S1: session 1; S2: session 2; S3: session 3  
 2 649 MVC: maximal voluntary contraction; RMS: root mean square; VA<sub>TMS</sub>: cortical voluntary  
 3 650 activation; SIT<sub>MVC</sub>: superimposed twitch amplitude evoked during MVC; ERT: estimated  
 4 651 resting twitch amplitude; M<sub>MVC-PP</sub>: M wave amplitude recorded during MVC; MEP: motor  
 5 652 evoked potential recorded at 50% MVC (MEP<sub>50</sub>), 75% MVC (MEP<sub>75</sub>) and 100% MVC  
 6 653 (MEP<sub>MVC</sub>). CSP: cortical silent period recorded at 50% MVC (CSP<sub>50</sub>), 75% MVC (CSP<sub>75</sub>) and  
 7 654 100% MVC (CSP<sub>MVC</sub>).  
 8  
 9 655 ICC: intraclass correlation coefficients; CV: coefficient of variation; SEM: standard error of  
 10 656 the mean.

11  
12 657  
13 658  
14  
15 659 **Table 2.** Intra- and inter-day reliabilities for input-output curves parameters.  
16 660

	Mean $\pm$ SD		Inter-day		
	S1	S3	ICC	CV (%)	SEM
<b>k</b>	<b>0.06 <math>\pm</math> 0.03</b>	<b>0.07 <math>\pm</math> 0.03</b>	<b>0.73</b>	<b>43.9 <math>\pm</math> 37.7</b>	<b>0.02</b>
<b>P (%M<sub>MVC</sub>)</b>	<b>55.8 <math>\pm</math> 12.3</b>	<b>49.0 <math>\pm</math> 12.6</b>	<b>0.71</b>	<b>14.8 <math>\pm</math> 11.3</b>	<b>6.7</b>
<b>I<sub>50</sub> (% stimulator output)</b>	<b>37.1 <math>\pm</math> 7.4</b>	<b>37.2 <math>\pm</math> 9.1</b>	<b>0.87</b>	<b>8.0 <math>\pm</math> 6.3</b>	<b>3.0</b>
<b>Peak slope</b>	<b>0.81 <math>\pm</math> 0.55</b>	<b>0.88 <math>\pm</math> 0.43</b>	<b>0.71</b>	<b>40.4 <math>\pm</math> 37.9</b>	<b>0.27</b>

28 661 Absolute data and CV are expressed as mean  $\pm$  standard deviation  
 29 662 S1: session 1; S3: session 3  
 30  
 31 663 K: slope parameter; P: function plateau; I<sub>50</sub>: stimulus intensity required to obtain a MEP  
 32 664 amplitude of half the size of the plateau; ICC: intraclass correlation coefficients; CV:  
 33 665 coefficient of variation; SEM: standard error of the mean.  
 34 666




Figure 2






Figure 3




Figure 4

