



**HAL**  
open science

## No evidence for behavioural syndrome and genetic basis for three personality traits in a wild bird population

Jennifer Morinay, Grégory Daniel, Lars Gustafsson, Blandine Doligez

### ► To cite this version:

Jennifer Morinay, Grégory Daniel, Lars Gustafsson, Blandine Doligez. No evidence for behavioural syndrome and genetic basis for three personality traits in a wild bird population. *Animal Behaviour*, 2019, 153, pp.69-82. 10.1016/j.anbehav.2019.05.001 . hal-02413241

**HAL Id: hal-02413241**

**<https://hal.science/hal-02413241>**

Submitted on 24 Nov 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **No evidence for behavioural syndrome and genetic basis for three**  
2 **personality traits in a wild bird population**

3

4

5 **Jennifer Morinay<sup>1,2,\*</sup>, Gregory Daniel<sup>1</sup>, Lars Gustafsson<sup>2</sup>, Blandine Doligez<sup>1</sup>**

6

7 <sup>1</sup> Université de Lyon, Université Lyon 1, CNRS, Laboratoire de Biométrie et Biologie  
8 Evolutive UMR 5558, Villeurbanne, France

9 <sup>2</sup> Department of Ecology and Evolution/Animal Ecology, Evolutionary Biology Centre,  
10 Uppsala University, Uppsala, Sweden

11

12 Received 3 September 2018

13 Initial acceptance 31 October 2018

14 Final acceptance 25 March 2019

15 MS number 18-00633R

16

17 \* *Correspondence:* J. Morinay, LBBE, CNRS UMR 5558, Univ Lyon–Université Claude  
18 Bernard Lyon 1, Bâtiment Gregor Mendel, 43 boulevard du 11 novembre 1918, 69622  
19 Villeurbanne Cedex, France.

20 E-mail address: [jennifer.morinay@gmail.com](mailto:jennifer.morinay@gmail.com)

21

22

23 **Keywords:** aggressiveness, between- and within-individual correlations, boldness, collared  
24 flycatcher, *Ficedula albicollis*, heritability, neophobia, parental effects, pedigree, quantitative  
25 genetics, repeatability

26 **ABSTRACT**

27

28 Personality traits and their correlations have been shown to be linked with life history  
29 strategies and fitness in various species. Between-individual correlations (i.e. behavioural  
30 syndromes) between personality traits can affect the evolutionary responses of these traits to  
31 environmental variation. Understanding the genetic and ecological determinants of  
32 personality traits and their interactions as behavioural syndromes in the wild is thus needed to  
33 shed light on the mechanisms shaping their evolution. Partitioning the observed (co)variance  
34 in these traits, however, requires large numbers of repeated behavioural measures on many  
35 individuals of known relatedness level. In the absence of such data, it is thus often assumed  
36 that phenotypic (co)variances inform about (1) underlying between-individual (co)variances  
37 (i.e. ignoring within-individual (co)variances) and (2) underlying genetic (co)variances. We  
38 tested these assumptions using three personality traits collected during 3 years on a long-term  
39 monitored breeding population of collared flycatchers, *Ficedula albicollis*. We partitioned the  
40 observed phenotypic (co)variance of aggressiveness, boldness and neophobia into genetic,  
41 permanent environment and parental components, and we estimated the repeatability and  
42 heritability of these traits and their between-individual correlations. All three traits were  
43 repeatable between years (at least on the latent scale) but none were heritable. Permanent  
44 environment effects explained 15% of the phenotypic variance in aggressiveness, and parental  
45 effects explained 25% of the phenotypic variance in neophobia, in line with previous studies  
46 in wild populations. The three traits showed phenotypic correlations but no between-  
47 individual correlations and no additive genetic covariance. Thus, our results did not support  
48 the assumptions that phenotypic covariance reflects behavioural syndromes and genetic  
49 covariance. We discuss the reasons for the absence of heritability and between-individual and

50 genetic covariance between these three personality traits in light of the possible selective  
51 pressures acting on this population.

52

53

## 54 **INTRODUCTION**

55

56 Over the past two decades, personality traits, that is, repeatable between-individual  
57 behavioural differences across time and contexts (Réale, Reader, Sol, McDougall, &  
58 Dingemanse, 2007), have received increasing attention in animal behavioural and  
59 evolutionary ecology studies (Bell, 2007; Bell, Hankison, & Laskowski, 2009; Carere &  
60 Maestriperi, 2013; Dingemanse & Réale, 2005; Réale et al., 2007; Sih, Bell, & Johnson,  
61 2004; Sih, Bell, Johnson, & Ziemba, 2004). Five ecologically important personality axes have  
62 been identified to characterize the behavioural responses of individuals when interacting with  
63 their environment (activity, exploration, boldness) and with others (aggressiveness,  
64 sociability; Réale et al., 2007). Personality traits have been shown to depend on ecological  
65 parameters (e.g. Réale et al., 2007; Sih, Cote, Evans, Fogarty, & Pruitt, 2012), to be heritable  
66 (e.g. van Oers, de Jong, van Noordwijk, & Drent, 2005; van Oers & Sinn, 2013), to be linked  
67 to life history traits or fitness (Dingemanse, Both, Drent, & Tinbergen, 2004; Dingemanse &  
68 Réale, 2013; Duckworth & Kruuk, 2009; Reale et al., 2010; Schuett, Tregenza, & Dall, 2010;  
69 Smith & Blumstein, 2008) and often to correlate with each other at the individual level,  
70 forming so-called behavioural syndromes (Garamszegi, Markó, & Herczeg, 2012; Andrew  
71 Sih, Bell, & Johnson, 2004; van Oers & Sinn, 2013). Such correlations may result from a  
72 functional integration of personality traits favoured by selection when interactions between  
73 these traits increase individuals' fitness in given environmental conditions (e.g. Dingemanse  
74 et al., 2007). Altogether, these various results reveal the crucial role that personality traits may

75 play in shaping evolutionary processes in wild populations (Dingemanse et al., 2004;  
76 Duckworth & Badyaev, 2007; Karlsson Green, Eroukhmanoff, Harris, Pettersson, &  
77 Svensson, 2016; Niemelä, Lattenkamp, & Dingemanse, 2015).

78         Understanding the evolution of personality traits and their associations in behavioural  
79 syndromes requires understanding the mechanisms underlying these between-individual  
80 differences in behaviour and their interactions, including their genetic basis. Phenotypic  
81 correlations between personality traits result from the addition of between-individual  
82 correlations (defining behavioural syndromes per se, whether genetic or nongenetic) and  
83 within-individual (or residual) correlations (Brommer, 2013; Dingemanse & Réale, 2013;  
84 Dingemanse, Kazem, Réale, & Wright, 2010). Assessing the relative contribution of between-  
85 and within-individual correlation components in observed phenotypic correlations can be  
86 crucial because between-individual correlations may constrain the independent evolution of  
87 the traits involved and thus may have major evolutionary consequences in the wild (Sih, Bell,  
88 & Johnson, 2004).

89         To reliably quantify this relative contribution of between- and within-individual  
90 correlations, multiple measurements of the personality traits considered must be collected on a  
91 large number of individuals. When only single measurements are available, it is often  
92 assumed that a phenotypic correlation observed between personality traits reflects an  
93 underlying between-individual correlation, i.e. the within-individual correlation is negligible  
94 (the so-called ‘individual gambit’, Brommer, 2013). Furthermore, to quantify genetic variance  
95 in personality traits and genetic covariance between them, the level of genetic relatedness  
96 between individuals measured has to be incorporated (e.g. via pedigree information) in the  
97 models. When relatedness information is unavailable, it is often assumed that the observed  
98 phenotypic (co)variance reflects the underlying genetic (co)variance (the so-called

99 'phenotypic gambit'; Grafen, 1984; Hadfield, Nutall, Osorio, & Owens, 2007; van Oers &  
100 Sinn, 2011).

101         These two crucial assumptions have been tested empirically in various species, and  
102 recent meta-analyses including over 30 studies, among which 25 are from wild populations,  
103 have confirmed their overall validity (Brommer & Class, 2017; Dochtermann, 2011;  
104 Dochtermann, Schwab, & Sih, 2015). Across these studies, the sign (and to a certain extent  
105 the magnitude) of the phenotypic correlations between personality traits reliably informed on  
106 the sign (and the magnitude) of the between-individual correlations (Brommer & Class, 2017)  
107 and of the genetic correlations (Dochtermann, 2011). Furthermore, 52% of the between-  
108 individual variation in personality traits taken separately was explained by additive genetic  
109 variance (Dochtermann et al., 2015). A recent empirical study on a wild population of yellow-  
110 bellied marmots, *Marmota flaviventris*, estimated the proportion of phenotypic (co)variance  
111 explained by genetic, permanent environment and maternal (co)variances between four  
112 different personality traits: docility, exploration, activity and sociability (Petelle, Martin, &  
113 Blumstein, 2015). Results showed additive genetic variations, as well as maternal and  
114 permanent environment variations, in all four traits and a positive genetic correlation between  
115 activity and sociability (Petelle et al., 2015). More of such integrative studies partitioning the  
116 observed phenotypic (co)variance in multiple personality traits simultaneously are needed in  
117 different biological models with contrasting life histories to better understand the mechanisms  
118 underlying and possibly constraining the evolution of correlated personality traits.

119         In this study, we assessed the genetic basis of three personality traits, together with the  
120 relative contribution of between- and within-individual variations in, and correlations  
121 between, these traits, chosen for their potentially important effects on crucial ecological  
122 processes (here, nest site acquisition and defence against competitors and predators) in a  
123 natural population of a small territorial, short-lived, migrant passerine bird, the collared

124 flycatcher, *Ficedula albicollis*. During 3 consecutive years, we measured for several hundreds  
125 of breeding pairs in the field (1) aggressiveness towards competitors (as the agonistic reaction  
126 to simulated territorial intrusions by intra- and interspecific competitors), (2) boldness  
127 towards predators (as the latency to resume nestling feeding after human disturbance) and (3)  
128 neophobia, possibly reflecting exploration (as the latency to resume nestling feeding in the  
129 presence of a novel object on the nest site; following Réale et al.'s 2007 definitions). To  
130 identify the mechanisms underlying the phenotypic (co)variation in these behavioural traits,  
131 we assessed to what extent (1) additive genetic, parental or permanent environment effects  
132 contributed to the observed phenotypic (co)variance, while accounting for fixed individual  
133 (sex, age) covariates, and (2) between-individual correlations explained phenotypic  
134 correlations between these three traits. Based on many previous studies on personality traits in  
135 populations of passerines of similar ecology (e.g. Dingemanse, Both, Drent, van Oers, & van  
136 Noordwijk, 2002; Drent, Oers, & Noordwijk, 2003; Duckworth & Badyaev, 2007;  
137 Garamszegi et al., 2015; Garamszegi, Rosivall, et al., 2012; van Oers, Drent, de Goede, & van  
138 Noordwijk, 2004), we expected heritable differences to partly explain variation in  
139 aggressiveness, boldness and neophobia in our study population. Furthermore, high  
140 aggressiveness, high boldness and low neophobia may allow individuals to efficiently secure  
141 and defend a breeding site when they are unfamiliar with the environment (e.g. for dispersers:  
142 Cote, Clobert, Brodin, Fogarty, & Sih, 2010; Duckworth & Kruuk, 2009; Korsten, van  
143 Overveld, Adriaensen, & Matthysen, 2013). Thus we predicted a functional integration and  
144 (possibly genetically based) between-individual correlations between these traits. In another  
145 population of collared flycatchers, male aggressiveness and boldness, but not neophobia, were  
146 found to be phenotypically correlated in some years (Garamszegi et al., 2015; Garamszegi,  
147 Eens, & Török, 2009). This population and ours, however, differ greatly in both demographic  
148 functioning (e.g. male age structure: Hegyi, Rosivall, & Török, 2006) and selective pressures

149 (e.g. nest predation: Doligez & Clobert, 2003; sexual selection: Qvarnström, 1997; see also  
150 Rosivall, Török, Hasselquist, & Bensch, 2004), which may affect the fitness consequences of  
151 interactions between personality traits, and thus their potential functional integration.

152

153

## 154 **METHODS**

155

### 156 *Study species and population monitoring*

157 Collared flycatchers are migratory cavity nesters and readily breed in artificial nestboxes,  
158 providing easy access to parents' identity and breeding data. Between 2011 and 2013, we  
159 conducted the behavioural tests (see below) on 1131 pairs breeding in nestboxes spread over  
160 14 to 22 forest patches in our study population located on the island of Gotland (Sweden,  
161 Baltic Sea). Each year since 1980, nests in boxes have been monitored at least weekly from  
162 late April until early July, allowing us to record major breeding variables (laying and hatching  
163 dates; clutch size; nestling number, condition and fledging success). Breeding pairs were  
164 captured, identified and ringed if previously unringed; females were caught during incubation  
165 and males while feeding nestlings. Nestlings were ringed between day 8 and day 13 after  
166 hatching; fledging typically occurs 16 days after hatching. Adult and nestling identification  
167 every year combined with a high return rate of both adults (approximately 40%) and juveniles  
168 (approximately 10%) for such a short-lived passerine bird (Gustafsson, 1989) allowed us to  
169 establish a high-quality social pedigree of the population which has previously been used in  
170 several quantitative genetic studies (e.g. Evans & Gustafsson, 2017; Merilä & Sheldon, 2000;  
171 Sheldon, Kruuk, & Merilä, 2003; Appendix Table A1). In this population, approximately 15%  
172 of all nestlings are extrapair (Sheldon & Ellegren, 1999), a percentage considered low enough  
173 for quantitative genetic models to provide valid (i.e. only slightly underestimated) heritability



174 estimates from the social pedigree (Charmantier & Réale, 2005; Firth, Hadfield, Santure,  
175 Slate, & Sheldon, 2015), even though no information is available yet on how extrapair  
176 paternity may affect genetic covariances. The clear sexual dimorphism in plumage coloration  
177 in this species allowed an easy discrimination of adult males (black and white plumage with a  
178 white forehead patch) from females (brown plumage; Svensson, 1992), even from several  
179 metres away during behavioural tests. Adults could also be aged by plumage criteria  
180 (yearlings versus older adults; Svensson, 1992).

181

### 182 *Aggressiveness score*

183 We measured the level of aggressiveness of breeding flycatchers soon after settlement, during  
184 nest building or early laying, that is, when the risk of losing a nestbox to a competitor is  
185 highest in this single-clutch species. During the breeding season, collared flycatchers compete  
186 for nest sites with conspecifics but also with great tits, *Parus major*, the second most abundant  
187 species breeding in nestboxes in the study area (Gustafsson, 1987). Aggressiveness towards  
188 conspecific intruders was shown to decrease after the start of incubation (Král & Bícík, 1989)  
189 even though aggressiveness towards great tit intruders remained high throughout the breeding  
190 cycle (Král & Bícík, 1992). To elicit an aggressive response from a focal flycatcher pair, we  
191 simulated the intrusion of competitors at the nest of the pair by attaching to its nestbox clay  
192 decoys mimicking either a flycatcher pair or a single (male) great tit. We used a pair (one  
193 male and one female) for flycatcher decoys to elicit and measure an aggressive response by  
194 both pair members, that is, to avoid a sex-specific response towards this intraspecific  
195 stimulus, while the response to the interspecific stimulus (great tit decoy) was not expected to  
196 differ depending on the sex of the decoy. In addition, we simultaneously broadcast male songs  
197 corresponding to the decoy(s) species with a loudspeaker placed just under the nestbox. To  
198 avoid pseudoreplication, we randomly used one of eight different sets of decoys and one of

199 five different song tracks per species for each test. After attaching the decoys to the nestbox  
200 and the loudspeaker under the box, the observer sat under a camouflage net approximately 8–  
201 10 m away from the nestbox and recorded the following behaviours for each pair member: (1)  
202 movements between perches and perching position (within 2 m, between 2 and 5 m or  
203 between 5 and 10 m away from the nestbox), (2) agonistic behaviours towards a decoy  
204 (attacks and stationary flights in front of the decoy) and (3) chases towards living birds  
205 attracted by the stimulus. A behavioural test started with an observation period of 15 min but  
206 we lengthened the test by up to 5 min when an individual arrived between 10 and 15 min after  
207 the start of the test, and up to 5 additional min if its partner arrived during this extra time, so  
208 that we could observe the behavioural response of each pair member for at least 5 min. Each  
209 test thus lasted between 15 and 25 min. If an individual was observed during less than 5 min  
210 before the end of the test, it was discarded from the analyses and these observations were  
211 therefore not used later on.

212         Aggressiveness level was measured as the sum of the number of movements within 2  
213 m of the nestbox, attacks, stationary flights and chases. We included this latter behaviour  
214 because chasing a live intruder may have prevented the focal flycatcher from interacting with  
215 the decoy, while reflecting an aggressive territory defence response. The number of each type  
216 of behavioural response (movements, agonistic behaviours and chases) was standardized by  
217 the time interval between the first observation of the individual and the end of the test,  
218 rescaled to 15 min. Using alternative scores did not qualitatively change the results (see  
219 Appendix and Table A2). We conducted aggressiveness tests two to four times per focal pair  
220 over a 5-day interval, with at most one test per day and tests on 2 days in a row. The stimulus  
221 used (intra- / interspecific decoys) was alternated between tests after a random assignment for  
222 the first test. An aggressiveness score was computed for each individual for each test. We  
223 obtained aggressiveness responses (for more than 5 min at least once per year) for 1974

224 individuals (including unidentified ones; 961 females and 1014 males in 1046 nests). Among  
225 those, 825 females and 667 males were later captured and identified, and thus used for  
226 heritability estimation. In 601 breeding pairs both partners were identified and responded to  
227 the tests and in 273 only one partner responded. We obtained repeated estimates for 502 and  
228 445 identified females and males, respectively.

229

### 230 *Boldness and neophobia scores*

231 During nestling rearing, we estimated (1) boldness level by measuring the individual's  
232 reaction towards the presence of a human observer near the nestbox and (2) neophobia level  
233 by measuring the reaction towards the presence of a novel object on the nestbox (i.e. in a  
234 familiar environment), following the definitions from Réale et al. (2007). As advised in  
235 Greenberg and Mettke-Hofmann (2001), we measured our behavioural responses as the  
236 latency to perform a highly motivating action (here, feeding their nestlings) after disturbance.  
237 We conducted one combined boldness/neophobia test per breeding pair when the nestlings  
238 were 5 days old, that is, at the beginning of the period of highest provisioning by parents (and  
239 before we caught the parents to avoid any behavioural interference). A test consisted of two  
240 consecutive periods of approximately 1 h each: the behaviour of the parents was recorded first  
241 without any change in the surroundings of the nestbox, that is, without the novel object, and  
242 second with a novel object (here a coloured figurine approximately 7 cm high) attached near  
243 the entrance hole of the nestbox. Both periods were video-recorded from a distance (6–8 m).  
244 At the beginning of each period, the observer checked the camouflaged video recorder,  
245 walked to the nestbox, opened it to check nestling satiety, closed it, and then left the area. The  
246 test was abandoned if the nestlings were very hungry to avoid them starving if the parents  
247 were too disturbed by the test.

248 We estimated boldness score using the latency to enter the nestbox after the departure  
249 of the observer in the first period (i.e. without the novel object). Reaction to disturbance by  
250 humans has previously been used in boldness tests in this species (e.g. Garamszegi et al.,  
251 2009). To ease interpretation (i.e. increasing values of boldness score corresponding to  
252 increasing level of boldness), we transformed the latency to enter the nestbox such that the  
253 boldness score of an individual was the maximum latency observed in the entire data set  
254 minus the latency for this individual. We estimated neophobia score based on the latency to  
255 enter the box after the departure of the observer in the second period (i.e. in the presence of  
256 the novel object). For both boldness and neophobia scores, individuals that did not enter the  
257 nestbox during the first period of the test were not used in the analysis (187 of 1251  
258 observations, i.e. 15%). Individuals that entered the nestbox during the first but not the second  
259 part of the test (411 of 1064 observations, i.e. 39%) were considered as the most neophobic  
260 ones but could not be assigned a latency. To include them in the analyses, we discretized the  
261 latency to enter the nestbox in the second period into four categories based on its quartiles,  
262 assigning values from 1 to 4 for increasing latencies, and adding a fifth category including  
263 individuals that did not enter in the second part of the test. Using alternative scores for  
264 boldness and neophobia did not qualitatively change the results (see Appendix and Tables A2  
265 and A3). We obtained boldness and neophobia estimates for 849 identified individuals (472  
266 females and 378 males). Over the 3 years, we assessed 403 unique breeding pairs where both  
267 identified partners responded to the tests and 185 pairs where only one partner responded. We  
268 obtained repeated boldness and neophobia scores (i.e. several years in a row) for 66 females  
269 and 65 males.

270

271 ***Repeatability and heritability of aggressiveness, boldness and neophobia***

272 We estimated the repeatability of aggressiveness, boldness and neophobia scores as well as  
273 their heritability by fitting three separate univariate animal models. The models included the  
274 following random effects: additive genetic effect (associated with the pedigree), individual  
275 identity for the repeated measures per individual (permanent environment effect), maternal  
276 and paternal identities, forest patch, observer identity (the person observing and reporting the  
277 behaviours onsite for the aggressiveness tests and the person extracting latencies from the  
278 video recording for the boldness and neophobia tests). The models of the aggressiveness score  
279 also included the broadcast song track and decoy set identifiers. In addition, the models  
280 included the following fixed effects, to control for potential confounding factors: sex, age  
281 (two levels: yearling versus older) and their interaction, as well as year (three levels: 2011,  
282 2012, 2013). The aggressiveness model also included stimulus type (two levels: flycatcher  
283 versus great tit decoys), the order of the test within a year (continuous variable: first to  
284 fourth), the presence of the partner during the test (binary variable: yes/no) and the presence  
285 of other live flycatchers or great tits (binary variable: yes/no). The boldness and neophobia  
286 models included the number of ringed nestlings as a proxy of the motivation to enter the  
287 nestbox to feed nestlings. The neophobia models included the boldness score to control for the  
288 effect of the human disturbance at the beginning of the period with the novel object. All  
289 continuous fixed terms were centred and standardized prior to analysis to allow comparisons  
290 between effects.

291 Repeatabilities ( $R$ ) were estimated as the ratio of the sum of the additive genetic ( $V_A$ ),  
292 permanent environment ( $V_{PE}$ ), maternal and paternal identities variances ( $V_M$  and  $V_F$ ,  
293 respectively) over the total phenotypic variance ( $V_P$ , sum of all variances; Falconer &  
294 Mackay, 1996; with possibly an additional term accounting for the distribution variance,  
295 Nakagawa & Schielzeth, 2010). Narrow-sense heritabilities ( $h^2$ ) were estimated as the ratio of  
296 the additive variance  $V_A$  over the phenotypic variance  $V_P$ . The presence of fixed effects in

297 models did not result in over- or underestimating repeatability and heritability estimates (as  
 298 warned against in Nakagawa & Schielzeth, 2010), because estimates were similar when  
 299 obtained from models with only the intercept (presented in the main text) and from models  
 300 with the previously described fixed effects (see Appendix Tables A1, A2, A3; except for a  
 301 slight overestimation of the boldness and neophobia repeatabilities on the latent scale). For  
 302 aggressiveness, we also estimated within-year repeatability by replacing the additive genetic  
 303 and permanent environment effects by a unique identifier per individual per breeding season.

304

305 *Correlations between aggressiveness, boldness and neophobia scores*

306 We estimated the between- and within-individual correlations between aggressiveness,  
 307 boldness and neophobia scores by fitting a trivariate mixed-effects model (Dingemanse &  
 308 Dochtermann, 2013). For this model, aggressiveness was averaged over all aggressiveness  
 309 scores of an individual in a given year (i.e. over up to four estimates). We chose this  
 310 averaging approach because aggressiveness score (1) differed depending on decoy species,  
 311 chosen at random for the first test, and (2) decreased with the order of the test (i.e. due to  
 312 habituation). Averaging all aggressiveness scores of an individual in a given year allowed us  
 313 to control for these differences and decrease the effect of varying environmental  
 314 (meteorological) conditions between tests. In this trivariate model, we included sex and year  
 315 as fixed effects and ring number as a random effect. The phenotypic correlation between two  
 316 traits  $A$  and  $B$ ,  $r_{P_A, P_B}$ , and the between-individual correlation between the traits  $A$  and  $B$ ,  
 317  $r_{ind_A, ind_B}$ , were estimated as follows (Snijders & Bosker 1999):

318 
$$r_{P_A, P_B} = \frac{COV_{ind_A, ind_B} + COV_{\epsilon_A, \epsilon_B}}{\sqrt{V_{P_A} \times V_{P_B}}} \quad (1)$$

319 
$$r_{ind_A, ind_B} = \frac{COV_{ind_A, ind_B}}{\sqrt{V_{ind_A} \times V_{ind_B}}} \quad (eq. 2)$$

320 where  $\text{COV}_{\text{ind}_A, \text{ind}_B}$  and  $\text{COV}_{\varepsilon_A, \varepsilon_B}$  are the between- and the within-individual covariances  
321 between traits  $A$  and  $B$ , and  $V_{P, A \text{ or } B}$  the total phenotypic variance (sum of the between- and  
322 within-individual variances) associated with trait  $A$  or  $B$ . Combining aggressiveness, boldness  
323 and neophobia score data for a given individual in a given year, we obtained 841 observations  
324 for which estimates for all three traits were available, and 152 observations for which only  
325 boldness and neophobia estimates were available. Among these 841 observations, 49 females  
326 and 52 males were repeatedly assessed over several years for all three traits, providing 100  
327 and 111 repeated observations, respectively.

328 To estimate the additive genetic correlations between traits, we fitted a model with the  
329 same fixed effects but with the additive genetic effect instead of the individual ring as a  
330 random factor. Fitting both additive genetic and individual (permanent environment) effects  
331 together in a single model indeed led to convergence failures. We thus fitted only one effect at  
332 a time.

333

### 334 *Implementation of Bayesian models*

335 All statistical analyses were performed within the Bayesian framework in R v.3.3.2 (R Core  
336 Team, 2016). Both univariate and trivariate models were fitted using the function  
337 MCMCglmm ('MCMCglmm' R package, Hadfield, 2010). The pedigree was prepared using  
338 the function fixPedigree ('pedantics' R package, Morrissey & Wilson, 2010) and pruned  
339 using the function prunePed ('nativ' R package, Wolak, 2012; see Appendix Table A1 for a  
340 description of the pedigree). Aggressiveness and boldness scores were fitted with a Poisson  
341 family (logit link), and neophobia scores with a Threshold family with the residual variance  
342 fixed to 10 (instead of the usual value of 1, to improve the mixing of the chains for low  
343 variances, which were expected from preliminary analyses; Hadfield, 2016). We used wide  
344 normally distributed priors for fixed effects (large variance  $V=10^8$ ; Hadfield, 2016) and

345 parameter expanded  $\chi^2$  distributed priors with 1 degree of freedom for random effects. For the  
346 univariate models, we adjusted the number of iterations, burn-in and thinning interval for each  
347 model so as to obtain an effective sample size over 1500 (see Appendix) and autocorrelations  
348 of posterior samples below 0.1 in all cases. For the trivariate models, we used  $4 \times 10^6$   
349 iterations, a burn-in of  $10^5$  and a thinning interval of 2000 to reach the same criteria. We  
350 visually assessed the convergence of each MCMC chain and compared three chains per model  
351 using the Gelman & Rubin approach (`gelman.diag` and `gelman.plot` functions, ‘coda’ R  
352 package, Plummer et al., 2016). Following Nakagawa and Schielzeth (2010), we retained in  
353 our univariate models data from individuals tested only once. For all three traits, estimates are  
354 presented as posterior modes with the associated 95% credible intervals (CI). Variance,  
355 repeatability and heritability estimates are presented on the latent scale ( $R_{\text{latent}}$ , following  
356 Nakagawa & Schielzeth, 2010;  $h^2_{\text{latent}}$  following de Villemereuil, Schielzeth, Nakagawa, &  
357 Morrissey, 2016). We also provide repeatability and heritability estimates on the observed  
358 scale ( $R_{\text{obs}}$  using the `QGicc` function,  $h^2_{\text{obs}}$  using the `QGparams` function, from the ‘`QGglmm`’  
359 R package; de Villemereuil, 2018; de Villemereuil, Schielzeth, Nakagawa, & Morrissey,  
360 2015). It is not possible here to discuss all results on the observed scale, even though it is the  
361 scale of the realized behaviour, that is, where natural selection can act (see de Villemereuil et  
362 al., 2016). Indeed, the estimation of  $R_{\text{obs}}$  for ordinal traits (here the neophobia score) is  
363 complex and is not currently implemented in `QGicc`. To allow us to compare estimates  
364 between traits on the observed scale, we computed repeatabilities for neophobia using two  
365 alternative scores with nonordinal distributions, namely (1) the latency to enter the nestbox in  
366 the presence of the novel object for individuals that entered the nestbox ( $N=653$ , fitted with a  
367 Poisson distribution), and (2) whether the individual entered the nestbox in the presence of the  
368 novel object ( $N=1064$ , binary variable fitted with the threshold family). Moreover, the  
369 estimation of  $h^2_{\text{obs}}$  for ordinal traits provides one heritability estimate per level, which in the



370 case of an artificial categorization as here is not biologically relevant (de Villemereuil, 2018).  
371 As the heritability estimates on the observed scale were fairly similar between neophobia  
372 levels, we compared the range of heritability estimates found for neophobia with the  
373 heritability estimates found on the observed scale for aggressiveness and boldness.  
374 Correlations are provided on the latent scale.

375

### 376 *Ethical note*

377 Permission for catching and ringing adults (here 838 yearlings, 1074 older birds) and  
378 nestlings (here 9750) with individually numbered aluminium rings was granted every year by  
379 the Ringing Centre from the Museum of Natural History in Stockholm (licence nb. 471: M015  
380 to B.D.). Adults were caught in the nest, either directly (females during incubation) or using  
381 swinging-door traps (both parents during nestling rearing). Traps were set for at most 30 - 60  
382 min depending on nestling age (30 min when nestlings were 5 days old or younger), to avoid  
383 nestling starvation if parents did not resume feeding during the catching period; traps were  
384 checked every 5–10 min, and removed as soon as the adults had been caught. Catching  
385 sessions started after 0600 hours to let birds feed and provision nestlings undisturbed for at  
386 least 2 h after the night period (sunrise occurs at approximately 0400 hours during spring).  
387 Adults were handled for 5 - 10 min and released straight after manipulation or (when catching  
388 both parents during nestling feeding) kept until the partner was captured (up to 40 min  
389 maximum). For nestling ringing, whole broods were taken directly from the nest and ringed  
390 just beside the nestbox (for approximately 10 min); nestlings that were not handled were kept  
391 warm using small heating packs. During the aggressiveness tests, we minimized disturbance  
392 by approaching the nestbox as quietly as possible and hiding below a camouflage net.  
393 Conversely, for the combined boldness/neophobia test, which aimed at measuring (or  
394 controlling for) the reaction towards human presence, we on purpose approached the nestbox

395 conspicuously. During the boldness/neophobia test, nestling satiety was checked at the  
396 beginning and in between the two parts of the test, and the test was aborted if nestlings were  
397 begging too strongly, to avoid any harmful effect of temporarily decreased provisioning by  
398 parents. All the manipulations were performed in accordance with the Swedish legislation  
399 applying at the time.

400

401

## 402 **RESULTS**

403

### 404 *Univariate models*

405 The level of repeatability for aggressiveness was 0.18 on the latent scale and 0.03 on the  
406 observed scale (Table 1). In addition, aggressiveness score was repeatable within years  
407 ( $R_{\text{latent}}=0.22$ , 95% CI = [0.18; 0.26];  $R_{\text{obs}}=0.04$ , 95% CI = [0.03; 0.06]) and between years  
408 (when averaging the aggressiveness score of 1 year;  $R_{\text{latent}}=0.26$ , 95% CI = [0.11; 0.38];  
409  $R_{\text{obs}}=0.11$ , 95% CI = [0.04; 0.20]). The level of repeatability for boldness was 0.11 on the  
410 latent scale and 0.10 on the observed scale (Table 1). Neophobia was slightly more repeatable  
411 than the other traits on the latent scale (Table 1) as well as on the observed scale when  
412 estimated from alternative nonordinal measures ( $R_{\text{latent}}=0.25$ , 95% CI = [0.06; 0.35] and  
413  $R_{\text{obs}}=0.14$ , 95%CI = [0.04; 0.23] for the latency to enter the nestbox in the presence of the  
414 novel object;  $R_{\text{latent}}=0.44$ , 95% CI = [0.22; 0.67] and  $R_{\text{obs}}=0.29$ , 95%CI = [0.12; 0.44] for  
415 whether the individual entered the nestbox during the test or not). None of the three  
416 behavioural scores, however, were heritable (all 95% CI for  $V_A$  and  $h^2$  values included zero;  
417 Table 1). Permanent environment effects explained 15% of the phenotypic variance in  
418 aggressiveness score (95% CI of  $V_{\text{PE}} = [0.28; 0.80]$ ; Appendix Table A2) and parental  
419 identities (i.e. maternal and paternal identities summed) explained 25% of the phenotypic

420 variance in neophobia score (95% CI of  $V_M + V_F = [0.38; 8.46]$ , even though the lower limit of  
421 the 95% CI for each parent identity separately was 0:  $[0.00; 6.49]$  for  $V_M$  and  $[0.00; 5.75]$  for  
422  $V_F$ ). When excluding the maternal (paternal) identity from the model, the paternal (maternal)  
423 identity explained 19% (18%) of the phenotypic variance. Fitting the neophobia model  
424 without the maternal and paternal identities did not change the heritability estimate, revealing  
425 that these effects were not confounded with the additive genetic effect (results not detailed).  
426 Observer identity explained 7% of the phenotypic variance for aggressiveness (95% CI of  
427  $V_{\text{Observer}} = [0.14; 0.55]$ ); paternal identity explained 11% of the phenotypic variance for  
428 aggressiveness but only when measuring aggressiveness as the first axis of a principal  
429 component analysis (see Appendix). All other variances were low (less than 4 % of the  
430 phenotypic variance) or not different from zero (Appendix Tables A1, A2 and A3).

431 Males were more aggressive than females, especially among yearlings (interaction  
432 sex\*age, with yearling males as reference: 95% CI =  $[0.44, 0.81]$ ; Appendix Fig. A1a). In  
433 addition, males were slightly shyer (longer latency to enter in the absence of a novel object)  
434 and less neophobic (shorter latency in the presence of a novel object) than females (with  
435 female as a reference 95% CI =  $[-0.12; -0.02]$  and  $[-2.78; -1.07]$  respectively; Appendix Fig.  
436 A1b), and this did not depend on age (see Appendix Tables A2 and A3 for sex\*age  
437 interactions). In addition, individuals with larger broods were bolder (95% CI =  $[0.002; 0.06]$ ;  
438 Table A3) and less neophobic (95% CI =  $[-1.03; -0.37]$ ; Table A4). Regarding environmental  
439 effects, aggressiveness, boldness and neophobia scores depended on the year: individuals  
440 were less aggressive and less neophobic in 2011 than 2012 (Appendix Tables A2 and A4),  
441 and shyer in 2011 than 2013 (Appendix Table A3). Finally, individuals were more aggressive  
442 in the presence of their partner or neighbouring tits attracted by the stimulus, and during the  
443 first tests of the sequence (Appendix Table A2).

444

445 *Trivariate model*

446 Aggressiveness and neophobia scores were phenotypically correlated: more aggressive  
447 individuals were less neophobic (Table 2, Appendix Fig. A2a). Boldness and neophobia  
448 scores were also phenotypically correlated: bolder individuals were less neophobic (Table 2,  
449 Appendix Fig. A2b). The corresponding within-individual correlations were negative (Table  
450 2). However, there was no phenotypic correlation between boldness and aggressiveness  
451 (Table 2; see Appendix Table A5 for the full model output) and none of the between-  
452 individual correlations differed from zero (Table 2). When accounting for an additive genetic  
453 effect, none of the additive genetic covariances differed from zero (Table 2; see Table A6 for  
454 the full model output).

455

456

457 **DISCUSSION**

458

459 In this study, we tested whether three personality traits (aggressiveness, boldness and  
460 neophobia) had a genetic basis in a wild population of collared flycatchers and formed  
461 (genetically based) behavioural syndromes during breeding, to shed light on constraints in  
462 their possible evolution. None of the three traits were heritable and their repeatability  
463 estimates were low (0.11–0.39 on the latent scale for all traits; 0.03–0.10 on the observed  
464 scale for aggressiveness and boldness; 0.14 and 0.29 on the observed scale for nonordinal  
465 measures of neophobia) compared to average estimates previously found for behavioural traits  
466 in two meta-analyses (0.37 in Bell et al., 2009; 0.41 in Holtmann et al., 2017), suggesting  
467 strong phenotypic plasticity in these traits. The repeatability originated mainly from  
468 permanent environment effects for aggressiveness and from parental effects for neophobia. In  
469 addition, we found that the three traits showed phenotypic covariance but no between-

470 individual covariance and no additive genetic covariance. The absence of behavioural  
471 syndromes among these personality traits may be due either to a lack of statistical power to  
472 detect between-individual covariances or to an absence of functional integration of these traits  
473 at the individual level and no genetic correlation at the population level.

474

#### 475 *Factors at the origin of behavioural trait repeatability*

476 Our levels of repeatability, estimated both within and between years for aggressiveness score  
477 and between years for boldness and neophobia scores, were lower than usually reported for  
478 such behaviours: around 0.50 for aggressiveness and exploration, and around 0.40 for  
479 antipredator behaviours (Bell et al., 2009). Interestingly, the repeatability level estimated here  
480 for aggressiveness score was similar within and between years, contrary to the usual decrease  
481 observed when the time interval between recordings increases (Bell et al., 2009; Chervet,  
482 Zöttl, Schürch, Taborsky, & Heg, 2011; Dingemanse et al., 2012; Garamszegi et al., 2015;  
483 Holtmann et al., 2017; Wuerz & Krüger, 2015; but see David, Auclair, & Cézilly, 2012 for  
484 differences between traits). Overall, our lower levels of repeatability, especially on the  
485 observed scale, suggest higher plasticity, both within and between years compared to studies  
486 on other species.

487         The observed repeatability in aggressiveness score resulted mostly from permanent  
488 environment effects, which explained 15% of the phenotypic variance and 72% of the  
489 repeatability in aggressiveness score. Here, because we controlled for the identity of the  
490 parents, permanent environment effects could be linked to differences in individual condition  
491 or experience. Some measures of personality traits have indeed been found to depend on  
492 individual condition or experience (reviewed in Stamps & Groothuis, 2010). In our  
493 population, condition and experience also affect breeding habitat choice depending on social  
494 cues (e.g. Doligez, Danchin, Clobert, & Gustafsson, 1999; Doligez, Pärt, Danchin, Clobert, &

495 Gustafsson, 2004; Kivelä et al., 2014), which could shape individuals' response to the risk of  
496 competition for nest sites. Permanent environment effects may also include a dominance  
497 effect (Kruuk & Hadfield, 2007; Wilson et al., 2010), which could not be directly modelled  
498 here because full- and half-sib links were too rare in our pruned pedigree for running such  
499 complex models (Wilson et al., 2010).

500 In turn, the observed repeatability in neophobia resulted mostly from parental  
501 identities, which accounted for 25% of the phenotypic variance and 55% of the between-  
502 individual variance. Both pre- and postnatal parental effects have been found to affect  
503 exploration and neophobia behavioural responses later in life (e.g. nestling provisioning and  
504 exploration in birds, Carere, Drent, Koolhaas, & Groothuis, 2005; maternal hormones early in  
505 life and neophobia, Spencer & Verhulst, 2007; see the review in Groothuis & Maestriperi,  
506 2013). However, parental identities did not explain between-individual differences in  
507 aggressiveness and boldness here, contrary to previous findings (e.g. Eising, Muller, &  
508 Groothuis, 2006; reviewed in Groothuis & Maestriperi, 2013). These behavioural responses  
509 may be more dependent on individual or local environmental conditions, in particular  
510 individual competitive ability and neighbour/predator presence or density, at the time of the  
511 test(s). To better understand how parental effects shape behavioural responses later in life in  
512 our study population, further experiments (e.g. nestling cross-fostering) would be necessary.

513

#### 514 *Personality traits with no genetic basis*

515 We found no genetic basis for our three personality traits. A meta-analysis on personality  
516 traits in wild animal populations estimated an average heritability level of 0.28 for  
517 aggressiveness, 0.31 for boldness and 0.58 for exploration-avoidance (including estimates  
518 from novel environment and novel object tests; van Oers & Sinn, 2013). The absence of  
519 heritability for our personality traits here was not due to a lack of statistical power to detect

520 significant additive genetic variance based on our sample and social pedigree, because based  
521 on the same sample with the same pedigree, we obtained positive heritability estimates for  
522 tarsus and wing length ( $h^2=0.59$ , 95% CI = [0.44; 0.69] for tarsus and  $h^2=0.30$ , 95% CI =  
523 [0.11; 0.50] for wing length, while accounting for maternal and permanent environment  
524 effects) which are consistent with previous estimates in this population ( $h^2 = 0.53$  and  $0.51$  for  
525 tarsus and wing length, respectively, in Merilä & Gustafsson, 1993). The absence of  
526 heritability in our personality traits was therefore likely to be the result of very low additive  
527 genetic variance combined with large environmental variance as illustrated, for instance, by  
528 between-year differences in behavioural scores, which reflected large variations in  
529 environmental conditions between the 3 years of our study (see Morinay, Forsman, Kivelä,  
530 Gustafsson, & Doligez, 2018 for differences between 2012 and 2013). Large environmental  
531 variance could originate from individuals being tested in different environments (including  
532 the social context) in different years, because between-year fidelity to the nestbox and/or  
533 partner is very low in this population (approximately 6.7% of 240 individuals bred in the  
534 same nestbox several years and 1.0% of 214 identified pairs were faithful over several years).  
535 This, however, limited the risk of pseudoreplication (Niemelä & Dingemans, 2017).

536         Because we measured personality traits at the nest during breeding, the reaction of the  
537 partner may have affected the reaction of the focal bird during a behavioural test, as found  
538 here with a higher aggressiveness score when the partner was present. To account for this  
539 effect, we could have included the partner's identity and genetic background (i.e. pedigree) as  
540 random effects in our models (see Morinay et al., 2018 for an example in the same  
541 population; and Wolf, Brodie III, Cheverud, Moore, & Wade, 1998 for so-called indirect  
542 genetic effects). However, this could have led to pseudoreplication, because the behavioural  
543 score of the partner itself was most of the time also analysed in this dataset (e.g. around 74%  
544 of the females and 90% of the males had their partner tested). To keep exploring a response at

545 the individual level (rather than combining behavioural scores at the pair level), a solution  
546 could be to fit a bivariate model of the two partners' responses and include both their  
547 pedigree, permanent environment and parental effects in the model. This would, however,  
548 require a larger data set than used here to reach sufficient statistical power to detect such  
549 effects with such complex models. Furthermore, because the focal bird chooses at least partly  
550 its partner (like its nest site), partner's effects can also be expected to be at least partly  
551 included in the individual's genetic and permanent environment effects. Disentangling such  
552 complex effects may require a more balanced sample of faithful and divorced pairs breeding  
553 in the same and different sites over several years than observed in our population.

554

#### 555 *No personality syndrome?*

556 Phenotypic correlations were observed between our personality traits, even though they did  
557 not constitute behavioural syndromes (i.e. there was no between-individual correlations): less  
558 neophobic individuals were more aggressive and bolder. This was in line with previous  
559 studies reporting bolder individuals to be more explorative in a novel environment (or less  
560 neophobic in a novel object test; e.g. Garamszegi et al., 2009; van Oers, De Jong, Drent, &  
561 van Noordwijk, 2004). Conversely, the absence of correlation between aggressiveness and  
562 boldness partly contrasts with previous results reporting more aggressive individuals to be  
563 bolder, as part of the proactive–reactive axis, in different species (Koolhaas et al., 1999; Sih,  
564 Bell, Johnson, et al., 2004) including the collared flycatcher (Garamszegi et al., 2015).

565 The observed phenotypic correlations resulted solely from correlated changes in  
566 behaviours between measurements for the same individuals, that is, within-individual  
567 correlations. Within-individual correlations could be due to micro-environmental effects (e.g.  
568 nestbox environment), to individual effects (e.g. long-term between-year plasticity but short-  
569 term within-year behavioural constraints, for instance due to experience) or to correlated



570 measurement errors (Dingemanse & Dochtermann, 2013). Error correlation, however, is more  
571 likely to occur between boldness and neophobia scores, which were extracted from the same  
572 test and might both be correlated with feeding rate (with birds investing more in nestling  
573 provisioning returning more rapidly to their nest in both situations), than between  
574 aggressiveness and neophobia scores, which were measured several weeks apart by different  
575 persons in different tests. To tease these sources of within-individual correlations apart,  
576 aggressiveness, boldness and neophobia scores need to be estimated several times during the  
577 same breeding season and possibly the same phase(s) of the reproductive cycle. The limited  
578 number of individuals measured several times here (211 observations of 101 individuals),  
579 however, is likely to be why we did not detect between-individual covariance (Dingemanse &  
580 Dochtermann, 2013 recommended sample sizes of at least 200 individuals tested twice; see  
581 also Garamszegi & Herczeg, 2012). Indeed, based on the same sample, we were not able to  
582 obtain positive genetic or between-individual covariances between tarsus and wing length,  
583 two morphological traits previously reported as genetically correlated in the same population  
584 (Merilä & Gustafsson, 1993).

585         Even though our limited statistical power does not allow us to conclude the absence of  
586 behavioural syndromes, this absence, if true, would suggest that selective pressures did not  
587 yield or maintain a functional integration between the personality traits investigated here. A  
588 true absence of behavioural syndromes among the traits we studied could be explained by  
589 specific breeding conditions in our population, possibly altering the selective regimes  
590 compared to other populations or species. In our population, the high availability of high-  
591 quality nest sites (i.e. nestboxes, provided in excess since the early 1980s) may have released  
592 joint selective pressures on exploration to find suitable nest sites and aggressiveness to  
593 acquire and defend this resource against dominant competitors (in particular tit species) in a  
594 natural context. In turn, providing nestboxes probably increased local breeding densities and

595 thereby competition for food resources during the nestling period, especially in a highly  
596 synchronous species such as the collared flycatcher. Furthermore, our population is subjected  
597 to very low nest predation rates, due to the absence of mustelid species on Gotland (Doligez  
598 & Clobert, 2003), which may have released selective pressures on boldness through the  
599 decrease in the need to defend the brood. Overall, these specific breeding conditions may  
600 have strongly modified the selective regime for personality traits and for a functional  
601 integration between them if they are costly.

602         In conclusion, we showed that aggressiveness, boldness and neophobia are repeatable  
603 but not heritable traits and do not seem to form behavioural syndromes in our population of  
604 collared flycatchers since only phenotypic correlations were observed between neophobia and  
605 the other two traits. Our study thus brings insights on the evolutionary potential of these  
606 personality traits alone and in interaction with each other during breeding in a wild population  
607 experiencing particular breeding conditions (low competition for nest sites, low nest predation  
608 rate). To understand the absence of individual covariance between, and heritable variations in,  
609 personality traits in our population, a first step would be to investigate the fitness benefits (i.e.  
610 reproductive success and survival) associated with each trait and their interactions. Plasticity  
611 in the associations between personality traits should be selected for if the fitness costs and  
612 benefits of expressing each trait relative to the others depend on the environmental (including  
613 social) context (e.g. competition level or predation risk), which remains to be explored for  
614 example by experimentally manipulating these environmental conditions.

615

#### 616 **Conflict of interest**

617 We declare no conflict of interest.

618

#### 619 **Acknowledgments**

620 We are grateful to the landowners from Gotland for allowing us to collect data on their  
621 properties, to all fieldworkers who collected the data between 2011 and 2013, and to all the  
622 students who contributed to the extraction of data from the audio and video recordings and to  
623 building the behavioural database. We also thank Ingrid Ahnesjö and Emmanuel Desouhant,  
624 and three anonymous referees for their critical comments on the manuscript. J.M. was funded  
625 by the Ministère de l'Enseignement Supérieur et de la Recherche (PhD grant), by Uppsala  
626 University (PhD grant and Stiftelsen research grant) and by the University of Lyon (IDEX  
627 mobility grant). G.D. was funded by the Région Auvergne Rhône-Alpes (PhD and  
628 Explora'Doc grants). L.G. was funded by the Swedish Research Council and B.D. by the  
629 Centre National pour la Recherche Scientifique.

630

## 631 **References**

- 632 Bell, A. M. (2007). Future directions in behavioural syndromes research. *Proceedings of the*  
633 *Royal Society B: Biological Sciences*, 274(1611), 755–761.  
634 <http://doi.org/10.1098/rspb.2006.0199>
- 635 Bell, A. M., Hankison, S. J., & Laskowski, K. L. (2009). The repeatability of behaviour: a  
636 meta-analysis. *Animal Behaviour*, 77(4), 771–783.  
637 <http://doi.org/10.1016/j.anbehav.2008.12.022>
- 638 Brommer, J. E. (2013). On between-individual and residual (co)variances in the study of  
639 animal personality: Are you willing to take the 'individual gambit'? *Behavioral Ecology*  
640 *and Sociobiology*, 67(6), 1027–1032. <http://doi.org/10.1007/s00265-013-1527-4>
- 641 Brommer, J. E., & Class, B. (2017). Phenotypic correlations capture between-individual  
642 correlations underlying behavioral syndromes. *Behavioral Ecology and Sociobiology*,  
643 71(3). <http://doi.org/10.1007/s00265-017-2278-4>
- 644 Carere, C., Drent, P. J., Koolhaas, J. M., & Groothuis, T. G. G. (2005). Epigenetic effects on

645 personality traits: Early food provisioning and sibling competition. *Behaviour*, 142(9),  
646 1329–1355. <http://doi.org/10.1163/156853905774539328>

647 Carere, C., & Maestriperi, D. (2013). *Animal personalities*. Chicago, IL: The University of  
648 Chicago Press.

649 Charmantier, A., & Réale, D. (2005). How do misassigned paternities affect the estimation of  
650 heritability in the wild? *Molecular Ecology*, 14(9), 2839–2850.  
651 <http://doi.org/10.1111/j.1365-294X.2005.02619.x>

652 Chervet, N., Zöttl, M., Schürch, R., Taborsky, M., & Heg, D. (2011). Repeatability and  
653 Heritability of Behavioural Types in a Social Cichlid. *International Journal of*  
654 *Evolutionary Biology*, 2011, 1–15. <http://doi.org/10.4061/2011/321729>

655 Cote, J., Clobert, J., Brodin, T., Fogarty, S., & Sih, A. (2010). Personality-dependent  
656 dispersal: characterization, ontogeny and consequences for spatially structured  
657 populations. *Philosophical Transactions of the Royal Society of London. Series B,*  
658 *Biological Sciences*, 365(1560), 4065–4076. <http://doi.org/10.1098/rstb.2010.0176>

659 David, M., Auclair, Y., & Cézilly, F. (2012). Assessing short- and long-term repeatability and  
660 stability of personality in captive zebra finches using longitudinal data. *Ethology*,  
661 118(10), 932–942. <http://doi.org/10.1111/j.1439-0310.2012.02085.x>

662 de Villemereuil, P. (2018). Quantitative genetic methods depending on the nature of the  
663 phenotypic trait. *Annals of the New York Academy of Sciences*, 1422(1), 29–47.  
664 <http://doi.org/10.1111/nyas.13571>

665 de Villemereuil, P., Schielzeth, H., Nakagawa, S., & Morrissey, M. (2016). General methods  
666 for evolutionary quantitative genetic inference from generalised mixed models. *Genetics*,  
667 204(3), 1–30. <http://doi.org/10.1534/genetics.115.186536>

668 Dingemanse, N. J., Both, C., Drent, P. J., & Tinbergen, J. M. (2004). Fitness consequences of  
669 avian personalities in a fluctuating environment. *Proceedings of the Royal Society B:*

670 *Biological Sciences*, 271(1541), 847–852. <http://doi.org/10.1098/rspb.2004.2680>

671 Dingemanse, N. J., Both, C., Drent, P. J., van Oers, K., & van Noordwijk, A. J. (2002).  
672 Repeatability and heritability of exploratory behaviour in great tits from the wild. *Animal*  
673 *Behaviour*, 64(6), 929–938. <http://doi.org/10.1006/anbe.2002.2006>

674 Dingemanse, N. J., Bouwman, K. M., van de Pol, M., van Overveld, T., Patrick, S. C.,  
675 Matthysen, E., & Quinn, J. L. (2012). Variation in personality and behavioural plasticity  
676 across four populations of the great tit *Parus major*. *Journal of Animal Ecology*, 81(1),  
677 116–126. <http://doi.org/10.1111/j.1365-2656.2011.01877.x>

678 Dingemanse, N. J., & Dochtermann, N. A. (2013). Quantifying individual variation in  
679 behaviour: Mixed-effect modelling approaches. *Journal of Animal Ecology*, 82(1), 39–  
680 54. <http://doi.org/10.1111/1365-2656.12013>

681 Dingemanse, N. J., Kazem, A. J. N., Réale, D., & Wright, J. (2010). Behavioural reaction  
682 norms: animal personality meets individual plasticity. *Trends in Ecology & Evolution*,  
683 25(2), 81–89. <http://doi.org/10.1016/j.tree.2009.07.013>

684 Dingemanse, N. J., & Réale, D. (2005). Natural selection and animal personality. *Behaviour*,  
685 142(9/10), 1165–1190. <http://doi.org/10.1163/156853905774539445>

686 Dingemanse, N. J., & Réale, D. (2013). What is the evidence that natural selection maintains  
687 variation in animal personalities? In C. Carere & D. Maestripieri (Eds.), *Animal*  
688 *personalities* (pp. 201–220). Chicago, IL: The University of Chicago Press.

689 Dingemanse, N. J., Wright, J., Kazem, A. J. N., Thomas, D. K., Hickling, R., & Dawnay, N.  
690 (2007). Behavioural syndromes differ predictably between 12 populations of three-  
691 spined stickleback. *Journal of Animal Ecology*, 76(6), 1128–1138.  
692 <http://doi.org/10.1111/j.1365-2656.2007.01284.x>

693 Dochtermann, N. A. (2011). Testing Cheverud’s conjecture for behavioral correlations and  
694 behavioral syndromes. *Evolution*, 65(6), 1814–1820. [28](http://doi.org/10.1111/j.1558-</a></p></div><div data-bbox=)

695 5646.2011.01264.x

696 Dochtermann, N. A., Schwab, T., & Sih, A. (2015). The contribution of additive genetic  
697 variation to personality variation: heritability of personality. *Proceedings of the Royal*  
698 *Society B: Biological Sciences*, 282(1798), 20142201.  
699 <http://doi.org/10.1098/rspb.2014.2201>

700 Doligez, B., & Clobert, J. (2003). Clutch size reduction as a response to increased nest  
701 predation rate in the collared flycatcher. *Ecology*, 84(10), 2582–2588.  
702 <http://doi.org/10.1890/02-3116>

703 Doligez, B., Danchin, E., Clobert, J., & Gustafsson, L. (1999). The use of conspecific  
704 reproductive success for breeding habitat selection in a non-colonial, hole-nesting  
705 species, the collared flycatcher. *Journal of Animal Ecology*, 68(6), 1193–1206.  
706 <http://doi.org/10.1046/j.1365-2656.1999.00362.x>

707 Doligez, B., Pärt, T., Danchin, E., Clobert, J., & Gustafsson, L. (2004). Availability and use  
708 of public information and conspecific density for settlement decisions in the collared  
709 flycatcher. *Journal of Animal Ecology*, 41, 75–87.

710 Drent, P. J., Oers, K. v., & Noordwijk, A. J. v. (2003). Realized heritability of personalities in  
711 the great tit (*Parus major*). *Proceedings of the Royal Society B: Biological Sciences*,  
712 270(1510), 45–51. <http://doi.org/10.1098/rspb.2002.2168>

713 Duckworth, R. A., & Badyaev, A. V. (2007). Coupling of dispersal and aggression facilitates  
714 the rapid range expansion of a passerine bird. *Proceedings of the National Academy of*  
715 *Sciences of the United States of America*, 104(38), 15017–22.  
716 <http://doi.org/10.1073/pnas.0706174104>

717 Duckworth, R. A., & Kruuk, L. E. B. (2009). Evolution of genetic integration between  
718 dispersal and colonization ability in a bird. *Evolution*, 63(4), 968–977.  
719 <http://doi.org/10.1111/j.1558-5646.2009.00625.x>

720 Eising, C. M., Muller, W., & Groothuis, T. G. . (2006). Avian mothers create different  
721 phenotypes by hormone deposition in their eggs. *Biology Letters*, 2(1), 20–22.  
722 <http://doi.org/10.1098/rsbl.2005.0391>

723 Evans, S. R., & Gustafsson, L. (2017). Climate change upends selection on ornamentation in a  
724 wild bird. *Nature Ecology and Evolution*, 1(2), 1–5. [http://doi.org/10.1038/s41559-016-](http://doi.org/10.1038/s41559-016-0039)  
725 0039

726 Falconer, D. S., & Mackay, T. F. C. (1996). *Introduction to quantitative genetics* (4th editio).  
727 London, U.K.: Prentice Hall.

728 Firth, J. A., Hadfield, J. D., Santure, A. W., Slate, J., & Sheldon, B. C. (2015). The influence  
729 of nonrandom extra-pair paternity on heritability estimates derived from wild pedigrees.  
730 *Evolution*, 69(5), 1336–1344. <http://doi.org/10.1111/evo.12649>

731 Garamszegi, L. Z., Eens, M., & Török, J. (2009). Behavioural syndromes and trappability in  
732 free-living collared flycatchers, *Ficedula albicollis*. *Animal Behaviour*, 77(4), 803–812.  
733 <http://doi.org/10.1016/j.anbehav.2008.12.012>

734 Garamszegi, L. Z., & Herczeg, G. (2012). Behavioural syndromes, syndrome deviation and  
735 the within- and between-individual components of phenotypic correlations: When reality  
736 does not meet statistics. *Behavioral Ecology and Sociobiology*, 66(12), 1651–1658.  
737 <http://doi.org/10.1007/s00265-012-1439-8>

738 Garamszegi, L. Z., Markó, G., & Herczeg, G. (2012). A meta-analysis of correlated  
739 behaviours with implications for behavioural syndromes: Mean effect size, publication  
740 bias, phylogenetic effects and the role of mediator variables. *Evolutionary Ecology*,  
741 26(5), 1213–1235. <http://doi.org/10.1007/s10682-012-9589-8>

742 Garamszegi, L. Z., Markó, G., Szász, E., Zsebők, S., Azcárate, M., Herczeg, G., & Török, J.  
743 (2015). Among-year variation in the repeatability, within- and between-individual, and  
744 phenotypic correlations of behaviors in a natural population. *Behavioral Ecology and*

745        *Sociobiology*, 69(12). <http://doi.org/10.1007/s00265-015-2012-z>

746 Garamszegi, L. Z., Rosivall, B., Rettenbacher, S., Markó, G., Zsebok, S., Szöllosi, E., ...

747        Török, J. (2012). Corticosterone, avoidance of novelty, risk-taking and aggression in a

748        wild bird: no evidence for pleiotropic effects. *Ethology*, 118(7), 621–635.

749        <http://doi.org/10.1111/j.1439-0310.2012.02049.x>

750 Grafen, A. (1984). Natural selection, kin selection and group selection. In J. R. Krebs & N. B.

751        Davies (Eds.), *Behavioural Ecology: an Evolutionary Approach* (pp. 62–84). Oxford,

752        U.K.: Blackwell Science.

753 Greenberg, R., & Mettke-Hofmann, C. (2001). Ecological aspects of neophobia and neophilia

754        in birds. *Current Ornithology*, 16I, 119–178. <http://doi.org/10.1007/978-1-4615-1211-0>

755 Groothuis, T. G. G., & Maestriperi, D. (2013). Parental influence on offspring personality

756        traits in oviparous and placental vertebrates. In C. Carere & D. Maestriperi (Eds.),

757        *Animal personalities: behaviour, physiology, and evolution* (pp. 317–352). Chicago, IL:

758        The University of Chicago Press.

759 Gustafsson, L. (1987). Interspecific competition lowers fitness in collared flycatchers

760        *Ficedula albicollis*: an experimental demonstration. *Ecology*, 68(2), 291–296.

761 Gustafsson, L. (1989). Collared flycatcher. In: I. Newton (ed.): *Lifetime reproduction in birds*

762        (pp. 75-88). London, U.K.: Academic Press.

763 Hadfield, J. D. (2010). MCMC methods for multi-response Generalized Linear Mixed

764        Models: The MCMCglmm R package. *Journal of Statistical Software*, 33(2), 1–22.

765        <http://doi.org/10.1002/ana.22635>

766 Hadfield, J. D. (2016). MCMCglmm course notes. [https://cran.r-](https://cran.r-project.org/web/packages/MCMCglmm/vignettes/CourseNotes.pdf)

767        [project.org/web/packages/MCMCglmm/vignettes/CourseNotes.pdf](https://cran.r-project.org/web/packages/MCMCglmm/vignettes/CourseNotes.pdf)

768 Hadfield, J. D., Nutall, A., Osorio, D., & Owens, I. P. F. (2007). Testing the phenotypic

769        gambit: Phenotypic, genetic and environmental correlations of colour. *Journal of*



770 *Evolutionary Biology*, 20(2), 549–557. <http://doi.org/10.1111/j.1420-9101.2006.01262.x>

771 Hegyi, G., Rosivall, B., & Török, J. (2006). Paternal age and offspring growth: Separating the  
772 intrinsic quality of young from rearing effects. *Behavioral Ecology and Sociobiology*,  
773 60(5), 672–682. <http://doi.org/10.1007/s00265-006-0211-3>

774 Holtmann, B., Lagisz, M., & Nakagawa, S. (2017). Metabolic rates, and not hormone levels,  
775 are a likely mediator of between-individual differences in behaviour: a meta-analysis.  
776 *Functional Ecology*, 31(685–696). <http://doi.org/10.1111/1365-2435.12779>

777 Karlsson Green, K., Eroukhmanoff, F., Harris, S., Pettersson, L. B., & Svensson, E. I. (2016).  
778 Rapid changes in genetic architecture of behavioural syndromes following colonisation  
779 of a novel environment. *Journal of Evolutionary Biology*, 29, 144–152.  
780 <http://doi.org/10.1111/jeb.12769>

781 Kivelä, S. M., Seppänen, J.-T., Ovaskainen, O., Doligez, B., Gustafsson, L., Mönkkönen, M.,  
782 & Forsman, J. T. (2014). The past and the present in decision-making: the use of  
783 conspecific and heterospecific cues in nest site selection. *Ecology*, 95, 3428–3439.  
784 <http://doi.org/10.1890/13-2103.1>

785 Koolhaas, J. ., Korte, S. ., De Boer, S. ., Van Der Vegt, B. ., Van Reenen, C. ., Hopster, H., ...  
786 Blokhuis, H. . (1999). Coping styles in animals: current status in behavior and stress-  
787 physiology. *Neuroscience & Biobehavioral Reviews*, 23(7), 925–935.  
788 [http://doi.org/10.1016/S0149-7634\(99\)00026-3](http://doi.org/10.1016/S0149-7634(99)00026-3)

789 Korsten, P., van Overveld, T., Adriaensen, F., & Matthysen, E. (2013). Genetic integration of  
790 local dispersal and exploratory behaviour in a wild bird. *Nature Communications*,  
791 4(2362). <http://doi.org/10.1038/ncomms3362>

792 Král, M., & Bícík, V. (1989). Intraspecific aggressive responses of male collared flycatchers  
793 (*Ficedula albicollis* temm.). *Acta Universitatis Palackianae Olomucensis, Facultas*  
794 *Rerum Naturalium, Biologica XXIX*, 96, 107–122.

795 Král, M., & Bícík, V. (1992). Nest defence by the collared flycatcher (*Ficedula albicollis*)  
796 against the great tit (*Parus major*). *Folia Zoologica*, 41(3), 263–269.

797 Kruuk, L. E. B., & Hadfield, J. D. (2007). How to separate genetic and environmental causes  
798 of similarity between relatives. *Journal of Evolutionary Biology*, 20(5), 1890–1903.  
799 <http://doi.org/10.1111/j.1420-9101.2007.01377.x>

800 Lê, S., Josse, J., & Husson, F. (2008). FactoMineR: an R package for multivariate analysis.  
801 *Journal of Statistical Software*, 25(1), 1–18.

802 Merilä, J., & Gustafsson, L. (1993). Inheritance of size and shape in a natural population of  
803 collared flycatchers, *Ficedula albicollis*. *Journal of Evolutionary Biology*, 6(3), 375–395.  
804 <http://doi.org/10.1046/j.1420-9101.1993.6030375.x>

805 Merilä, J., & Sheldon, B. C. (2000). Lifetime Reproductive Success and Heritability in  
806 Nature. *The American Naturalist*, 155(3), 301–310. <http://doi.org/10.1086/303330>

807 Morinay, J., Forsman, J. T., Kivelä, S. M., Gustafsson, L., & Doligez, B. (2018).  
808 Heterospecific nest site copying behavior in a wild bird: Assessing the influence of  
809 genetics and past experience on a joint breeding phenotype. *Frontiers in Ecology and*  
810 *Evolution*, 5(January), 1–14. <http://doi.org/10.3389/fevo.2017.00167>

811 Morrissey, M. B., & Wilson, A. J. (2010). Pedantics: an R package for pedigree-based genetic  
812 simulation and pedigree manipulation, characterization and viewing. *Molecular Ecology*  
813 *Resources*, 10(4), 711–719. <http://doi.org/10.1111/j.1755-0998.2009.02817.x>

814 Nakagawa, S., & Schielzeth, H. (2010). Repeatability for Gaussian and non-Gaussian data: a  
815 practical guide for biologists. *Biological Reviews*, 85, 935–956.  
816 <http://doi.org/10.1111/j.1469-185X.2010.00141.x>

817 Niemelä, P. T., & Dingemanse, N. J. (2017). Individual versus pseudo-repeatability in  
818 behaviour: Lessons from translocation experiments in a wild insect. *Journal of Animal*  
819 *Ecology*, 86(5), 1033–1043. <http://doi.org/10.1111/1365-2656.12688>

820 Niemelä, P. T., Lattenkamp, E. Z., & Dingemanse, N. J. (2015). Personality-related survival  
821 and sampling bias in wild cricket nymphs. *Behavioral Ecology*, 26(3), 936–946.  
822 <http://doi.org/10.1093/beheco/arv036>

823 Petelle, M. B., Martin, J. G. A., & Blumstein, D. T. (2015). Heritability and genetic  
824 correlations of personality traits in a wild population of yellow-bellied marmots  
825 (*Marmota flaviventris*). *Journal of Evolutionary Biology*, 28(10), 1840–1848.  
826 <http://doi.org/10.1111/jeb.12700>

827 Plummer, M., Best, N., Cowles, K., & Vines, K (2006). CODA: Convergence Diagnosis and  
828 Output Analysis for MCMC, *R News*, 6, 7-11.

829 Qvarnström, A. (1997). Experimentally increased badge size increases male competition and  
830 reduces male parental care in the collared flycatcher. *Proceedings of the Royal Society B:*  
831 *Biological Sciences*, 264(1385), 1225–1231. <http://doi.org/10.1098/rspb.1997.0169>

832 R Core Team. (2016). *R: a language and environment for statistical computing*. Vienna,  
833 Austria: R Foundation for Statistical Computing. <https://www.R-project.org/>.

834 Reale, D., Garant, D., Humphries, M. M., Bergeron, P., Careau, V., & Montiglio, P.-O.  
835 (2010). Personality and the emergence of the pace-of-life syndrome concept at the  
836 population level. *Philosophical Transactions of the Royal Society B: Biological Sciences*,  
837 365(1560), 4051–4063. <http://doi.org/10.1098/rstb.2010.0208>

838 Réale, D., Reader, S. M., Sol, D., McDougall, P. T., & Dingemanse, N. J. (2007). Integrating  
839 animal temperament within ecology and evolution. *Biological Reviews*, 82(2), 291–318.  
840 <http://doi.org/10.1111/j.1469-185X.2007.00010.x>

841 Rosivall, B., Török, J., Hasselquist, D., & Bensch, S. (2004). Brood sex ratio adjustment in  
842 collared flycatchers (*Ficedula albicollis*): results differ between populations. *Behavioral*  
843 *Ecology and Sociobiology*, 56(4), 346–351. <http://doi.org/10.1007/s00265-004-0796-3>

844 Schuett, W., Tregenza, T., & Dall, S. R. X. (2010). Sexual selection and animal personality.

845 *Biological Reviews*, 85, 217–246. <http://doi.org/10.1111/j.1469-185X.2009.00101.x>

846 Sheldon, B. C., & Ellegren, H. (1999). Sexual selection resulting from extrapair paternity in  
847 collared flycatchers. *Animal Behaviour*, 57(2), 285–298.  
848 <http://doi.org/10.1006/anbe.1998.0968>

849 Sheldon, B. C., Kruuk, L. E. B., & Merilä, J. (2003). Natural selection and inheritance of  
850 breeding time and clutch size in the collared flycatcher. *Evolution*, 57(2), 406–420.  
851 <http://doi.org/10.1111/j.0014-3820.2003.tb00274.x>

852 Sih, A., Bell, A. M., & Johnson, J. C. (2004). Behavioral syndromes: an ecological and  
853 evolutionary overview. *Trends in Ecology & Evolution*, 19(7), 372–378.  
854 <http://doi.org/10.1016/j.tree.2004.04.009>

855 Sih, A., Bell, A. M., Johnson, J. C., & Ziemba, R. E. (2004). Behavioral syndromes: an  
856 integrative overview. *The Quarterly Review of Biology*, 79(3), 241–277.  
857 <http://doi.org/10.1086/422893>

858 Sih, A., Cote, J., Evans, M., Fogarty, S., & Pruitt, J. (2012). Ecological implications of  
859 behavioural syndromes. *Ecology Letters*, 15(3), 278–289. [http://doi.org/10.1111/j.1461-](http://doi.org/10.1111/j.1461-0248.2011.01731.x)  
860 [0248.2011.01731.x](http://doi.org/10.1111/j.1461-0248.2011.01731.x)

861 Smith, B. R., & Blumstein, D. T. (2008). Fitness consequences of personality: A meta-  
862 analysis. *Behavioral Ecology*, 19(2), 448–455. <http://doi.org/10.1093/beheco/arm144>

863 Spencer, K. A., & Verhulst, S. (2007). Delayed behavioral effects of postnatal exposure to  
864 corticosterone in the zebra finch (*Taeniopygia guttata*). *Hormones and Behavior*, 51(2),  
865 273–280. <http://doi.org/10.1016/j.yhbeh.2006.11.001>

866 Stamps, J., & Groothuis, T. G. G. (2010). The development of animal personality: Relevance,  
867 concepts and perspectives. *Biological Reviews*, 85(2), 301–325.  
868 <http://doi.org/10.1111/j.1469-185X.2009.00103.x>

869 Svensson, L. (1992). *Identification guide to european passerines* (Märstatryc). Stockholm,

870 Sweden: Svensson.

871 van Oers, K., De Jong, G., Drent, P. J., & van Noordwijk, A. J. (2004). A genetic analysis of  
872 avian personality traits: Correlated response to artificial selection. *Behavior Genetics*,  
873 *34*(6), 611–619. <http://doi.org/10.1007/s10519-004-5588-z>

874 van Oers, K., de Jong, G., van Noordwijk, A. J., & Drent, P. J. (2005). Contribution of  
875 genetics to the study of animal personalities: a review of case studies. *Behaviour*, *142*(9),  
876 1185–1206. <http://doi.org/10.1163/156853905774539364>

877 van Oers, K., Drent, P. J., de Goede, P., & van Noordwijk, A. J. (2004). Realized heritability  
878 and repeatability of risk-taking behaviour in relation to avian personalities. *Proceedings*  
879 *of the Royal Society B: Biological Sciences*, *271*, 65–73.  
880 <http://doi.org/10.1098/rspb.2002.2168>

881 van Oers, K., & Sinn, D. L. (2011). Towards a basis for the phenotypic gambit: advances in  
882 the evolutionary genetics of animal personality. In M. Inoue-Murayama, S. Kawamura,  
883 & A. Weiss (Eds.), *From genes to behavior* (pp. 165–183). Tokyo, Japan: Springer.

884 van Oers, K., & Sinn, D. L. (2013). Quantitative and molecular genetics of animal  
885 personality. In C. Carere & D. Maestripieri (Eds.), *Animal personalities: behaviour,*  
886 *physiology, and evolution* (pp. 149–200). Chicago, IL: The University of Chicago Press.

887 Wilson, A. J., Réale, D., Clements, M. N., Morrissey, M. M., Postma, E., Walling, C. A., ...  
888 Nussey, D. H. (2010). An ecologist's guide to the animal model. *Journal of Animal*  
889 *Ecology*, *79*(1), 13–26. <http://doi.org/10.1111/j.1365-2656.2009.01639.x>

890 Wolak, M. E. (2012). nadiv: an R package to create relatedness matrices for estimating non-  
891 additive genetic variances in animal models. *Methods in Ecology and Evolution*, *3*(5),  
892 792–796. <http://doi.org/10.1111/j.2041-210X.2012.00213.x>

893 Wolf, J. B., Brodie III, E. D. B., Cheverud, J. M., Moore, A. J., & Wade, M. J. (1998).  
894 Indirect genetic effects. *Trends in Ecology & Evolution*, *13*(2), 64–69.

895 [http://doi.org/10.1016/S0169-5347\(97\)01233-0](http://doi.org/10.1016/S0169-5347(97)01233-0)

896 Wuerz, Y., & Krüger, O. (2015). Personality over ontogeny in zebra finches: Long-term  
897 repeatable traits but unstable behavioural syndromes. *Frontiers in Zoology*, 12(1).  
898 <http://doi.org/10.1186/1742-9994-12-S1-S9>

899

## 900 **Appendix**

### 901 Aggressiveness score

902 We estimated aggressiveness score using the following alternative measures and modelled  
903 them using the parameters given in parentheses. (1) Number of aggressive behaviours  
904 (movements < 2 m from the nestbox, stationary flights and attacks towards the decoys, chases  
905 of live birds) standardized per 15 min (Poisson family; number of iterations =  $10^6$ ; burn-in =  
906  $10^4$ ; thinning interval = 400). (2) First axis of the PCA presented below (Table A7; Gaussian  
907 family; number of iterations =  $10^6$ ; burn-in =  $10^4$ ; thinning interval = 400). (3) Discrete score  
908 (threshold family, residual variance  $V_R = 10$ ; number of iterations =  $13 \times 10^5$ ; burn-in =  $8 \times$   
909  $10^4$ ; thinning interval = 700). This score was based on the distinction between activity  
910 (number of movements/min, including stationary flights and chases) performed far from (> 2  
911 m) and close to (< 2 m) the nestbox, and on attacks, subdivided into six categories (Fig. A3a):  
912 **0**: individuals that performed no movements (either far from or close to the nestbox); 1:  
913 individuals that performed no attack or movements close to the nestbox and performed less  
914 than 0.440 movements/min far from (> 5 m) the nestbox; 2: individuals that performed no  
915 attack or movements close to the nestbox and performed more than 0.440 movements/min far  
916 from (> 5 m) the nestbox; 3: individuals that performed no attacks and less than 0.282  
917 movements/min close to (< 2 m) the nestbox; 4: individuals that performed no attacks and  
918 between 0.282 and 0.784 movements/min close to (< 2 m) the nestbox; 5: individuals that  
919 performed no attacks and above 0.784 movements/min close to (< 2 m) the nestbox; 6:

920 individuals that performed attacks towards decoys(s). The thresholds were chosen so as to  
921 distribute individuals equally among categories for scores 1 and 2 on the one hand and scores  
922 3, 4 and 5 on the other.

923

#### 924 Boldness score

925 We estimated boldness score using the following alternative measures and modelled them  
926 using the parameters given in parentheses. (1) Maximum latency to enter the nestbox after  
927 human departure (from all individuals) minus the same latency for the focal individual on the  
928 focal test (Poisson family; number of iterations =  $1 \times 10^6$ ; burn-in =  $10^4$ ; thinning interval =  
929 500). (2) Inverse logarithmic ratio of the latency to enter the nestbox after human departure  
930 divided by the feeding rate for the remaining time after the first entrance in the nestbox. We  
931 divided by the feeding rate for this alternative measurement because the average feeding  
932 interval might have affected the latency to return to the nestbox, for example if individuals  
933 feeding more frequently entered the nestbox faster. The feeding rate after the first entrance  
934 was estimated as the average time interval between two feeding events by the focal birds,  
935 after the first entrance; it was thus only computable for individuals that fed at least twice in  
936 the period; for technical reasons, further individuals could not be used for this variable,  
937 leading to a final used data set of 641 observations (Gaussian family; number of iterations =  
938  $25 \times 10^4$ ; burn-in =  $10^4$ ; thinning interval = 100). (3) Discrete score (threshold family,  $V_R =$   
939 10; number of iterations =  $3 \times 10^6$ ; burn-in =  $2 \times 10^5$ ; thinning interval = 1000). The score was  
940 based on entrance in the nestbox during the first part of the boldness-neophobia test (no novel  
941 object) and latency to enter after human disturbance; individuals that did not enter were given  
942 a score of 0, and individuals that entered the nestbox were given a score of 1 - 5 based on five  
943 quantiles of the inverse latency to enter (Fig. A3b).

944

945

946 Neophobia score

947 We estimated neophobia score for individuals that entered the nestbox during the first period  
948 of the test, using the following alternative measures and modelled them using the parameters  
949 given in parentheses. (1) Discrete score based on the latency to enter the nestbox in the  
950 presence of a novel object (second period of the test), discretized in four quantiles, the fifth  
951 category including individuals that did not enter the nestbox at all in the presence of the novel  
952 object (threshold family;  $V_R = 10$ ; number of iterations =  $15 \times 10^5$ ; burn-in =  $2 \times 10^5$ ; thinning  
953 interval = 500; Fig. A3c); (2) Binary variable separating individuals that did and did not enter  
954 during the second period of the test (threshold family;  $V_R = 10$ ; number of iterations =  $10^6$ ;  
955 burn-in =  $10^5$ ; thinning interval = 500). (3) Latency to enter the nestbox in the presence of the  
956 novel object, excluding the individuals that did not enter the nestbox during the second period  
957 of the test (Poisson family; number of iterations =  $15 \times 10^5$ ; burn-in =  $15 \times 10^4$ ; thinning  
958 interval = 500).

959 Results obtained for these alternative scores for the three behavioural traits are given in  
960 Tables A2 to A4, and the main text presents the first score in each case.



961 **TABLES**

962

963 **Table 1.** Between-year repeatability and heritability estimates for aggressiveness, boldness  
 964 and neophobia scores

	$R_{\text{latent}}$	$R_{\text{obs}}$	$h^2_{\text{latent}}$	$h^2_{\text{obs}}$
Aggressiveness	0.18 *	0.03 *	0.00	0.00
	[0.15; 0.23]	[0.02; 0.04]	[0.00; 0.08]	[0.00; 0.008]
Boldness	0.11 *	0.10 *	0.00	0.00
	[0.01; 0.21]	[0.01; 0.19]	[0.00; 0.10]	[0.00; 0.09]
Neophobia	0.39 *		0.00	0.00 for all scores
	[0.25; 0.54]		[0.00; 0.15]	From [0.00; 0.00] to [0.00; 0.09]

965

966 Repeatabilities and heritabilities (posterior modes and 95% credible intervals) are given on  
 967 the latent scale ( $R_{\text{latent}}$ ,  $h^2_{\text{latent}}$ ) and on the observed scale ( $R_{\text{obs}}$  obtained with the QGicc function  
 968 and  $h^2_{\text{obs}}$  with the QGparams function from ‘QGglmm’ R package; de Villemereuil et al.  
 969 2016; de Villemereuil 2018). Asterisks indicate estimates whose 95% CI do not encompass  
 970 zero. For aggressiveness, estimates are given using all scores. For neophobia, we provide a  
 971 range of heritability values on the observed scale, since one value is provided per neophobia  
 972 score level (i.e. five values in total); however, we could not derive repeatability estimates on  
 973 the observed scale for ordinal variables.

974

975 **Table 2.** Phenotypic, between- and within-individual, and additive genetic correlations  
 976 between aggressiveness, boldness and neophobia scores  
 977

	Aggressiveness	Boldness	Aggressiveness
Correlation level	-	-	-
	Neophobia	Neophobia	Boldness
Phenotypic	-0.20*	-0.30*	0.02
	[-0.25; -0.12]	[-0.38; -0.23]	[-0.04; 0.10]
Between-individual	0.00	0.00	-0.02
	[-0.06; 0.06]	[-0.07; 0.05]	[-0.06; 0.06]
Within-individual	-0.28*	-0.43*	0.03
	[-0.38; -0.18]	[-0.53; -0.32]	[-0.04; 0.12]
Additive genetic	0.00	-0.01	0.01
	[-0.06; 0.006]	[-0.06; 0.06]	[-0.07; 0.06]

978  
 979 Posterior modes and 95% credible intervals (CI) on the latent scale are shown. Asterisks  
 980 indicate estimates whose 95% CI do not encompass zero.

981 **Table A1.** Detailed description of the collared flycatcher pedigree from the Gotland

982 Island population

---

Pedigree statistics	
Records	2 218
Founders	1423
Maternities	728
Paternities	781
Mothers with $\geq 2$ offspring	120
Fathers with $\geq 2$ offspring	133
Full sibs	130
Maternal sibs	222
Maternal half sibs	92
Paternal sibs	230
Paternal half sibs	100
Maternal grandmothers	249
Maternal grandfathers	268
Paternal grandmothers	220
Paternal grandfathers	236
Maximum pedigree depth	15
Mean relatedness	$5.29 \times 10^{-4}$

---

983

984 The pedigree statistics were obtained from all identified individuals involved in either  
985 aggressiveness, boldness or neophobia assays, and were extracted using the pedigreeStats and  
986 pedStatSummary functions from ‘pedantics’ R package (Morrissey & Wilson, 2010).

987

1 **Table A2.** Output of the univariate models fitting aggressiveness score

	Number of aggressive behaviours/15 min				PC1		D
	Without fixed effect		With fixed effects				
	Posterior mode	95% CI	Posterior mode	95% CI	Posterior mode	95% CI	
<b>Fixed effects</b>							
Intercept	1.22	[ 0.91 ; 1.50 ] *	-0.29	[ -0.72 ; 0.26 ]	-0.98	[ -1.30 ; -0.65 ] *	5
Sex (male)			0.63	[ 0.44 ; 0.81 ] *	0.50	[ 0.35 ; 0.62 ] *	1
Age (young)			0.23	[ -0.01 ; 0.40 ]	0.06	[ -0.05 ; 0.25 ]	0
Presence of tits (present)			0.34	[ 0.19 ; 0.50 ] *	0.05	[ -0.07 ; 0.15 ]	0
Presence of flycatchers (present)			0.27	[ -0.02 ; 0.55 ]	0.29	[ 0.05 ; 0.49 ] *	0
Presence of the partner (present)			0.62	[ 0.44 ; 0.80 ] *	0.34	[ 0.20 ; 0.44 ] *	1
Test number			-0.20	[ -0.27 ; -0.14 ] *	-0.16	[ -0.27 ; -0.05 ] *	-1

							0.20	0.11	
Dummy type (flycatcher)	0.08	[ -0.23 ; 0.34 ]		0.16	[ - ; 0.38 ]		0.01		0
Year (2012)	0.48	[ 0.08 ; 0.87 ] *		0.19	[ - ; 0.50 ]		0.05		0
Year (2013)	0.29	[ -0.09 ; 0.67 ]		0.10	[ - ; 0.46 ]		0.09		0
Sex*age (male*young)	0.28	[ 0.03 ; 0.61 ] *		0.24	[ 0.09 ; 0.50 ] *				0
<b>Random effects</b>									
$V_A$	0.00	[ 0.00 ; 0.32 ]		0.00	[ 0.00 ; 0.41 ]		0.00	[ 0.00 ; 0.16 ]	0
$V_{PE}$	0.60	[ 0.28 ; 0.80 ] *		0.52	[ 0.11 ; 0.71 ] *		0.19	[ 0.03 ; 0.28 ] *	1
$V_M$	0.00	[ 0.00 ; 0.20 ]		0.00	[ 0.00 ; 0.17 ]		0.00	[ 0.00 ; 0.12 ]	0
$V_F$	0.00	[ 0.00 ; 0.20 ]		0.00	[ 0.00 ; 0.24 ]		0.23	[ 0.06 ; 0.37 ] *	0
$V_{plot}$	0.07	[ 0.02 ; 0.16 ] *		0.05	[ 0.01 ; 0.15 ] *		0.01	[ 0.00 ; 0.04 ]	0
$V_{decoy}$	0.00	[ 0.00 ; 0.03 ]		0.00	[ 0.00 ; 0.04 ]		0.00	[ 0.00 ; 0.03 ]	0
$V_{song}$	0.02	[ 0.00 ; 0.09 ]		0.01	[ 0.00 ; 0.09 ]		0.00	[ 0.00 ; 0.03 ]	0

$V_{\text{obs}}$	0.25	[ 0.14 ; 0.55 ] *	0.30	[ 0.15 ; 0.63 ] *	0.06	[ 0.02 ; 0.16 ] *	0
$V_{\epsilon}$	2.56	[ 0.91 ; 1.50 ] *	2.11	[ 1.97 ; 2.34 ] *	1.49	[ 1.40 ; 1.57 ] *	1
<b>Derived estimates</b>							
$R_{\text{latent}}$	0.18	[ 0.15 ; 0.23 ] *	0.18	[ 0.13 ; 0.22 ] *	0.23	[ 0.17 ; 0.27 ] *	0
$R_{\text{obs}}$	0.03	[ 0.02 ; 0.04 ] *	0.03	[ 0.02 ; 0.04 ] *	0.21	[ 0.16 ; 0.25 ] *	
$h^2_{\text{latent}}$	0.00	[ 0.00 ; 0.08 ]	0.00	[ 0.00 ; 0.12 ]	0.00	[ 0.00 ; 0.08 ]	0
$h^2_{\text{obs}}$	0.00	[ 0.00 ; 0.01 ]	0.00	[ 0.00 ; 0.01 ]	0.00	[ 0.00 ; 0.07 ]	P
$\overline{N}$	4680		3271		3271		
Effective sample size	> 2357		> 2234		> 2209		

2

3 Models for the general aggressiveness score, based on the number of aggressive behaviours standardized per 15 min, are shown without and with  
4 fixed effects. We also present the posterior modes and 95% credible intervals (CI) of models for alternative aggressiveness scores: the first axis  
5 of the principal component analysis (PC1) and the discrete score (see Appendix). Asterisks indicate estimates whose 95% CI do not encompass  
6 zero. For categorical fixed terms, estimates refer to the category indicated in parentheses.  $V_A$ ,  $V_{PE}$ ,  $V_M$ ,  $V_F$  and  $V_e$  refer to the additive genetic,  
7 permanent environment, maternal, paternal and residual variances, respectively.  $V_{\text{plot}}$ ,  $V_{\text{decoy}}$ ,  $V_{\text{song}}$ ,  $V_{\text{obs}}$  refer to the variances associated with the  
8 plot, the decoy set used, the song track played and observer identity, respectively.  $N$  is the sample size of the data set used in the model.

- 9 Repeatability and heritability estimates are given both on the latent scale ( $R_{\text{latent}}, h^2_{\text{latent}}$ ) and on the observed scale ( $R_{\text{obs}}, h^2_{\text{obs}}$ ) whenever these
- 10 could be estimated.

11 **Table A3 Output of the models fitting boldness scores.**

	Maximum latency–individual latency				Latency/feeding rate		Discrete score
	Without fixed effects		Without fixed effects		Posterior mode	95% CI	Posterior mode
	Posterior mode	95% CI	Posterior mode	95% CI			
<b>Fixed effects</b>							
Intercept	8.00	[ 7.97 ; 8.04 ] *	8.05	[ 7.96 ; 8.10 ] *	-3.78	[ -4.07 ; 3.46 ] *	4.80
Sex (male)			-0.04	[ -0.12 ; 0.02 ]	-0.10	[ -0.29 ; 0.13 ]	-1.33
Age (young)			-0.02	[ -0.10 ; 0.05 ]	-0.12	[ -0.39 ; 0.13 ]	-0.65
Year (2012)			-0.03	[ -0.10 ; 0.04 ]	0.05	[ -0.19 ; 0.43 ]	-0.38
Year (2013)			0.08	[ 0.00 ; 0.16 ] *	0.47	[ 0.10 ; 0.73 ] *	1.46
No. of chicks			0.03	[ 0.00 ; 0.06 ] *	0.26	[ 0.15 ; 0.33 ] *	0.36
Sex*age (male*young)			0.03	[ -0.08 ; 0.13 ]	0.30	[ -0.18 ; 0.55 ]	0.60
<b>Random effects</b>							
$V_A$	0.00	[ 0.00 ; 0.02 ]	0.00	[ 0.00 ; 0.03 ]	0.00	[ 0.00 ; 0.20 ]	0.01



$V_{PE}$	0.00	[ 0.00 ; 0.02 ]	0.00	[ 0.00 ; 0.03 ]	0.00	[ 0.00 ; 0.25 ]	0.04
$V_M$	0.00	[ 0.00 ; 0.02 ]	0.00	[ 0.00 ; 0.02 ]	0.00	[ 0.00 ; 0.13 ]	0.01
$V_F$	0.00	[ 0.00 ; 0.01 ]	0.00	[ 0.00 ; 0.01 ]	0.00	[ 0.00 ; 0.15 ]	0.01
$V_{plot}$	0.00	[ 0.00 ; 0.00 ]	0.00	[ 0.00 ; 0.01 ]	0.00	[ 0.00 ; 0.07 ]	0.00
$V_{obs}$	0.00	[ 0.00 ; 0.00 ]	0.00	[ 0.00 ; 0.00 ]	0.00	[ 0.00 ; 0.03 ]	0.00
$V_{\epsilon}$	0.16	[ 0.14 ; 0.17 ] *	0.13	[ 0.11 ; 0.16 ] *	1.20	[ 0.93 ; 1.37 ] *	10.00

**Derived estimates**

$R_{latent}$	0.11	[ 0.01 ; 0.21 ] *	0.19	[ 0.07 ; 0.31 ] *	0.12	[ 0.01 ; 0.29 ] *	0.28
$R_{obs}$	0.10	[ 0.01 ; 0.19 ] *	0.12	[ 0.02 ; 0.23 ] *	0.04	[ 0.01 ; 0.14 ] *	
$h^2_{latent}$	0.00	[ 0.00 ; 0.10 ]	0.00	[ 0.00 ; 0.18 ]	0.00	[ 0.00 ; 0.14 ]	0.00
$h^2_{obs}$	0.00	[ 0.00 ; 0.09 ]	0.00	[ 0.00 ; 0.09 ]	0.00	[ 0.00 ; 0.06 ]	Posterior m
$N$	1064		914		641		
Effective sample size	> 1718		> 1979		> 2053		

12 Models for the general boldness estimate (maximum latency to enter the nestbox after human disturbance observed in the entire data set minus  
13 the individual latency) are shown without and with fixed effects. We also present the posterior modes and 95% credible intervals (CI) of models

14 for alternative boldness estimates: the log-transformed and inverse ratio of the latency to enter the nestbox after human disturbance over the  
15 feeding rate during the time remaining and the discrete score based on the latency (see Appendix). Asterisks indicate estimates whose 95% CI do  
16 not encompass zero. For categorical fixed terms, estimates refer to the category indicated in parentheses.  $V_A$ ,  $V_{PE}$ ,  $V_M$ ,  $V_F$  and  $V_e$  refer to the  
17 additive genetic, permanent environment, maternal, paternal and residual variances, respectively.  $V_{plot}$  and  $V_{obs}$  refer to the variances associated  
18 with the plot and the observer identity, respectively.  $N$  is the sample size of the data set used in the model. Repeatability and heritability estimates  
19 are given both on the latent scale ( $R_{latent}$ ,  $h^2_{latent}$ ) and on the observed scale ( $R_{obs}$ ,  $h^2_{obs}$ ) whenever these could be estimated. Repeatability and  
20 heritability estimates for the latency/feeding rate model (fitted with a Gaussian distribution) differ between the latent and observed scales because  
21 we accounted for the inverse log transformation in the `QGicc` and `QGparams` functions (`'QGglmm'` R package; de Villemereuil et al. 2016).

22 **Table A4. Output of the models fitting neophobia scores**

	Discrete score				Binary		La
	Without fixed effect		With fixed effects		Posterior mode	95% CI	
	Posterior mode	95% CI	Posterior mode	95% CI			
<b>Fixed effects</b>							
Intercept	4.22	[ 3.40 ; 4.94 ] *	5.08	[ 3.63 ; 6.38 ] *	-1.67	[ -3.39 ; -0.30 ] *	6.
Sex (male)			-1.89	[ -2.78 ; -1.07 ] *	-1.45	[ -2.44 ; -0.41 ] *	-0.
Age (young)			-0.18	[ -1.01 ; 0.76 ]	0.07	[ -1.26 ; 0.92 ]	-0.
Year (2012)			1.62	[ 0.72 ; 2.93 ] *	2.19	[ 0.59 ; 3.85 ] *	0.
Year (2013)			1.21	[ -0.05 ; 2.37 ]	1.84	[ -0.06 ; 3.34 ]	0.
Boldness			-1.58	[ -1.93 ; -1.12 ] *	-1.34	[ -1.95 ; -0.93 ] *	-0.

No. of chicks			-0.71	[ -1.03 ; -0.37 ] *	-0.71	[ -1.15 ; -0.30 ] *	-0.71	[ -1.15 ; -0.30 ] *	-0.71
---------------	--	--	-------	---------------------	-------	---------------------	-------	---------------------	-------

Sex*age (male*young)			-1.02	[ -2.44 ; 0.11 ]	-0.79	[ -2.75 ; 0.69 ]	-0.79	[ -2.75 ; 0.69 ]	-0.79
----------------------	--	--	-------	------------------	-------	------------------	-------	------------------	-------

**Random effects**

$V_A$	0.02	[ 0.00 ; 2.90 ]	0.03	[ 0.00 ; 5.76 ]	0.08	[ 0.00 ; 8.41 ]	0.08	[ 0.00 ; 8.41 ]	0.08
-------	------	-----------------	------	-----------------	------	-----------------	------	-----------------	------

$V_{PE}$	0.04	[ 0.00 ; 5.77 ]	0.06	[ 0.00 ; 9.03 ]	0.05	[ 0.00 ; 10.70 ]	0.05	[ 0.00 ; 10.70 ]	0.05
----------	------	-----------------	------	-----------------	------	------------------	------	------------------	------

$V_M$	0.04	[ 0.00 ; 6.49 ]	0.06	[ 0.00 ; 8.81 ]	0.12	[ 0.00 ; 22.53 ]	0.12	[ 0.00 ; 22.53 ]	0.12
-------	------	-----------------	------	-----------------	------	------------------	------	------------------	------

$V_F$	0.02	[ 0.00 ; 5.75 ]	0.03	[ 0.00 ; 7.85 ]	0.04	[ 0.00 ; 11.33 ]	0.04	[ 0.00 ; 11.33 ]	0.04
-------	------	-----------------	------	-----------------	------	------------------	------	------------------	------

$V_{plot}$	0.98	[ 0.26 ; 2.50 ] *	1.01	[ 0.27 ; 3.42 ] *	0.80	[ 0.00 ; 3.08 ]	0.80	[ 0.00 ; 3.08 ]	0.80
------------	------	-------------------	------	-------------------	------	-----------------	------	-----------------	------

$V_{obs}$	0.28	[ 0.02 ; 1.01 ] *	0.38	[ 0.00 ; 1.41 ]	0.80	[ 0.14 ; 3.14 ] *	0.80	[ 0.14 ; 3.14 ] *	0.80
-----------	------	-------------------	------	-----------------	------	-------------------	------	-------------------	------

$V_{\epsilon}$	10.00	[ 10.00 ; 10.00 ]	10.00	[ 10.00 ; 10.00 ]	10.00	[ 10.00 ; 10.00 ]	10.00	[ 10.00 ; 10.00 ]	10.00
----------------	-------	-------------------	-------	-------------------	-------	-------------------	-------	-------------------	-------

**Derived estimates**

$R_{latent}$	0.39	[ 0.25 ; 0.54 ] *	0.50	[ 0.33 ; 0.63 ] *	0.56	[ 0.35 ; 0.75 ] *	0.56	[ 0.35 ; 0.75 ] *	0.56
--------------	------	-------------------	------	-------------------	------	-------------------	------	-------------------	------

$R_{obs}$					0.29	[ 0.17 ; 0.48 ] *	0.29	[ 0.17 ; 0.48 ] *	0.29
-----------	--	--	--	--	------	-------------------	------	-------------------	------

$h^2_{latent}$	0.00	[ 0.00 ; 0.15 ]	0.00	[ 0.00 ; 0.22 ]	0.00	[ 0.00 ; 0.26 ]	0.00	[ 0.00 ; 0.26 ]	0.00
----------------	------	-----------------	------	-----------------	------	-----------------	------	-----------------	------

$h^2_{\text{obs}}$	Posterior mode range: 0.00-0.00	Posterior mode range: 0.00-0.00	0.00	[ 0.00 ; 0.14 ]	0.
$N$	1 064	914	914		
Effective sample size	> 2302	> 2317	> 1583		

23 Models for the general neophobia discrete score, based on the latency to enter the nestbox in presence of the novel object, are shown without and  
24 with fixed effects We also present the posterior modes and 95% credible intervals (CI) of models for alternative neophobia estimates: a binary  
25 variable (individual entered versus did not enter in the presence of the novel object) and a continuous latency to enter the nestbox, for individuals  
26 that entered in the presence of the novel object (see Appendix). Asterisks indicate estimates whose 95% CI do not encompass zero. For  
27 categorical fixed terms, estimates refer to the category indicated in parentheses.  $V_A$ ,  $V_{PE}$ ,  $V_M$  and  $V_F$  refer to the additive genetic, permanent  
28 environment, maternal and paternal variances, respectively. The residual variance  $V_\epsilon$  was set to 10.  $V_{\text{plot}}$  and  $V_{\text{obs}}$  refer to the variances associated  
29 with the plot and the observer identity, respectively.  $N$  is the sample size of the data set used in the model. Repeatability and heritability estimates  
30 are given both on the latent scale ( $R_{\text{latent}}$ ,  $h^2_{\text{latent}}$ ) and the observed scale ( $R_{\text{obs}}$ ,  $h^2_{\text{obs}}$ ) whenever these could be estimated.

31

32 **Table A5.** Output of the trivariate model fitting aggressiveness, boldness and neophobia scores with an individual random effect

33

	Posterior mode	95% CI	
<b>Fixed effects</b>			
Aggressiveness	1.73	[ 1.65 ; 1.82 ]	*
Boldness	7.97	[ 7.93 ; 8.02 ]	*
Neophobia	4.44	[ 3.79 ; 5.00 ]	*
Sex (male)	-0.02	[ -0.06 ; 0.04 ]	
Year (2012)	0.04	[ -0.02 ; 0.09 ]	
Year (2013)	0.09	[ 0.03 ; 0.15 ]	*
<b>Random effects</b>			
$V_{\text{ind, aggressiveness}}$	0.42	[ 0.16 ; 0.66 ]	*
$V_{\text{ind, boldness}}$	0.02	[ 0.00 ; 0.04 ]	
$V_{\text{ind, neophobia}}$	6.14	[ 2.96 ; 10.27 ]	*
$V_{\epsilon, \text{aggressiveness}}$	1.36	[ 1.09 ; 1.62 ]	*

$V_{\varepsilon, \text{boldness}}$	0.13	[ 0.12 ; 0.16 ]	*
$V_{\varepsilon, \text{neophobia}}$	10.00	[ 10.00 ; 10.00 ]	
$\text{COV}_{\text{ind}, \text{aggressiveness-neophobia}}$	0.00	[ -0.09 ; 0.11 ]	
$\text{COV}_{\text{ind}, \text{aggressiveness-boldness}}$	0.00	[ -0.01 ; 0.01 ]	
$\text{COV}_{\text{ind}, \text{boldness-neophobia}}$	0.00	[ -0.03 ; 0.02 ]	
$\text{COV}_{\varepsilon, \text{aggressiveness-neophobia}}$	-1.03	[ -1.37 ; -0.63 ]	*
$\text{COV}_{\varepsilon, \text{aggressiveness-boldness}}$	0.02	[ -0.02 ; 0.05 ]	
$\text{COV}_{\varepsilon, \text{boldness-neophobia}}$	-0.49	[ -0.62 ; -0.36 ]	*
$N$	1689		
Effective sample size:	>1740		

34

35  $V$  stands for variance terms and Cov for covariance terms (posterior mode and 95% credible interval (CI)). The individual effect was the only  
36 random term included; ‘ind’ and ‘ $\varepsilon$ ’ stand for between-individual and residual terms, respectively. The residual variance for neophobia score was  
37 fixed to 10 (see text for the distributions used for the three scores). Asterisks indicate estimates whose 95% CI do not encompass zero.

38 **Table A6.** Output of the trivariate model fitting aggressiveness, boldness and neophobia scores with an additive genetic effect

39

	Posterior mode	95% CI	
<b>Fixed effects</b>			
Aggressiveness	1.71	[ 1.62 ; 1.79 ]	*
Boldness	7.97	[ 7.93 ; 8.02 ]	*
Neophobia	4.12	[ 3.55 ; 4.62 ]	*
Sex (male)	-0.01	[ -0.05 ; 0.04 ]	
Year (2012)	0.04	[ -0.02 ; 0.09 ]	
Year(2013)	0.09	[ 0.03 ; 0.15 ]	*
<b>Random effects</b>			
$V_A$ , aggressiveness	0.15	[ 0.00 ; 0.38 ]	
$V_A$ , boldness	0.02	[ 0.00 ; 0.04 ]	
$V_A$ , neophobia	4.81	[ 2.07 ; 7.52 ]	*
$V_\epsilon$ , aggressiveness	1.46	[ 1.29 ; 1.78 ]	*



$V_{\epsilon, \text{boldness}}$	0.14	[ 0.12 ; 0.16 ]	*
$V_{\epsilon, \text{neophobia}}$	10.00	[ 10.00 ; 10.00 ]	
$\text{COV}_A, \text{aggressiveness-neophobia}$	0.00	[ -0.06 ; 0.07 ]	
$\text{COV}_A, \text{aggressiveness-boldness}$	0.00	[ 0.00 ; 0.00 ]	
$\text{COV}_A, \text{boldness-neophobia}$	0.00	[ -0.02 ; 0.02 ]	
$\text{COV}_{\epsilon}, \text{aggressiveness-neophobia}$	-1.01	[ -1.35 ; -0.64 ]	*
$\text{COV}_{\epsilon}, \text{aggressiveness-boldness}$	0.01	[ -0.02 ; 0.05 ]	
$\text{COV}_{\epsilon}, \text{boldness-neophobia}$	-0.48	[ -0.61 ; -0.37 ]	*
$N$	1686		
Effective sample size	> 1816		

40

41  $V$  stands for variance terms and Cov for covariance terms (posterior mode and 95% credible interval (CI)). The additive genetic effect was the  
42 only random term included; ‘A’ and ‘ $\epsilon$ ’ stand for additive genetic and residual terms, respectively. The residual variance for neophobia score was  
43 fixed to 10 (see text for the distributions used for the three scores). Asterisks indicate estimates whose 95% CI do not encompass zero. Positive  
44 additive genetic variances are found here because permanent environment and parental effects are not taken into account.

- 1 **Table A7.** Output of a principal component analysis of behaviours recorded during the
- 2 aggressiveness assays

	Coordinates		Contribution	
	PC1	PC2	PC1	PC2
Moves < 2 m from the nestbox	0.83	-0.12	36.12	0.95
Stationary flights towards the decoy	0.73	-0.21	27.74	2.99
Attacks towards the decoy	0.67	-0.24	23.24	4.06
Moves between 2 and 5 m from the nestbox	0.42	0.65	9.40	29.96
Moves between 5 and 10 m from the nestbox	-0.03	0.74	0.03	38.49
Chases of live birds	0.26	0.58	3.47	23.55

3

- 4 We used the function PCA from the ‘FactoMineR’ R package (Lê, Josse, & Husson, 2008).

5 **FIGURE CAPTIONS**

6

7 Figure A1. Sex differences in aggressiveness, boldness and neophobia scores (means  $\pm$  95%  
8 confidence interval). (a) Between-sexes differences in average aggressiveness score for a  
9 given individual in a given year depending on age (yearling versus older). (b) Between-sexes  
10 differences in the latency to return after human disturbance for the period without a novel  
11 object, as a proxy of (inverse) boldness, and for the period with the novel object, as a proxy of  
12 neophobia. Number of observations is indicated near each estimate.

13

14 Figure A2. (a) Aggressiveness and (b) boldness scores depending on the neophobia score  
15 (means  $\pm$  95% confidence interval). See text for the definitions of the scores. Aggressiveness  
16 is here the averaged value of all scores for a given individual in a given year. Number of  
17 observations is indicated near each estimate.


18

19 Figure A3. Distribution of the discrete scores for (a) aggressiveness, (b) boldness and (c)  
20 neophobia. Number of observations is indicated above each bar.


21

22 **Figure A1**

23



24 **Figure A2**



25

26 **Figure A3**

