



Corrigendum to

“Impact of rainfall spatial distribution on rainfall-runoff modelling efficiency and initial soil moisture conditions estimation” published in Nat. Hazards Earth Syst. Sci., 11, 157–170, 2011

Y. Tramblay¹, C. Bouvier¹, P.-A. Ayrat², and A. Marchandise³

¹Hydrosociences Montpellier, UMR5569, CNRS-IRD-UM1-UM2, Université Montpellier 2, Maison des Sciences de l’Eau, Place Eugène Bataillon, 34095 Montpellier Cedex 5, France

²Ecole Des Mines d’Alès, 6 Avenue de Clavières, 30319 Alès Cedex, France

³SCHAPI, 42 Avenue Gaspard Coriolis, 31 057 Toulouse Cedex 1, France

Correspondence to: Y. Tramblay (ytramblay@gmail.com)

An error occurred in the computation of the radar rainfall amounts, causing a +20 % bias on the computed amounts. All the radar rainfall values must therefore be corrected, and reduced by 20 %. Thus, the precipitations measured by the radar are in good agreement with the rain gauges (with only

+3 % in average for radar data). Consequently, the S parameter values obtained with radar data (UR or SR) are of the same order of magnitude as the S parameter values obtained with rain gauge data (UG or SG). Since the bias is constant, it does not imply modifications of the main conclusions of the paper.