



HAL
open science

Field-induced phase separation of a magnetic colloid under AC fields

Maxime Raboisson-Michel, J. Queiros Campos, Pavel Kuzhir, Grégory
Verger-Dubois

► **To cite this version:**

Maxime Raboisson-Michel, J. Queiros Campos, Pavel Kuzhir, Grégory Verger-Dubois. Field-induced phase separation of a magnetic colloid under AC fields. International Conference on Magnetic Fluids, 2019, Paris, France. hal-02426137

HAL Id: hal-02426137

<https://hal.science/hal-02426137v1>

Submitted on 1 Jan 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Field-induced phase separation of a magnetic colloid under AC fields

UNIVERSITÉ CÔTE D'AZUR

INPHYNI
INSTITUT DE PHYSIQUE DE NICE

Δxλσpιθζ
BIOMEDICAL

M. Raboisson-Michel^{1,2}, J. Campos Queiros¹, P. Kuzhir¹, G. Verger-Dubois²

¹Université Côte d'Azur, CNRS UMR 7010 Institut de Physique de Nice, Parc Valrose 06100 Nice, FRANCE

²Axlepios Biomedical, 1^{ère} avenue, 5^{ème} rue, 06510 Carros, FRANCE


Introduction

This work is devoted to the experimental study of kinetics of phase separation of a magnetic colloid under AC rotating magnetic fields. To this purpose, we use citrate-coated maghemite nanoparticles (medium size 8 nm) dispersed in distilled water and slightly destabilized addition of salts screening electrostatic repulsion between nanoparticles and leading to primary aggregates. In the presence of an AC unidirectional sinusoidal magnetic field (5-10 kA/m), the primary isotropic aggregates are attracted to each other and form long needle-shape aggregates of a typical length of 0.1-0.5 mm during a timescale related to a translational diffusivity of nanoparticles and the suspension supersaturation. Once they reach a maximal length related to homogeneity of the chemical potential across the suspension, some aggregates coalesce with each other in order to decrease the surface energy, but the coalescence is strongly slowed down by repulsive magnetic dipolar interactions, such that in practice the phase separation is not accomplished during a few hours. However, when a rotating sinusoidal magnetic field of the same amplitude is applied, the needle-like aggregates (appearing a few minutes after switching on the field) grow very quickly due to their collisions with free nanoparticles as they synchronously rotate with the field. We focus on fundamental understanding of this phenomenon tuned by Mason and Péclet numbers, as well as by magnetic nanoparticle concentration and suspension supersaturation. From the practical point of view, the obtained results open new perspectives for controlling the field-induced aggregation which may be applied in co-operative magnetophoresis, immuno-agglutination assays and magnetically assisted thrombolysis.


Methods

Experimental system


The system used for the experimental study consists of two nested Helmholtz coil pairs powered by a suitable alternating current to generate a rotating magnetic field and a microfluidic circuit within which the formation of rotating needle-like aggregates under rotating magnetic fields are observed.




Suspension of nanoparticles



Characterisation of Iron Oxide nanoparticles in solution through DLS. Particle volumic fraction of 0,16%, addition of NaCl with $[Na^+] = 0,350mM$ and $pH = 5,50$. Aggregation time without rotating field \gg Aggregation time with rotating field



Theory



Peclet's Number :

$$Pe \sim \frac{j_{conv}}{j_{diff}} = \frac{L \omega d_a}{D_p}$$

$$\square \text{ Particle flux } J = \phi_i \frac{dV_a}{dt}$$

$$\square \text{ Diffusive Boundary Layer Theory: } J \sim D_p Pe^{1/3} (\phi_\infty - \phi') L$$

$$\square \text{ Particle Volume Conservation: } \phi_\infty = \phi_0 - \phi_i n_a V_a$$

(number of agregates per unit volume n_a constant with time)


$$\square \text{ Agregate Volume: } V_a = \frac{\pi L^3}{6 r_a^2}$$

$$\square \text{ Minimum of free energy of agregate: } V_a \sim d_p^3 \frac{r_a^7}{ln^3 r_a}$$


$$\left. \begin{aligned} \bar{V}_a &\sim \frac{V_a}{V_m}, \bar{V}_m \sim \frac{\phi_0 - \phi'}{\phi_i n_a d_p^3} \\ \frac{d\bar{V}_a}{dt} &\sim \frac{\phi_0 - \phi'}{\phi_i \bar{V}_m^{1/3}} \bar{V}_a^{2/3} (1 - \bar{V}_a) \\ \bar{t} &= \frac{D_p^{2/3} \omega^{1/3}}{d_p^{4/3}} t \end{aligned} \right\} \text{ we find } \bar{V}_a(\bar{t})$$

Results



Sequence of shots showing kinetics of aggregation of nanoparticles with different rotating field frequencies




Phase diagram with NaCl constant, $[Na^+] = 350mM$ and $pH = 5,50$



Analysis of the size distribution at a given time, Time $t = 1s$, Time $t = 150s$, Time $t = 300s$



Experimental dependencies of the aggregate mean length (obtained from the size distribution) on the elapsed time.
(a) 4 raw experimental curves (dashed) and smoothed (continuous) for a frequency $f = 5Hz$,
(b) Length averaged over 4 measurements for each field frequency. The error bars are presented for $f = 5Hz$ and $f = 20Hz$. For (a) and (b) the amplitude of the field is $B_0 = 11mT$



Characteristic aggregation time:

$$t_{agr} = \frac{\phi_i V_{max}^{1/3} d_a^{1/3}}{D_p^{2/3} \omega^{1/3} (\phi_\infty - \phi')} \sim 100s$$

Conclusions

From the experiment:

At $f=5 Hz$: Aggregates longer than for any of $f > 5 Hz$ and a faster aggregation rate
Explanation: Largest aggregates size at low frequency for the hydrodynamic forces exerted on the aggregates are smaller as their rotation speed is low.
Extremely fast aggregates creation (1min) (length 50-80 μm) rotating with the field

From theory :

$$\frac{dL}{dt} \sim \frac{L_{max}}{t_{agr}} \sim L_{max}^{2/9} \omega^{1/3}$$

Aggregation speed is decreasing with the frequency as $L_{max}^{2/9}$ decreases faster than $\omega^{1/3}$ increases

Bibliography

Gabymo, J. L. E. Liu, D. W. Chang, M., & Lin, Y. H. "Controlled manipulation of Fe₃O₄ nanoparticles in an oscillating magnetic field for fast ablation of microchannel occlusion." *Nanoscale*, 7(9), 3947-3953, (2015).
Levich, V. G. *Physicochemical hydrodynamics*. Englewood Cliffs, N.J., Prentice-Hall, (1962).
Sandre, O., Browaes, J., Perzynski, R., Bacri, J. C., Cabuil, V., Rosenweig, R. E. "Assembly of microscopic highly magnetic droplets: Magnetic alignment versus viscous drag". *Physical Review E*, 59(2), 1736, (1999).
Adamczyk, Z. and Van de Ven, T. G. M. "Deposition of Brownian particles onto cylindrical collectors". *Journal of Colloid and Interface Science*, 84(2), 497-518, (1981).
Jeffery, G. B. "The motion of ellipsoidal particles immersed in a viscous fluid". *Proceedings of the Royal Society of London. Series A*, 102(715), 161-179, (1922).
Ezzaier, H., Alves Martins, J., Razvin, I., Abbas, M., Ben Haj Amara, A., Zubarev, A., & Kuzhir, P. "Two-stage kinetics of field-induced aggregation of medium-sized magnetic nanoparticles". *J.Chem.Phys.*146, 114902, (2017).