



Hair mercury levels in Amazonian populations: spatial distribution and trends

Flavia Barbieri, J. Gardon

► To cite this version:

Flavia Barbieri, J. Gardon. Hair mercury levels in Amazonian populations: spatial distribution and trends. *International Journal of Health Geographics*, 2009, 8 (1), pp.71. 10.1186/1476-072x-8-71 . hal-02429034

HAL Id: hal-02429034

<https://hal.science/hal-02429034>

Submitted on 3 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Review

Open Access

Hair mercury levels in Amazonian populations: spatial distribution and trends

Flavia L Barbieri*¹ and Jacques Gardon²

Address: ¹IRD - SELADIS, La Paz, Bolivia and ²IRD - HSM, La Paz, Bolivia

Email: Flavia L Barbieri* - flbarbieri@gmail.com; Jacques Gardon - jacques.gardon@ird.fr

* Corresponding author

Published: 21 December 2009

Received: 1 September 2009

Accepted: 21 December 2009

International Journal of Health Geographics 2009, **8**:71 doi:10.1186/1476-072X-8-71

This article is available from: <http://www.ij-healthgeographics.com/content/8/1/71>

© 2009 Barbieri and Gardon; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Mercury is present in the Amazonian aquatic environments from both natural and anthropogenic sources. As a consequence, many riverside populations are exposed to methylmercury, a highly toxic organic form of mercury, because of their intense fish consumption. Many studies have analysed this exposure from different approaches since the early nineties. This review aims to systematize the information in spatial distribution, comparing hair mercury levels by studied population and Amazonian river basin, looking for exposure trends.

Methods: The reviewed papers were selected from scientific databases and online libraries. We included studies with a direct measure of hair mercury concentrations in a sample size larger than 10 people, without considering the objectives, approach of the study or mercury speciation. The results are presented in tables and maps by river basin, displaying hair mercury levels and specifying the studied population and health impact, if any.

Results: The majority of the studies have been carried out in communities from the central Amazonian regions, particularly on the Tapajós River basin. The results seem quite variable; hair mercury means range from 1.1 to 34.2 µg/g. Most studies did not show any significant difference in hair mercury levels by gender or age. Overall, authors emphasized fish consumption frequency as the main risk factor of exposure. The most studied adverse health effect is by far the neurological performance, especially motricity. However, it is not possible to conclude on the relation between hair mercury levels and health impact in the Amazonian situation because of the relatively small number of studies.

Conclusions: Hair mercury levels in the Amazonian regions seem to be very heterogenic, depending on several factors. There is no obvious spatial trend and there are many areas that have never been studied. Taking into account the low mercury levels currently handled as acceptable, the majority of the Amazonian populations can be considered exposed to methylmercury contamination. The situation for many of these traditional communities is very complex because of their high dependence on fish nutrients. It remains difficult to conclude on the Public Health implication of mercury exposure in this context.

Background

Since the Minamata tragedy as well as the Basra incident, methylmercury contamination has been a source of concern worldwide. There has been extensive work attempting to assess the human health impact of mercury contamination through fish, seafood or marine mammal consumption, such as the Seychelles and Faroe Islands birth cohorts and the New Zealand study [1-8].

In the Amazonian aquatic environments, mercury is present in soils, water and food chains from complex sources [9-16]. On one hand, these soils have accumulated mercury naturally through time [9]. Human activities such as deforestation and agricultural land use can mobilize mercury from soils and vegetation [14,15,17,18]. Also, the use of metallic mercury in the gold mining process can contribute to mercury contamination in these areas [10-13,16].

According to specific methylation rates, mercury compounds in certain aquatic environments can be transformed into methylmercury, the most toxic mercury compound. This organic form of mercury is easily assimilated and accumulated into the food chains, with biomagnification along the trophic levels [10]. This is the main pathway for human exposure via fish consumption.

The most deleterious and studied health effects of methylmercury are neurological dysfunctions [19-22], especially from *in utero* exposure [4,19,23-29]. Also, immunotoxic and citotoxic damages have been shown [30-33], as well as cardiovascular deleterious effects [5,34-37]. It was not easy to reach a consensus regarding safe values for methylmercury exposure [38]. Overall, the tendency through time has been to lower the recommended level as much as possible, in order to minimize the health risk [39,40]. Based on *in utero* neurological development, the Joint FAO/WHO Expert Committee on Food Additives suggested a Benchmark Dose Limit (BMDL) of 14 µg/g of mercury in maternal hair, recommending a daily mercury intake lower than 1.5 µg/kg of body weight [41]. The Environmental Protection Agency (EPA) stated the Reference Dose (RfD) at 0.1 µg/kg/day, expecting no more than 11 µg/g of mercury in maternal hair [42]. Most researchers have handled for Amazonian populations a BMDL of 10 µg/g of hair mercury [36,43].

Amazonian riverside populations, which are amongst the highest fish consumers in the world, are exposed to mercury because of their alimentary habits. This situation has been widely studied in Brazil and French Guyana since the late eighties, showing quite complete information about the determinants of mercury exposure in these riverside populations [38,44-51]. However, there is an evident lack

of data in the rest of the Amazonian countries, where there have been a few isolated studies.

We aimed to systematize the available data and extract relevant information regarding hair mercury levels and observed health effects in Amazonian populations. We focused on the spatial distribution trends, in order to compare hair mercury levels by river basin and studied population.

Methods

We used scientific databases and online libraries such as PubMed, ISI Web of Knowledge, ScienceDirect, Springer-Link, SciELO, Horizon, selecting articles with "Mercury, methylmercury, Amazon, hair, human exposure, fish consumption" as main keywords. We included the websites of international public institutions such as the Environmental Protection Agency (EPA), the Agency for Toxic Substances and Disease Registry (ATSDR), the World Health Organization (WHO) and the Food and Agriculture Organization (FAO).

The articles were included in this review if a measure of the hair mercury concentration was presented from the general population or a target group of an Amazonian region. We also considered similar basins in Brazil such as the Tocantins River and Guyana Plateau, because of their geographic and ecological interfaces with the Amazon regions.

Exclusion criteria were sample sizes less than 10 people, and the absence of a measure of central tendency (mean, geometric mean or median). Also, we excluded studies where the hair mercury concentration presented was an estimate calculated from the mercury concentration in fish tissue and the frequency and amount of fish consumption. In the cases where we found two or more articles corresponding to the same study and in the same population, we chose one to be represented on the maps.

We presented the results from all the selected articles in tables, showing author, study sites, sample sizes, hair mercury levels and fish consumption measures when available. The tables were organized by regions and using the same continuity given by the maps, starting with studies from Andean Amazonian countries (Table 1) and French Guyana (Table 2). Brazilian studies are presented in the following tables, divided according to the studied population or group (Tables 3, 4, 5). The studies conducted in the Tapajós River basins were presented in independent tables, also separated according to the studied population or group (Tables 6, 7, 8, 9). Finally, the studies from the Madeira River basin were displayed separately in the last table (Table 10).

Table 1: Hair mercury levels in Andean Amazonian Countries

Studied population	Study	N	Hg mean $\mu\text{g/g}$	Range	> 10 $\mu\text{g/g}^a$	Fish
Country, River Basin			(median) [SD]	$\mu\text{g/g}$		
General population						
Barbieri et al. 2009 Bolivia, Beni River	Cachuela Esperanza	150	3.76 (3.01) [2.52]	0.42-15.65	3%	10.5 meals/week
Monrroy et al. 2008 Bolivia, Beni River	Upper Beni River	556	5.3 (4.0) [4.3]	0.08-34.	$\approx 14.0\%$	
	Children	393	5.2 (3.9) [4.4]	0.08-34.1		
	Mothers	163	5.5 (4.4) [4.1]	0.15-20.0		
	Pregnant women	18	3.2 (3.3) [2.1]	0.2-7.8		
	Breastfeeding (BF)	57	6.2 (5.5) [4.1]	0.5-18.3		
	Non pregnant, non BF	93	5.4 (4.1) [4.1]	0.15-20.0		
Maurice-Bourgoin et al. 2000 Bolivia, Beni River	Rurrenabaque	80	9.8	4.3-19.5		
	Esse-Ejjas indigenous	37				
Webb et al. 2004 Ecuador, Napo River	Coca	45	1.9(1.5)	0.03-10.0		7.5 meals/month
	Añangu	27	8.7 (7.8)	2.2-20.5		17.2 meals/month
	Pañacocha	27	5.3 (5.0)	1.5-13.6		33.9 meals/month
Children						
Counter et al. 2005 Ecuador, Nambija River	Nambija gold-mining settlement	80	2.8 ^b (2.0) [17.5]	1.0-135.0	<10%	

^aPercentage of the population with hair mercury levels higher than 10 $\mu\text{g/g}$. ^bGeometric mean.

In the cases when an article studied populations from different river basins, we presented them separately, in the corresponding basin tables.

For the design of the maps, we located each Amazonian study site using their longitude and latitude data when available in the article. Otherwise, we searched longitude and latitude data in geographic databases and using Google Earth, which allowed us to find and successfully locate the majority of the study sites. When it was not possible to identify the coordinates, we used the maps and/or site descriptions presented by the authors. We marked and named each study site on the map displaying a representation of the hair mercury levels found in this population in a six colour scale. Each individual square on the map represents a hair mercury measure and the target group used for the study, as well as the reference number. We prepared separated maps for the Madeira River, the Tapajós River and French Guiana. We also designed a map illustrating the studies that assessed health outcomes in relation to hair mercury levels in the studied populations, using a three colour scale for the different results.

Results and Discussion

We found 58 articles meeting our criteria for the elaboration of the maps. The majority of the studies were carried out in Brazil (86%), while there are only 3 studies in French Guyana, 3 in Bolivia and 2 in Ecuador (Figures 1, 2). In Brazil, 30 studies are focused on the Tapajós River basin (Figure 3), 10 on the Madeira River basin (Figure 4) and the remaining 20% of these Brazilian studies are from other basins, such as the Negro River, the Tocantins River or the Xingu River (Figure 1).

The approaches and strategies are quite variable between studies. Several studies have used a large sample size, selected randomly from the general population (Tables 1, 2, 3, 6, 10), and usually comparing two or three sites from the same basin. Some of them used smaller sample sizes, accepting the variability induced by the sampling fluctuation in order to extend to several populations geographically close to each other. In the Madeira River (Figure 4; Table 10), the study by Bastos et al. (2006) covered from the upper basin (near the Bolivian border) to a point downstream on the Amazon River [52]. Even if some of the populations had sample sizes of less than 10 people,

Table 2: Hair mercury levels in French Guyana

Studied population	River basin	Location	N	Hg mean $\mu\text{g/g}$ (median) [SD]	Range $\mu\text{g/g}$	> 10 $\mu\text{g/g}$ ^a	Fish consumption
General population							
BASAG 2007	Maroni River	Sinnamary	285	1.8		5%	
	Oyapok River	Lower Maroni River	740	1.7-3.6		2.4%	
		Lower Oyapok River	144	1.5-3.4		1 person	
		Upper Oyapok River	181	4.6-7.2		7.6-18.2%	
Cordier et al. 1998	Maroni River Oyapok River	11 health centers					
		Adults	255	3.4 ^b	0.2-22.0	12.2%	
		Pregnant women	109	1.6 ^b	0.2-13.0	4.6%	
		Children	136	2.5 ^b	0.2-31.0	11.8	
Fréry et al. 2001	Maroni River	Cayodé, Twenké, Taluhen and Antécume-Pata	235	11.4 [4.2]	1.9-27.2	57.4%	20-317 g/day
Mothers and their infants							
Cordier et al. 2002	Maroni River Oyapok River	Upper Maroni River					
		Children	156	10.2 ^b		79%	2 meals/day
		Mothers	90	12.7 ^b			
		Oyapok River					
		Children	69	6.5 ^b			
		Mothers	63	6.7 ^b			
		Lower Maroni River					
		Children	153	1.4 ^b			
		Mothers	77	2.8 ^b			

^aPercentage of the population with hair mercury levels higher than 10 $\mu\text{g/g}$. ^bGeometric mean.

the locations evaluated were 44, distributed along the largest part of the Madeira basin (Figure 4).

In French Guiana the studies have used both approaches, covering almost the totality of the Maroni and Oyapok River basins, with large sample sizes from the general population (Figure 2; Table 2) [47,53,54].

Hair mercury levels in the Amazon

In the Andean Amazonian regions (Figure 1; Table 1), mercury levels were found below 10 $\mu\text{g/g}$ in Ecuador [55,56] and Bolivia [57-59].

In French Guiana (Figure 2; Table 2), most study sites showed hair mercury levels below 10 $\mu\text{g/g}$ in the Oyapok River and the lower Maroni River basins [53,60]. The only

populations with mercury levels above 10 $\mu\text{g/g}$ were located in an Amerindian reservation in the upper Maroni River basin [47,53,54].

In Brazilian Amazonia (Figures 1, 3, 4; Tables 3, 4, 5) hair mercury levels are very variable between basins and study sites. Without considering the studies from the Tapajós and Madeira Rivers, which will be explained in detail below, the highest mercury levels can be found in the Negro River basin [61,62] and at the lakes from the Amapá State (Figure 1; Tables 3, 4, 5) [63], with means above 20 $\mu\text{g/g}$ and even higher than 25 $\mu\text{g/g}$. These high values can also be found in two other sites from the Xingu and Tocantins Rivers, both from Amerindian reservations (Figure 1) [51,64-68]. These levels can be considered of a high risk for the populations and would merit further

Table 3: Hair mercury levels in Brazilian River Basins, General Population

Study River Basin	Location	N	Hg mean $\mu\text{g/g}$ (median) [SD]	Range $\mu\text{g/g}$	> 10 $\mu\text{g/g}$ ^a	Fish consumption
Akagi et al. 1995 Araguari River	Três Bocas	11	28.1	8.4-54.0		
Castro et al. 1991 Branco River	Surucucus, Paapiu and Mujacai areas	162	3.61	1.4-8.1		
Forsberg et al. 1995 Negro River	Various sites between Marie and Paduari Rivers	154	75.5 [35.2]	5.8-171.2		
Kehrig et al. 1998 Negro River	Balbina Reservoir	53	6.5 [5.4]	1.2-22.0		110 g
Leino & Lodenius 1995 Tocantins River	Tucuruí area	125	35.0 (29.0)	0.9-240.0		11 meals/week
Pinheiro, Nakanishi et al. 2000 Tocantins River	Belém	13	2.0			
Santos, Camara et al. 2002 Amazon River	Caxiuanã	214	8.6 [6.3]	0.6-46.0		12.3 meals/week
Santos et al. 2003 Mamoré River	Lower Mamoré: Pakaás Novos Indigenous Areas	910	8.4 (6.9) [6.4]	0.5-83.9		
Soares et al. 2002 Mamoré River	Doutor Tanajura	13	(6.1)	1.4-11.7		
Vasconcellos et al. 1994 Xingu River	Indigenous Xingu Park	27	18.5 (18) [5.9]	6.9-34.0		
	Billings Dam	28	0.9 (0.7) [0.7]	0.3-3.0		
	Controls	25	1.1 (1.0) [0.6]	0.3-2.5		
Vasconcellos et al. 2000 Xingu River	Xingu Park (13 groups)			1.2-57.3		
	Highest values		21.8 (20.8) [6.1]			
	Lowest values		3.6 (2.6) [2.4]			

The Tapajós and Madeira Rivers basins are not displayed in this table (see Tables 5, 6, 7, 8, 9, 10). ^aPercentage of the population with hair mercury levels higher than 10 $\mu\text{g/g}$.

investigations about the health impact of this exposure. The rest of these study sites show hair mercury levels below 10 $\mu\text{g/g}$, with the exception of an Amerindian reservation on the Mamoré River (13.1 $\mu\text{g/g}$) [69].

Tapajós River basin

In the Tapajós River basin (Figure 3; Tables 6, 7, 8, 9) there is a wide difference in mercury levels between the various populations. Most locations present hair mercury levels

above 10 $\mu\text{g/g}$, especially in some populations such as Rainha, Barreiras, Brasília Legal or São Luís do Tapajós [34,70-73], where exposure levels seem to be alarming, even reaching a mean over 30 $\mu\text{g/g}$ in the Apiacás Reservation study [74]. In very few sites the hair mercury levels remained below 5 $\mu\text{g/g}$ [75,76]. The populations living in urban or suburban areas (Santarém and Santana do Itiquí) were used as control populations by the authors and showed lower exposure levels attributed to their food

Table 4: Mercury levels in Brazilian River Basins, Target Groups

Studied population River Basin	Location	N	Hg mean $\mu\text{g/g}$ (median) [SD]	Range $\mu\text{g/g}$	> 10 $\mu\text{g/g}$ ^a	Fish consumption
Adults						
Barbosa et al. 2001 Negro River	Negro River shores	76	21.4 (17.8) [12.7]	1.7-59.0	79%	
	Men	17	26.2 [13.7]			
	Women	31	18.3 [11.1]			
Pinheiro et al. 2006	Panacauera	22	≈ 7.0			
Tocantins River	Pindobal Grande	43	≈ 3.0			
Silva et al. 2004 Amazon River, lakes	Tabatinga	98	(6.4)	1.2-17.0		
Yokoo et al. 2003 Cuiabá River	Pantanal region	129	4.2 (3.7) [2.4]	0.6-13.6		
Children						
Barbosa et al. 2001 Negro River	Negro River shores	73	18.5 (16.4) [10.0]	0.5-45.9	79%	
Pinheiro et al. 2006 Tocantins River	Pindobal Grande	88	≈ 3.0			
Pinheiro et al. 2007 Tocantins River	Panacauera	36	2.3	0.4-9.5	0%	
Santos-Filho et al. 1993 Cubatão River	Cubatão Municipality	217	0.8 [0.5]	0.2-3.0	0%	
Tavares et al. 2005 Cuiabá River	Barão de Melgaço	114	2.1 (1.8) [1.4]	0.4-7.6	0%	4.6 meals/week
	Riverside communities	72	5.4 (4.7) [3.4]	0.6-17.1		7.8 meals/week
Women						
Pinheiro et al. 2008 Tocantins River	Panacauera	20	3.3	1.3-6.0	0%	
Mothers and their infants						
Barbosa et al. 1998 Fresco River (Xingu basin)	Garimpo Maria Bonita Mothers	28	8.1 (8.3) [3.2]	0.8-13.7		
	Infants	54	7.3 (6.6) [3.5]	2.0-20.4		

The Tapajós and Madeira Rivers basins are not displayed in this table (see Tables 5, 6, 7, 8, 9, 10). ^aPercentage of the population with hair mercury levels higher than 10 $\mu\text{g/g}$.

diversification [33,77-79]. In the same way, some authors used study sites from different river basins to compare their exposure levels, such as Panacuera or the city of Belém (Figure 1) [65,73].

Unexpectedly, studies carried out in the same site and target group obtained quite different results. For example, in Jacareacanga the studies on the general population published in 1995 [77,80] showed much higher levels (means of 16.6 and 25.0 µg/g) than those carried out in the following decade (mean = 8.6 µg/g and median = 8.0 µg/g) [33,81]. The sample sizes and study designs were very different. In 1995, Jacareacanga was a village with approximately 3000 inhabitants, and the two studies published that year took samples of 10 and 48 people. In 2002, the population of Jacareacanga was around 2000 inhabitants, and the studies published in 2002 and 2004 took samples of 140 and 205 people. Other than that, there is not enough information to explain the observed difference in hair mercury levels. There is no evidence of a change in the fish consumption frequency, deforestation, colonization, agricultural land use or gold mining activities.

Madeira River basin

In the Madeira River basin (Figure 4; Table 10) most study sites showed hair mercury levels between 11 µg/g and 15 µg/g [52,82-85]. A couple of sites (very close to each other) presented hair mercury levels above 20 µg/g [52]. A few sites showed levels between 16 µg/g and 20 µg/g and the others presented hair mercury levels under 10 µg/g [52,82,83,85]. This shows a mercury exposure distribution that appears to be less heterogenic than the Tapajós River basin. The higher mercury levels seem to be found in small isolated villages on the middle of the basin, downstream from Humaitá, near Manicoré (Figure 4).

Mercury levels and fish consumption

In the Brazilian Amazon, many studies include especially fish consumption in the population, even though not all of them show these results (Tables 1, 2, 3, 4, 5, 6, 7, 8, 9, 10). Fish consumption is not measured in a uniform man-

ner in all the studies. Each research team chose their indicators according to their objectives, so this measure can be found in grams per day, percentage of meals composed by fish, meals per week or per day and/or times per week. No matter which indicator for fish consumption the authors chose, there was always a positive relation between fish consumption and hair mercury levels [19,46,47,49,64,75,76]. Most of these study sites are usually small and traditional riverside villages, some of them Amerindian reservations, without the proper roads connecting them to larger villages and cities. It is possible to hypothesize that this situation makes those populations more dependent on fish as source of protein intake.

In spite of those considerations, some populations with high fish consumption showed hair mercury levels below 10 µg/g, even if there was a significant relation with fish consumption frequency [55-57,59,78,79]. In those particular six studies, it is important to consider the different geographical and socio-economic situation those populations are in: three of those studies were conducted in the Andean piedmont Amazonian regions, the two Ecuadorian studies [55,56] and one of the Bolivian studies [57]. Also, those populations are not solely dependent on fish because they are also farmers or hunters, even if fish remains the most important source of nutrients. The Bolivian population of Cachuela Esperanza, located near the Brazilian border, consumes large amounts of fish (10 fish meals per week), but only during the dry season, preferring game meat during the rainy season, which lasts from October to April [57].

The other two cases with intense fish consumption and low hair mercury levels are the studies conducted in Aldeia do Lago Grande, Santana do Ituí and Vila do Tabatinga. These communities are located on the Amazon River shore, near the confluence with the Tapajós River. They are small but not isolated because they are connected by roads to the cities of Monte Alegre, Santarém and Juruti. The low hair mercury levels cannot be explained by food diversification, because these populations consume

Table 5: Hair mercury levels in Brazilian River Basins, Occupational Groups

Study	Location	N	Hg mean µg/g	Range	> 10 µg/g ^a	Fish consumption
River Basin			(median) [SD]			
Guimarães et al 1999	Pracuúba Lake	15	16.7			
Tartarugal Grande River (Fishermen)	Duas Bocas Lake	15	28.0		87%	14 meals/week (200 g per day)
	Groups					
Palheta & Taylor 1995	Garimpeiros	20		0.4-32.		
Gurupi River	Cachoeira Villagers	5		0.8-4.6		
	River dwellers	10		0.2-15		

The Tapajós and Madeira Rivers basins are not displayed in this table (see Tables 5, 6, 7, 8, 9, 10). ^aPercentage of the population with hair mercury levels higher than 10 µg/g.

Table 6: Hair mercury levels in the Tapajós River Basin, General Population

Study	Location	N	Hg mean $\mu\text{g/g}$ (median) [SD]	Range $\mu\text{g/g}$	> 10 $\mu\text{g/g}$ ^a	Fish consumption
Akagi et al. 1995	Rainha	11	15.8	2.4-31.0		
	Brasília Legal	56	22.6	3.5-151.0		
	Ponta de Pedra	10	10.2	6.2-12.6		
	Jacareacanga	48	16.6	1.5-46.0		
Barbosa et al. 1997	Apiacás Reservation	55	34.2 (42.8)	?-128	93%	≈ 6 times/week
Crompton et al. 2002	Jacareacanga	205	8.6	0.3-83.2		
Dorea et al. 2005	Kaburuá	89	2.5 [1.4]			22 g/day
	Cururu Mission	138	3.7 [1.6]			32 g/day
	Terra Preta	22	6.0 [2.9]			52 g/day
	Kayabi	47	12.8 [7.0]			110 g/day
Malm et al. 1995	Jacareacanga	10	25.0	5.7-52.0		
	Brasília Legal	13-29	26.0	4.7-151.0		
	Ponta de Pedra	4-26	12.0			
	Santarem	11	2.7			
Pinheiro, Guimarães et al. 2000	Rainha	29	17.2			
	Barreiras	111	18.9			
	São Luís do Tapajós	30	25.3			
	Paraná-Mirim	21	9.2			
Pinheiro, Nakanishi et al. 2000	Rainha	29	17.6			
	Barreiras	78	19.1			
Santos et al. 2000	Brasília Legal	220	11.8 [8.0]	0.5-50.0		10 meals/week
	São Luís do Tapajós	327	19.9 [12.0]	0.1-94.5		13 meals/week
	Santana do Itaquí	321	4.3 [1.9]	0.4-11.6		13 meals/week
Santos, Camara et al 2002	Santana do Itaquí	321	4.3 [2.2]	0.4-12.0		12.7 meals/week
	Aldeia do Lago Grande	316	4.0 [2.1]	0.4-12.0		12.0 meals/week
	Vila do Tabatinga	499	5.4 [3.1]	0.4-17.0		10.5 meals/week
Santos, de Jesus et al. 2002	Sai Cinza	324	16.0 [18.9]	4.5-90.4		
Silva et al. 2004	Jacareacanga	140	(8.0)	0.3-58.5		
	Rio-Rato	98		0.01-81.4		

^a Percentage of the population with hair mercury levels higher than 10 $\mu\text{g/g}$.

10 to 13 fish meals per week. However, the fish they consume is collected from local lakes and minor rivers, and the mercury concentrations in fish tissue was found to be much lower than in other exposed populations [52,78,79].

On the Madeira River basin, where most hair mercury levels remained under 10 $\mu\text{g/g}$, the fish consumption in the majority of the populations was around seven meals per week (Table 10). On the contrary, on the Tapajós River basin the fish consumption in the majority of the populations was higher than 10 meals per week (Table 6).

Some studies measured mercury concentrations in fish tissue, finding positive relations between fish mercury concentrations, fish consumption and human hair mercury levels [13,75,77,80,86].

Studied populations

Due to the importance of mercury exposure and its health effects in children, as well as *in utero* exposure, many studies have chosen fertile age women and children as target groups (Tables 1, 2, 4, 8, 10). These studies are generally consistent with studies on the general population, with comparable hair mercury levels. When studied together as mother-infant pairs, there was always a strong relation

Table 7: Hair mercury levels in the Tapajós River Basin, Adults

Study	Location	N	Hg mean $\mu\text{g/g}$ (median) [SD]	Range $\mu\text{g/g}$	> 10 $\mu\text{g/g}$ ^a	Fish consumption
Amorim et al. 2000	Brasília Legal Women Men	98	(13.5) (10.8) (17.1)	0.6-71.8	>50%	
Dolbec et al. 2000	Cametá Women Men	68 41 27	10.8 (9.0) [6.1] 9.9 (8.0) [5.6] 12.2 (10.8) [6.8]		>25% >50%	61.8% of total meals
Fillion et al. 2006	São Luís do Tapajós, Nova Canaã, Santo Antônio, Mussum, Vista Alegre and Açaituba	251	17.8	0.2-77.2	69.7%	6.8 meals/week
Lebel et al. 1997	Brasília Legal Women Men	96	(12.9) (11.2) (15.7)		>50%	44.7% of total meals 43.9% of total meals
Lebel et al. 1998	Brasília Legal Men Women	34 46	14.3 [9.4] 12.6 [7.0]			
Passos et al. 2007	SLTapajós, Nova Canaã, Santo Antônio, Ipaupixuna, Novo Paraíso, Teça, Timbu, Açaituba, Campo Alegre, Samauma, Mussum, Vista Alegre and Santa Cruz	457	16.8 (15.7) [10.3]	0.2-58.3	>50%	6.6 meals/week
Pinheiro et al. 2006	São Luís do Tapajós Barreiras	32 37	\approx 15.0 \approx 15.5			
Silva et al. 2004	Vila do Tabatinga	98	(6.4)	1.2-17.0		

^aPercentage of the population with hair mercury levels higher than 10 $\mu\text{g/g}$.

between maternal hair mercury and the exposure in their infants; the mothers always had higher mercury levels than their infants [26,60,82,83,85,87].

The majority of the studies did not find any significant relationship between age and hair mercury levels [13,30,33,44,48,61,66,77,78,85,88-92]. Some studies found that hair mercury levels increase with age [74,78,79,81,84]. Nevertheless, a couple of studies found the opposite results, showing higher hair mercury levels in younger people, children and infants [58,69].

Several studies found that men have higher hair mercury levels than women [19,30,34,61,65,69,71,74,84,87]. In two articles from the Tapajós River basin, only the fishermen had hair mercury levels significantly higher than the women, while the other men from the same village presented hair mercury levels similar to those of the women (Tables 7, 8). Those differences corresponded to their fish consumption, which was also significantly higher for the fishermen [20,49]. Only one study found women to have higher mercury levels than men [69].

No study revealed physiological basis that could lead to hypothesize a difference between female and male mercury metabolism. In fact, most studies (78%) did not find any significant relation between mercury exposure and gender.

A few studies focused their attention on fishermen, because of their obvious access to fish as main nutrient [49,63,70]. Their hair mercury means ranged from 16 to more than 25 $\mu\text{g/g}$, regardless of the river basin. For instance, the studies conducted by Lebel et al. (1997, 1998) compared fishermen and other adults from the Brasília Legal population. Both men and women had mercury levels lower than 15 $\mu\text{g/g}$, while fishermen presented levels of 27.3 $\mu\text{g/g}$ and 23.9 $\mu\text{g/g}$ (Figure 3; Tables 7, 8) [20,49].

The particular case of gold miners (called *garimpeiros* in Brazil and Bolivia) has been documented in a couple of studies. It is important to remark that occupational mercury exposure is very different from the environmental exposure. Artisanal *garimpeiros* are occupationally exposed to metallic mercury vapours, which are rapidly trans-

Table 8: Hair mercury levels in the Tapajós River Basin Women and Children

Studied population	Location	N	Hg mean $\mu\text{g/g}$ (median) [SD]	Range $\mu\text{g/g}$	> 10 $\mu\text{g/g}$ ^a	Fish consumption
Mothers and their Children						
Grandjean et al. 1999	Mothers	114	11.6 (14.0)			
	Children		11.0 (12.8)	0.5-83.5		2 meals/day
	Brasília Legal	76	11.9	0.7-35.8	76%	
	São Luís do Tapajós	71	25.4	0.6-83.5	91%	
	Sai-Cinza	87	17.7	7.3-63.8	92%	
	Santana do Itaquí	105	3.8	0.5-12.4	2%	
Children						
Barbosa et al. 1997	Apiacás Reservation	28	29		86%	
Dorea, Barbosa et al. 2005	Kaburuá	77	2.9 [2.1]			
	Cururu Mission	86	4.8 [2.1]			
	Kayabi	40	16.6 [11.4]			
Pinheiro et al. 2007	São Luís do Tapajós	48	10.9	1.3-53.8	52%	
	Barreiras	84	6.1	1.4-23.6	21%	
Women						
Barbosa et al. 1997	Apiacás Reservation	13	41.2		100%	
Dolbec et al. 2001	Cametá	98	(12.5)	2.9-27.0	>50%	
Hacon et al. 2000	Alta Floresta	75	1.12 [1.2]	0.05-8.2	0%	8-20 g/day
Passos et al. 2003	Brasília Legal	26	10.0 (9.1)	4.0-20.0	≈ 50%	8 meals/week
Pinheiro et al. 2007	São Luís do Tapajós					5-14 meals/week
	Rainha, and Barreiras					
	Pregnant	19	8.2	1.5-19.4	37%	
	Non-pregnant	21	9.4	5.2-21.0	28%	
Pinheiro et al. 2005	São Luís do Tapajós	28	13.7	3.2-30.04	36%	
	Barreiras	39	12.1	3.04-33.4	38%	

^aPercentage of the population with hair mercury levels higher than 10 $\mu\text{g/g}$.

formed in the human body into inorganic mercury, best measured in urine or plasma, while total hair mercury corresponds mainly to methylmercury exposure in high fish eating populations [36,93]. Generally, the authors found that *garimpeiros* had lower hair mercury levels than the *ribeirinhos* [58,94], concluding that because of their better economical situation, they were able to diversify their food, consuming less fish than the rest of the population [13]. However, in a few studies the *garimpeiros* or gold miners had higher total mercury levels in hair than the general population, without apparent relation with fish consumption frequency [45,57,95,96]. Consistently, in those three studies the hair mercury means in the general population were lower than those found in most central

Amazonian studies. In both the Bolivian study and the Colombian study, hair mercury means remained below 6 $\mu\text{g/g}$, even in the most exposed groups [57,95]. It would be possible to assume that metallic mercury exposure can acquire significance in hair mercury levels in some specific situations of low methylmercury exposure. However, considering the differences between those studies and regions, and without hair mercury speciation, it is not possible to reach valid conclusions.

Multidisciplinary studies

Five studies carried out a multidisciplinary approach, measuring as well mercury in air, water, sediments, fish and human samples [13,52,58,63,96]. In those studies,

Table 9: Hair mercury levels in the Tapajós River Basin, Occupational Groups

Study	Locations and groups	N	Hg mean $\mu\text{g/g}$ (median) [SD]	Range $\mu\text{g/g}$	> 10 $\mu\text{g/g}^a$	Fish consumption
Harada et al. 2001	Fishermen and families					
	Barreiras	76	16.4 [10.6]	1.8-53.8	75%	
	Rainha	12	14.1 [9.3]	3.1-34.5	67%	
	São Luís do Tapajós	44	20.8 [10.6]	5.1-42.2	86%	
Lebel et al. 1997	Brasília Legal Fishermen	14	(27.3)		> 50%	68.8% of total meals
Lebel et al. 1998	Brasília Legal Fishermen	11	23.9 [9.3]			
Santos, de Jesus et al. 2002	Sai Cinza	324	16.0 [18.9]	4.5-90.4		
	Agriculture	127	17.3	6.8-90.4		
	Gold mining	6	13.8	10.7-18.5		
	Both of the above	25	13.8	6.9-25.7		
	Others	11	12.3	6.6-20.6		
	Children up to 6 y/o	93	16.8	4.5-66.6		
	Students	38	15.1	9.0-38.7		
	Without information	24	17.0	9.6-30.8		

^aPercentage of the population with hair mercury levels higher than 10 $\mu\text{g/g}$.

Table 10: Hair mercury levels in the Madeira River Basin

Studied population	Location	N	Hg mean $\mu\text{g/g}$ (median) [SD]	Range $\mu\text{g/g}$	> 10 $\mu\text{g/g}^a$	Fish consumption
General population						
Bastos et al. 2006	Various populations along the Madeira River	713	15.2 (12.5) [9.6]	0.36-150.0	>50%	7 meals/week
Boischio & Barbosa 1993	Near Porto Velho	311	\approx (10)	?-303.1	51%	200 g/day ^b
Mothers and their infants						
Barbosa & Dorea 1998	Near Porto Velho Mothers Infants	98	14.1 (12.8) [10.7]	2.6-94.7	>50%	
		71	10.8 (7.8) [8.5]	0.8-44.4		
Boschio & Cernichiari 1998	Near Porto Velho Mothers Infants	12		4.0-41.0		
		12		8.2-50.4		
Boischio & Henshel 2000	Near Porto Velho Mothers Infants	90	12.6 [6.5]	15.0-45.0		
		89	10.2 [7.2]	1.0-34.2		
Marques et al. 2007	Porto Velho city Mothers Infants	82	(5.4)	0.4-62.4		1 meal/week
		82	(1.8)	0.02-32.9		

^aPercentage of the population with hair mercury levels higher than 10 $\mu\text{g/g}$. ^bFrom the Letter to the Editor Boischio Environ Research 2000; 82:91-92




Figure 1
Hair mercury levels in the Amazon River basin.

human exposure seemed to be considered as a part of the totality of the environmental contamination. Therefore, those hair mercury levels were useful as a reference value for those specific study sites, and need to be interpreted in the whole context rather than as a Public Health assessment. These studies showed in a precise manner the interactions within various environmental components, showing a direct relation between mercury concentrations at different levels of the ecosystem and the food chains, including the human being as the last receptor of this pollution.

An example of an ecosystem approach is the CARUSO project [14,15,17-20,30,34,49,71,88,97-100]. This project started in 1994, and consists on a series of studies conducted by research teams of different disciplines, attempting to assess the various aspects of mercury pollution and human exposure in the Tapajós River basin.

Hair mercury levels and health impact

There were 15 studies conducted in Brazil and one in French Guyana (Figure 5). Most of them (73%) focused on neurological effects.




Figure 2
Hair mercury levels in French Guyana.

In the Amazonian context it is not easy to identify a clear relation between hair mercury levels and patent neurological abnormalities. Observing the spatial distribution of the studies, we can see that in populations with low hair mercury levels, such as Santana do Itaquí, Vila do Tabatinga or the city of Porto Velho, it was not possible to confirm a relation between mercury levels and neurological performance [26,78,79,101]. Also, in the populations with higher mercury exposure, the authors found an impact on the nervous system, especially motricity [19,20,60].

However, in São Luís do Tapajós or Brasília Legal the information is difficult to interpret. There were two studies assessing neurotoxicity in the general population and one in adults. While one of those studies documented cases of mild Minamata disease [70], the other did not find any relation between hair mercury levels and neurological outcomes [79]. On the contrary, the study con-

ducted in adults found a dose-dependent relation between hair mercury levels and motor abnormalities [20]. There was also a study that found an association between exposure and psychomotor performance in children, but the authors concluded that the outcomes observed were also influenced by socioeconomic factors, such as maternal educational level or nutritional status [87].

Two studies have evaluated the impact of mercury exposure on the immune system *in vitro*. The authors found positive relations between mercury levels and the presentation of auto-antibodies [33] and cytogenetic abnormalities in peripheral lymphocytes [30], especially in the most exposed Amazonian populations.

There are a couple of studies about mercury exposure and blood pressure on the Tapajós River basin. One of them supports a negative impact of mercury exposure on the cardiovascular system [34], while the other did not find a significant relation between mercury levels and blood pressure [76]. This second study was conducted in four Amerindian populations located on rivers tributaries to the Tapajós, with low hair mercury levels. The authors found that the community with higher hair mercury levels (12.8 µg/g) seemed to be protected against age-related increase of blood pressure, because of their intense fish consumption [76].

Two studies were focused on children's growth. One of these studies was carried out in an Amerindian community from the Tapajós River [75]. No significant correlation was found between growth or nutritional status and hair mercury levels. On the contrary, a study in the Bolivian Amazonia found a positive relation between hair mercury levels and a better nutritional status in children ranging from 5 to 10 years of age [102]. Considering the characteristics of that population, the authors hypothesized that the relation observed was caused by the nutritional contents in fish, mostly important at that age group. Furthermore, the same study found the opposite result in the mothers, with worse nutritional indices in the women with higher mercury levels. In fact, while some authors worry about the health impact of mercury exposure in high fish eating populations [38,99,103], others consider that the nutritional value of fish probably compensate an exposure to low mercury levels [46,75,102,104,105]. In absence of large cohort studies, as the ones developed in the Faeroe Islands and Seychelles, the issue of health impact in the Amazonian context still remains open for debate.

Other interactions

Interest has recently been focused on the relation between mercury exposure and antioxidant defences, suggesting




Figure 3
Hair mercury levels in the Tapajós River basin, Brazil.




Figure 4
Hair mercury levels in the Madeira River basin, Brazil.

that long term mercury exposure induced a depletion in the antioxidant enzymatic activity [67].

Interactions between fish and fruit consumption were studied in a population from the Tapajós River, finding an inverse relationship between fruit consumption and Hg levels [71,100]. The authors recommended further investigation to establish new nutritional strategies aiming to

limit mercury exposure while maintaining fish consumption.

Recently, some studies also emphasized the interactions between mercury and selenium from dietary sources, especially fish consumption [68,90,92,98]. The authors generally found high concentrations of selenium in the populations exposed to methylmercury, with molar ratios




Figure 5
Hair mercury levels and health impact in the Amazon River basin.

consistently close to 1. Based on their observations, the authors suggest the need of further research regarding the protective role of selenium against mercury toxicity.

Variability in exposure assessment

As seen on the maps and tables, there is a considerable spatial and temporal variability in the measure of mercury exposure. Fish consumption frequency alone does not explain the differences in hair mercury levels in some study sites. There are several factors involved: the characteristics of the soils, methylation rates in different aquatic environments, the food chain structures, specific human habits and diets, social and economic characteristics, mercury interactions with other elements, and also strong local variations without a clear explanation. Moreover, many of these small populations are subjects to intense changes through time. Human migration, land use and deforestation are usually associated with erosion peaks, which could temporarily mobilize mercury from the soils [14,15,17,18].

It is also possible to attribute some of the variations observed to the use of hair as biomarker. It has been stated that methylmercury corresponds to more than 90% of total hair mercury [19,93,106]. Therefore, total mercury in hair has been widely accepted as a well-validated methylmercury exposure marker, reflecting non-occupational exposure by seafood or fish consumption. Nevertheless, a study by Barbosa et al. (2001) in a Negro River basin population found important variations in the percentage of methylmercury in total hair mercury, ranging from 34% to 100% [61]. This variation did not seem to be influenced by age, gender, body mass index, frequency of fish consumption or total hair mercury levels. Besides, it is worth mentioning that the use of this biomarker for toxicological purposes has been criticized for its lack of reproducibility [107,108].

That sum of factors makes it difficult to assess a certain situation with exactitude in a determined population. Also, it is important to consider the variability induced by the different sampling, target populations and methodologies used by the researchers according to their objectives. Therefore, we assume that not every hair mercury value represented on the maps and tables can be completely or accurately comparable to each other for epidemiological purposes.

Conclusions

Considering the current recommendations from the environmental agencies, it is patent that almost all the Amazonian riverside populations are exposed to mercury contamination through alimentary habits. There is no evident spatial trend, even if the highest hair mercury levels were found in the central Amazonian regions, especially

the Tapajós River basin. Small and isolated communities with traditional lifestyles seem to be the most exposed to mercury, regardless of the river basin. This situation is very complex for these populations, given that many of them depend on fish for economic support as well as almost unique source of dietary protein.

The available information about the health impact of this situation is not conclusive. Therefore, it becomes difficult to assess accurately the Public Health implication of mercury exposure in this particular context.

Besides, there are numerous Amazonian regions with lacking data and also with discordant results. Thus, a harmonized assessment of mercury human exposure based on a standardized approach would be valuable in order to spatially identify the most contaminated populations. This assessment would also allow prospective follow-ups of exposed populations, especially considering that these small communities are in constant evolution and subject of global changes.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Both authors conceived the review, drafted the manuscript and designed the maps.

References

- Davidson PW, Myers GJ, Cox C, Wilding GE, Shamlaye CF, Huang LS, Cernichiari E, Sloane-Reeves J, Palumbo D, Clarkson TW: **Methylmercury and neurodevelopment: longitudinal analysis of the Seychelles child development cohort.** *Neurotoxicol Teratol* 2006, **28**:529-535.
- Davidson PW, Palumbo D, Myers GJ, Cox C, Shamlaye CF, Sloane-Reeves J, Cernichiari E, Wilding GE, Clarkson TW: **Neurodevelopmental outcomes of Seychellois children from the pilot cohort at 108 months following prenatal exposure to methylmercury from a maternal fish diet.** *Environ Res* 2000, **84**:1-11.
- Davidson PW, Strain JJ, Myers GJ, Thurston SW, Bonham MP, Shamlaye CF, Stokes-Riner A, Wallace JM, Robson PJ, Duffy EM, Georger LA, Sloane-Reeves J, Cernichiari E, Canfield RL, Cox C, Huang LS, Janicuras J, Clarkson TW: **Neurodevelopmental effects of maternal nutritional status and exposure to methylmercury from eating fish during pregnancy.** *Neurotoxicology* 2008, **29**:767-775.
- Debes F, Budtz-Jorgensen E, Weihe P, White RF, Grandjean P: **Impact of prenatal methylmercury exposure on neurobehavioral function at age 14 years.** *Neurotoxicol Teratol* 2006, **28**:363-375.
- Grandjean P, Murata K, Budtz-Jorgensen E, Weihe P: **Cardiac autonomic activity in methylmercury neurotoxicity: 14-year follow-up of a Faroese birth cohort.** *J Pediatr* 2004, **144**:169-176.
- Grandjean P, Weihe P, Needham LL, Burse VW, Patterson DG Jr, Sampson EJ, Jorgensen PJ, Vahter M: **Relation of a seafood diet to mercury, selenium, arsenic, and polychlorinated biphenyl and other organochlorine concentrations in human milk.** *Environ Res* 1995, **71**:29-38.
- Hightower JM, Moore D: **Mercury levels in high-end consumers of fish.** *Environ Health Perspect* 2003, **111**:604-608.
- Kjellström T: **Sverige. Statens naturvårdsverk: Physical and mental development of children with prenatal exposure to mercury from fish. Stage I, Preliminary tests at age 4.** City: National Swedish Environmental Protection Board [Statens naturvårdsverk]; 1986:96.

9. Carmouze J, Boudou A, Lucotte M: **Le mercure en Amazonie. Rôle de l'homme et de l'environnement, risques sanitaires.** IRD, Paris; 2000.
10. Clarkson TW: **Mercury: major issues in environmental health.** *Environ Health Perspect* 1993, **100**:31-38.
11. Lacerda LD, de Souza M, Ribeiro MG: **The effects of land use change on mercury distribution in soils of Alta Floresta, Southern Amazon.** *Environ Pollut* 2004, **129**:247-255.
12. Malm O: **Gold mining as a source of mercury exposure in the Brazilian Amazon.** *Environ Res* 1998, **77**:73-78.
13. Malm O, Castro MB, Bastos WR, Branches FJP, Guimaraes JRD, Zuffo CE, Pfeiffer WC: **An assessment of Hg pollution in different goldmining areas, Amazon Brazil.** *Science of the Total Environment* 1995, **175**:127-140.
14. Roulet M, Lucotte M, Canuel R, Farella N, Courcelles M, Guimaraes JRD, Mergler D, Amorim M: **Increase in mercury contamination recorded in lacustrine sediments following deforestation in the central Amazon.** *Chemical Geology* 2000, **165**:243-266.
15. Roulet M, Lucotte M, Saint-Aubin A, Tran S, Rheault I, Farella N, De Jesus Da Silva E, Dezencourt J, Sousa Passos CJ, Santos Soares G, Guimaraes JR, Mergler D, Amorim M: **The geochemistry of mercury in central Amazonian soils developed on the Alter-do-Chao formation of the lower Tapajos River Valley, Para state, Brazil.** *Sci Total Environ* 1998, **223**:1-24.
16. Veiga MM, Maxson PA, Hylander LD: **Origin and consumption of mercury in small-scale gold mining.** *Journal of Cleaner Production* 2006, **14**:436-447.
17. Beliveau A, Lucotte M, Davidson R, Lopes LO, Paquet S: **Early Hg mobility in cultivated tropical soils one year after slash-and-burn of the primary forest, in the Brazilian Amazon.** *Sci Total Environ* 2009, **407**:4480-4489.
18. Sampaio da Silva D, Lucotte M, Paquet S, Davidson R: **Influence of ecological factors and of land use on mercury levels in fish in the Tapajos River basin, Amazon.** *Environ Res* 2009, **109**:432-446.
19. Dolbec J, Mergler D, Sousa Passos CJ, Sousa de Moraes S, Lebel J: **Methylmercury exposure affects motor performance of a riverine population of the Tapajos river, Brazilian Amazon.** *Int Arch Occup Environ Health* 2000, **73**:195-203.
20. Lebel J, Mergler D, Branches F, Lucotte M, Amorim M, Larribe F, Dolbec J: **Neurotoxic effects of low-level methylmercury contamination in the Amazonian Basin.** *Environ Res* 1998, **79**:20-32.
21. Steuervald U, Weihe P, Jorgensen PJ, Bjerve K, Brock J, Heinzow B, Budtz-Jorgensen E, Grandjean P: **Maternal seafood diet, methylmercury exposure, and neonatal neurologic function.** *J Pediatr* 2000, **136**:599-605.
22. Yokoo EM, Valente JG, Grattan L, Schmidt SL, Platt I, Silbergeld EK: **Low level methylmercury exposure affects neuropsychological function in adults.** *Environ Health* 2003, **2**:8.
23. Axelrad DA, Bellinger DC, Ryan LM, Woodruff TJ: **Dose-response relationship of prenatal mercury exposure and IQ: an integrative analysis of epidemiologic data.** *Environ Health Perspect* 2007, **115**:609-615.
24. Grandjean P, Budtz-Jorgensen E, Jorgensen PJ, Weihe P: **Umbilical cord mercury concentration as biomarker of prenatal exposure to methylmercury.** *Environ Health Perspect* 2005, **113**:905-908.
25. Grandjean P, Weihe P, Burse VW, Needham LL, Storr-Hansen E, Heinzow B, Debes F, Murata K, Simonsen H, Ellefsen P, Budtz-Jorgensen E, Keiding N, White RF: **Neurobehavioral deficits associated with PCB in 7-year-old children prenatally exposed to seafood neurotoxins.** *Neurotoxicol Teratol* 2001, **23**:305-317.
26. Marques RC, Garrofe Dorea J, Rodrigues Bastos W, de Freitas Rebelo M, de Freitas Fonseca M, Malm O: **Maternal mercury exposure and neuro-motor development in breastfed infants from Porto Velho (Amazon), Brazil.** *Int J Hyg Environ Health* 2007, **210**:51-60.
27. Murata K, Weihe P, Budtz-Jorgensen E, Jorgensen PJ, Grandjean P: **Delayed brainstem auditory evoked potential latencies in 14-year-old children exposed to methylmercury.** *J Pediatr* 2004, **144**:177-183.
28. Ramirez GB, Cruz MC, Pagulayan O, Ostrea E, Dalisay C: **The Tagum study I: analysis and clinical correlates of mercury in maternal and cord blood, breast milk, meconium, and infants' hair.** *Pediatrics* 2000, **106**:774-781.
29. Ramirez GB, Pagulayan O, Akagi H, Rivera AF, Lee LV, Berroya A, Cruz MCV, Casintahan D: **Tagum study II: Follow-up study at two years of age after prenatal exposure to mercury.** *Pediatrics* 2003, **111**:.
30. Amorim MI, Mergler D, Bahia MO, Dubeau H, Miranda D, Lebel J, Burbano RR, Lucotte M: **Cytogenetic damage related to low levels of methyl mercury contamination in the Brazilian Amazon.** *An Acad Bras Cienc* 2000, **72**:497-507.
31. Shenker BJ, Guo TL, Shapiro IM: **Mercury-induced apoptosis in human lymphoid cells: evidence that the apoptotic pathway is mercurial species dependent.** *Environ Res* 2000, **84**:89-99.
32. Silbergeld EK, Silva IA, Nyland JF: **Mercury and autoimmunity: implications for occupational and environmental health.** *Toxicol Appl Pharmacol* 2005, **207**:282-292.
33. Silva IA, Nyland JF, Gorman A, Perisse A, Ventura AM, Santos EC, Souza JM, Burek CL, Rose NR, Silbergeld EK: **Mercury exposure, malaria, and serum antinuclear/antinucleolar antibodies in Amazon populations in Brazil: a cross-sectional study.** *Environ Health* 2004, **3**:11.
34. Fillion M, Mergler D, Sousa Passos CJ, Larribe F, Lemire M, Guimaraes JR: **A preliminary study of mercury exposure and blood pressure in the Brazilian Amazon.** *Environ Health* 2006, **5**:29.
35. Guallar E, Sanz-Gallardo MI, van't Veer P, Bode P, Aro A, Gomez-Aracena J, Kark JD, Riemersma RA, Martin-Moreno JM, Kok FJ: **Mercury, fish oils, and the risk of myocardial infarction.** *N Engl J Med* 2002, **347**:1747-1754.
36. **National Research Council.** In *Toxicological effects of methylmercury* 1st edition. Washington, DC: National Academy Press; 2000.
37. Salonen JT, Seppanen K, Nyyssönen K, Korpela H, Kauhanen J, Kanto M, Tuomilehto J, Esterbauer H, Tatzber F, Salonen R: **Intake of mercury from fish, lipid peroxidation, and the risk of myocardial infarction and coronary, cardiovascular, and any death in eastern Finnish men.** *Circulation* 1995, **91**:645-655.
38. Mergler D, Anderson HA, Chan LH, Mahaffey KR, Murray M, Sakamoto M, Stern AH: **Methylmercury exposure and health effects in humans: a worldwide concern.** *Ambio* 2007, **36**:3-11.
39. Dourson ML, Wullenweber AE, Poirier KA: **Uncertainties in the reference dose for methylmercury.** *Neurotoxicology* 2001, **22**:677-689.
40. Rice DC: **The US EPA reference dose for methylmercury: sources of uncertainty.** *Environ Res* 2004, **95**:406-413.
41. Joint FAO/WHO Expert Committee on Food Additives: **Evaluation of Certain Food Additives and Contaminants.** Geneva 2004.
42. **United States Environmental Protection Agency. Office of Science and Technology.** In *Water quality criterion for the protection of human health: methylmercury: final* Washington, DC: Office of Science and Technology, Office of Water, U.S. Environmental Protection Agency; 2001.
43. **UNEP Chemicals, Inter-Organization Programme for the Sound Management of Chemicals.** In *Global mercury assessment* Geneva, Switzerland: UNEP Chemicals; 2002.
44. Castro MB, Albert B, Pfeiffer WC: **Mercury levels in Yanomami Indians hair from Roraima, Brazil.** *Proc 8th Intern Conf Heavy Metals in the Environment* 1991:367-370.
45. Couto RCS, Câmara VM, Sabroza PC: **Intoxicação mercurial: resultados preliminares em duas áreas garimpeiras - PA.** *Cad Saúde Pública* 1988, **4**:301-315.
46. Dorea JG: **Fish are central in the diet of Amazonian riparians: should we worry about their mercury concentrations?** *Environ Res* 2003, **92**:232-244.
47. Frery N, Maury-Brachet R, Maillot E, Deheeger M, de Merona B, Boudou A: **Gold-mining activities and mercury contamination of native amerindian communities in French Guiana: key role of fish in dietary uptake.** *Environ Health Perspect* 2001, **109**:449-456.
48. Hacon S, Yokoo E, Valente J, Campos RC, da Silva VA, de Menezes ACC: **Exposure to mercury in pregnant women from Alta Floresta-Amazon Basin, Brazil.** *Environmental Research* 2000, **84**:204-210.
49. Lebel J, Roulet M, Mergler D, Lucotte M, Larribe F: **Fish diet and mercury exposure in a riparian Amazonian population.** *Water Air and Soil Pollution* 1997, **97**:31-44.
50. Passos CJ, Mergler D: **Human mercury exposure and adverse health effects in the Amazon: a review.** *Cad Saúde Pública* 2008, **24**(Suppl 4):503-520.

51. Vasconcellos MBA, Saiki M, Paletti G, Pinheiro RMM, Baruzzi RG, Spindel R: **Determination of mercury in head hair of Brazilian population groups by neutron-activation analysis.** *Journal of Radioanalytical and Nuclear Chemistry-Articles* 1994, **179**:369-376.
52. Bastos WR, Gomes JP, Oliveira RC, Almeida R, Nascimento EL, Bernardi JV, de Lacerda LD, da Silveira EG, Pfeiffer WC: **Mercury in the environment and riverside population in the Madeira River Basin, Amazon, Brazil.** *Sci Total Environ* 2006, **368**:344-351.
53. Bulletin d'Alerte et de Surveillance Antilles Guyane: **Mercury in French Guiana** Cellule Inter Régionale d'Epidémiologie Antilles Guyane - Institut de Veille Sanitaire; 2007.
54. Cordier S, Grasmick C, Paquier-Passelaigue M, Mandereau L, Weber JP, Jouan M: **Mercury exposure in French Guiana: levels and determinants.** *Arch Environ Health* 1998, **53**:299-303.
55. Counter SA, Buchanan LH, Ortega F: **Mercury levels in urine and hair of children in an Andean gold-mining settlement.** *Int J Occup Environ Health* 2005, **11**:132-137.
56. Webb J, Nicolas Mainville N, Mergler D, Lucotte M, Betancourt O, Davidson R, Cueva E, Quizhpe E: **Mercury in Fish-eating Communities of the Andean Amazon, Napo River Valley, Ecuador.** *Eco Health* 2004, **1**:SU59-SU71.
57. Barbieri FL, Cournil A, Gardon J: **Mercury exposure in a high fish eating Bolivian Amazonian population with intense small-scale gold-mining activities.** *Int J Environ Health Res* 2009, **19**:267-277.
58. Maurice-Bourgoin L, Quiroga I, Chincheros J, Courau P: **Mercury distribution in waters and fishes of the upper Madeira rivers and mercury exposure in riparian Amazonian populations.** *Sci Total Environ* 2000, **260**:73-86.
59. Monrroy SX, Lopez RW, Roulet M, Benefice E: **Lifestyle and mercury contamination of Amerindian populations along the Beni river (lowland Bolivia).** *J Environ Health* 2008, **71**:44-50.
60. Cordier S, Garel M, Mandereau L, Morcel H, Doineau P, Gosme-Seguret S, Josse D, White R, Amiel-Tison C: **Neurodevelopmental investigations among methylmercury-exposed children in French Guiana.** *Environ Res* 2002, **89**:1-11.
61. Barbosa AC, Jardim WV, Dorea JG, Fosberg B, Souza J: **Hair mercury speciation as a function of gender, age, and body mass index in inhabitants of the Negro River basin, Amazon, Brazil.** *Arch Environ Contam Toxicol* 2001, **40**:439-444.
62. Forsberg BR, Forsberg MCS, Padovani CR, Sargentini E, Malm O, Kato H: **High levels of mercury in fish and human hair from the Rio Negro basin (Brazilian Amazon): Natural background or anthropogenic contamination?** In *Proceedings of the International Workshop on Environmental Mercury Pollution and its Health Effects in the Amazon River Basin*; National Institute for Minamata Disease/UFRJ Kato HaWCP; 1995:33-40.
63. Guimaraes JRD, Fostier AH, Forti MC, Melfi JA, Kehrig H, Mauro JBN, Malm O, Krug JF: **Mercury in human and environmental samples from two lakes in Amapá, Brazilian Amazon.** *Ambio* 1999, **28**:296-301.
64. Leino T, Lodenius M: **Human hair mercury levels in Tucuruí area, State of Pará, Brazil.** *Sci Total Environ* 1995, **175**:119-125.
65. Pinheiro MC, Crespo-Lopez ME, Vieira JL, Oikawa T, Guimaraes GA, Araujo CC, Amoras WW, Ribeiro DR, Herculanio AM, do Nascimento JL, Silveira LC: **Mercury pollution and childhood in Amazon riverside villages.** *Environ Int* 2007, **33**:56-61.
66. Pinheiro MC, Oikawa T, Vieira JL, Gomes MS, Guimaraes GA, Crespo-Lopez ME, Muller RC, Amoras WW, Ribeiro DR, Rodrigues AR, Cortes MI, Silveira LC: **Comparative study of human exposure to mercury in riverside communities in the Amazon region.** *Braz J Med Biol Res* 2006, **39**:411-414.
67. Pinheiro MCN, Macchi BM, Vieira JLF, Oikawa T, Amoras WW, Guimaraes GA, Costa CA, Crespo-Lopez ME, Herculanio AM, Silveira LCL, do Nascimento JLM: **Mercury exposure and antioxidant defenses in women: A comparative study in the Amazon.** *Environmental Research* 2008, **107**:53-59.
68. Vasconcellos MBA, Bode P, Paletti G, Catharino MGM, Ammerlaan AK, Saiki M, Favaro DIT, Byrne AR, Baruzzi R, Rodrigues DA: **Determination of mercury and selenium in hair samples of Brazilian Indian populations living in the Amazonian region by NAA.** *Journal of Radioanalytical and Nuclear Chemistry* 2000, **244**:81-85.
69. Santos EC, Camara Vde M, Brabo Eda S, Loureiro EC, de Jesus IM, Fayal K, Sagica F: **Mercury exposure evaluation among Pakaanova Indians, Amazon Region, Brazil.** *Cad Saúde Pública* 2003, **19**:199-206.
70. Harada M, Nakanishi J, Yasoda E, Pinheiro MC, Oikawa T, de Assis Guimaraes G, da Silva Cardoso B, Kizaki T, Ohno H: **Mercury pollution in the Tapajós River basin, Amazon: mercury level of head hair and health effects.** *Environ Int* 2001, **27**:285-290.
71. Passos CJ, Mergler D, Fillion M, Lemire M, Mertens F, Guimaraes JR, Philibert A: **Epidemiologic confirmation that fruit consumption influences mercury exposure in riparian communities in the Brazilian Amazon.** *Environ Res* 2007, **105**:183-193.
72. Pinheiro MC, Guimaraes GA, Nakanishi J, Oikawa T, Vieira JL, Quaresma M, Cardoso B, Amoras W: **Total mercury in hair samples of inhabitants of Tapajós River, Para State, Brazil.** *Rev Soc Bras Med Trop* 2000, **33**:181-184.
73. Pinheiro Md, Nakanishi J, Oikawa T, Guimaraes G, Quaresma M, Cardoso B, Amoras WW, Harada M, Magno C, Vieira JL, Xavier MB, Bacelar DR: **Methylmercury human exposure in riverside villages of Tapajós basin, Para State, Brazil.** *Rev Soc Bras Med Trop* 2000, **33**:265-269.
74. Barbosa AC, Garcia AM, deSouza JR: **Mercury contamination in hair of riverine populations of Apiacas Reserve in the Brazilian Amazon.** *Water Air and Soil Pollution* 1997, **97**:1-8.
75. Dorea JG, Barbosa AC, Ferrari I, De Souza JR: **Fish consumption (hair mercury) and nutritional status of Amazonian Amer-Indian children.** *Am J Hum Biol* 2005, **17**:507-514.
76. Dorea JG, de Souza JR, Rodrigues P, Ferrari I, Barbosa AC: **Hair mercury (signature of fish consumption) and cardiovascular risk in Mundurucu and Kayabi Indians of Amazonia.** *Environ Res* 2005, **97**:209-219.
77. Malm O, Branches FJ, Akagi H, Castro MB, Pfeiffer WC, Harada M, Bastos WR, Kato H: **Mercury and methylmercury in fish and human hair from the Tapajós river basin, Brazil.** *Sci Total Environ* 1995, **175**:141-150.
78. Santos EC, Camara VM, Jesus IM, Brabo ES, Loureiro EC, Mascarenhas AF, Fayal KF, Sa Filho GC, Sagica FE, Lima MO, Higuchi H, Silveira IM: **A contribution to the establishment of reference values for total mercury levels in hair and fish in Amazonia.** *Environ Res* 2002, **90**:6-11.
79. Santos EC, Jesus IM, Brabo ES, Loureiro EC, Mascarenhas AF, Weirich J, Camara VM, Cleary D: **Mercury exposures in riverside Amazon communities in Pará, Brazil.** *Environ Res* 2000, **84**:100-107.
80. Akagi H, Malm O, Kinjo Y, Harada M, Branches FJP, Pfeiffer WC, Kato H: **Methylmercury pollution in the Amazon, Brazil.** *Science of the Total Environment* 1995, **175**:85-95.
81. Crompton P, Ventura AM, de Souza JM, Santos E, Strickland GT, Silbergeld E: **Assessment of mercury exposure and malaria in a Brazilian Amazon riverine community.** *Environmental Research* 2002, **90**:69-75.
82. Barbosa AC, Dorea JG: **Indices of mercury contamination during breast feeding in the Amazon Basin.** *Environmental Toxicology and Pharmacology* 1998, **6**:71-79.
83. Barbosa AC, Silva SR, Dorea JG: **Concentration of mercury in hair of indigenous mothers and infants from the Amazon basin.** *Arch Environ Contam Toxicol* 1998, **34**:100-105.
84. Boischoia AA, Barbosa A: **Exposure to organic mercury in riparian populations on the Upper Madeira river, Rondonia, Brazil, 1991: preliminary results.** *Cad Saúde Pública* 1993, **9**:155-160.
85. Boischoia AA, Henshel DS: **Linear regression models of methyl mercury exposure during prenatal and early postnatal life among riverside people along the Upper Madeira river, Amazon.** *Environmental Research* 2000, **83**:150-161.
86. Malm O, Guimaraes JRD, Castro MB, Bastos WR, Viana JP, Branches FJP, Silveira EG, Pfeiffer WC: **Follow-up of mercury levels in fish, human hair and urine in the Madeira and Tapajós basins, Amazon, Brazil.** *Water Air and Soil Pollution* 1997, **97**:45-51.
87. Grandjean P, White RF, Nielsen A, Cleary D, de Oliveira Santos EC: **Methylmercury neurotoxicity in Amazonian children downstream from gold mining.** *Environ Health Perspect* 1999, **107**:587-591.
88. Dolbec J, Mergler D, Larribe F, Roulet M, Lebel J, Lucotte M: **Sequential analysis of hair mercury levels in relation to fish diet of an Amazonian population, Brazil.** *Sci Total Environ* 2001, **271**:87-97.
89. Hacon S, Barrocas PR, Vasconcellos AC, Barcellos C, Wasserman JC, Campos RC, Ribeiro C, Azevedo-Carlioni FB: **An overview of mer-**

- cury contamination research in the Amazon basin with an emphasis on Brazil. *Cad Saude Publica* 2008, **24**:1479-1492.
90. Pinheiro MC, Muller RC, Sarkis JE, Vieira JL, Oikawa T, Gomes MS, Guimaraes GA, do Nascimento JL, Silveira LC: **Mercury and selenium concentrations in hair samples of women in fertile age from Amazon riverside communities.** *Sci Total Environ* 2005, **349**:284-288.
 91. Santos Filho E, Silva Rde S, Sakuma AM, Scorsafava MA: **Lead and mercury in the hair of children living in Cubatao, in the southeast region of Brazil.** *Rev Saude Publica* 1993, **27**:81-86.
 92. Soares MC, Sarkis JE, Muller RC, Brabo ES, Santos EO: **Correlation between mercury and selenium concentrations in Indian hair from Rondonia State, Amazon region, Brazil.** *Sci Total Environ* 2002, **287**:155-161.
 93. Berglund M, Lind B, Bjornberg KA, Palm B, Einarsson O, Vahter M: **Inter-individual variations of human mercury exposure biomarkers: a cross-sectional assessment.** *Environ Health* 2005, **4**:20.
 94. Santos EC, de Jesus IM, Camara Vde M, Brabo E, Loureiro EC, Mascarenhas A, Weirich J, Luiz RR, Cleary D: **Mercury exposure in Mundurucu Indians from the community of Sai Cinza, State of Para, Brazil.** *Environ Res* 2002, **90**:98-103.
 95. Olivero J, Mendonza C, Mestre J: **Hair mercury levels in different occupational groups in a gold mining zone in the north of Colombia.** *Rev Saude Publica* 1995, **29**:376-379.
 96. Palheta D, Taylor A: **Mercury in environmental and biological samples from a gold mining area in the Amazon region of Brazil.** *Sci Total Environ* 1995, **168**:63-69.
 97. Fillion M, Passos CJ, Lemire M, Fournier B, Mertens F, Guimaraes JR, Mergler D: **Quality of life and health perceptions among fish-eating communities of the Brazilian Amazon: an ecosystem approach to well-being.** *Ecohealth* 2009, **6**:121-134.
 98. Lemire M, Mergler D, Fillion M, Passos CJ, Guimaraes JR, Davidson R, Lucotte M: **Elevated blood selenium levels in the Brazilian Amazon.** *Sci Total Environ* 2006, **366**:101-111.
 99. Mergler D: **Review of neurobehavioral deficits and river fish consumption from the Tapajo's (Brazil) and St. Lawrence (Canada).** *Environmental Toxicology and Pharmacology* 2002, **12**:93-99.
 100. Passos CJ, Mergler D, Gaspar E, Morais S, Lucotte M, Larribe F, Davidson R, de Grosbois S: **Eating tropical fruit reduces mercury exposure from fish consumption in the Brazilian Amazon.** *Environ Res* 2003, **93**:123-130.
 101. Tavares LM, Camara VM, Malm O, Santos EC: **Performance on neurological development tests by riverine children with moderate mercury exposure in Amazonia, Brazil.** *Cad Saude Publica* 2005, **21**:1160-1167.
 102. Benefice E, Monrroy SJ, Rodriguez RW: **A nutritional dilemma: fish consumption, mercury exposure and growth of children in Amazonian Bolivia.** *Int J Environ Health Res* 2008, **18**:415-427.
 103. Boischio AA, Henshel D: **Human biomonitoring to optimize fish consumption advice: reducing uncertainty when evaluating benefits and risks.** *Environmental Research* 2000, **84**:108-126.
 104. Arnold SM, Lynn TV, Verbrugge LA, Middaugh JP: **Human biomonitoring to optimize fish consumption advice: reducing uncertainty when evaluating benefits and risks.** *Am J Public Health* 2005, **95**:393-397.
 105. Dorea JG: **Persistent, bioaccumulative and toxic substances in fish: human health considerations.** *Sci Total Environ* 2008, **400**:93-114.
 106. Kehrig HA, Malm O, Akagi H, Guimaraes JR, Torres JP: **Methylmercury in fish and hair samples from the Balbina Reservoir, Brazilian Amazon.** *Environ Res* 1998, **77**:84-90.
 107. Harkins DK, Susten AS: **Hair analysis: exploring the state of the science.** *Environ Health Perspect* 2003, **111**:576-578.
 108. Miekeley N, Dias Carneiro MT, da Silveira CL: **How reliable are human hair reference intervals for trace elements?** *Sci Total Environ* 1998, **218**:9-17.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

