


Version auteure de l'article :

Are cleaning activities a source of exposure to crystalline silica in women with rheumatoid arthritis? A case–control study

Publié dans: *RMD Open. Rheumatic & Musculoskeletal Diseases* 2023;9:e003205. doi:10.1136/rmdopen-2023-003205

Johanna Sigaux ^{1,2}, Catherine Cavalin,^{3,4} Alain Lescoat,^{5,6} Sarah El Rharras,² Odile Macchi,⁷ Pierre-Yves Brillet,⁸ Lucile Sesé,⁹ Hilario Nunes,¹⁰ Marie-Christophe Boissier,^{1,2} Paul-André Rosental,¹¹ Luca Semerano^{1,2}

ABSTRACT

Introduction Inhalation of crystalline silica (silicon dioxide, SiO₂) is associated with a wide range of acute and chronic diseases, including rheumatoid arthritis (RA). The objectives of this work were to identify the main sources of exposure to SiO₂ in a series of patients with RA not selected on the basis of their professional activity, compared with a representative sample of the French general population, and to assess the association between silica exposure and disease features.

Methods The Dust Exposure Life-Course Questionnaire (DELCQ) is a tool that enables retrospective quantification of both occupational and non-occupational lifetime exposure to SiO₂. DELCQ—previously validated in a large representative sample of the French general population—was administered to 97 consecutive RA patients, and exposure scores were compared between cases and age, gender and smoking status-matched controls (1:4). The main sources of SiO₂ exposure were identified in cases and controls, and source-specific exposure levels were compared. The association between DELCQ scores and disease variables in cases was tested via univariable and multivariable analyses.

Results In women with RA, the main sources of SiO₂ exposure were cleaning activities and dusty clothes laundry, with higher exposure levels from these sources versus the general population ($p < 0.005$). Across the whole series of RA patients, high SiO₂ exposure was independently associated with mediastinal lymphadenopathy (OR 6.3, 95% CI 1.4 to 27.7).

Conclusion Cleaning activities and dusty clothes laundry may be underestimated sources of SiO₂ exposure in women with RA.

INTRODUCTION

Inhalation of crystalline silica (silicon dioxide, SiO₂) is associated with a wide range of acute and chronic diseases,¹ including rheumatoid arthritis (RA).² In a recent meta-analysis of epidemiological studies, occupational exposure to SiO₂ was associated with a 2.59-fold

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Inhalation of crystalline silica (silicon dioxide, SiO₂) is associated with rheumatoid arthritis (RA).
- ⇒ Research on the role of SiO₂ in human disease has been traditionally focused on occupational exposures, with male workers as the main target population.
- ⇒ Substantial exposure from both occupational and non-occupational sources may occur in both men and women over the life course.

WHAT THIS STUDY ADDS

- ⇒ The Dust Exposure Life-Course Questionnaire (DELCQ), is a novel tool that allows to quantify retrospectively both occupational and non-occupational lifetime exposure to SiO₂.
- ⇒ Women with RA have high lifetime exposure to SiO₂ from sources that are currently neglected in research: cleaning activities and dusty clothes laundry.
- ⇒ High occupational DELCQ scores were associated with mediastinal lymphadenopathy, a proxy of crystalline silica exposure.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ This study highlights the importance of daily-life sources of exposure to crystalline silica, particularly in household products, and their association with RA in women.

higher odds of RA.³ Exposure to SiO₂ has been identified as a risk factor for incident RA in the Swedish population,⁴ showing a closer association with seropositive RA^{5,6} and positive interaction with smoking.

SiO₂ is present in diverse polymorphs like quartz, cristobalite and tridymite; quartz is the most common form found in nature, and the main component of the crystalline silica dust that may be emitted in manufacturing processes.⁵ SiO₂ is used in a wide range of

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/rmdopen-2023-003205>).



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Johanna Sigaux;
johanna.sigaux@aphp.fr

industrial activities. High exposure to crystalline silica dust may occur in blast furnace, electronics glass, cement and ceramic manufacturing. It may also occur in other sectors such as mining (historically the main sector of exposure to SiO₂ and subsequent silicosis), farming and construction.⁷

Since these activities may involve intense exposure for workers, research on SiO₂ in human diseases has historically focused on occupational exposure and on male workers as the main target population.⁸ However, SiO₂ is ubiquitous in the environment. Over a lifetime, both men and women may experience substantial exposure across many professional activities and even beyond occupational contexts.

SiO₂ is present in a wide range of products (scouring powders, cat litter, rubber tires, etc) that may emit airborne particles. Thus, an accurate measure of actual lifetime exposure to crystalline silica remains a challenging task.⁹

The Dust Exposure Life-Course Questionnaire (DELCO) is a novel tool that enables retrospective quantification of both occupational and non-occupational lifetime exposure to SiO₂.¹⁰

A multidisciplinary team of epidemiologists, social scientists, occupational health and medicine collaborators developed the questionnaire to identify situations likely to expose people to SiO₂. The exposures were based on an extensive list of products and activities summarised by the International Agency for Research on Cancer¹¹ and complemented by medical and statistical surveys of the general population^{12–14} as well as data from the literature about exposure to SiO₂ and inorganic particles in occupational or environmental settings.^{9,15}

DELCO was administered in 2016 to a representative sample of the French general population (ELIPSS cohort, ELIPSSilice2 survey) selected from the French national rolling census by the French National Institute of Statistics and Economic Studies (INSEE). This population serves as a control for case–control studies of lifetime exposure to SiO₂ in patients with specific diseases such as RA and systemic sclerosis.¹⁰ Using DELCO, we recently found that both male and female RA patients had higher occupational exposure to SiO₂ than their respective controls, and that women with RA also had higher non-occupational exposure than controls.¹⁰

In this case–control study, we sought to identify the actual sources of SiO₂ exposure in patients with RA, and to explore the association between the level of SiO₂ exposure and major disease features and outcomes.

PATIENTS AND METHODS

Patients

The case and control populations and SiO₂ exposure assessment are detailed elsewhere.¹⁰ In short, 97 patients who were diagnosed with RA according to 2010 ACR/EULAR classification criteria and followed by the Department of Rheumatology of Avicenne Teaching Hospital

(GHUPSSD, APHP, Bobigny, France) were consecutively recruited in a retrospective observational study between June and December 2016.

A specially trained interviewer administered DELCO to all patients. The interviews typically lasted 45–60 min and were conducted in person for 56 patients and by phone for 41.

Clinical, laboratory and radiological data were retrieved from patient files.

For a subgroup of patients, high-resolution CT (HRCT) images of the chest were available. HRCT was not performed on purpose for this study, and had been carried out in the setting of patient usual care or prior initiating a targeted treatment. Two investigators evaluated HRCT images of the chest while blinded to the results of the SiO₂ exposure evaluations (LSesé, P-YB). The investigators determined the presence and extent of interstitial lung disease (ILD), and the presence of emphysema, mediastinal lymphadenopathy (LA), nodules and micronodules (number and shape). They assessed the presence and extent of ILD according to the Fleischner Society criteria.¹⁶ The most relevant CT signs of ILD are reticular patterns, traction bronchiectasis, ground-glass opacity and honeycombing. We also included ILAs (interstitial lung abnormalities) in our definition of ILD. ILAs are described as non-dependent abnormalities affecting more than 5% of any lung zone.¹⁷ Mediastinal LAs were measured along their short axis and considered significant if >10 mm.

Controls

For each patient, up to four gender, age and smoking-status matched controls were randomly sampled from the ELIPSSilice2 cohort (online supplemental figure 1). Briefly, ELIPSSilice2 is composed of subjects sampled by INSEE, based on national rolling census data and representative of the French general population (aged 18–76 at the time of the census), that is, independent of any health considerations. Subjects (n=2 739) completed the DELCO questionnaire in 2016. The questionnaire was self-administered on tablets.

Patient and public involvement

Patients and the public were neither involved in the design and conduct of the study, nor the choice of outcome measures, nor the recruitment.

Assessment of silica exposure

The design and the content of DELCO are thoroughly described elsewhere.¹⁰

Briefly, DELCO addresses both occupational and non-occupational exposure over a lifetime through more than 130 questions (online supplemental data 1 and 2).

The questionnaire was conducted by phone or in-person interview by a trained interviewer (OM or SER) who was unaware of the results of clinical, immunological and pulmonary assessments. A score from 0 to 5 (based on the cumulative exposure duration and level

of protection) was given for each specific exposure. Both the occupational (OES) and non-occupational exposure scores (NOES) were calculated by adding up the scores for each specific exposure. The two scores were analysed separately.

Statistical analysis

The occupational and non-occupational DELCQ scores related to each single source of exposure were compared between women with RA and controls using the Wilcoxon test. The Bonferroni correction was applied to avoid alpha-risk inflation due to multiple comparisons. The associations between DELCQ score quartiles and patient variables (for both male and female patients) were analysed with the Chi-square test or Fisher's exact test and expressed with OR and 95% CIs. A logistic regression model including age, gender, disease duration, anticitrullinated protein antibody (ACPA)/rheumatoid factor (RF) positivity and smoking status (defined as never, current or former smoker) was used to assess the association between SiO₂ exposure and HRCT abnormalities at multivariable analysis. Patients with an exposure score in the fourth quartile were considered to be highly exposed, and were compared with patients in the first to third quartiles. For all tests, the significance level was set at $p < 0.05$. All statistical analyses were performed with R and RStudio V.2022.07.0 software.

Role of the funding source

The SILICOSIS project was funded with an Advanced Grant from the European Research Council. As such, its principal investigator Paul-André Rosental of the Centre for European Studies and Comparative Politics (Sciences Po, Paris) and his team are solely responsible for the design of the study; the data collection, analysis and interpretation; the writing of publications; and decisions to submit written outputs for publication in journals or books.

RESULTS

A total of 97 patients with RA (mean age: 59.4 ± 13.2 years; 76 women and 21 men) completed the DELCQ questionnaire (table 1).

Mean RA duration was 13.1 ± 11 years, most patients were RF and/or ACPA positive, and 61.9% were never smokers (19% of male vs 73% of female patients, $p < 0.05$). At least one HRCT scan of the chest was available for 63.9% of patients.

As previously reported,¹⁰ RA patients had significantly higher OES than their matched controls, whereas no difference appeared for non-occupational exposures. However, women with RA had both higher occupational and non-occupational exposures to SiO₂ than their female matched controls. We therefore sought to uncover the most relevant sources of SiO₂ exposure in female patients by studying and comparing the distribution of occupational and non-occupational sources of SiO₂ exposure in women with RA and their controls.

Sources and levels of occupational and non-occupational silica exposure in women with RA and controls

Cleaning activities are the most important source of occupational exposure, accounting for 31% of women's total occupational exposure (15.7% from 'proper' cleaning activities+15.3% from laundry for dusty work clothes). Then came farming (9.1%), steel welding (4.2%) and construction (3.0%) activities (data not shown). The sources of occupational SiO₂ exposures in controls were also primarily cleaning activities (15.2% from 'proper' cleaning activities+6.2% from laundry for dusty work clothes=21.4%) and farming activities (19.6%) (data not shown).

We then compared the mean score allocated to each main occupational and non-occupational source for women with RA versus controls (table 2).

Within occupational activities, women with RA had higher exposure scores for cleaning activities (1.03 ± 1.71 vs 0.46 ± 1.32 , $p = 0.02$) and laundry for dusty work clothes (1.00 ± 1.86 vs 0.19 ± 0.86 , $p = 0.01$).

Among non-occupational SiO₂ exposure sources, women with RA had higher scores than controls for cleaning activities (8.03 ± 4.29 vs 3.92 ± 4.48 , $p < 0.0001$).

Association between silica exposure and patient variables

Ever-smoker RA patients (active or former) were more likely to be in the highest quartile of SiO₂ OES than never-smoker patients ($p = 0.002$), but no difference was observed for NOES ($p = 0.18$) (table 3).

Being in the highest quartile of SiO₂ exposure (occupational or non-occupational) showed no association with ACPA or RF status, radiographic erosion, or the need for biologic disease-modifying antirheumatic drugs (DMARDs). Conversely, high occupational exposure to SiO₂ was associated with ILD, emphysema and mediastinal LA (all $p < 0.05$, table 3).

The association between high occupational SiO₂ exposure, ILD and mediastinal LA at HRCT persisted after adjustment for smoking status (ie, ever-smoker or never-smoker), with a sixfold increased risk for patients in the highest OES quartile (table 4). Further adjusting for ACPA and/or RF status, gender and disease duration had no bearing on the associations.

DISCUSSION

This study shows that women with RA, not selected on the basis of their professional activity, have high lifetime exposure to SiO₂ from sources that are currently neglected in research: cleaning activities and dusty clothes laundry. The use of an accurate questionnaire derived from social sciences enabled identification of these unconventional sources of SiO₂ exposure.

Cleaning activities in particular were a relevant source of exposure to SiO₂ in both occupational and non-occupational contexts. Given that around 60% of the women in the series were retired or officially unemployed, our study's retrospectivity allowed us to identify

Table 1 RA patient characteristics

| General characteristics | Overall RA population (n=97) | Female RA (n=76) | Male RA (n=21) | P value |
|---|------------------------------|------------------|----------------|---------|
| Age (years) | 59.4±13.2 | 57.4±13.0 | 66.4±12.2 | <0.01 |
| RA duration (years) | 13.1±11.0 | 13.4±10.9 | 12.1±11.6 | NS |
| ACPA positive | 78 (83.0) | 57 (75.0) | 21 (100) | <0.05 |
| RF positive | 76 (84.4) | 58 (76.3) | 18 (85.7) | NS |
| Erosive RA | 80 (84.2) | 61 (80.3) | 19 (90.5) | NS |
| Tobacco users | | | | |
| Never a smoker | 60 (61.9) | 56 (73.7) | 4 (19.0) | <0.0001 |
| Current smoker | 14 (14.4) | 9 (11.8) | 5 (23.8) | NS |
| Former smoker | 23 (23.7) | 11 (14.5) | 12 (57.1) | <0.001 |
| Occupation | | | | |
| Never worked or student | 6 (6.4) | 6 (7.9) | 0 (0) | NS |
| Active worker | 29 (30.8) | 23 (30.3) | 6 (28.6) | NS |
| Non-active worker (retired or unemployed) | 59 (62.8) | 48 (63.2) | 11 (52.4) | NS |
| RA treatments | | | | |
| MTX | 53 (54.6) | 42 (55.3) | 11 (52.5) | NS |
| bDMARDs | 54 (55.7) | 44 (57.9) | 10 (47.6) | NS |
| Never bDMARDs user | 30 (30.9) | 23 (30.3) | 7 (33.3) | NS |
| bDMARD 1st line | 37 (38.1) | 31 (40.8) | 6 (28.6) | NS |
| bDMARD 2nd line | 18 (18.6) | 13 (17.1) | 5 (23.8) | NS |
| bDMARD 3rd line | 6 (6.2) | 5 (6.6) | 1 (4.8) | NS |
| bDMARD >3rd line | 6 (6.2) | 4 (5.3) | 2 (9.5) | NS |
| Chest HRCT scan | | | | |
| Normal HRCT chest scan | 12 (19.3) | 12 (24.5) | 0 (0) | NS |
| ILD | 11 (18.0) | 7 (14.3) | 4 (16.7) | NS |
| Emphysema | 12 (19.7) | 5 (10.2) | 7 (58.3) | <0.001 |
| Mediastinal lymph nodes | 17 (27.4) | 12 (24.5) | 5 (41.7) | NS |
| Pulmonary nodules* | 18 (29.0) | 13 (26.5) | 5 (41.7) | NS |

Data are expressed as absolute number and percentage (%) or mean (SD).

*Nodules or micronodules.

ACPA, anticitrullinated protein antibody; bDMARD, biologic disease-modifying antirheumatic drug; HRCT, high-resolution CT; ILD, interstitial lung disease; MTX, methotrexate; NS, not significant; RA, rheumatoid arthritis; RF, rheumatoid factor.

substantial exposure to SiO₂ in a population that would not usually be considered at risk.

Like cigarette smoke, SiO₂ is an airborne agent that may initiate an inflammatory response in lung tissue. In predisposed individuals, local inflammation is conducive to the activation of adaptive immunity, production of autoantibodies and progression towards systemic disease.^{5 6}

The association between high occupational exposure to SiO₂ and incident RA is supported by several observational studies.^{3 18} A Swedish nationwide study also found an association between the risk of incident RA and a job-matrix score based on occupations with established high exposure to SiO₂.⁸ However, all these studies focused on specific jobs known to expose workers to high levels of SiO₂.

Conversely, the novelty of this study was the assessment and quantification of SiO₂ exposure in an unselected sample of consulting RA patients. In a previous study, we showed that patients (men and women with RA) primarily differ from their (respective) matched controls on the level of lifetime occupational exposure.¹⁰ Consistent with the literature, men and women with RA had higher occupational exposure to SiO₂ than both (respective) gender-matched controls. Occupational exposure was mainly attributable to male-dominated activities such as construction, glass, mining, agriculture, ceramics and electronics.⁷

The novelty of this work lies in its measurement of non-occupational exposure, and of occupational exposure from occupational activities that are not necessarily identified as at-risk. In the case of women, we had

Table 2 Main sources of occupational and non-occupational silica exposure in women with RA versus controls (expressed as the mean and SD of exposure score)

| Activities | RA women n=76 | | Control women n=308 | | P value* | Bonferroni correction† |
|---|---------------|------|---------------------|------|----------|------------------------|
| | Mean | SD | Mean | SD | | |
| Main occupational activities (OES) | | | | | | |
| Construction | 0.19 | 1.71 | 0.12 | 1.22 | 0.99 | NS |
| Farming | 0.60 | 2.64 | 0.60 | 2.52 | 0.72 | NS |
| Stone | 0.09 | 0.57 | 0.03 | 0.32 | 0.13 | NS |
| Wood | 0.00 | 0.00 | 0.10 | 1.05 | 0.39 | NS |
| Cleaning | 1.03 | 1.71 | 0.46 | 1.32 | <0.001 | † |
| Metallurgy | 0.05 | 0.46 | 0.03 | 0.51 | 0.29 | NS |
| Painting | 0.06 | 0.41 | 0.06 | 0.40 | 0.99 | NS |
| Plumbing | 0.00 | 0.00 | 0.00 | 0.00 | – | NS |
| Steel welding | 0.27 | 1.48 | 0.03 | 0.33 | 0.02 | NS |
| Dusty work clothes laundry | 1.00 | 1.86 | 0.19 | 0.86 | <0.001 | † |
| Main non-occupational activities (NOES) | | | | | | |
| Farming | 2.30 | 4.07 | 1.06 | 1.88 | 0.13 | NS |
| DIY | 1.05 | 1.61 | 0.80 | 1.40 | 0.19 | NS |
| Wood | 0.32 | 0.97 | 0.29 | 0.96 | 0.73 | NS |
| Cleaning | 8.03 | 4.29 | 3.92 | 4.48 | <0.001 | † |
| Dusty work clothes laundry | 2.55 | 3.71 | 1.76 | 3.30 | 0.064 | NS |
| Cat litter | 0.47 | 0.79 | 0.82 | 0.95 | 0.003 | † |
| Mud bath | 0.49 | 0.64 | 0.81 | 0.94 | 0.03 | NS |

*Wilcoxon test, level of significance p<0.05.

†Bonferroni correction, level of significance p<0.004 for occupational activities, p<0.006 for non-occupational activities. NOES, non-occupational exposure score; OES, occupational exposure score; RA, rheumatoid arthritis.

previously shown that our sample of female RA patients were more exposed to SiO₂ from both occupational and non-occupational than their matched controls randomly

sampled from the general population.¹⁰ Moreover, while tobacco plays a role in the pathogenesis of RA, since the vast majority of these female patients were non-smokers,

Table 3 Association between SiO₂ exposure (occupational and non-occupational, by quartile) and patient characteristics

| RA patient variable | Occupational exposure score | | | Non-occupational exposure score | | |
|----------------------------|-----------------------------|-----------|--------------------|---------------------------------|-----------|----------|
| | Q1–Q3 | Q4 | P value* | Q1–Q3 | Q4 | P value* |
| Ever-smoking, n (%) | 21 (28.8) | 16 (66.7) | 0.002 | 25 (33.8) | 12 (55.2) | 0.18 |
| ACPA-positivity, n (%) | 56 (78.9) | 22 (95.6) | 0.16 | 61 (85.9) | 17 (73.9) | 0.31 |
| Erosive RA patients, n (%) | 61 (84.7) | 19 (82.6) | 0.93 | 62 (84.9) | 18 (81.8) | 0.98 |
| bDMARD treatment n (%) | 44 (60.3) | 10 (41.7) | 0.17 | 38 (51.3) | 16 (69.6) | 0.19 |
| ILD, n (%) | 5 (10.9) | 6 (37.5) | 0.04 [†] | 9 (19.1) | 2 (13.3) | 0.9 |
| Emphysema, n (%) | 5 (10.9) | 7 (43.7) | 0.004 [†] | 8 (17.4) | 4 (25.0) | 0.51 |
| Mediastinal LA, n (%) | 9 (19.6) | 8 (50.0) | 0.04 [†] | 14 (28.0) | 3 (25.0) | 0.88 |
| Lung micronodules, n (%) | 4 (8.7) | 3 (18.7) | 0.27 | 5 (10.6) | 2 (13.3) | 0.77 |

*χ² test, level of significance p<0.05.

ACPA, anticitrullinated protein antibody; bDMARD, biologic disease-modifying antirheumatic drug; ILD, interstitial lung disease; LA, lymphadenopathy; RA, rheumatoid arthritis.

Table 4 Association between occupational silica exposure (by quartile) and HRCT lung abnormalities

| HRCT abnormality | OES (quartiles) | Unadjusted OR (95% CI) | Adjusted OR (95% CI) |
|-----------------------------|-----------------|------------------------|----------------------|
| Mediastinal lymphadenopathy | Q1–Q3 | (ref) | (ref) |
| | Q4 | 4.1 (1.2 to 13.9) | 6.3 (1.4 to 27.7) |
| Interstitial lung disease | Q1–Q3 | (ref) | (ref) |
| | Q4 | 6.3 (1.5 to 26.6) | 6.5 (1.3 to 32.6) |
| Emphysema | Q1–Q3 | (ref) | (ref) |
| | Q4 | 6.4 (1.6 to 24.7) | 2.9 (0.6 to 13.7) |

ORs adjusted for smoking status, gender, disease duration, anticitrullinated protein antibody and rheumatoid factor status.
HRCT, high-resolution CT; OES, occupational exposure score.

SiO₂ may be a major contributor to the pathogenesis of the disease in this subgroup.

The thoroughness of DELCQ enables the identification of specific sources of SiO₂ exposure in women with RA. The main sources are related to cleaning activities. Contact with SiO₂ comes from household cleaning products¹⁹ and from the manipulation of clothes belonging to workers directly exposed to high levels of SiO₂ (dusty work clothes laundry). As those activities (mainly performed by women) are not considered to be associated with SiO₂ hazards, specific protections are rarely (if ever) implemented. Even fewer protections are considered when these activities are practiced at home. Hence, the absence of protection during house chores impacted the DELCQ score and led us to identify hitherto unrecognised sources of non-occupational exposure to SiO₂.

According to DELCQ, people who perform housekeeping activities are exposed to SiO₂, which many detergents contain. However, the use of cleansing agents has not been demonstrated to result in substantial SiO₂ exposure, and the association with RA may be due to other compounds contained in the detergents. In the Swedish Epidemiological Investigation of Rheumatoid Arthritis cohort, a strong association between occupational exposure to detergents and the risk of both ACPA-positive and ACPA-negative RA was found. Detergents were the most prevalent occupational inhalant for men (percentage exposed: 31%) and women (51%) with RA.²⁰

In a Swedish cross-sectional study of 40 000 workers, the prevalence of RA was higher in women working as cleaners; doctors and nurses; nurse assistants; textile workers; shop assistants; bank, postal and telegraph personnel; and housewives.²¹ The study hypothesised the action of an environmental factor, but SiO₂ was not mentioned as a possible one. Another epidemiological study found a looser association with RA in women working as hairdressers or beauticians, especially those exposed to hair products. No information was provided about the composition of the products, including any possible SiO₂ content.²⁰

Interestingly, Colinet's patients (whose description was used for the Caplan-Colinet syndrome²²) were employees

of a soap factory, involved in silica-scouring powder packing.²³

Our female patient sample includes a vast majority of housewives from low-income households, partially reflecting Avicenne hospital's location in a poor urban area. Their cumulative exposure may be compared with an occupational one, even though housecleaning products may differ from those used in occupational contexts. A documented association between low socioeconomic status and RA exists,^{24 25} but no study has ever considered the potential role of SiO₂ exposure in this association.

The association of high occupational DELCQ scores with HRCT abnormalities, notably mediastinal LA, provides external validity to the questionnaire. Since mediastinal LA is a common characteristic of silicosis, it can be considered a proxy for high SiO₂ exposure.^{26 27} A similar association with high occupational DELCQ scores was demonstrated in a series of scleroderma patients.²⁸

High occupational SiO₂ exposure was also independently associated with ILD after adjusting for potential confounders. This association is intriguing as it links SiO₂ exposure with a type of rheumatoid lung involvement. However, this association may only be due to chance and needs to be confirmed in larger series. In addition, we may have overestimated the proportion of ILD and the strength of the association. As chest HRCT had been performed only for some patients in the setting of their usual care or prior initiating a targeted treatment, a selection bias based on the presence of pulmonary involvement or on the severity of RA is likely present.

No association was found between the DELCQ and other markers of disease severity like radiographic erosion, ACPA/RF positivity and level, or the need for biologic DMARDs therapy.

The main limitation of the study is the limited sample size and the recruitment of patients from a single centre located in a poor urban area, which may result in under or overestimating relevant sources of exposure among RA patients. This may reduce the generalisability of the results and larger studies in more heterogeneous populations are needed to confirm this finding. However, the patients were recruited from a 2 million inhabitants

territory that stretches over two different administrative areas, one of which displays a sociodemographic profile corresponding to that of the French society. In fact, the main sources of exposure that we have identified did not greatly differ from those of controls extracted from a representative sample of the French general population. Another limitation is that the questionnaire administration method differed between the controls (self-administered questionnaire on digital tablets) and the cases (administered by a trained interviewer by phone or in person), potentially introducing an evaluation bias. We estimated and discussed the impact of that potential bias elsewhere,^{10 29} as we compared the DELCQ scores of our hospital cohort of patients with RA to those of self-reported RA patients inside the general population panel from which the healthy controls were selected (ELLIP-SILICE 1 and 2). Although occupational (but not non-occupational) scores were lower in self-declared patients, the difference was not major and may be partly explained by higher diagnostic accuracy in a hospital setting. More importantly, inside the general population sample, the DELCQ score was able to discriminate between healthy subjects and self-reported patients both on occupational and non-occupational scores. Thus, although the different mode of administration might have led to overestimate the size of the difference, this is unlikely to affect the validity of the analysis.

The study's main strength lies in the tool we used to assess SiO₂ exposure. DELCQ is a detailed questionnaire that is not excessively time-consuming but enables quantification of lifetime SiO₂ exposure, accounting for the source of SiO₂, context of exposure (occupational vs non-occupational), cumulative duration of exposure and use of protection.

In conclusion, this study identifies underestimated sources of potential SiO₂ exposure in women with RA. Both mechanistic and larger epidemiological studies are warranted to explore the relevance of SiO₂ exposure from those sources and the possible need for preventive strategies.

Author affiliations

¹UMR Inserm U1125, Université Sorbonne Paris Nord, Bobigny, France

²Rheumatology Department, Assistance Publique-Hôpitaux de Paris, GH HUPSSD, Bobigny, France

³UMR CNRS-INRAE 7170-1427, Université Paris Dauphine, Paris, France

⁴Laboratoire interdisciplinaire d'évaluation des politiques publiques (LIEPP), Paris, France

⁵University Hospital Centre Rennes, Rennes, France

⁶Institut de Recherche en Santé, Environnement et Travail, INSERM, Paris, France

⁷Centre d'études des mouvements sociaux, Ecole des Hautes Etudes en Sciences Sociales (EHESS), Paris, France

⁸Radiology Department, Assistance Publique-Hôpitaux de Paris, GH HUPSSD, Bobigny, France

⁹Physiology Department, Assistance Publique-Hôpitaux de Paris, GH HUPSSD, Bobigny, France

¹⁰Respiratory Department, Assistance Publique-Hôpitaux de Paris, GH HUPSSD, Bobigny, France

¹¹Centre d'histoire, Sciences Po, Paris, France

Contributors JS, CC, P-AR and LSeerano conceptualised the study. JS, CC and LSeerano created the methodology for the study. CC, SER, OM, P-YB, LSeésé and HN carried out the data collection. JS and LSeerano did the data analysis and verified the data. LSeerano and AL supervised the data collection. JS and CC wrote the original draft. JS is the guarantor. All authors reviewed and edited the manuscript. All authors had full access to all of the data and the final responsibility to submit for publication.

Funding SILICOSIS Project, led by Paul-André Rosental, Centre for European Studies and Comparative Politics, Sciences Po (Paris, France), sponsored by the European Research Council (ERC); (grant number ERC-2011-ADG_20110406, project ID 295817).

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and was approved by CLEA (Comité local éthique Avicenne), N° 2016-256. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Johanna Sigaux <http://orcid.org/0000-0002-6371-740X>

REFERENCES

- 1 Marinaccio A, Consonni D, Mensi C, *et al*. Association between asbestos exposure and Pericardial and Tunica vaginalis Testis malignant Mesothelioma: a case-control study and Epidemiological remarks. *Scand J Work Environ Health* 2020;46:609–17.
- 2 Klareskog L, Rönnelid J, Saevarsdottir S, *et al*. The importance of differences; on environment and its interactions with genes and immunity in the causation of rheumatoid arthritis. *J Intern Med* 2020;287:514–33.
- 3 Mehri F, Jenabi E, Bashirian S, *et al*. The association between occupational exposure to silica and risk of developing rheumatoid arthritis: A meta-analysis. *Saf Health Work* 2020;11:136–42.
- 4 Stolt P, Källberg H, Lundberg I, *et al*. Silica exposure is associated with increased risk of developing rheumatoid arthritis: results from the Swedish EIRA study. *Ann Rheum Dis* 2005;64:582–6.
- 5 Stolt P, Yahya A, Bengtsson C, *et al*. Silica exposure among male current smokers is associated with a high risk of developing ACPA-positive rheumatoid arthritis. *Ann Rheum Dis* 2010;69:1072–6.
- 6 Zeng P, Chen Z, Klareskog L, *et al*. Amount of smoking, duration of smoking cessation and their interaction with silica exposure in the risk of rheumatoid arthritis among males: results from the Swedish Epidemiological investigation of rheumatoid arthritis (EIRA) study. *Ann Rheum Dis* 2018;77:1238–41.
- 7 Yassin A, Yebesi F, Tingle R. Occupational exposure to crystalline silica dust in the United States, 1988–2003. *Environ Health Perspect* 2005;113:255–60.
- 8 Ilar A, Gustavsson P, Wiebert P, *et al*. Occupational exposure to organic dusts and risk of developing rheumatoid arthritis: findings from a Swedish population-based case-control study. *RMD Open* 2019;5:e001049.
- 9 U.S. EPA. *Health Effects Of Inhaled Crystalline And Amorphous Silica*. U.S. Environmental Protection Agency, Office of Research and Development, National Center For Environmental Assessment, Research Triangle Park Office, Research Triangle Park, NC, EPA/600/R-95/115. 1996.

- 10 Cavalin C, Lescoat A, Sigaux J, *et al.* Crystalline silica exposure in patients with rheumatoid arthritis and systemic sclerosis: a nationwide cross-sectional survey. *Rheumatology (Oxford)* 2022;keac675.
- 11 IARC publications Website - pharmaceuticals. Available: <https://publications.iarc.fr/Book-And-Report-Series/IARC-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Pharmaceuticals-2012> [Accessed 26 Mar 2021].
- 12 Matinet B, Rosankis É, Léonard M. Les expositions aux Risques Professionnels. LES Produits chimiques. *Synthèse-Stat DARES* 2020:323.
- 13 Couraud S, Souquet P-J, Paris C, *et al.* Biocast/IFCT-1002: Epidemiological and molecular features of lung cancer in never-smokers. *Eur Respir J* 2015;45:1403–14.
- 14 Parks CG, Cooper GS. Occupational exposures and risk of systemic lupus erythematosus: a review of the evidence and exposure assessment methods in Population- and clinic-based studies. *Lupus* 2006;15:728–36.
- 15 Drent M, Wijnen PA, Boots AW, *et al.* Cat litter is a possible trigger for Sarcoidosis. *Eur Respir J* 2012;39:221–2.
- 16 Hansell DM, Goldin JG, King TE, *et al.* CT staging and monitoring of Fibrotic interstitial lung diseases in clinical practice and treatment trials: a position paper from the Fleischner society. *Lancet Respir Med* 2015;3:483–96.
- 17 Hatabu H, Hunninghake GM, Richeldi L, *et al.* Interstitial lung abnormalities detected incidentally on CT: a position paper from the Fleischner society. *Lancet Respir Med* 2020;8:726–37.
- 18 Biton J, Saldenber-Kermanac'h N, Decker P, *et al.* The Exposome in rheumatoid arthritis. *Joint Bone Spine* 2022;89:105455.
- 19 Dumontet C, Vincent M, Laennec E, *et al.* Silicosis due to inhalation of domestic cleaning powder. *Lancet* 1991;338:1085.
- 20 Tang B, Liu Q, Ilar A, *et al.* Occupational Inhalable agents constitute major risk factors for rheumatoid arthritis, particularly in the context of genetic predisposition and smoking. *Ann Rheum Dis* 2023;82:316–23.
- 21 Hellgren L. The prevalence of rheumatoid arthritis in occupational groups. *Scand J Rheumatol* 1987;16:106–13.
- 22 CAPLAN A. Certain unusual radiological appearances in the chest of coal-miners suffering from rheumatoid arthritis. *Thorax* 1953;8:29–37.
- 23 Ronsmans S, Blanc PD. Colinet-Caplan syndrome: history of an outbreak of autoimmune disease in scouring powder workers. *Ann Intern Med* 2023;176:260–5.
- 24 Bengtsson C, Nordmark B, Klareskog L, *et al.* Socioeconomic status and the risk of developing rheumatoid arthritis: results from the Swedish EIRA study. *Ann Rheum Dis* 2005;64:1588–94.
- 25 Mackie SL, Taylor JC, Twigg S, *et al.* Relationship between area-level socio-economic deprivation and autoantibody status in patients with rheumatoid arthritis: Multicentre cross-sectional study. *Ann Rheum Dis* 2012;71:1640–5.
- 26 Satija B, Kumar S, Ojha UC, *et al.* Spectrum of high-resolution computed tomography imaging in occupational lung disease. *Indian J Radiol Imaging* 2013;23:287–96.
- 27 Antao V dos S, Pinheiro GA, Terra-Filho M, *et al.* High-resolution CT in Silicosis: correlation with radiographic findings and functional impairment. *J Comput Assist Tomogr* 2005;29:350–6.
- 28 Ballerie A, Cavalin C, Lederlin M, *et al.* Association of silica exposure with chest HRCT and clinical characteristics in systemic sclerosis. *Semin Arthritis Rheum* 2020;50:949–56.
- 29 Cavalin C, Catinon M, Macchi O, *et al.* Expositions aux Particules Inorganiques: comment poser La question? in Duwez Emmanuelle, Mercklé Pierre (Dir.). In: *Un panel français L'Étude longitudinale par internet pour les sciences sociales (ELIPSS)*. Paris: Éditions de l'INED, 2021: 185–212.