



HAL
open science

Longest increasing paths with Lipschitz constraints

Anne-Laure Basdevant, Lucas Gerin

► **To cite this version:**

Anne-Laure Basdevant, Lucas Gerin. Longest increasing paths with Lipschitz constraints. 2020.
hal-02437174v1

HAL Id: hal-02437174

<https://hal.science/hal-02437174v1>

Preprint submitted on 13 Jan 2020 (v1), last revised 16 Dec 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Longest increasing paths with Lipschitz constraints

BASDEVANT A.-L.* GERIN L.†

January 17, 2020

Abstract

The Hammersley problem asks for the maximal number of points in a monotonous path through a Poisson point process. It is exactly solvable and notoriously known to belong to the KPZ universality class, with a cube-root scaling for the fluctuations.

Here we introduce and analyze a variant in which we impose a Lipschitz condition on paths. Thanks to a coupling with the classical Hammersley problem we observe that this variant is also exactly solvable. It allows us to derive first and second orders asymptotics. It turns out that the cube-root scaling only holds for certain choices of the Lipschitz constants.

MSC 2010 Classification: 60K35, 60F15.

Keywords: combinatorial probability, last-passage percolation, longest increasing subsequences, longest increasing paths, Hammersley's process, cube-root fluctuations.

1 Introduction

The focus of the present paper is the Hammersley (or Ulam-Hammersley) problem. It asks for the asymptotic behaviour of the maximal number ℓ_n of points that belong to a monotonous path through n uniform points in a square (or more precisely a Poissonized version thereof, see details below). It is among the few exactly solvable models which are known to lie in the *Kardar-Parisi-Zhang (KPZ) universality class*, another exactly solvable model is the corner-growth model (which is itself closely related to last-passage percolation). The main feature of the KPZ class is a universal relation $\chi = 2\xi - 1$ between the *fluctuation exponent* χ and the *wandering exponent* ξ (also called *dynamic scaling exponent* or *transversal exponent*). We refer the reader to [11] for an introduction to the KPZ class.

For the Hammersley problem, critical exponents have been shown to exist and we have $\chi = 1/3$ ([3]) and $\xi = 2/3$ ([16]). The proof of $\chi = 1/3$ requires *hard* analysis and the deep connection between ℓ_n and the shape of random Young tableaux (see [17]). So-called *soft* arguments were gathered since ([1, 8, 9]) and lead to a weaker form of $\chi = 1/3$ but they are of course still non-trivial.

Various authors have studied variants of the Hammersley problem and last-passage percolation with an additional geometric constraint on paths. In a non-exhaustive way one can mention: convexity constraint [2], Hölder constraint [6, 7], gaps constraint [4], localization constraint [13], area below the path [5].

In particular some of the above works have studied the robustness of the membership to KPZ under such modifications of the constraints. Our goal here is mainly to illustrate the (non)-robustness of $\chi = 1/3$ for the Hammersley process when we impose a Lipschitz condition on

*Laboratoire Modal'X, UPL, Univ. Paris Nanterre, France. email: anne.laure.basdevant@normalesup.org. Work partially supported by ANR PPPP, ANR Malin and Labex MME-DII.

†CMAP, Ecole Polytechnique, France. email: gerin@cmap.polytechnique.fr Work partially supported by ANR PPPP and ANR GRAAL.

paths. It turns out that this problem is easily seen to be exactly solvable, thanks to a coupling with the classical Hammersley problem.

We now formally introduce our model. Let Ξ be a homogeneous Poisson process with intensity one in the quarter-plane $[0, +\infty)^2$. Hammersley [15] studied the problem of the maximal number of points $L(x, t)$ in an increasing path of points of $\Xi \cap ([0, x] \times [0, t])$. He used subadditivity to prove the existence of a constant $\pi/2 \leq c \leq e$ such that $L(x, t)/t \rightarrow c$ in probability and conjectured $c = 2$. This result was first obtained by Vershik-Kerov [20]. We give here a formulation due to Aldous-Diaconis [1] which provided a more probabilistic proof by exploiting the geometric construction of Hammersley:

Proposition 1 ([1], Th.5 (a)). *Let $a, b > 0$. Then*

$$f(a, b) := \lim_{t \rightarrow +\infty} \frac{L(at, bt)}{t} = 2\sqrt{ab}. \quad (1)$$

The convergence holds a.s. and in L^1 .

Let $0 < \alpha < \beta$ be fixed parameters, we introduce the partial order relation $\preceq^{\alpha, \beta}$ in $[0, +\infty)^2$ defined by

$$(x, y) \preceq^{\alpha, \beta} (x', y') \Leftrightarrow \begin{cases} x \leq x', \\ \alpha \leq \frac{y' - y}{x' - x} \leq \beta. \end{cases}$$

In the extremal case $\alpha = 0, \beta = +\infty$ then $\preceq^{0, +\infty}$ is the classical partial order on $[0, +\infty)^2$ defined by $\{x \leq x', y \leq y'\}$. For any domain $D \subset [0, +\infty)^2$, we consider the random variable $L^{\alpha, \beta}(D)$ given by the length of the longest chain in $\Xi \cap D$ with respect to $\preceq^{\alpha, \beta}$:

$$L^{\alpha, \beta}(D) = \max \left\{ L; \text{there are } (u_1, v_1), \dots, (u_L, v_L) \in \Xi \cap D \text{ such that} \right. \\ \left. (u_1, v_1) \preceq^{\alpha, \beta} (u_2, v_2) \preceq^{\alpha, \beta} \dots \preceq^{\alpha, \beta} (u_L, v_L) \right\}.$$

When D is a rectangle of the form $[0, x] \times [0, t]$, we just write $L^{\alpha, \beta}(x, t)$ instead of $L^{\alpha, \beta}(D)$. In the case $\alpha = 0, \beta = +\infty$, we recover the usual quantity $L^{0, +\infty}(x, t) := L(x, t)$ studied by Hammersley.

The first result of this paper is that we can also explicitly compute the limit of $L^{\alpha, \beta}(at, bt)/t$.

Theorem 2 (Limiting shape). *For every $a, b > 0$, every $\beta > \alpha \geq 0$, we have the following almost-sure limit:*

$$f^{\alpha, \beta}(a, b) := \lim_{t \rightarrow \infty} \frac{L^{\alpha, \beta}(at, bt)}{t} = \begin{cases} a\sqrt{\beta - \alpha} & \text{if } \frac{b}{a} > \frac{\alpha + \beta}{2}, \\ \frac{2\sqrt{(\beta a - b)(b - a\alpha)}}{\sqrt{\beta - \alpha}} & \text{if } \frac{\alpha + \beta}{2} \geq \frac{b}{a} \geq \frac{2}{1/\beta + 1/\alpha}, \\ b\sqrt{\frac{1}{\alpha} - \frac{1}{\beta}} & \text{if } \frac{b}{a} < \frac{2}{1/\beta + 1/\alpha}. \end{cases}$$

(Basic calculus shows that we always have $\frac{\alpha + \beta}{2} \geq \frac{2}{1/\beta + 1/\alpha}$.)

In the sequel, the condition $\frac{\alpha + \beta}{2} \geq \frac{b}{a} \geq \frac{2}{1/\beta + 1/\alpha}$ will be referred to as the *central case*.

A key consequence of Theorem 2 is that the limiting shape is strictly convex exactly within the central case. The proof of Theorem 2 is only simple calculus once we notice a coupling with the classical Hammersley problem (see Proposition 6 below). What is more appealing is that we obtain non-trivial dichotomies for the shapes of paths and the fluctuations. The following result




Figure 1: An example where $L^{\alpha,\beta}(x,y) = 4$. Every point (u,v) of Ξ (represented by \bullet 's) is drawn with its set of majorants for $\preceq^{\alpha,\beta}$. One maximizing path is drawn in blue.




Figure 2: The maximazing paths through $n = 1000$ points (red: $\alpha = 1/6, \beta = 6$, green: $\alpha = 1/2, \beta = 2$.)

states that in the central case all the maximizing paths are localized along the main diagonal $\{y = xb/a\}$ (asymptotically with high probability) while in the two non-central cases, optimal paths have slope respectively $\frac{\alpha+\beta}{2}$ or $\frac{2}{1/\beta+1/\alpha}$. In order to state the result, we set for $R > 0$ and a slope $\theta \in \mathbb{R}$

$$D_{R,\theta} = \{(x,y) \in \mathbb{R}_+^2 ; |y - \theta x| \leq R\}.$$

Theorem 3 (Localization of paths).

i). If $\frac{\alpha+\beta}{2} \geq \frac{b}{a} \geq \frac{2}{1/\beta+1/\alpha}$ then for every $\delta > 0$

$$\mathbb{P} \left(\begin{array}{c} \text{There is an optimal path for } L^{\alpha,\beta}(at, bt) \\ \text{not wholly contained in } D_{t\delta, \frac{b}{a}} \end{array} \right) \xrightarrow{t \rightarrow +\infty} 0.$$

ii). If $\frac{b}{a} > \frac{\alpha+\beta}{2}$ then for every $\delta > 0$

$$\mathbb{P} \left(\begin{array}{c} \text{There is an optimal path for } L^{\alpha,\beta}(at, bt) \\ \text{not wholly contained in } (0, k) + D_{t\delta, \frac{\alpha+\beta}{2}} \text{ for some } k \geq 0 \end{array} \right) \xrightarrow{t \rightarrow +\infty} 0.$$

iii). If $\frac{b}{a} < \frac{2}{1/\beta+1/\alpha}$ then for every $\delta > 0$

$$\mathbb{P} \left(\begin{array}{c} \text{There is an optimal path for } L^{\alpha,\beta}(at, bt) \\ \text{not wholly contained in } (k, 0) + D_{t\delta, \frac{2}{1/\beta+1/\alpha}} \text{ for some } k \geq 0 \end{array} \right) \xrightarrow{t \rightarrow +\infty} 0.$$

(Three cases of Theorem 3 are illustrated in the rightmost column of table p.10.)

This a folklore result that in the classical Hammersley problem (*i.e.* item (i) above with $\alpha = 0, \beta = +\infty$) asymptotically all the optimizing paths concentrate on the diagonal $y = xb/a$. We cannot date back the first appearance of this result but it can be seen as a consequence of a more general result by Deuschel-Zeitouni [12, Th.2 (ii)]. In the case $b/a > \beta$ the diagonal is not




Figure 3: Illustration of Theorem 3. Left: A maximizing path through $n = 1000$ points with $\alpha = 0.2, \beta = 3$ (central case). Right: A maximizing path with $\alpha = 0.2, \beta = 1.1$ (non-central case). In blue: a straight line with slope $(\alpha + \beta)/2$.

an admissible path so it is obvious that optimizing paths will not concentrate on it. However in the intermediate case $\beta > b/a > \frac{\alpha + \beta}{2}$ the diagonal is admissible but Theorem 3 shows that paths close to the diagonal are not optimal.

We now discuss the fluctuations of $L^{\alpha, \beta}(at, bt)$. In 1999, Baik-Deift-Johansson have identified the fluctuations of $L(at, bt)$:

Proposition 4 (Cube root fluctuations for the Hammersley problem ([3] Th.1.1)). *For every $a, b > 0$, we have the convergence in law*

$$\frac{L(at, bt) - 2\sqrt{abt}}{(\sqrt{abt})^{1/3}} \xrightarrow{t \rightarrow +\infty} \text{TW},$$

where TW is the standard Tracy-Widom distribution.

We show that the same asymptotic behaviour is true exactly within the central case:

Theorem 5 (Cube root fluctuations for $L^{\alpha, \beta}$).

i). *If $\frac{\alpha + \beta}{2} \geq \frac{b}{a} \geq \frac{2}{1/\beta + 1/\alpha}$, we have the convergence in law*

$$\frac{L^{\alpha, \beta}(at, bt) - t f^{\alpha, \beta}(a, b)}{\sigma_{a, b, \beta, \alpha}^{1/3} \times t^{1/3}} \xrightarrow{t \rightarrow +\infty} \text{TW},$$

where TW is the standard Tracy-Widom distribution and

$$\sigma_{a, b, \beta, \alpha} = \frac{1}{2} f^{\alpha, \beta} = \frac{\sqrt{(\beta a - b)(b - \alpha)}}{\sqrt{\beta - \alpha}}.$$

ii). *If $\frac{b}{a} > \frac{\beta + \alpha}{2}$ or $\frac{b}{a} < \frac{2}{1/\beta + 1/\alpha}$, then*

$$\begin{aligned} \frac{L^{\alpha, \beta}(at, bt) - t f^{\alpha, \beta}(a, b)}{t^{1/3}} &\xrightarrow{t \rightarrow +\infty} +\infty \text{ in prob.}, \\ \frac{L^{\alpha, \beta}(at, bt) - t f^{\alpha, \beta}(a, b)}{t^{1/3 + \varepsilon}} &\xrightarrow{t \rightarrow +\infty} 0 \text{ in prob. for every } \varepsilon. \end{aligned}$$




Figure 4: The picture of Fig.1 under map ϕ . In particular, the image of the maximizing path for α, β is mapped onto one maximal path in the parallelogram for the classical order \preceq .

2 Proof of the limiting shape

As mentioned earlier, the proof of Theorem 2 follows from a coupling argument between $L^{\alpha, \beta}$ and $L^{0, +\infty}$. Thanks to this coupling we can split the proof of Theorem in two: a probabilistic part (Proposition 7) and simple lemma of optimization (Lemma 8). Proofs of Proposition 7 and Lemma 8 rely on quite usual and simple arguments but we detail all the computations as they will be needed in the (more involved) proofs of Theorems 3 and 5.

Proposition 6 (Coupling with the classical Hammersley problem). *For every $x, y \geq 0$ and $\beta > \alpha \geq 0$ we have the identity*

$$L^{\alpha, \beta}([0, x] \times [0, y]) \stackrel{(d)}{=} L(\phi([0, x] \times [0, y])),$$

where $\phi(x, y)$ is defined by

$$\phi(x, y) = \frac{\beta^{1/4} \alpha^{1/4}}{\sqrt{\beta - \alpha}} \left(\beta^{1/2} x - \beta^{-1/2} y, -\alpha^{1/2} x + \alpha^{-1/2} y \right). \quad (2)$$

(In fact, we have equality in distribution of all the processes $(L^{\alpha, \beta}([0, x] \times [0, y]))_{x>0, y>0}$ and $(L(\phi([0, x] \times [0, y])))_{x>0, y>0}$, not only marginals.)

Proof of Proposition 6. The linear function ϕ has determinant 1 and is constructed so that

$$(x, y) \stackrel{\alpha, \beta}{\preceq} (x', y') \Leftrightarrow \phi(x, y) \preceq \phi(x', y').$$

Namely, straight lines $\{y = \alpha x\}$ (resp. $\{y = \beta x\}$) are mapped onto horizontal (resp. vertical) lines. Therefore the maximizing paths for the classical order \preceq in $\phi([0, x] \times [0, y])$ are exactly the images by ϕ of the maximizing paths in $[0, x] \times [0, y]$ for the order $\stackrel{\alpha, \beta}{\preceq}$.

Besides, the fact that ϕ preserves areas implies that the image of Ξ by ϕ is still a Poisson process with intensity one. \square

We now solve the classical Hammersley problem in a parallelogram:

Proposition 7. *Let $p, q, r, s > 0$ and denote P_t, Q_t, R_t, S_t the points of the plane of respective coordinates $(-pt, qt)$, $(0, 0)$, $(rt, -st)$, $(rt - pt, qt - st)$. Denote \mathcal{P}_t be the parallelogram $P_t Q_t R_t S_t$. Then*

$$\lim_{t \rightarrow \infty} \frac{L(\mathcal{P}_t)}{t} = \sup\{2\sqrt{\text{Area}(\mathcal{R})}, \text{ such that the rectangle } \mathcal{R} := [u, u'] \times [v, v'] \subset \mathcal{P}_1\}, \quad (3)$$

where $\text{Area}(\mathcal{R})$ denotes the area of \mathcal{R} .

Proof. For any u, u', v, v' such that $[u, u'] \times [v, v'] \subset \mathcal{P}_1$, we have

$$L(\mathcal{P}_t) \geq L([ut, u't] \times [vt, v't]). \quad (4)$$

By Proposition 1 we have that

$$\lim_{t \rightarrow \infty} \frac{L([ut, u't] \times [vt, v't])}{t} = 2\sqrt{(u' - u)(v' - v)} = 2\sqrt{\text{Area}(\mathcal{R})}.$$

Thus we obtain the lower bound of (3).

Let us now prove the upper bound. We discretize the boundary of \mathcal{P}_t as follows. Let $(x_i, y_i)_{1 \leq i \leq t^3}$ (resp. (x_i, y'_i)) be points of the plane such that

- For all i , (x_i, y_i) lies on the segment $[P_t, Q_t]$ or on $[Q_t, R_t]$ (resp. (x_i, y'_i) lies on the segment $[P_t, S_t]$ or on $[S_t, R_t]$)
- $x_1 = -pt$ and $x_{i+1} - x_i = (p + r)/t^2$. Thus $x_{t^3} = rt$.

Let

$$\mathcal{I} = \{(i, j), i < j \text{ and } y_i \leq y'_j\}.$$

For $(i, j) \in \mathcal{I}$, denote $R_{i,j}$ the rectangle $[x_i, x_j] \times [y_i, y'_j]$ and $\mathcal{Q}_t = \bigcup_{(i,j) \in \mathcal{I}} R_{i,j}$. Let us notice that $\mathcal{P}_t \setminus \mathcal{Q}_t$ is composed of $2t^3$ right triangles of width $(p + r)/t^2$ and of height bounded by C/t^2 for some constant C depending of p, q, r, s . Thus the area of $\mathcal{P}_t \setminus \mathcal{Q}_t$ is of order C/t . In particular

$$\lim_{t \rightarrow \infty} \mathbb{P}(\text{no point of } \Xi \text{ lies in } \mathcal{P}_t \setminus \mathcal{Q}_t) = 1.$$

Thus

$$\lim_{t \rightarrow \infty} \mathbb{P}(L(\mathcal{P}_t) = L(\mathcal{Q}_t)) = 1.$$

Due to the particular form of \mathcal{Q}_t , any increasing path included in \mathcal{Q}_t is necessarily included in a rectangle $R_{i,j}$. Thus

$$L(\mathcal{Q}_t) = \max\{L(R_{i,j}), (i, j) \in \mathcal{I}\}.$$

Hence for $\gamma > 0$

$$\begin{aligned} \mathbb{P}(L(\mathcal{Q}_t) \geq \gamma t) &= \mathbb{P}(\exists (i, j) \in \mathcal{I}, L(R_{i,j}) \geq \gamma t) \\ &\leq \sum_{(i,j) \in \mathcal{I}} \mathbb{P}(L(R_{i,j}) \geq \gamma t) \\ &\leq t^6 \max_{(i,j) \in \mathcal{I}} \mathbb{P}(L(R_{i,j}) \geq \gamma t) \end{aligned} \quad (5)$$

We conclude using, that if $\gamma > \sup\{2\sqrt{\text{Area}(\mathcal{R})}, \text{ such that the rectangle } \mathcal{R} := [u, u'] \times [v, v'] \subset \mathcal{P}_1\}$ then there exists some constant C , such that, for any $(i, j) \in \mathcal{I}$ and $t \geq 0$,

$$\mathbb{P}(L(R_{i,j}) \geq \gamma t) \leq \frac{C}{t^7}, \quad (6)$$

Indeed, this inequality is a direct consequence of [3, Th.1.2] (see also Proposition 9 below). This concludes the proof of the upper bound. \square

Hence, to obtain Theorem 2, it remains to determine the rectangle of maximal area in the parallelogram $\phi([0, a] \times [0, b])$.

Lemma 8. *Let $\mu > 1$, $c, c' > 0$ and denote P, Q, R, S the points of the plane of respective coordinates $(-c, c\mu)$, $(0, 0)$, $(c'\mu, -c')$, (σ, ρ) with $\sigma := -c + c'\mu$ and $\rho := -c' + c\mu$ so that the quadrilateral $\mathcal{P} : PQRS$ is a parallelogram. Then*

$$\max\{\text{Area}(\mathcal{R}) \text{ such that } \mathcal{R} := [u, u'] \times [v, v'] \subset \mathcal{P}\} = \begin{cases} \frac{(\rho\mu + \sigma)^2}{4\mu} & \text{if } \rho < \mu^{-1}\sigma \\ \frac{(\sigma\mu + \rho)^2}{4\mu} & \text{if } \rho > \mu\sigma, \\ \sigma\rho & \text{otherwise.} \end{cases} \quad (7)$$

More precisely,

- In the first case $\rho < \mu^{-1}\sigma$ a maximizing rectangle is given by $\mathcal{R} = [0, \mu\xi] \times [0, \xi]$ where $\xi = \frac{\rho\mu + \sigma}{2\mu}$. Every translation of \mathcal{R} by a vector $u(\mu, -1)$ for some $u \in [0, (\sigma - \mu\rho)/2\mu]$ is also maximizing (a similar result holds in the second case).
- In the third case $\mu^{-1}\sigma < \rho < \mu\sigma$ there is a unique rectangle \mathcal{R} with maximal area and $\mathcal{R} = [0, \sigma] \times [0, \rho]$.

Proof of Lemma 8. Set


$$M := \max\{\text{Area}(\mathcal{R}) \text{ such that the rectangle } \mathcal{R} := [u, u'] \times [v, v'] \subset \mathcal{P}\}.$$

We are going to identify a rectangle $\mathcal{R} := [u, u'] \times [v, v'] := ABCD$ inside \mathcal{P} with the largest area. Clearly, such rectangle must have two opposite vertices on sides of \mathcal{P} , for instance A and C , with A on $[PQ]$ or $[QR]$ and C on $[PS]$ or $[SR]$. With this convention, note first that if $ABCD \subset \mathcal{P}$, then the image by a translation of vector \overrightarrow{AQ} of this rectangle is a rectangle inside \mathcal{P} with a vertex at $(0, 0)$. Thus, we get

$$M = \max\{\text{Area}(\mathcal{R}) \text{ such that the rectangle } \mathcal{R} := [0, u] \times [0, v] \subset \mathcal{P}\}.$$

Assume first that $\sigma, \rho > 0$, so that the vertex S is in the upper right quarter plane. Then, they are two possible cases (drawn in red and blue in the picture below):

- Either $C \in [PS]$ (drawn in blue)
- Or $C \in [SR]$ (drawn in red)



We now distinguish the two cases. If $C \in [PS]$, then $C := (-c + xc'\mu, c\mu - xc')$ for some $x \in [c/(c'\mu), 1]$ since C must be in $(\mathbb{R}_+)^2$. Thus

$$\text{Area}(\mathcal{R}) = (-c + xc'\mu) \times (c\mu - xc') \quad \text{which is maximal for } x = \min\left(\frac{c}{2c'}\left(\mu + \frac{1}{\mu}\right), 1\right).$$

If $C \in [SR]$, then $C := (c'\mu - xc, -c' + xc\mu)$ for some $x \in [c'/(c\mu), 1]$ since C must be in $(\mathbb{R}_+)^2$. Thus

$$\text{Area}(\mathcal{R}) = (c'\mu - xc) \times (-c' + xc\mu) \quad \text{which is maximal for } x = \min\left(\frac{c'}{2c}\left(\mu + \frac{1}{\mu}\right), 1\right).$$

Note that, assuming $\rho, \sigma > 0$,


$$\frac{\rho}{\sigma} > \mu \iff c\mu - c' > \mu(-c + c'\mu) \iff 1 - \frac{c'}{c\mu} > -1 + \frac{c'\mu}{c} \iff \frac{c'}{2c}\left(\mu + \frac{1}{\mu}\right) < 1$$

Symmetrically, we have

$$\frac{\rho}{\sigma} < \mu^{-1} \iff \frac{c}{2c'}\left(\mu + \frac{1}{\mu}\right) < 1$$

Hence, if $\mu^{-1} < \rho/\sigma < \mu$, the maximum are in both cases obtained for $x = 1$, *i.e.* $C = S$. On the other hand, if this condition does not hold, the maximum is obtained for some $C \neq S$, thus the image of $ABCD$ by a translation of vector \overrightarrow{CS} yields another rectangle inside \mathcal{P} with the same area. Then, simple calculus yields the claimed expression for M .

Assume now that the vertex S is not in the upper right quarter plane, for example we have $\sigma > 0$ but $\rho < 0$ as in the picture below.



Then, if $A = Q = (0, 0)$, necessarily $C \in [PS]$, *i.e.* $C := (-c + xc'\mu, c\mu - xc')$ for some $x \in [0, 1]$. In this case, $\text{Area}(\mathcal{R}) = (-c + xc'\mu) \times (c\mu - xc')$ is always maximal for $x = \frac{c}{2c'}\left(\mu + \frac{1}{\mu}\right) < 1$ and we still get $M = \frac{c^2(\mu^2-1)^2}{4\mu} = \frac{(\rho\mu+\sigma)^2}{4\mu}$.

Similar arguments give the case $\sigma < 0$ but $\rho > 0$. Note that we can not have simultaneously $\sigma < 0$ and $\rho < 0$ since $\mu > 1$. \square

We now can proceed to the proof of the limiting shape. We set

$$c = \frac{\alpha^{1/4}\beta^{-1/4}b}{\sqrt{\beta-\alpha}}, \quad c' = \frac{\beta^{1/4}\alpha^{3/4}a}{\sqrt{\beta-\alpha}}, \quad \mu = \sqrt{\frac{\beta}{\alpha}} > 1 \quad (8)$$

and

$$\sigma = -c + c'\mu, \quad \rho = -c' + c\mu. \quad (9)$$

We easily check that

$$\rho/\sigma < \mu^{-1} \iff \frac{b}{a} < \frac{2}{\frac{1}{\alpha} + \frac{1}{\beta}}, \quad \rho/\sigma > \mu \iff \frac{b}{a} > \frac{\alpha + \beta}{2}. \quad (10)$$

Proof of Theorem 2. Combining Proposition 7 and Proposition 6 we obtain that

$$f^{\alpha,\beta}(a,b) = \sup\{2\sqrt{\text{Area}(\mathcal{R})}, \text{ such that the rectangle } \mathcal{R} = [u, u'] \times [v, v'] \subset \phi([0, a] \times [0, b])\}$$

where ϕ is defined by (2). Hence, $\phi([0, a] \times [0, b])$ is a parallelogram with vertices

$$P : c(-1, \mu), \quad Q : (0, 0), \quad R : c'(\mu, -1), \quad S : (\sigma, \rho),$$

where c, c', μ, σ, ρ are defined by (8),(9). We first collect the results of Lemma 8 in the case $\rho, \sigma > 0$. If we have $\frac{\rho}{\sigma} < \mu^{-1}$ then Lemma 8 yields

$$f^{\alpha,\beta}(a,b) = 2\sqrt{\frac{(\rho\mu + \sigma)^2}{4\mu}} = b\sqrt{\frac{1}{\alpha} - \frac{1}{\beta}}.$$

If $\frac{\rho}{\sigma} > \mu$ then again by Lemma 8

$$f^{\alpha,\beta}(a,b) = 2\sqrt{\frac{(\sigma\mu + \rho)^2}{4\mu}} = a\sqrt{\beta - \alpha}.$$

Finally, in the central case (which corresponds to $\mu^{-1} \leq \rho/\sigma \leq \mu$) we obtain that

$$f^{\alpha,\beta}(a,b) = 2\sqrt{\sigma\rho} = 2\sqrt{\frac{(-b + \beta a)(-\alpha a + b)}{\beta - \alpha}}.$$

For the case where σ or ρ is negative we observe that

$$\sigma < 0, \rho > 0 \iff \frac{b}{a} > \beta \left(\geq \frac{\alpha + \beta}{2} \right), \quad \sigma > 0, \rho < 0 \iff \frac{b}{a} < \alpha \left(\leq \frac{2}{1/\alpha + 1/\beta} \right)$$

so the formula claimed in Theorem 2 is also valid. \square

For the sake of the reader we sum up notation and qualitative results for the three cases when $\rho, \sigma > 0$:

After transformation ϕ		Before transformation ϕ	
	Maximizing rectangle(s) (drawn in red)		Shape of maximizing path(s)
$\rho/\sigma < \mu^{-1}$		$\frac{b}{a} < \frac{2}{1/\beta+1/\alpha}$	
$\rho/\sigma > \mu$		$\frac{b}{a} > \frac{\alpha+\beta}{2}$	
$\mu^{-1} \leq \rho/\sigma \leq \mu$		central case	

2.1 Localization of paths

This Section is devoted to the proof of Theorem 3. Thanks to the coupling of Proposition 6, it suffices to prove that in the classical Hammersley problem in a parallelogram $t \times \mathcal{P}$, any optimal path concentrates along the diagonal of one of the largest rectangles included in $t \times \mathcal{P}$ (asymptotically with high probability).

The proof relies on a simple convexity argument for the limiting shape. In the central case the proof is almost identical to that of ([18], Theorem 2 eq.(1.7)) so we only write the details in the non-central case. (This case is more involved as we know from Lemma 8 that there are several maximizing rectangles).

Proof of Theorem 3. We fix δ throughout the proof. As in Lemma 8, let $\mu > 1$, $c, c' > 0$ and denote P, Q, R, S the points of the plane of respective coordinates $(-c, c\mu)$, $(0, 0)$, $(c'\mu, -c')$, (σ, ρ) with $\sigma := -c + c'\mu$ and $\rho := -c' + c\mu$ so that the quadrilateral $\mathcal{P}_1 : PQRS$ is a parallelogram. Denote \mathcal{P}_t be the homothetic transformation of \mathcal{P}_1 by a factor t . By symmetry, we can assume without loss of generality that $\rho < \mu^{-1}\sigma$. Recall from Lemma 8 that in that case there are




Figure 5: Discretization of \mathcal{P}_t into squares of size $t\varepsilon$. In yellow, the small squares indexed by the set $\mathcal{I}_{c,\delta}^1$ and in blue the one indexed by the set $\mathcal{I}_{c,\delta}^2$. The three parallelograms $\mathcal{G}_{i,j}$, $\tilde{\mathcal{G}}_{k,l}$ and $\mathcal{R}_{i,j,k,l}$ are drawn in red.

several maximizing rectangles in \mathcal{P}_1 . For all such maximizing rectangles the diagonal has slope $1/\mu$. Let us recall some notation of Proposition 7.

Denote \mathcal{M}_t the set of maximizing paths for the classical Hammersley problem in \mathcal{P}_t . For $\varepsilon > 0$ which will be chosen later, we decompose \mathcal{P}_1 into disjoint squares of size $\varepsilon \times \varepsilon$:

$$\mathcal{P}_1 \subset \bigcup_{(i,j) \in \mathcal{I}} \left\{ [\varepsilon i, \varepsilon(i+1)) \times [\varepsilon j, \varepsilon(j+1)) \right\},$$

for some finite set \mathcal{I} . For a real c , let $D_{c,\delta}$ be the diagonal strip $D_{c,\delta} = \{|\mu y - x - c| \leq \delta/2\}$. Hence, $\mathcal{P}_1 \setminus D_{c,\delta}$ is composed of (at most) two connected components,

$$\begin{aligned} \mathcal{Q}_{c,\delta}^1 &:= \{(x, y) \in \mathcal{P}_1, \mu y - x - c > \delta/2\}, \\ \mathcal{Q}_{c,\delta}^2 &:= \{(x, y) \in \mathcal{P}_1, \mu y - x - c < -\delta/2\}. \end{aligned}$$

Let $\mathcal{I}_{c,\delta}^\ell$ for $\ell \in \{1, 2\}$ be the smallest subset of \mathcal{I} such that

$$\mathcal{Q}_{c,\delta}^\ell \subset \bigcup_{(i,j) \in \mathcal{I}_{c,\delta}^\ell} \left\{ [\varepsilon i, \varepsilon(i+1)) \times [\varepsilon j, \varepsilon(j+1)) \right\}$$

(notations are summarized in Fig. 5). Note that by homogeneity

$$\mathcal{P}_t \setminus (t \times D_{c,\delta}) \subset (t \times \mathcal{Q}_{c,\delta}^1) \cup (t \times \mathcal{Q}_{c,\delta}^2).$$

If there exists some path $\pi \in \mathcal{M}_t$ not contained in any $tD_{c,\delta}$ for any $c \in \mathbb{R}$, this implies in particular that there exists some c such that the path π intersects $\mathcal{Q}_{c,\delta}^1$ and $\mathcal{Q}_{c,\delta}^2$. Besides,

because of discretization, there exists a finite number of real numbers c_1, \dots, c_n such that, for any $c \in \mathbb{R}$, $\mathcal{I}_{c,\delta}^\ell = \mathcal{I}_{c_a,\delta}^\ell$, for some $a \in \{1, \dots, n\}$. Thus we have

$$\begin{aligned} & \left\{ \exists \pi \in \mathcal{M}_t \text{ not contained in } tD_{c,\delta} \text{ for some } c \right\} \\ \subset & \bigcup_{a=1}^n \bigcup_{(i,j) \in \mathcal{I}_{c_a,\delta}^1} \bigcup_{(k,l) \in \mathcal{I}_{c_a,\delta}^2} \left\{ \exists \pi \text{ intersecting } [t\varepsilon i, t\varepsilon(i+1)] \times [t\varepsilon j, t\varepsilon(j+1)] \text{ and } [t\varepsilon k, t\varepsilon(k+1)] \times [t\varepsilon l, t\varepsilon(l+1)] \right\}. \end{aligned} \quad (11)$$

Let $a \in \{1, \dots, n\}$, $(i, j) \in \mathcal{I}_{c_a,\delta}^1$ and $(k, l) \in \mathcal{I}_{c_a,\delta}^2$. We need to prove that the probability that there exists a maximizing path which intersects the squares $[t\varepsilon i, t\varepsilon(i+1)] \times [t\varepsilon j, t\varepsilon(j+1)]$ and $[t\varepsilon k, t\varepsilon(k+1)] \times [t\varepsilon l, t\varepsilon(l+1)]$ tends to 0 as t tends to infinity. By symmetry, we can assume that $i \leq k$ and $j \leq l$ as in Figure 5.

Let $\mathcal{G}_{i,j}$ be the parallelogram with opposite vertices $(0, 0)$ and $\varepsilon(i+1, j+1)$ and with sides parallel to the sides of \mathcal{P} . Let $\tilde{\mathcal{G}}_{k,l}$ be the parallelogram with opposite vertices $\varepsilon(k, l)$ and (σ, ρ) and with sides parallel to the sides of \mathcal{P} . Denote $\mathcal{R}_{i,j,k,l}$ the rectangle with sides parallel to the axes and with opposite vertices $\varepsilon(i, j)$ and $\varepsilon(k+1, l+1)$. Thus we get

$$\begin{aligned} & \left\{ \exists \pi \text{ intersecting } [t\varepsilon i, t\varepsilon(i+1)] \times [t\varepsilon j, t\varepsilon(j+1)] \text{ and } [t\varepsilon k, t\varepsilon(k+1)] \times [t\varepsilon l, t\varepsilon(l+1)] \right\} \\ & \subset \left\{ L(t\mathcal{G}_{i,j}) + L(t\tilde{\mathcal{G}}_{k,l}) + L(t\mathcal{R}_{i,j,k,l}) \geq L(\mathcal{P}_t) \right\}. \end{aligned}$$

We can compute the almost sure limit of $(L(t\mathcal{G}_{i,j}) + L(t\tilde{\mathcal{G}}_{k,l}) + L(t\mathcal{R}_{i,j,k,l})) / t$ using Proposition 7 and Lemma 8. We let the reader check that, if ε is chosen small enough, for any $(i, j) \in \mathcal{I}_{c,\delta}^1$ and $(k, l) \in \mathcal{I}_{c,\delta}^2$, this limit is strictly smaller than the limit of $L(\mathcal{P}_t) / t$. This computation is very similar to the proof of Lemma 8. It is a consequence of the fact that the function

$$\begin{aligned} (\mathcal{P}_t)^2 & \rightarrow \mathbb{R}^2 \\ (a, b), (\alpha, \beta) & \mapsto \tilde{f}(a, b) + \tilde{f}(\alpha - a, \beta - b) + \tilde{f}(\sigma - \alpha, \rho - \beta) \end{aligned}$$

(where $\tilde{f}(a, b) = 2\sqrt{ab}\mathbf{1}_{a,b>0}$) is strictly maximized when (a, b) and (α, β) belong to a straight line with slope μ^{-1} . Thus, we deduce that

$$\lim_{t \rightarrow \infty} \mathbb{P} \left\{ L(t\mathcal{G}_{i,j}) + L(t\tilde{\mathcal{G}}_{k,l}) + L(t\mathcal{R}_{i,j,k,l}) \geq L(\mathcal{P}_t) \right\} = 0.$$

Therefore the event on the LHS of (11) is included in a finite union of events (and the number of those events does not depend on t) whose probabilities tend to zero.

This proves that in the classical Hammersley problem in a parallelogram optimal paths concentrate along one of the maximizing diagonals. The proof of Theorem 3 follows from the coupling of Proposition 6. \square

3 Fluctuations

3.1 The standard case : Proof of Theorem 5 (i)

We first recall some known facts regarding the fluctuations for the classical Hammersley problem.

Proposition 9. *For every $m > 0$, there exists $C_m > 0$ such that for every rectangle \mathcal{R} with sides parallel to the axes,*

$$\mathbb{P}\left(L(\mathcal{R}) \geq 2\sqrt{\text{Area}(\mathcal{R})} + \text{Area}(\mathcal{R})^{1/6+\varepsilon}\right) \leq \frac{C_m}{\text{Area}(\mathcal{R})^{\varepsilon m}}. \quad (12)$$

Proof. The claimed inequality is obtained by applying Markov's inequality to ([3, Th.1.2]). \square

With the help of Proposition 4, we will have proved Theorem 5 as soon as we establish that, inside a large parallelogram \mathcal{P} with a unique optimal rectangle \mathcal{R} , the fluctuations of $L(\mathcal{P})$ are equal to the fluctuations of $L(\mathcal{R})$, *i.e.* non-optimal rectangles do not modify them. To this end, we show the following proposition.

Proposition 10. *Let $\mu > 1$ and $\sigma, \rho > 0$. Let denote $\mathcal{P}_1 = PQRS$ the parallelogram with $Q = (0, 0)$, $S = (\sigma, \rho)$ and such that $P = c(-1, \mu)$ and $R = c'(\mu, -1)$ for some $c, c' > 0$. For $t \geq 0$, let \mathcal{P}_t be the homothetic transformation of \mathcal{P}_1 by a factor t . Assume $\mu^{-1} \leq \rho/\sigma \leq \mu$. Then*

$$\lim_{t \rightarrow \infty} \mathbb{P}(L(\mathcal{P}_t) \geq L(\sigma t, \rho t) + t^{\frac{1}{3} - \frac{1}{32}}) = 0.$$

(Observe that we always have the inequality $L(\mathcal{P}_t) \geq L(\sigma t, \rho t)$.)

Let us first explain why this proposition implies Theorem 5.

Proof of Proposition 10 \Rightarrow Theorem 5. Using Proposition 6, we know that $L^{\alpha, \beta}(at, bt)$ has the law of $L(\mathcal{P}_t)$ where \mathcal{P}_t is the homothetic transformation of $PQRS$ by a factor t , where

$$P : c(-1, \mu), \quad Q : (0, 0), \quad R : c'(\mu, -1), \quad S : (\sigma, \rho),$$

and c, c', μ, σ, ρ are defined by (8),(9).

Recall from (10) that the central case $\frac{2}{1/\alpha + 1/\beta} \leq \frac{b}{a} \leq \frac{\beta + \alpha}{2}$ corresponds to $0 < \mu^{-1} < \rho/\sigma < \mu$. Thus, \mathcal{P}_t is an homothetic transformation of a parallelogram satisfying the condition of Proposition 10. Hence, we get that

$$\frac{L^{\alpha, \beta}(at, bt) - L(\sigma t, \rho t)}{t^{1/3}}$$

converges in probability to 0. We conclude using Proposition 4 and noticing that $2\sqrt{\sigma\rho} = f^{\alpha, \beta}(a, b)$. \square

Proof of Proposition 10. To prove Proposition 10 we need to bound, for any rectangle of the form

$$\mathcal{R}_{uv} := [-u, \sigma t + v] \times [\mu^{\text{sign}(u)}u, \rho t - \mu^{\text{sign}(v)}v]$$

inside \mathcal{P}_t (and with two vertices on the boundary of \mathcal{P}_t), the probability that $L(\mathcal{R}_{uv}) - L(\sigma t, \rho t)$ is larger than $t^{1/3-1/32}$. In fact, we will establish the following equation

$$\lim_{t \rightarrow \infty} \mathbb{P}\left(\sup_{(u, v) \in \mathcal{U}} L(\mathcal{R}_{uv}) - L(\sigma t, \rho t) \geq t^{1/3-1/32}\right) = 0 \quad (13)$$

where \mathcal{U} is the set of value (u, v) for which \mathcal{R}_{uv} is inside \mathcal{P}_t . To this end, we set

$$\kappa = \frac{1}{3} + \frac{1}{32}$$

and we treat separately two cases.

- (i) Either $|u| > t^\kappa$ or $|v| > t^\kappa$. Then the rectangle \mathcal{R}_{uv} is sufficiently smaller than the optimal rectangle $[0, \sigma t] \times [0, \rho t]$ such that the law of large number for $L(\mathcal{R}_{uv})$ will insure that it is very unlikely that $L(\mathcal{R}_{uv}) \geq 2\sqrt{\sigma\rho}t - t^{1/3+\varepsilon}$.
- (ii) Or $|u| < t^\kappa$ and $|v| < t^\kappa$. Then \mathcal{R}_{uv} and $[0, \sigma t] \times [0, \rho t]$ are so closed to each other than necessarily $L(\mathcal{R}_{uv}) - L(\sigma t, \rho t)$ can not be too large.

The case (i): $|u| > t^\kappa$ or $|v| > t^\kappa$

By symmetry, we only consider here the case $u > t^\kappa$ and $v \geq 0$ but the reader can easily adapt the proof to the other cases. As in the proof of Proposition 7, we discretize the problem. More precisely, let $u_i = v_i = i/t^2$ and denote \mathcal{I} the set of indices (i, j) such that $\mathcal{R}_{u_i v_j}$ satisfies $u_i > t^\kappa$ and $v_j \geq 0$ and is inside \mathcal{P}_t namely

$$\mathcal{I} = \{(i, j) \in \mathbb{Z}_+^2, u_i > t^\kappa, (u_i, v_j) \in \mathcal{U}\}.$$

As in the proof of Proposition 7, we have, with probability tending to 1 as t tend to infinity

$$\sup\{L(\mathcal{R}_{uv}), (u, v) \in \mathcal{U}, u > t^\kappa, v \geq 0\} = \sup\{L(\mathcal{R}_{u_i v_j}), (i, j) \in \mathcal{I}\} \quad (14)$$

and we have $|\mathcal{I}| \leq Ct^6$ for some constant $C := C(\sigma, \rho, \mu)$. Besides, we have

$$\text{Area}(\mathcal{R}_{u_i v_j}) = \sigma\rho t^2 - t(u_i + v_j)(\sigma\mu - \rho) - \mu(u_i + v_j)^2 \leq \sigma\rho t^2 - t^{\kappa+1}(\sigma\mu - \rho).$$

Hence

$$\sqrt{\sigma\rho}t - \sqrt{\text{Area}(\mathcal{R}_{u_i v_j})} = \frac{\sigma\rho t^2 - \text{Area}(\mathcal{R}_{u_i v_j})}{\sqrt{\sigma\rho}t + \sqrt{\text{Area}(\mathcal{R}_{u_i v_j})}} \geq \frac{\sigma\rho t^2 - \text{Area}(\mathcal{R}_{u_i v_j})}{2\sqrt{\sigma\rho}t} \geq \frac{\sigma\mu - \rho}{2\sqrt{\sigma\rho}}t^\kappa.$$

Thus

$$\mathbb{P}(L(\mathcal{R}_{u_i v_j}) \geq 2\sqrt{\sigma\rho}t - t^{1/3+\varepsilon}) \leq \mathbb{P}(L(\mathcal{R}_{u_i v_j}) - 2\sqrt{\text{Area}(\mathcal{R}_{u_i v_j})} \geq \frac{\sigma\mu - \rho}{\sqrt{\sigma\rho}}t^\kappa - t^{1/3+\varepsilon}).$$

Take $\varepsilon = 1/65$ such that $1/3+2\varepsilon < \kappa$ and so that t^κ is asymptotically larger than $\text{Area}(\mathcal{R}_{u_i v_j})^{1/6+\varepsilon}$. Then Proposition 9 yields the existence of a constant $C := C(\sigma, \rho, \mu)$ such that

$$\forall (i, j) \in \mathcal{I}, \quad \mathbb{P}(L(\mathcal{R}_{u_i v_j}) \geq 2\sqrt{\text{Area}(\mathcal{R}_{u_i v_j})} + \frac{\sigma\mu - \rho}{\sqrt{\sigma\rho}}t^\kappa) \leq \frac{C}{t^7}$$

and finally

$$\mathbb{P}(\sup_{(i, j) \in \mathcal{I}} L(\mathcal{R}_{u_i v_j}) \geq 2\sqrt{\sigma\rho}t - t^{1/3+\varepsilon}) \leq |\mathcal{I}| \frac{C}{t^7} \leq \frac{\tilde{C}}{t}.$$

Case $u \geq t^\kappa, v \leq 0$ is treated in the same way and the cases where $u < -t^\kappa$ hold by symmetry. Besides, Proposition 9 implies that

$$\lim_{t \rightarrow \infty} \mathbb{P}(L(\sigma t, \rho t) \leq 2\sqrt{\sigma\rho}t - t^{1/3+\varepsilon}) = 0.$$

Combining this with (14) we get

$$\lim_{t \rightarrow \infty} \mathbb{P}(\sup\{L(\mathcal{R}_{uv}), (u, v) \in \mathcal{U}, |u| > t^\kappa \text{ or } |v| > t^\kappa\} - L(\sigma t, \rho t) \geq 0) = 0.$$




Figure 6: Summary of notations: sources, sinks, Hammersley lines. On this sample we have $L_\lambda(x, y) = 6$ and an optimal path (drawn in dark red) passes through two sources.

The case (ii): $|u| < t^\kappa$ and $|v| < t^\kappa$

This case is more delicate than the previous one. We discretize again the boundary of \mathcal{P}_t but with a different scaling than before. All along case (ii) we put

$$\delta = \frac{1}{3} - \frac{1}{16} \quad \text{and} \quad u_i = v_i = it^\delta \quad \text{for} \quad |i|, |j| \leq t^{\kappa-\delta}.$$

The main ideas of the proof of case (ii) are inspired by the proof of Theorem 2 in [10] and require some important tools regarding the Hammersley process. In particular, we will use

- the graphical representation of *Hammersley lines* (implicitly introduced by Hammersley [15]) ;
- the *sinks/sources* approach introduced in [8].

Sinks and sources. We introduce the Hammersley process with *sinks* and *sources*. For $\lambda > 0$ we consider the point process Ξ^λ on the plane defined by

$$\Xi^\lambda := \Xi \cup \mathcal{M}_\lambda \cup \mathcal{N}_\lambda$$

where \mathcal{M}_λ is a one dimensional Poisson point process of intensity λ on the x -axis, \mathcal{N}_λ is a one dimensional Poisson point process of intensity λ^{-1} on the y -axis (and $\Xi, \mathcal{M}_\lambda, \mathcal{N}_\lambda$ are independent). A point on \mathcal{M}_λ (resp. on \mathcal{N}_λ) is called a source (resp. a sink).

We will modify L by considering additional points of $\mathcal{M}_\lambda \cup \mathcal{N}_\lambda$:

$$L_\lambda(x, y) = \max \left\{ L; \text{there are } (u_1, v_1), \dots, (u_L, v_L) \in \Xi \cup \mathcal{M}_\lambda \cup \mathcal{N}_\lambda \text{ such that} \right. \\ \left. (0, 0) \preceq (u_1, v_1) \preceq (u_2, v_2) \preceq \dots \preceq (u_L, v_L) \preceq (x, y) \right\}.$$

Hammersley lines. For $1 \leq \ell \leq L_\lambda(x, y)$ the ℓ -th *Hammersley line* \mathcal{H}_λ^ℓ is defined as the lowest level set of level ℓ for the function $(x, y) \mapsto L_\lambda(x, y)$ (see Fig. 6, Hammersley lines are drawn in

blue). It is easy to see that by construction every Hammersley line is a simple curve composed of an alternation of South/East straight lines.

It was shown and exploited by Cator and Groeneboom [14, 8] that for the choice of intensities λ (sources) and λ^{-1} (sinks) the process of Hammersley lines is stationary either seen from bottom to top or from left to right. As a consequence we obtain the following non-asymptotic estimate.

Lemma 11 ([8], Theorem 3.1). *For every $0 < x_0 < x_1$ and $0 < y_0 < y_1$ we have:*

- **Stationarity of sources.**

$$\mathcal{S}_1 := L_\lambda(x_1, y_0) - L_\lambda(x_0, y_0) \stackrel{(d)}{=} \text{Poisson r.v. with mean } \lambda(x_1 - x_0).$$

- **Stationarity of sinks.** *For every $x > 0$ and every $y_0 < y_1$,*

$$\mathcal{S}_2 := L_\lambda(x_0, y_1) - L_\lambda(x_0, y_0) \stackrel{(d)}{=} \text{Poisson r.v. with mean } \frac{1}{\lambda}(y_1 - y_0).$$

- *Furthermore, \mathcal{S}_1 and \mathcal{S}_2 are independent.*

We now introduce the random variable $Z_\lambda(x, y)$ which represents the maximal length that an optimal path spends on the x -axis:

$$Z_\lambda(x, y) = \sup \{ \xi \geq 0; L_\lambda(x, y) = \text{card}(\mathcal{M}_\lambda \cap [0, \xi]) + L([\xi, x] \times [0, y]) \},$$

where $\sup \emptyset = 0$.

Lemma 12. *Let $\lambda_+^* = \sqrt{\rho/\sigma}(1+t^{-1/4})$. There exists some constant C_1 such that, for any $t > 0$,*

$$\mathbb{P}(Z_{\lambda_+^*}(\sigma t, \rho t) = 0) \leq \frac{C_1}{t^{1/4}}.$$

Proof. Lemma 6 of [10] states that there exists some constant $C > 0$ such that, for any $t \geq 0$,

$$\mathbb{P}(Z_{1+t^{-1/4}}(t, t) = 0) \leq \frac{C}{t^{1/4}}.$$

We just need here to establish an analogue result for a rectangle. Using the invariance of a Poisson process under the transformation $(x, y) \mapsto (x\sqrt{\rho/\sigma}, y\sqrt{\sigma/\rho})$, we get the equality in distribution:

$$Z_{\lambda_+^*}(\sigma t, \rho t) \stackrel{\text{law}}{=} Z_{\lambda_+^* \sqrt{\sigma/\rho}}(\sqrt{\sigma \rho t}, \sqrt{\sigma \rho t}).$$

Since $\lambda_+^* = \sqrt{\rho/\sigma}(1+t^{-1/4})$ we get

$$\mathbb{P}(Z_{\lambda_+^*}(\sigma t, \rho t) = 0) \leq \mathbb{P}(Z_{1+t^{-1/4}}(\sqrt{\sigma \rho t}, \sqrt{\sigma \rho t}) = 0) \leq \frac{C}{(\sigma \rho)^{1/8} t^{1/4}}.$$


□

We now state a deterministic lemma.

Lemma 13. *For any $y \geq x$, $s \leq t$ and $\lambda > 0$, if the event $Z_\lambda(x, t) > 0$ occurs then we have*

$$L(y, s) - L(x, t) \leq L_\lambda(y, s) - L_\lambda(x, t).$$

Proof. For $s = t$ this is Lemma 2 in [10]. The diagram below explains how it can be extended for every $s \leq t$: the dashed line represents an optimal path for $L_\lambda(x, t)$ while the plain line represents an optimal path for $L(y, s)$. On the event $Z_\lambda(x, t) > 0$ the dashed line crosses the plain line and these paths can be decomposed respectively as ②+④ and ①+③ as follows:



We have that

$$L(y, s) = \textcircled{1} + \textcircled{3}, \quad L(x, t) \geq \textcircled{1} + \textcircled{4}$$

from which we deduce $L(y, s) - L(x, t) \leq \textcircled{3} - \textcircled{4}$. On the other hand

$$L_\lambda(y, s) \geq \textcircled{2} + \textcircled{3}, \quad L_\lambda(x, t) = \textcircled{2} + \textcircled{4}$$

from which we deduce $L_\lambda(y, s) - L_\lambda(x, t) \geq \textcircled{3} - \textcircled{4}$. \square

We will use the following exponential concentration inequalities (see Sect.2.4.1 in [19]):

Lemma 14. *If X is a Poisson random variable with mean θ and $\varepsilon > 0$ then*

$$\mathbb{P}(X \geq \theta + \varepsilon) \leq \exp\left(-\frac{\varepsilon^2}{2(\theta + \varepsilon)}\right), \quad \mathbb{P}(X \leq \theta - \varepsilon) \leq \exp\left(-\frac{\varepsilon^2}{2(\theta + \varepsilon)}\right). \quad (15)$$

As a consequence, there exists $C' > 0$ such that if X, Y are Poisson random variables with respective means $\theta < \theta'$ then for every $10(\theta' - \theta) \leq \varepsilon \leq \theta/10$ then

$$\mathbb{P}(|X - Y| \geq \varepsilon) \leq 2 \exp\left(-\frac{C' \varepsilon^2}{\theta'}\right). \quad (16)$$

We first establish Proposition 10 outside the critical cases $\rho/\sigma \in \{\mu^{-1}, \mu\}$ by proving the following lemma.

Lemma 15. *Assume $\mu^{-1} < \rho/\sigma < \mu$. There exists C_1 (depending on σ, ρ, μ) such that, for any $i, j \in [-t^{\kappa-\delta}, t^{\kappa-\delta}]$ and $t \geq 1$,*

$$\mathbb{P}(L(\mathcal{R}_{u_i, v_j}) \geq L(\sigma t, \rho t)) \leq \frac{C_2}{t^{1/4}}. \quad (17)$$

Proof. We begin by proving the lemma for $i = 0$ and $j \geq 0$ i.e. showing that there exists C_2 such that, for any $j \in \mathbb{Z}_+$ and $t \geq 1$,

$$\mathbb{P}\left(L(\sigma t + jt^\delta, \rho t - \mu jt^\delta) \geq L(\sigma t, \rho t)\right) \leq \frac{C_2}{t^{1/4}}. \quad (18)$$

As in Lemma 12, let $\lambda_+^* = \sqrt{\rho/\sigma}(1 + t^{-1/4})$. Using Lemma 13, we have

$$\begin{aligned} \mathbb{P}(L(\mathcal{R}_{0, v_j}) \geq L(\sigma t, \rho t)) &\leq \mathbb{P}(Z_{\lambda_+^*}[\sigma t, \rho t] = 0) + \mathbb{P}\left(L_{\lambda_+^*}(\mathcal{R}_{0, v_j}) \geq L_{\lambda_+^*}(\sigma t, \rho t)\right) \\ &\leq \frac{C_1}{t^{1/4}} + \mathbb{P}\left(L_{\lambda_+^*}(\mathcal{R}_{0, v_j}) \geq L_{\lambda_+^*}(\sigma t, \rho t)\right). \end{aligned} \quad (19)$$

In order to bound the last term in the RHS of (19) we write

$$L_{\lambda_+^*}(\mathcal{R}_{0, v_j}) - L_{\lambda_+^*}(\sigma t, \rho t) = \left(L_{\lambda_+^*}(\mathcal{R}_{0, v_j}) - L_{\lambda_+^*}(\sigma t, \rho t - \mu jt^\delta)\right) - \left(L_{\lambda_+^*}(\sigma t, \rho t) - L_{\lambda_+^*}(\sigma t, \rho t - \mu jt^\delta)\right).$$

Using the stationarity of the Hammersley process with sources and sink (Lemma 11), we know that both terms in the RHS of the above equation has a Poisson distribution. More precisely,

$$\mathbb{P}\left(L_{\lambda_+^*}(\mathcal{R}_{0v_j}) \geq L_{\lambda_+^*}(\sigma t, \rho t)\right) = \mathbb{P}\left(\text{Poisson}(\lambda_+^* j t^\delta) \geq \text{Poisson}\left(\frac{\mu}{\lambda_+^*} j t^\delta\right)\right), \quad (20)$$

where the two Poisson random variables on the RHS are independent.

For t large enough, λ_+^* tends to $\sqrt{\rho/\sigma}$. Therefore the mean of the first Poisson r.v. in (20) becomes strictly smaller than the second mean (here we use $\rho/\sigma < \mu$). Hence, (15) implies the existence of a constant $C_2 := C_2(\sigma, \rho, \mu, j)$ satisfying (18).

This concludes the proof of Lemma 15 for $i = 0, j > 0$. The proof of the case $i = 0, j < 0$ is identical (using $\mu^{-1} < \rho/\sigma$).

To extend the formula to $i \neq 0$, we write

$$L(\mathcal{R}_{u_i v_j}) - L(\sigma t, \rho t) = \left(L(\mathcal{R}_{u_i v_j}) - L(\mathcal{R}_{0v_j})\right) + \left(L(\mathcal{R}_{0v_j}) - L(\sigma t, \rho t)\right)$$

Using the same argument has before in the rectangle \mathcal{R}_{0v_j} instead of \mathcal{R}_{00} we still have that λ_+^* (which depends on j) is $\sqrt{\rho/\sigma} + o(1)$ uniformly in j (here we use that $|j t^\delta| \leq t^\kappa \ll t$).

Therefore we can bound the probability of the event $\{L(\mathcal{R}_{u_i v_j}) - L(\mathcal{R}_{0v_j}) \geq 0\}$ uniformly for $j \in [-t^{\kappa-\delta}, t^{\kappa-\delta}]$. □

Since $\text{Card}\{|i|, |j| \leq t^{\kappa-\delta}\} \sim 4t^{2\kappa-2\delta} = 4t^{3/16} = o(t^{1/4})$, we get using Lemma 15 that

$$\lim_{t \rightarrow \infty} \mathbb{P}\left(\sup_{|i|, |j| \leq t^{\kappa-\delta}} L(\mathcal{R}_{u_i v_j}) \geq L(\sigma t, \rho t)\right) = 0. \quad (21)$$

Now, for $i, j \geq 0$, we have

$$\begin{aligned} \sup_{\substack{u \in [u_i, u_{i+1}] \\ v \in [v_j, v_{j+1}]}} L(\mathcal{R}_{uv}) &\leq \max(L(\mathcal{R}_{u_i v_j}), L(\mathcal{R}_{u_i v_{j+1}}), L(\mathcal{R}_{u_{i+1} v_j}), L(\mathcal{R}_{u_{i+1} v_{j+1}})) \\ &+ L([-u_{i+1}, -u_i] \times [\mu u_i, \mu u_{i+1}]) + L([\sigma t + v_i, \sigma t + v_{i+1}] \times [\rho t - \mu v_{i+1}, \rho t - \mu v_i]). \end{aligned}$$

The random variables $L([-u_{i+1}, -u_i] \times [\mu u_i, \mu u_{i+1}])$ and $L([\sigma t + v_i, \sigma t + v_{i+1}] \times [\rho t - \mu v_{i+1}, \rho t - \mu v_i])$ has both the same distribution as $L(t^\delta, \mu t^\delta)$. Since $\delta < 1/3 - 1/32$, using Proposition 9, we get

$$\mathbb{P}(L([-u_{i+1}, -u_i] \times [\mu u_i, \mu u_{i+1}]) \geq t^{1/3-1/32}) \leq \frac{C}{t}.$$

Thus, an union bound implies Proposition 10 in the case $\mu^{-1} < \rho/\sigma < \mu$. □

The case $\rho/\sigma \in \{\mu^{-1}, \mu\}$. We now explain how to modify the previous proof in the critical cases where $\rho/\sigma = \mu$. Using the same argument as above, Proposition 10 will still be proved if, instead of (17), we show that

$$\mathbb{P}\left(L(\mathcal{R}_{u_i v_j}) \geq L(\sigma t, \rho t) + t^{1/3-1/32}\right) \leq \frac{C_2}{t^{1/4}}.$$

With the same sinks/sources strategy we get that

$$\begin{aligned} \mathbb{P}\left(L(\mathcal{R}_{u_i v_j}) \geq L(\sigma t, \rho t) + t^{1/3-1/32}\right) &\leq \\ &\frac{C_1}{t^{1/4}} + \mathbb{P}\left(\text{Poisson}(\lambda_+^* j t^\delta) \geq \text{Poisson}(\mu j t^\delta / \lambda_+^*) + t^{1/3-1/32}\right), \\ &= \frac{C_1}{t^{1/4}} + \mathbb{P}\left(\text{Poisson}(\sqrt{\rho/\sigma} j t^\delta (1 + t^{-1/4})) \geq \text{Poisson}(\sqrt{\rho/\sigma} j t^\delta (1 + t^{-1/4})^{-1}) + t^{1/3-1/32}\right) \end{aligned}$$




Figure 7: Example of a parallelogram with $\rho/\sigma < \mu^{-1}$.

which is indeed a $\mathcal{O}(t^{-1/4})$ using (16) with:

$$\begin{aligned}\theta' - \theta &\leq C_3 j t^{\delta-1/4} = \mathcal{O}(t^{\kappa-\delta+\delta-1/4}) = \mathcal{O}(t^{1/12+1/32}), \\ \theta' &\geq \sqrt{\rho/\sigma} j t^\delta \geq \sqrt{\rho/\sigma} t^{1/3-1/16}, \\ \varepsilon &= t^{1/3-1/32}.\end{aligned}$$

3.2 Fluctuations: The non-central case. Proof of Theorem 5 (ii).

Fluctuations are of order $\gg t^{1/3}$. We will show that, in the non-central case, for all $x \in \mathbb{R}$

$$\lim_{t \rightarrow \infty} \mathbb{P} \left(\frac{L^{\alpha,\beta}(at, bt) - t f^{\alpha,\beta}(a, b)}{t^{1/3}} \leq x \right) = 0.$$

The key argument is to exploit a result by Johansson [16] for the *transversal fluctuations* in the Hammersley problem. We make the assumption that

$$\frac{b}{a} < \frac{2}{\frac{1}{\alpha} + \frac{1}{\beta}} \Leftrightarrow \rho/\sigma < \mu^{-1}$$

where, as in the proof of Theorem 2, ρ, σ, μ are defined in (8),(9). Let $\mathcal{P}_1 = PQRS$ be the parallelogram with $Q = (0, 0)$, $S = (\sigma, \rho)$ and such that $P = c(-1, \mu)$ and $R = c'(\mu, -1)$ for some $c, c' > 0$. Recall Lemma 8: if $\xi := \frac{\rho\mu + \sigma}{2\mu}$, the rectangle $\mathcal{R} := [0, \mu\xi] \times [0, \xi]$ is a maximizing rectangle inside \mathcal{P}_1 . Any translation of this rectangle by a vector $u(\mu, -1)$ with $u \in [0, (\sigma - \mu\rho)/2\mu]$ remains a maximizing rectangle. Denote \mathcal{R}_t a homothetic transformation by a factor t of \mathcal{R} and let

$$\mathcal{S}_t := \left\{ (x, y) \in \mathcal{R}_t, |\mu y - x| \leq t^{3/4} \right\}$$

be a "small cylinder" around the diagonal of \mathcal{R}_t . The following Lemma is a weak consequence of [16, Th.1.1] (take $\gamma = 3/4$ in eq.(1.6) in [16]).

Lemma 16 (Johansson (2000)).

$$\lim_{t \rightarrow \infty} \mathbb{P}(L(\mathcal{S}_t) = L(\mathcal{R}_t)) = 1.$$

In particular,

$$N_t := \frac{L(\mathcal{S}_t) - 2\sqrt{\mu\xi t}}{(\sqrt{\mu\xi t})^{1/3}}$$

tends to a Tracy-Widom distribution. Let $\mathcal{S}_t(u)$ be the image of \mathcal{S}_t after a translation of vector $ut(\mu, -1)$. We clearly have

$$L(\mathcal{P}_t) \geq \sup_{0 \leq u \leq (\sigma - \mu\rho)/2\mu} L(\mathcal{S}_t(u)).$$

Moreover, the random variables $L(\mathcal{S}_t(u))$ and $L(\mathcal{S}_t(u'))$ have the same distribution and are independent if $|u - u'| \geq 2t^{-1/4}$, since, in this case, $\mathcal{S}_t(u) \cap \mathcal{S}_t(u') = \emptyset$. Hence, we get, with $c_0 := \frac{\sigma - \mu\rho}{4\mu}$, the stochastic lower bound

$$\frac{L(\mathcal{P}_t) - 2\sqrt{\mu\xi t}}{(\sqrt{\mu\xi t})^{1/3}} \succcurlyeq \max_{1 \leq i \leq c_0 t^{1/4}} N_t^i$$

where $(N_t^i)_{i \geq 1}$ are i.i.d. random variables with common distribution N_t .

Therefore, for every real x ,

$$\mathbb{P}\left(\frac{L(\mathcal{P}_t) - 2\sqrt{\mu\xi t}}{(\sqrt{\mu\xi t})^{1/3}} \geq x\right) \leq \mathbb{P}(N_t^1 \geq x)^{c_0 t^{1/4}} \xrightarrow{t \rightarrow +\infty} 0.$$

Fluctuations are of order $\ll t^{1/3+\varepsilon}$. Here we can re-use the strategy of the proof of the limiting shape. If we take (5),(6) with

$$\gamma t = t f^{\alpha, \beta}(a, b) + At^{1/3+\varepsilon}$$

for a given real A then we obtain with Proposition 9 that

$$\mathbb{P}\left(L(\mathcal{P}_t) \geq t f^{\alpha, \beta}(a, b) + At^{1/3+\varepsilon}\right) \rightarrow 0.$$

Concluding remarks

- There is a little room left between $t^{1/3}$ and $t^{1/3+\varepsilon}$ in the non-central case of Theorem 5. We leave open the question of the exact order of fluctuations.
- Proposition 7 (Hammersley problem in a parallelogram) can be generalized to an arbitrary convex domain D . We would get that

$$\lim_{t \rightarrow \infty} \frac{L(t \times D)}{t} = \sup\{2\sqrt{(v' - v)(u' - u)}\}, \quad (u, v) \preccurlyeq (u', v') \in D\}.$$

Indeed, the proof of the upper bound is easily generalized. Regarding the lower bound, (4) is not relevant anymore as the whole rectangle $[u, v] \times [u', v']$ may not be included in \mathcal{P}_1 . In order to generalize the proof we need to use the localization of the longest path in $[u, v] \times [u', v']$ along the diagonal.

References

- [1] D.Aldous, P.Diaconis. Hammersley's interacting particle process and longest increasing subsequences. *Probability Theory and Related Fields* vol.103 (1995), n.2, p.199-213.

- [2] G. Ambrus and I. Bárány. Longest convex chains. *Random Struct. Alg.*, vol. 35 (2009), n.2, p.137-162.
- [3] J. Baik, P. A. Deift and K. Johansson, On the distribution of the length of the longest increasing subsequence of random permutations. *Journal of the American Mathematical Society*, vol.12 (1999) n.4, p.1119-1178.
- [4] A.-L. Basdevant, L.Gerin. Longest increasing paths with gaps. *ALEA, Lat. Am. J. Probab. Math. Stat.* vol.16 (2019), p.1141-1163.
- [5] R.Basu, S.Ganguly, A.Hammond. The competition of roughness and curvature in area-constrained polymer models. *Communications in Mathematical Physics*, vol.364 (2018), n.3, p.1121-1161.
- [6] Q. Berger, N. Torri. Beyond Hammersley’s Last-Passage Percolation: a discussion on possible local and global constraints. *arXiv:1802.04046* (2018).
- [7] Q. Berger, N. Torri. Entropy-controlled Last-Passage Percolation. *Ann. Appl. Probab.* vol.29 (2019), n.3, p.1878-1903.
- [8] E. Cator, P. Groeneboom. Hammersley’s process with sources and sinks. *Annals of Probability* vol.33 (2005), n.3, p.879-903.
- [9] E.Cator, P.Groeneboom. Second class particles and cube root asymptotics for Hammersley’s process. *Annals of Probability* vol.34 (2006) n.4, p.1273-1295.
- [10] E.Cator, L.P.R. Pimentel, On the local fluctuations of last-passage percolation models. *Stochastic Processes and their Applications*, vol.125 (2015) n.2 p.538-551.
- [11] I.Corwin. Kardar-Parisi-Zhang universality. *Notices of the AMS*, vol.63 (2016) n.3, p.230-239.
- [12] J.D.Deuschel, O.Zeitouni. Limiting curves for i.i.d. records. *Annals of Probability* vol.23 (1995), n.2, p.852-878.
- [13] P.Dey, M.Joseph, R.Peled. Longest increasing path within the critical strip. *arXiv:1808.08407* (2018).
- [14] P. Groeneboom. Hydrodynamical methods for analyzing longest increasing subsequences. *J. Comput. Appl. Math.* vol. 142 (2002), p.83-105
- [15] J.M.Hammersley. A few seedlings of research. *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability*, vol.1 (1972), p.345-394.
- [16] K.Johansson. Transversal fluctuations for increasing subsequences on the plane. *Probability theory and related fields*, vol. 116 (2000) n.4, p.445-456.
- [17] D. Romik. *The Surprising Mathematics of Longest Increasing Subsequences*. Cambridge University Press (2015).
- [18] T.Seppäläinen. Increasing sequences of independent points on the planar lattice. *Annals of Applied Probability* vol.7 (1997) n.4 p.886-898.
- [19] R.Van Der Hofstad. *Random graphs and complex networks* (2009). Available on <http://www.win.tue.nl/rhofstad/>

- [20] A. M. Veršik and S. V. Kerov. Asymptotic behavior of the Plancherel measure of the symmetric group and the limit form of Young tableaux. *Dokl. Akad. Nauk SSSR*. 233 (1977) n.6 p.1024-1027.