



HAL
open science

Areas of muscle tissue alteration can differ from activated regions during electrically-induced isometric contractions

Alexandre Fouré, Arnaud Le Troter, Augustin C. Ogier, Maxime Guye, Julien Gondin, David Bendahan

► **To cite this version:**

Alexandre Fouré, Arnaud Le Troter, Augustin C. Ogier, Maxime Guye, Julien Gondin, et al.. Areas of muscle tissue alteration can differ from activated regions during electrically-induced isometric contractions. *MYO-MRI, Imaging in Neuromuscular Disease*, Nov 2019, Montréal, Canada. 2019. hal-02438631

HAL Id: hal-02438631

<https://hal.science/hal-02438631>

Submitted on 24 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

AREAS OF MUSCLE TISSUE ALTERATION CAN DIFFER FROM ACTIVATED REGIONS DURING ELECTRICALLY-INDUCED ISOMETRIC CONTRACTIONS

Fouré A^{1,2,3}, Le Troter A^{1,2}, Ogier A⁴, Guye M^{1,2}, Gondin J^{1,5}, Bendahan D^{1,2}



¹Aix-Marseille Univ, CNRS, CRMBM, Marseille, France, ²APHM, Hôpitaux de la Timone, CEMEREM, Marseille, France, ³Université de Lyon, LIBM, Villeurbanne, France, ⁴Aix-Marseille Univ, Université de Toulon, CNRS, LIS, Marseille, France, ⁵Institut NeuroMyoGène, UCBL1, CNRS, INSERM, Lyon, France

Introduction

Neuromuscular electrical stimulation (NMES) has been shown to induce severe muscle damage under isometric conditions. Structural alterations [1] have been assessed in the muscles located beneath the stimulation electrodes. These findings have been supported by muscle transverse relaxation time (T_2) mapping using magnetic resonance imaging (MRI)[2].

Although muscle activation is considered to be mainly superficial using NMES [3], MRI investigations have demonstrated a relative spatial heterogeneity regarding T_2 changes [4]. Therefore, two main contributing causes could be linked to isometric NMES-induced muscle damage. A direct damage of weaker sarcomeres as suggested by the *popping sarcomere hypothesis* [5] in the activated muscle areas during the NMES exercise and/or an indirect damage of less activated muscle areas associated to intramuscular shear strain during the NMES exercise.

This study aimed at comparing the localization of activated and damaged muscle areas after a NMES session in order to distinguish these potential causes related to isometric NMES-induced muscle damage.

Materials & Methods

Population. Thirteen subjects (26 ± 3 y, 173 ± 8 cm, 70 ± 9 kg, 4 women) volunteered to participate in this study. They were fully informed about the nature and the aim of the study and gave their informed written consent to participate in this study conducted in conformity with the Declaration of Helsinki

Study design. All testing sessions were performed at the same time of day before (PRE) and immediately after the damaging exercise (POST) and then seven (D7), fourteen (D14) and twenty-one (D21) days after the first exploration as detailed in Figure 1.




Figure 1: Schematic representation of the experimental design




Figure 2: Subject position during the neuromuscular electrostimulation session

Exercise. During NMES exercise inducing muscle damage, subjects were positioned on *vastus lateralis* (VL) and *vastus medialis* (VM) as shown in Figure 2. NMES exercise consisted of 40 isometric contractions (5s on and 35s off) evoked with pulses delivered at 100Hz (pulse duration of 400 μ s). Intensity was increased until maximally tolerable level.

MRI. MR images were acquired at rest using a 1.5T scanner. T_1 -weighted images were assessed for anatomical muscle delineations of the thigh. T_2 maps were determined with a segmented spin-echo echo-planar imaging sequence with the following parameters: 20 slices, slice thickness = 6mm, slice spacing = 6mm, TR = 4800ms, TE_s = 15/25/35/45/55ms.

T_2 quantification in deep and superficial parts of the VM and VL muscles. Two depth levels (*i.e.*, superficial and deep) were considered for VM and VL the most damaged muscles located beneath the stimulation electrodes. As previously described [6], a polar coordinate system (with the bone center used as the reference) was used to determine the thickness of damaged muscles at different angles and the frontiers between superficial and deep parts was then characterized in each slice [6]. Regarding muscle length, three levels were considered (*i.e.*, distal from slice #1 to #7, central from slice 8 to #13 and proximal from slice #14 to #20). T_2 was then quantified for each region and at each time point

Statistics. Normality of the data distribution was initially investigated using Shapiro-Wilk test. One-way ANOVA (time) were used (Statistica, Statsoft, Tulsa, USA) to assess immediate (*i.e.*, PRE vs. POST) and remaining (*i.e.*, PRE vs. D7 to D21) changes in volume and T_2 of each muscle. In addition, two-way ANOVAs (muscle depth * time) were used to assess T_2 differences between superficial and deep parts over time for both VL and VM and so for each muscle length level (*i.e.*, proximal, mid and distal). A Tukey's HSD post hoc analysis was performed when appropriate.

Conclusion

On the basis of immediate and long lasting T_2 changes recorded after an isometric NMES session, the present results clearly demonstrated that muscle areas activated during the contractions can be spatially different than the damaged muscle regions.

The discrepancies between activation areas and alteration regions in the *vastus lateralis* implies that tissues other than active skeletal muscle fibers were altered during unaccustomed NMES-induced isometric contractions.

Results

Significant decrease in MVC was found at POST ($-26 \pm 9\%$) and in delayed time at D7 ($-20 \pm 6\%$) and D14 ($-12 \pm 5\%$).

Although activation of muscle was statistically detected through T_2 increase at POST in superficial parts of the two muscles located beneath the stimulation electrodes (*i.e.*, *vastus lateralis* and *vastus medialis*), alterations quantified in a delayed time from increased T_2 were mainly located in the deep muscle region of the *vastus lateralis* ($+57 \pm 24\%$) and superficial area of the *vastus medialis* ($+24 \pm 16\%$) at D7 and were still observed in whole muscle at D21 (Table 1).

Table 1: Muscle T_2 values in deep and superficial parts along the *vastus lateralis* and *vastus medialis* before (PRE), immediately after (POST) and seven days (D7), fourteen days (D14) and twenty-one days (D21) after NMES-induced muscle damage.

Mean \pm SD (ms)	VL		VM		
	Sup	Deep	Sup	Deep	
PRE	Proximal	32.5 \pm 1.2	32.5 \pm 1.2	33.0 \pm 1.0	32.4 \pm 0.8
	Central	31.2 \pm 0.8	31.9 \pm 0.8	32.4 \pm 0.6	32.3 \pm 0.8
	Distal	32.4 \pm 1.0	33.1 \pm 1.0	33.6 \pm 0.5	33.5 \pm 0.8
	Whole muscle	32.2 \pm 1.0	32.5 \pm 1.0	33.3 \pm 0.6	32.9 \pm 0.7
POST	Proximal	38.2 \pm 1.7 ^{a,#}	36.1 \pm 1.4 ^a	34.5 \pm 1.5 ^a	33.3 \pm 1.0 ^a
	Central	37.0 \pm 2.0 ^a	36.7 \pm 1.7 ^a	35.6 \pm 2.0 ^{a,#}	33.8 \pm 1.6 ^a
	Distal	37.6 \pm 2.1 ^a	38.4 \pm 2.2 ^a	38.4 \pm 2.1 ^{a,#}	36.1 \pm 1.7 ^a
	Whole muscle	37.8 \pm 1.7 ^a	36.6 \pm 1.4 ^a	36.9 \pm 1.9 ^{a,#}	34.9 \pm 1.4 ^a
D7	Proximal	45.3 \pm 8.0 ^a	51.2 \pm 8.9 ^a	37.1 \pm 3.6 ^{a,#}	33.5 \pm 1.1
	Central	44.3 \pm 7.1 ^{a,#}	52.8 \pm 6.8 ^a	40.9 \pm 5.7 ^{a,#}	34.7 \pm 2.4
	Distal	44.3 \pm 6.7 ^a	50.3 \pm 10.0 ^a	42.7 \pm 5.6 ^{a,#}	37.6 \pm 3.8 ^a
	Whole muscle	44.6 \pm 7.0 ^{a,#}	51.1 \pm 7.8 ^a	41.1 \pm 5.0 ^{a,#}	35.8 \pm 2.6 ^a
D14	Proximal	37.7 \pm 3.0 ^{a,b}	40.9 \pm 3.7 ^{a,b}	35.7 \pm 2.2 ^a	33.5 \pm 1.4
	Central	36.9 \pm 3.1 ^{a,b}	41.3 \pm 3.4 ^{a,b}	36.9 \pm 2.5 ^{a,b}	33.8 \pm 1.5
	Distal	37.1 \pm 3.3 ^{a,b}	39.4 \pm 4.4 ^{a,b}	38.6 \pm 2.0 ^{a,b}	36.0 \pm 2.2
	Whole muscle	37.4 \pm 2.8 ^{a,b}	40.7 \pm 3.4 ^{a,b}	37.6 \pm 1.9 ^{a,b}	34.8 \pm 1.6
D21	Proximal	36.4 \pm 2.5 ^{a,b}	38.4 \pm 3.0 ^{a,b}	34.4 \pm 1.3 ^b	32.9 \pm 1.0
	Central	35.3 \pm 2.4 ^b	38.6 \pm 2.8 ^{a,b}	35.5 \pm 2.4 ^{a,b}	33.1 \pm 1.1
	Distal	35.6 \pm 1.8 ^{a,b}	37.9 \pm 3.1 ^{a,b}	37.1 \pm 1.9 ^{a,b}	35.3 \pm 1.7
	Whole muscle	36.0 \pm 2.3 ^b	38.4 \pm 2.8 ^{a,b}	36.2 \pm 1.8 ^{a,b}	34.1 \pm 1.2

VL: *vastus lateralis*; VM: *vastus medialis*. Sup: superficial part, Deep: deep part

a: significantly different from PRE ($P < 0.05$), b: significantly different from D7 ($P < 0.05$), #: significantly different from deep part ($P < 0.05$).

Bibliography

- [1] Mackey AL *et al.* (2008). Evidence of skeletal muscle damage following electrically stimulated isometric muscle contractions in humans. *J Appl Physiol* 105, 1620-1627.
- [2] Fouré A *et al.* (2015a). Heterogeneity of muscle damage induced by electrostimulation: a multimodal MRI study. *Med Sci Sports Exerc* 47, 166-175.
- [3] Maffiuletti NA. (2010). Physiological and methodological considerations for the use of neuromuscular electrical stimulation. *Eur J Appl Physiol* 110, 223-234.
- [4] Adams GR, Harris RT, Woodard D & Dudley GA. (1993). Mapping of electrical muscle stimulation using MRI. *J Appl Physiol* 74, 532-537.
- [5] Morgan DL & Proske U. Popping sarcomere hypothesis explains stretch-induced muscle damage. *Clinical and experimental pharmacology & physiology*. 2004; 31(8):541-545.
- [6] Fouré A *et al.* (2015). Localization and quantification of intramuscular damage using statistical parametric mapping and skeletal muscle parcellation. *Sci Rep* 5, 18580.