



**HAL**  
open science

## Global marine biodiversity in the context of achieving the Aichi Targets: ways forward and addressing data gaps

Hanieh Saeedi, James Davis Reimer, Miriam Brandt, Philippe-Olivier Dumais, Anna Maria Jazdzewska, Nicholas W. Jeffery, Peter M. Thielen, Mark John Costello

### ► To cite this version:

Hanieh Saeedi, James Davis Reimer, Miriam Brandt, Philippe-Olivier Dumais, Anna Maria Jazdzewska, et al. Global marine biodiversity in the context of achieving the Aichi Targets: ways forward and addressing data gaps. PeerJ, 2019, 7, pp.e7221. 10.7717/peerj.7221 . hal-02439858

**HAL Id: hal-02439858**

**<https://hal.science/hal-02439858>**

Submitted on 4 Jun 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# Global marine biodiversity in the context of achieving the Aichi Targets: ways forward and addressing data gaps

Hanieh Saeedi<sup>1,2,3</sup>, James Davis Reimer<sup>4</sup>, Miriam I. Brandt<sup>5</sup>, Philippe-Olivier Dumais<sup>6</sup>, Anna Maria Jazdzewska<sup>7</sup>, Nicholas W. Jeffery<sup>8</sup>, Peter M. Thielen<sup>9</sup> and Mark John Costello<sup>10</sup>

<sup>1</sup> Senckenberg Research Institute and Natural History Museum, Frankfurt am Main, Germany

<sup>2</sup> FB 15 Biological Sciences Institute for Ecology, Diversity and Evolution Biologicum, Goethe University of Frankfurt, Frankfurt am Main, Germany

<sup>3</sup> Senckenberg Research Institute and Natural History Museum, OBIS Data Manager, Deep-sea Node, Frankfurt am Main, Germany

<sup>4</sup> Marine Invertebrate Systematics & Ecology Laboratory, Faculty of Science, University of the Ryukyus, Nishihara, Okinawa, Japan

<sup>5</sup> IMARBEC, fremer, IRD, CNRS, Univ. Montpellier, Sète, France

<sup>6</sup> Benthic Ecology Laboratory, Biology Department, Université Laval, Québec, Canada

<sup>7</sup> Laboratory of Polar Biology and Oceanobiology, Department of Invertebrate Zoology and Hydrobiology, Faculty of Biology and Environmental Protection, University of Lodz, Lodz, Poland

<sup>8</sup> Fisheries and Oceans Canada, Bedford Institute of Oceanography, Dartmouth, Nova Scotia, Canada

<sup>9</sup> Research and Exploratory Development Department, Johns Hopkins Applied Physics Laboratory, Laurel, MD, United States of America

<sup>10</sup> Institute of Marine Science, University of Auckland, Auckland, New Zealand

## ABSTRACT

In 2010, the Conference of the Parties of the Convention on Biological Diversity agreed on the Strategic Plan for Biodiversity 2011–2020 in Aichi Prefecture, Japan. As this plan approaches its end, we discussed whether marine biodiversity and prediction studies were nearing the Aichi Targets during the 4th World Conference on Marine Biodiversity held in Montreal, Canada in June 2018. This article summarises the outcome of a five-day group discussion on how global marine biodiversity studies should be focused further to better understand the patterns of biodiversity. We discussed and reviewed seven fundamental biodiversity priorities related to nine Aichi Targets focusing on global biodiversity discovery and predictions to improve and enhance biodiversity data standards (quantity and quality), tools and techniques, spatial and temporal scale framing, and stewardship and dissemination. We discuss how identifying biodiversity knowledge gaps and promoting efforts have and will reduce such gaps, including via the use of new databases, tools and technology, and how these resources could be improved in the future. The group recognised significant progress toward Target 19 in relation to scientific knowledge, but negligible progress with regard to Targets 6 to 13 which aimed to safeguard and reduce human impacts on biodiversity.

Submitted 5 February 2019

Accepted 31 May 2019

Published 29 October 2019

Corresponding author

Hanieh Saeedi,  
hanieh.saeedi@senckenberg.de,  
hanieh.saeedi@gmail.com

Academic editor

Pablo Marquet

Additional Information and  
Declarations can be found on  
page 11

DOI 10.7717/peerj.7221

© Copyright  
2019 Saeedi et al.

Distributed under  
Creative Commons CC-BY 4.0

OPEN ACCESS

**Subjects** Biodiversity, Marine Biology, Molecular Biology, Taxonomy, Data Mining and Machine Learning

**Keywords** Aichi targets, Marine biodiversity, Prediction, Discovery, Biodiversity tools and pipelines, Biogeography, Data standard, Stewardship and dissemination, Stewardship, Data standards, Dissemination, Tools and pipelines, Marine biodiversity

## INTRODUCTION

The ‘Strategic Plan for Biodiversity 2011–2020’ of the Convention on Biological Diversity was agreed during the 10th Conference of the Parties, held from 18 to 29 October 2010, in Aichi Prefecture, Japan. The strategic plan included five main “Strategic Goals” that were divided into 20 targets. Each ‘Aichi Target’ was designed to better understand and predict biodiversity dynamics, such as how biological diversity underpins ecosystem function, and how the provision of ecosystem services is essential for human well-being. Meeting the Aichi Targets would ultimately benefit local livelihoods and economic development, and is essential for biodiversity maintenance and poverty reduction (*Shepherd et al., 2016; Tittensor et al., 2010*). In this paper, we report on the findings of a working group that discussed how the targets related to marine biodiversity were being achieved.

## SURVEY METHODOLOGY

The 4th World Conference on Marine Biodiversity (WCMB 2018) in Montreal organised a working group to review and evaluate how the Aichi Targets have been met by the global community. In particular, the review group was asked to focus on Target 19 regarding scientific knowledge about biodiversity as this was the area of most expertise of the participants. To identify and reduce biodiversity knowledge gaps we examined how marine biodiversity discoveries and their predictions need to be redirected to better understand and predict how marine biodiversity will change within the next 10 years. Here, we identify seven important priorities for this topic to support the Aichi Targets (*Table 1*). These foci arose iteratively from discussion between the group members and other groups at the conference. The priorities address issues of (1) data standards, (2) education in data management, (3) taxonomic expertise, (4) genetic tools, (5) international collaboration, (6) identifying knowledge gaps and understanding biogeography, and (7) the need to reduce human pressures on marine biodiversity.

## RESULTS AND DISCUSSION

### Reviewed priorities

#### ***Developing, improving and enhancing biodiversity data standards, exchange, and analytical tools, via standardized techniques***

Recent marine biodiversity discoveries have been greatly enhanced by standardised open-access taxonomic and biogeographic data repositories such as the World Register of Marine Species (WoRMS) and the <http://www.iobis.org/> (OBIS) (*Ahyong et al., 2018; Costello et al., 2013a*). Large-scale development of marine biodiversity data standards and exchange started when the Census of Marine Life (2000–2010) established OBIS (*O’Dor et al., 2012*).

Biodiversity data standards, such as “Darwin Core”, a data schema that provides stable terms and vocabularies for universal sharing of biodiversity data, and management techniques have been improved to ensure that published data have high quality. For example, there are now available taxonomic and geographic data management tools and R packages, e.g., rOBIS (*Provoost, 2018*) (*Fig. 1*). The taxon match tool in WoRMS can automatically check the correct spelling, authority, classification and validity of species

**Table 1 The examined Aichi Targets.** The Aichi Targets examined in this study and their relationship to the scientific priorities identified by participants of the 2018 World Conference on Marine Biodiversity. The Targets were grouped into Strategic Goals by the Convention on Biological Diversity. \* negligible progress, \*\* notable progress, \*\*\* good progress.

Aichi target	Scientific priority	Progress
<p><i>Strategic Goal E: Enhance implementation through participatory planning, knowledge management and capacity building</i></p> <p><b>Target 19</b> By 2020, knowledge, the science base and technologies relating to biodiversity, its values, functioning, status and trends, and the consequences of its loss, are improved, widely shared and transferred, and applied.</p> <p><i>Strategic Goals A: Address the underlying causes of biodiversity loss by mainstreaming biodiversity across government and society [1] B: Reduce the direct pressures on biodiversity and promote sustainable use, Targets 6 to 10 [2] and C: To improve the status of biodiversity by safeguarding ecosystems, species and genetic diversity, Targets 11–13 [3]</i></p>	Developing, improving and enhancing biodiversity data standards, exchange, and analytical tools, via standardized techniques.	***
	Educational activities to increase data mobilisation by taxonomists, data users, and/or wider audiences.	***
	Promoting synergy of biodiversity research efforts via increased collaboration at all levels.	***
	Utilization and promotion of taxonomic expertise and species identification tools to better recognize and catalogue biodiversity.	**
	Improvement and standardization of genetic, genomic, and other “omics” tools to aid in discovery, assessment, description, and cataloging of biodiversity.	**
	Identifying biodiversity and biogeographic knowledge gaps and promoting efforts to reduce such gaps.	***
	Control of anthropogenic pressures on vulnerable ecosystems impacted by climate change or ocean acidification to maintain their integrity and functioning.	*


#### Notes.

[1] **Target 3:** By 2020, at the latest, incentives, including subsidies, harmful to biodiversity are eliminated, phased out or reformed in order to minimize or avoid negative impacts, and positive incentives for the conservation and sustainable use of biodiversity are developed and applied, consistent and in harmony with the Convention and other relevant international obligations, taking into account national socio economic conditions. **Target 4:** By 2020, at the latest, Governments, business and stakeholders at all levels have taken steps to achieve or have implemented plans for sustainable production and consumption and have kept the impacts of use of natural resources well within safe ecological limits.

[2] **Target 6:** By 2020 all fish and invertebrate stocks and aquatic plants are managed and harvested sustainably, legally and applying ecosystem-based approaches, so that over-fishing is avoided, recovery plans and measures are in place for all depleted species, fisheries have no significant adverse impacts on threatened species and vulnerable ecosystems and the impacts of fisheries on stocks, species and ecosystems are within safe ecological limits. **Target 7:** By 2020 areas under agriculture, aquaculture and forestry are managed sustainably, ensuring conservation of biodiversity. **Target 8:** By 2020, pollution, including from excess nutrients, has been brought to levels that are not detrimental to ecosystem function and biodiversity. **Target 9:** By 2020, invasive alien species and pathways are identified and prioritized, priority species are controlled or eradicated, and measures are in place to manage pathways to prevent their introduction and establishment. **Target 10:** By 2015, the multiple anthropogenic pressures on coral reefs, and other vulnerable ecosystems impacted by climate change or ocean acidification are minimized, so as to maintain their integrity and functioning.

[3] **Target 11:** By 2020, at least 17 per cent of terrestrial and inland water, and 10 per cent of coastal and marine areas, especially areas of particular importance for biodiversity and ecosystem services, are conserved through effectively and equitably managed, ecologically representative and well-connected systems of protected areas and other effective area-based conservation measures and integrated into the wider landscapes and seascapes. **Target 12:** By 2020 the extinction of known threatened species has been prevented and their conservation status, particularly of those most in decline, has been improved and sustained. **Target 13:** By 2020, the genetic diversity of cultivated plants and farmed and domesticated animals and of wild relatives, including other socio-economically as well as culturally valuable species, is maintained, and strategies have been developed and implemented for minimizing genetic erosion and safeguarding their genetic diversity.

names uploaded to a webpage (Costello et al., 2013a). A marine gazetteer matches place names to geographic coordinates and polygon (shapefiles) of localities (Claus et al., 2014). The standardization and open storage of metadata, taxonomic, genetic, and geographic data also allows for greater stewardship by stakeholders, enhanced public awareness and education, and importantly, the ability to easily share data among institutions (Fig. 1).



**Figure 1 Biodiversity data processing.** Biodiversity data processing using novel analytical standardized techniques and technologies.

Full-size DOI: [10.7717/peerj.7221/fig-1](https://doi.org/10.7717/peerj.7221/fig-1)

Developing biodiversity data standards and data exchange protocols enables both data users and providers to benefit from the high-quality data that later allow for more reliable and precise biodiversity analyses. The expansion of the OBIS data schema to include additional information associated with sampling events, including sampling methods and environmental data, is a significant recent advance (De Pooter *et al.*, 2017). The open-access publication of thousands of data sets integrated into OBIS has enabled major advances in our understanding of global patterns of biodiversity. For example, several studies have utilized open-access marine species distribution records to discover and confirm large-scale biodiversity patterns. These findings include observations that global species richness is bimodal with latitude, and that species richness decreases with depth (Chaudhary, Saeedi & Costello, 2016; Chaudhary, Saeedi & Costello, 2017; Costello & Chaudhary, 2017; Saeedi, Basher & Costello, 2016; Saeedi & Costello, 2019a; Saeedi & Costello, 2019b; Saeedi, Dennis & Costello, 2017; Saeedi *et al.*, *in press*; Hillebrand, 2004). We conclude that there has been noticeable progress towards achieving the potential of Aichi Target 19 through establishing global databases and taxonomic resources. These are leading to better quality control, data management efficiencies, and new insights into our understanding of biodiversity.

However, these benefits, as well as contributions to these resources, are not realised in all countries.

### ***Educational activities to increase data mobilisation by taxonomists, data users, and/or wider audiences***

Many scientists and the general public around the world are unaware of the presence and advantages of large-scale open-access databases. Even if scientists are familiar with these facilities, data preparation and submission can be complex for contributors unfamiliar with data publication protocols. These issues are more pronounced for scientists in developing countries or non-native English speakers. As a result of these perceived and real data publication obstacles, significant biodiversity and biogeography knowledge remains in personal databases and non-digital archives. These logistical hurdles and data ownership perceptions frequently stand in the way of data publication. To expose researchers to these resources, initiatives like OceanTeacher Global Academy (OTGA) (<https://classroom.oceanteacher.org>) or Ocean School (<https://oceanschool.nfb.ca/>) provide a valuable educational platform that aids sustainable development. OceanTeacher is part of the <http://www.iode.org/> Programme of the <http://ioc.unesco.org/>. It has trained nearly 2,000 students from 120 countries since 2005 (<https://classroom.oceanteacher.org/mod/page/view.php?id=2033>). OTGA has hosted OBIS training workshops to train data providers on how to prepare, standardize, and submit their data to OBIS, where their data is cited and safely secured. Organisations such as OTGA need to be financially supported by governments in order to actively educate and train data keepers and encourage them to share their data with the global community. However, as the scientific community is relatively small and financial sources for training are limited, the future of large-scale biodiversity studies is likely to rely on the well-designed application of citizen science in addition to technological advances (*Stuart-Smith et al., 2017; Thiel et al., 2014*).

### ***Promoting synergy of biodiversity research efforts via increased collaboration at all levels***

In order to predict and discover biodiversity on a global scale, collaborative approaches among institutions and nations are necessary. *Guralnick, Hill & Lane (2007)* proposed a framework to use online databases and tools to improve and standardize geographic data, and to validate and highlight taxonomic data and misidentifications. They also suggested that a global infrastructure for web-based tools would enhance the quality of visualizing and standardizing raw biodiversity data and lead to a higher degree of collaboration and accessibility of knowledge (*Guralnick, Hill & Lane, 2007*).

The decade long Census of Marine Life was the largest global collaboration amongst marine biologists (*O'Dor et al., 2012*). Its legacy continues in OBIS with regard to data publication, but also continued international collaboration amongst polar, deep-sea and other researchers. The International Association for Biological Oceanography (IABO) is the organisation officially responsible for coordinating the marine biodiversity community (*Costello et al., 2015a*). It runs the MARINE-B email list with over 1,000 subscribers, and holds a World Conference on Marine Biodiversity every three years. Many others, often more specialist-focused, conferences also serve to bring marine biodiversity researchers

together. These serve to make introductions and help build collaborative relationships among researchers. However, most research funding is for topics of national rather than international importance. For example, not every country will have specialist expertise in every taxonomic group. Thus, sharing of taxonomic expertise can alleviate funding deficits, allow the transfer of knowledge, and lead to international partnerships.

The number of marine species formally described each year has never been greater, and aside from naming these species, more work is required to understand their life histories and ecology, biogeography, and evolution ([Appeltans et al., 2012](#)). [Costello, Vanhoorne & Appeltans \(2015\)](#) recommended the use of collaborative online databases, taxonomic effort improved and increased through communication, easier access to specimens, engagement of non-specialists, and international collaboration ([Costello, Vanhoorne & Appeltans, 2015](#)). Further, [Costello et al. \(2013b\)](#) advocated abandoning “data-sharing” and instead suggested requiring data publication within a journal or to online infrastructures such as OBIS, WoRMS, and/or the Global Biodiversity Information Facility (GBIF) ([Costello et al., 2013b](#)). A fundamental aspect of Aichi Target 19, namely discovering the full extent of biodiversity in the world’s oceans, is not possible without international collaboration.

#### ***Utilization and promotion of taxonomic expertise and species identification tools to better recognize and catalogue biodiversity***

“Good” taxonomy is an absolute necessity for biodiversity recognition and management ([Thomson et al., 2018](#)). It is very important to pass on the knowledge of experienced taxonomists to others. In this regard, field-specific training workshops can be of great importance. As an example, two ‘IceAGE (Icelandic marine Animals: Genetics and Ecology) amphipod identification workshops’ were recently held, consisting of two weeks of work of a group of taxonomists accompanied by students. This resulted in the identification of more than 20,000 amphipod specimens, and the publication of seven research papers dealing with the taxonomy, diversity and ecology of this group around Iceland ([Brix et al., 2018](#)).

Another problem in recognizing biodiversity is the complexity of access to information and images that can help end-users in identifying the organisms collected in their samples. To solve this requires comprehensive online identification guides to all species on Earth ([Costello, May & Stork, 2013b](#)). It is surprising that despite the numerous online digital initiatives regarding biodiversity resources for this most practical need (how to identify species) lack leadership and remain scarce. Targeted funding to support such resources is urgently needed to help the wider community identify species quickly and accurately, including species that may be invasive or pathogenic. Despite great publicity and interest, publically available DNA libraries, such as the Barcode of Life ([Ratnasingham & Hebert, 2007](#)), are still far from having complete coverage of the Tree of Life. Moreover, DNA is often only useful if the species has already been formally described, specimens from which DNA has been sampled have been correctly identified, and the DNA sequence(s) published in an open access database. Other tools that can help in species identification for conservation and resource management include interactive keys. Unfortunately, there are

only such keys for a few marine taxa, and these mostly concern higher taxonomic levels (family-level or higher) (Dallwitz, 1997; Nimbs, 2017).

Along with Aichi Target 19, the exchange of knowledge between experienced scientists and young researchers as well as the use of different identification tools (e.g., interactive keys, barcode databases) will help in biodiversity recognition. These targets need to be prioritised by governments and funding agencies as they are fundamental to quality assurance in other areas of biology, ecology, and resource management.

### ***Improvement and standardization of genetic, genomic, and other “omics” tools to aid in discovery, assessment, description, and cataloging of biodiversity***

Countless studies have used genomic tools to study marine biodiversity, connectivity, and functional diversity (Carvalho *et al.*, 2010). While data sharing, standardized sampling, metadata collection, and sequencing protocols still require significant standardization, the use of repositories such as GenBank, Dryad, and Sequence Read Archive have made published data much more accessible. We expect this trend to continue as new tools, such as the Genomic Observatories Metadata database (GeOMe), streamline sequencing data and metadata submission (Deck *et al.*, 2017).

The classification and phylogeny of eukaryotic organisms benefits from the use of genetic markers that help delineate species in combination with morphological, ecological, and/or physiological information (Hebert, Ratnasingham & De Waard, 2003; Palumbi, 2003). Currently, no uniform threshold value has been established for species delineation, and there is no single “universal” DNA barcode that captures all eukaryotic life. Even within a single animal order there can be large differences in this value between families (Tempestini, Rysgaard & Dufresne, 2018). Despite these issues, DNA barcoding and next-generation metabarcoding can reveal genetic diversity and are useful tools for the description and cataloguing of biodiversity.

Modern high-throughput sequencing (HTS) technologies have advanced DNA barcoding methods by producing millions of individual sequences per analyzed sample, enabling DNA metabarcoding from environmental DNA (eDNA) and complex community mixtures. Community and environmental metabarcoding are both useful tools to discover cryptic (undetected until genetic analyses) diversity in the marine realm, and assess ocean biodiversity in a non-invasive and high-throughput manner (Goodwin *et al.*, 2017).

In order to obtain robust and reproducible metabarcoding results, critical methodological aspects remain to be improved (Dickie *et al.*, 2018; Goldberg *et al.*, 2016; Nichols *et al.*, 2018). Studies are needed to address the effects of alternative protocols on sampling, molecular, and bioinformatic processing level in order to develop standardized and reliable techniques for applying these new methods. Furthermore, because large amounts of marine organisms have not been genetically characterized, integrative approaches should be supported in order to fill database gaps. The continued improvement and standardization of genetic, genomic, and other “-omics” tools (e.g., proteomics, transcriptomics) will continue to be valuable components in the discovery of new marine



prokaryotes and eukaryotes, as well as in monitoring biodiversity, thereby contributing to Aichi Target 19.

### ***Identifying biodiversity and biogeographic knowledge gaps and promoting efforts to reduce such gaps***

Deep-sea ecosystems include about 65% of the world's surface but are far less studied and sampled than shallower depths (Costello *et al.*, 2018; Costello, Cheung & Hauwera, 2010). Although deep-sea studies have increased rapidly in recent decades, there are large gaps in global sampling coverage, for example in the Indian and Pacific Oceans, and major efforts are needed to continue to be directed into offshore research (Saeedi *et al.*, 2019; Barroso *et al.*, 2018). The distribution and diversity of deep-sea fauna thus still remains poorly-known due to the size and remoteness of deep-sea ecosystems. For example, recent studies have shown that the global latitudinal marine species richness gradient follows a bi-modal pattern related to temperature and habitat availability (Chaudhary, Saeedi & Castello, 2016; Chaudhary, Saeedi & Costello, 2017; Saeedi, Basher & Costello, 2016; Saeedi & Costello, 2019a; Saeedi & Costello, 2019b; Saeedi, Dennis & Costello, 2017). This finding is supported by the fossil record, which shows reduced species richness at the equator in warm periods (Kiessling & Aberhan, 2007; Yasuhara *et al.*, 2012). As such, the peaks in this bimodal distribution may become further separated under future climate change and ocean warming. However, there is still no consensus about this bimodal pattern in the deep sea, where food supply may be more important than temperature in defining species distribution. For example, Woolley *et al.* (2016) examined 165,000 distribution records of Ophiuroidea and revealed that biogeographic patterns in species richness in the deep sea are associated with chemical energy and proximity to slope habitats. However, these patterns require investigation in other taxa, from micro- to mega-fauna, epifauna and infauna (Woolley *et al.*, 2016; Clarke & Gaston, 2006).

One issue in studying present and future global deep-sea biodiversity patterns are the few publicly available distribution records and environmental data. Deep-sea expeditions began over 100 years ago, but distribution data are often still retained in inaccessible archives, sometimes in local languages of that country, and are not publically available to the global community. Additionally, acquisition of data describing the distributions of deep-sea species is often limited by prohibitive costs and logistical difficulties in surveying the deep ocean. Environmental suitability modeling has thus become a cost-effective tool for identifying potential locations of deep-sea species, particularly for areas that have never been explored (Assis *et al.*, 2018; Basher & Costello, 2016; Danovaro *et al.*, 2017; Serrano *et al.*, 2017).

Understanding how abiotic drivers influence species distributions can contribute to filling spatial gaps of biodiversity hotspots and endangered areas (McHenry, Steneck & Brady, 2017; Saeedi, Basher & Costello, 2016; Saeedi & Costello, 2019a; Saeedi & Costello, 2019b; Saeedi, Dennis & Costello, 2017). Since some of these drivers can be observed by satellite imagery, it is possible to model some community assemblages in difficult-to-access locations. The development of models of species richness and cumulative anthropogenic impact distributions could be useful for conservation purposes and/or other spatial

planning applications (Selig *et al.*, 2014). Costello *et al.* (2018) proposed that sampling of the oceans should be stratified in relation to environmental variability, with more variable environments receiving more sampling focus in space and time (Costello *et al.*, 2018). Coupled with a recent objective (data driven) delineation of marine biodiversity into 30 biogeographic realms based on the endemisms of marine plants and animals (Costello *et al.*, 2018; Costello, Cheung & Hauwera, 2010; Costello, Vanhoorne & Appeltans, 2015), this provides a framework for more representative sampling of the oceans. These realms include 18 continental-shelf and 12 offshore realms, including unique seas, such as the Baltic and Black seas, and subdivisions of the Indian, Atlantic, and Pacific oceans, and polar waters. These are broad scale patterns, and application of stratified sampling at local levels requires finer spatial resolution data; such as benthic habitat maps and observations of movements of threatened species to know where they occur.

### ***Control of anthropogenic pressures on vulnerable ecosystems impacted by climate change or ocean acidification to maintain their integrity and functioning***

Fisheries and marine mammal hunting have had large impacts on biodiversity at local scales for centuries, and at global scales for the past two centuries (Pauly *et al.*, 2002). Clearly fishery management measures struggle to prevent overfishing, and trawling that destroys seabed habitats is widely permitted, while bycatch of seabirds, turtles and marine mammals is pushing some species to extinction (McCauley *et al.*, 2015; Mittelbach *et al.*, 2007). Progress in reducing bycatch is compromising reaching the achievement of Aichi Target 12 related to preventing species extinctions. A proven solution to reversing some negative trajectories are marine reserves (no-take MPAs) (Costello, 2014). However, about two-thirds of coastal countries lack even one marine reserve, and over 90% of MPAs allow fishing and thus prevent the recovery of biodiversity to natural conditions (Costello & Ballantine, 2015). This failure to conserve and help fisheries recover, despite the potential benefits of MPAs to nature, education, science (they act as controls for effects of fishing outside them), tourism, and fish populations defies what is best for society. With less than 3% of the ocean in reserves, there seems little hope that Aichi Target 11's goal of 10% of the oceans being protected in MPAs by 2020 will be reached. In addition, there appears to be negligible progress towards more sustainable use of the oceans, as called for in Targets 4, 6 and 7. Target 3, the reduction of harmful subsidies, has also seen little progress and too many fisheries still receive indirect and/or direct subsidies from governments that enable further unsustainable overfishing.

Aichi Target 10 calls for reduced anthropogenic impacts on coral reefs. Coral reefs suffered global-scale bleaching events in 2015–2017, even within MPAs, resulting in massive damage to these ecosystems, including mass mortality of hermatypic corals and other zooxanthellate organisms (Hughes *et al.*, 2017), and associated reduced ecosystem functioning (Hughes, Kerry & Simpson, 2018; Hughes *et al.*, 2018). Additionally, such events have economic effects such as reduced tourism (Prideaux, Carmody & Pabel, 2017). Overall, the trajectory of coral reefs continues to be one of downward degradation in the face of

increasing anthropogenic pressures, including climate change and continued exploitation (Heery *et al.*, 2018).

Other anthropogenic impacts on marine biodiversity include excess nutrient input, oxygen depletion, and invasive species. Levels of these impacts are to be reduced and their management improved as part of Aichi Targets 8 and 9. Progress in management of introduced and invasive marine species has been made with the establishment of the World Register of Introduced Marine Species (WRiMS) (Ahyong *et al.*, 2018). Because of the nature of invasive species, management of their information is most cost-effectively done at a global rather than local scale. The next steps should include access to species identification resources and a dynamic online reporting and early warning system.

Both global warming and ocean acidification are closely linked with the anthropogenic input of CO<sub>2</sub> and other greenhouse gases into the atmosphere, and without controlling these issues, the future of coral reefs looks bleak IPCC2018. Minimizing anthropogenic impacts such as increased runoff from coastal development and reducing overfishing can help delay the degradation of coral reef ecosystems, but it is estimated more than half of all coral reefs now experience medium to high anthropogenic pressures (Halpern *et al.*, 2008) and the extirpation of species from many coral reefs due to climate change is predicted (Molinos *et al.*, 2016).

There are some success stories, such as Palau, which has passed stringent legislation protecting coral reef diversity, including the world's first no-take zone for sharks (Vianna *et al.*, 2012), stringent legal protection (Gouezo *et al.*, 2017), and a visitor's pledge and public awareness campaign (<https://palaupledge.com/>). Other regions or countries following the lead of these exemplars could help buy time for coral reef ecosystems. For instance, the Australian government implemented the Great Barrier Reef (GBR) Zoning Plan 2003 in 2004, which set aside one-third of the GBR as a no-take zone (McCook *et al.*, 2010). This resulted in a significantly lower proportion of reefs being affected by Crown-of-thorns starfish outbreaks in no-take zones than in fished zones (McCook *et al.*, 2010), but the trajectory of GBR coral reef ecosystems remains bleak due to warming-associated coral bleaching (Hughes, Kerry & Simpson, 2018; Hughes *et al.*, 2018).

## CONCLUSIONS

While there has been considerable progress in addressing many of the priorities of the Aichi Targets, including the development and application of biodiversity tools and higher standards, as well as increased educational activity and increasing standardization of genetic and genomic tools, progress towards sustainable use is very limited (Table 1). Of the seven priorities for marine biodiversity to achieve the Aichi Targets we reviewed here, we judged six have seen some progress. However, other goals such as reducing anthropogenic stressors on vulnerable ecosystems have clearly not been met and seem certain to fall short of the 2020 Targets, as previously concluded (Tittensor *et al.*, 2014). We recommend continued efforts regarding international cooperation in marine biodiversity informatics to increase efficiencies of data management and accessibility, paralleled by field observations stratified by environmental conditions and societal needs, synergised

by online access to taxonomic information and species identification guides, and research into more cost-efficient methods for field sampling that minimise impacts on biodiversity (e.g., sensors, video, eDNA). This surveillance of marine biodiversity should occur in areas where human impacts, particularly fishing, occur, but also in fully-protected Marine Protected Areas where no fishing occurs (i.e., marine reserves). The lack of progress towards establishing the latter reference areas compromises our understanding of natural conditions and how to judge the sustainability of human activities in the ocean and their impacts on marine biodiversity.

## ACKNOWLEDGEMENTS

We would like to thank David Beauchesne, Rémi M. Daigle, Jéssica Goldsmit, Philippe Archambault, Anna Metaxas, and Paul Snelgrove for their thoughts and leadership of the mentorship program organizing committee and workshop facilitators. Special thanks also to Rémi M. Daigle and Jéssica Goldsmit for reviewing the current paper. Comments from Martin Thiel and Naomi Kingston greatly improved an earlier version of this paper.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This research is sponsored by the NSERC Canadian Healthy Oceans Network and its Partners: Department of Fisheries and Oceans Canada and INREST (representing the Port of Sept-Îles and City of Sept-Îles). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

NSERC Canadian Healthy Oceans Network and its Partners: Department of Fisheries and Oceans Canada.

INREST (representing the Port of Sept-Îles and City of Sept-Îles).

### Competing Interests

James D. Reimer and Mark J. Costello are Academic Editors for PeerJ.

### Author Contributions

- Hanieh Saeedi and Mark John Costello conceived and designed the experiments, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft, draft and revised some paragraphs of the current paper.
- James Davis Reimer conceived and designed the experiments, authored or reviewed drafts of the paper, approved the final draft, draft and revised some paragraphs of the current paper.
- Miriam I. Brandt, Philippe-Olivier Dumais, Anna Maria Jazdzewska, Nicholas W. Jeffery and Peter M. Thielen conceived and designed the experiments, draft and revised some paragraphs of the current paper.

## Data Availability

The following information was supplied regarding data availability:

We do not have any raw data as this is a review article.

## REFERENCES

- Ahyong SCMJ, Galil BS, Gollasch S, Hutchings P, Katsanevakis S, Lejeune C, Marchini A, Occhipinti A, Pagad S, Poore G, Rius M, Robinson TB, Sterrer W, Turon X, Willan RC, Zhan A. 2018. World Register of Introduced Marine Species (WRiMS). Available at <http://www.marinespecies.org/introduced> (accessed on 14 October 2018).
- Appeltans W, Ahyong ST, Anderson G, Angel MV, Artois T, Bailly N, Bamber R, Barber A, Bartsch I, Berta A. 2012. The magnitude of global marine species diversity. *Current Biology* 22:2189–2202 DOI 10.1016/j.cub.2012.09.036.
- Assis J, Tyberghein L, Bosch S, Verbruggen H, Serrão EA, De Clerck O. 2018. Bio-ORACLE v2.0: extending marine data layers for bioclimatic modelling. *Global Ecology and Biogeography* 27:277–284 DOI 10.1111/geb.12693.
- Barroso R, Kudenov JD, Halanych KM, Saeedi H, Sumida PYG, Bernardino AF. 2018. A new species of xylophylic fireworm (Annelida: Amphinomidae: Cryptonome) from deep-sea wood falls in the SW Atlantic. *Deep Sea Research Part I: Oceanographic Research Papers* 137:66–75 DOI 10.1016/j.dsr.2018.05.005.
- Basher Z, Costello MJ. 2016. The past, present and future distribution of a deep-sea shrimp in the Southern Ocean. *PeerJ* 4:e1713 DOI 10.7717/peerj.1713.
- Brix S, Lörz A-N, Jazdzewska AM, Hughes L, Tandberg AHS, Pabis K, Stransky B, Krapp-Schickel T, Sorbe JC, Hendrycks E, Vader W, Frutos I, Horton T, Jazdzewski K, Peart R, Beermann J, Coleman CO, Buhl-Mortensen L, Corbari L, Havermans C, Tato R, Jimenez Campean A. 2018. Amphipod family distributions around Iceland. *ZooKeys* 731:1–53.
- Carvalho G, Creer S, Allen MJ, Costa F, Tsigenopoulos C, Le Goff-Vitry M, Magoulas A, Medlin L, Metfies K. 2010. Genomics in the discovery and monitoring of marine biodiversity. In: Cock J, Tessmar-Raible K, Boyen CC, Viard F, eds. *Introduction to marine genomics. Advances in marine genomics, vol 1*. Dordrecht: Springer.
- Chaudhary C, Saeedi H, Costello MJ. 2016. Bimodality of latitudinal gradients in marine species richness. *Trends in Ecology & Evolution* 31:670–676 DOI 10.1016/j.tree.2016.06.001.
- Chaudhary C, Saeedi H, Costello MJ. 2017. Marine species richness is bimodal with latitude: a reply to Fernandez and Marques. *Trends in Ecology & Evolution* 32:234–237 DOI 10.1016/j.tree.2017.02.007.
- Clarke A, Gaston KJ. 2006. Climate, energy and diversity. *Proceedings Biological Sciences* 273:2257–2266 DOI 10.1098/rspb.2006.3545.
- Claus S, De Hauwere N, Vanhoorne B, Deckers P, Souza Dias F, Hernandez F, Mees J. 2014. Marine regions: towards a global standard for georeferenced marine names and boundaries. *Marine Geodesy* 37:99–125 DOI 10.1080/01490419.2014.902881.

- Costello MJ. 2014.** Long live marine reserves: a review of experiences and benefits. *Biological Conservation* 176:289–296 DOI 10.1016/j.biocon.2014.04.023.
- Costello MJ, Archambault P, Chavanich S, Miloslavich P, Paterson DM, Phang S-M, Pinto IS, Pierrot-Bults A, Song S, Soto EH. 2015a.** Organizing, supporting and linking the world marine biodiversity research community. *Journal of the Marine Biological Association of the United Kingdom* 95:431–433 DOI 10.1017/S0025315414001969.
- Costello MJ, Ballantine B. 2015.** Biodiversity conservation should focus on no-take marine reserves: 94% of marine protected areas allow fishing. *Trends in Ecology & Evolution* 30:507–509 DOI 10.1016/j.tree.2015.06.011.
- Costello MJ, Basher Z, Sayre R, Breyer S, Wright DJ. 2018.** Stratifying ocean sampling globally and with depth to account for environmental variability. *Scientific Reports* 8:11259 DOI 10.1038/s41598-018-29419-1.
- Costello MJ, Bouchet P, Boxshall G, Fauchald K, Gordon D, Hoeksema BW, Poore GC, Van Soest RW, Stöhr S, Walter TC. 2013a.** Global coordination and standardisation in marine biodiversity through the World Register of Marine Species (WoRMS) and related databases. *PLOS ONE* 8:e51629 DOI 10.1371/journal.pone.0051629.
- Costello MJ, Chaudhary C. 2017.** Marine biodiversity, biogeography, deep-sea gradients, and conservation. *Current Biology* 27:R511–R527 DOI 10.1016/j.cub.2017.04.060.
- Costello MJ, Cheung A, De Hauwere N. 2010.** Surface area and the seabed area, volume, depth, slope, and topographic variation for the world's seas, oceans, and countries. *Environmental Science & Technology* 44:8821–8828 DOI 10.1021/es1012752.
- Costello MJ, May RM, Stork NE. 2013.** Can we name Earth's species before they go extinct? *Science* 339:413–416 DOI 10.1126/science.1230318.
- Costello MJ, Michener WK, Gahegan M, Zhang Z-Q, Bourne PE. 2013b.** Biodiversity data should be published, cited, and peer reviewed. *Trends in Ecology & Evolution* 28:454–461 DOI 10.1016/j.tree.2013.05.002.
- Costello MJ, Vanhoorne B, Appeltans W. 2015.** Conservation of biodiversity through taxonomy, data publication, and collaborative infrastructures. *Conservation Biology* 29:1094–1099 DOI 10.1111/cobi.12496.
- Dallwitz M. 1997.** Descriptions, illustrations, interactive identification, and information retrieval from DELTA databases. DELTA databases. Available at <https://www.delta-intkey.com/>.
- Danovaro R, Corinaldesi C, Dell'Anno A, Snelgrove PVR. 2017.** The deep-sea under global change. *Current Biology* 27:R461–R465 DOI 10.1016/j.cub.2017.02.046.
- De Pooter D, Appeltans W, Bailly N, Bristol S, Deneudt K, Eliezer M, Fujioka E, Giorgetti A, Goldstein P, Lewis M, Lipizer M, Mackay K, Marin M, Moncoiffé G, Nikolopoulou S, Provoost P, Rauch S, Roubicek A, Torres C, Van de Putte A, Vandepitte L, Vanhoorne B, Vinci M, Wambiji N, Watts D, Klein Salas E, Hernandez F. 2017.** Toward a new data standard for combined marine biological and environmental datasets—expanding OBIS beyond species occurrences. *Biodiversity Data Journal* 5:e10989 DOI 10.3897/BDJ.5.e10989.
- Deck J, Gaither MR, Ewing R, Bird CE, Davies N, Meyer C, Riginos C, Toonen RJ, Crandall ED. 2017.** The Genomic Observatories Metadatabase (GeOMe): a new

- repository for field and sampling event metadata associated with genetic samples. *PLOS Biology* 15:e2002925 DOI 10.1371/journal.pbio.2002925.
- Dickie IA, Boyer S, Buckley H, Duncan RP, Gardner P, Hogg ID, Holdaway RJ, Lear G, Makiola A, Morales SE. 2018. Towards robust and repeatable sampling methods in eDNA based studies. *Molecular Ecology Resources* 18:940–952.
- Goldberg CS, Turner CR, Deiner K, Klymus KE, Thomsen PF, Murphy MA, Spear SF, McKee A, Oyler-McCance SJ, Cornman RS. 2016. Critical considerations for the application of environmental DNA methods to detect aquatic species. *Methods in Ecology and Evolution* 7:1299–1307 DOI 10.1111/2041-210X.12595.
- Goodwin KD, Thompson LR, Duarte B, Kahlke T, Thompson AR, Marques JC, Caçador I. 2017. DNA sequencing as a tool to monitor marine ecological status. *Frontiers in Marine Science* 4:Article 107 DOI 10.3389/fmars.2017.00107.
- Gouezo M, Nestor V, Olsudong D, Marino L, Mereb G, Jonathan R, Golbuu Y. 2017. 15 years of coral reef monitoring demonstrates the resilience of Palau’s coral reefs. PICRC Technical Report 17-01. Available at [http://picrc.org/picrcpage/wp-content/uploads/2016/01/Gouezo-et-al-2017-coral-reef-status-and-trends\\_15-years\\_FINAL\\_August18th.pdf](http://picrc.org/picrcpage/wp-content/uploads/2016/01/Gouezo-et-al-2017-coral-reef-status-and-trends_15-years_FINAL_August18th.pdf).
- Guralnick RP, Hill AW, Lane M. 2007. Towards a collaborative, global infrastructure for biodiversity assessment. *Ecology Letters* 10:663–672 DOI 10.1111/j.1461-0248.2007.01063.x.
- Halpern BS, Walbridge S, Selkoe KA, Kappel CV, Micheli F, D’agrosa C, Bruno JF, Casey KS, Ebert C, Fox HE. 2008. A global map of human impact on marine ecosystems. *Science* 319(5865):948–952 DOI 10.1126/science.1149345.
- Hebert PD, Ratnasingham S, De Waard JR. 2003. Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. *Proceedings of the Royal Society B: Biological Sciences* 270:S96–S99.
- Heery EC, Hoeksema BW, Browne NK, Reimer JD, Ang PO, Huang D, Friess DA, Chou LM, Loke LH, Saksena-Taylor P. 2018. Urban coral reefs: degradation and resilience of hard coral assemblages in coastal cities of East and Southeast Asia. *Marine Pollution Bulletin* 135:654–681 DOI 10.1016/j.marpolbul.2018.07.041.
- Hillebrand H. 2004. On the generality of the latitudinal diversity gradient. *The American Naturalist* 163:192–211 DOI 10.1086/381004.
- Hughes TP, Kerry JT, Baird AH, Connolly SR, Dietzel A, Eakin CM, Heron SF, Hoey AS, Hoogenboom MO, Liu G. 2018. Global warming transforms coral reef assemblages. *Nature* 556:492–496 DOI 10.1038/s41586-018-0041-2.
- Hughes TP, Kerry JT, Álvarez Noriega M, Álvarez Romero JG, Anderson KD, Baird AH, Babcock RC, Beger M, Bellwood DR, Berkelmans R. 2017. Global warming and recurrent mass bleaching of corals. *Nature* 543(7645):373–377 DOI 10.1038/nature21707.
- Hughes T, Kerry J, Simpson T. 2018. Large-scale bleaching of corals on the Great Barrier Reef. *Ecology* 99:501–501 DOI 10.1002/ecy.2092.

- Kiessling W, Aberhan M. 2007. Geographical distribution and extinction risk: lessons from Triassic-Jurassic marine benthic organisms. *Journal of Biogeography* 34:1473–1489 DOI [10.1111/j.1365-2699.2007.01709.x](https://doi.org/10.1111/j.1365-2699.2007.01709.x).
- McCauley DJ, Pinsky ML, Palumbi SR, Estes JA, Joyce FH, Warner RR. 2015. Marine defaunation: animal loss in the global ocean. *Science* 347(6219):1255641 DOI [10.1126/science.1255641](https://doi.org/10.1126/science.1255641).
- McCook LJ, Ayling T, Cappo M, Choat JH, Evans RD, De Freitas DM, Heupel M, Hughes TP, Jones GP, Mapstone B. 2010. Adaptive management of the Great Barrier Reef: a globally significant demonstration of the benefits of networks of marine reserves. *Proceedings of the National Academy of Sciences of the United States of America* 107:18278–18285 DOI [10.1073/pnas.0909335107](https://doi.org/10.1073/pnas.0909335107).
- McHenry J, Steneck RS, Brady DC. 2017. Abiotic proxies for predictive mapping of nearshore benthic assemblages: implications for marine spatial planning. *Ecological Applications* 27:603–618 DOI [10.1002/eap.1469](https://doi.org/10.1002/eap.1469).
- Mittelbach GG, Schemske DW, Cornell HV, Allen AP, Brown JM, Bush MB, Harrison SP, Hurlbert AH, Knowlton N, Lessios HA, McCain CM, McCune AR, McDade LA, McPeck MA, Near TJ, Price TD, Ricklefs RE, Roy K, Sax DF, Schluter D, Sobel JM, Turelli M. 2007. Evolution and the latitudinal diversity gradient: speciation, extinction and biogeography. *Ecology Letters* 10:315–331 DOI [10.1111/j.1461-0248.2007.01020.x](https://doi.org/10.1111/j.1461-0248.2007.01020.x).
- Molinos JG, Halpern BS, Schoeman DS, Brown CJ, Kiessling W, Moore PJ, Pandolfi JM, Poloczanska ES, Richardson AJ, Burrows MT. 2016. Climate velocity and the future global redistribution of marine biodiversity. *Nature Climate Change* 6:83–88 DOI [10.1038/nclimate2769](https://doi.org/10.1038/nclimate2769).
- Nichols RV, Vollmers C, Newsom LA, Wang Y, Heintzman PD, Leighton M, Green RE, Shapiro B. 2018. Minimizing polymerase biases in metabarcoding. *Molecular Ecology Resources* 18(5):927–939 DOI [10.1111/1755-0998.12895](https://doi.org/10.1111/1755-0998.12895).
- Nimbs MJ. 2017. NudiKey: an illustrated, interactive identification key to the families of Australian heterobranch sea-slugs (Mollusca: Gastropoda). *Australian Zoologist* 38:537–546 DOI [10.7882/AZ.2017.004](https://doi.org/10.7882/AZ.2017.004).
- O’Dor R, Boustany AM, Chittenden CM, Costello MJ, Moustahfid H, Payne J, Steinke D, Stokesbury MJ, Vanden Berghe E. 2012. A census of fishes and everything they eat: how the census of marine life advanced fisheries science. *Fisheries* 37:398–409 DOI [10.1080/03632415.2012.714323](https://doi.org/10.1080/03632415.2012.714323).
- Palumbi SR. 2003. Population genetics, demographic connectivity, and the design of marine reserves. *Ecological Applications* 13:146–158 DOI [10.1890/1051-0761\(2003\)013\[0146:PGDCAT\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2003)013[0146:PGDCAT]2.0.CO;2).
- Pauly D, Christensen V, Guénette S, Pitcher T, Sumaila U, Walters C, Watson R, Zeller D. 2002. Towards sustainability in world fisheries. *Nature* 418(6898):689–695 DOI [10.1038/nature01017](https://doi.org/10.1038/nature01017).
- Prideaux B, Carmody J, Pabel A. 2017. Impacts of the 2016 and 2017 mass coral bleaching events on the Great Barrier Reef tourism industry and tourism-dependent



coastal communities of Queensland. Report to the reef and rainforest research centre limited cairns. Reef and Rainforest Research Centre Limited.

- Provoost PBS. 2018.** robis: R Client to access data from the OBIS API. Ocean Biogeographic Information System. R package version 1.0.1. Intergovernmental Oceanographic Commission of UNESCO. Available at <https://cran.r-project.org/package=robis>.
- Ratnasingham S, Hebert PD. 2007.** BOLD: the barcode of life data system (<http://www.barcodinglife.org>). *Molecular Ecology Notes* 7:355–364 DOI 10.1111/j.1471-8286.2007.01678.x.
- Saeedi H, Basher Z, Costello MJ. 2016.** Modelling present and future global distributions of razor clams (Bivalvia: Solenidae). *Helgoland Marine Research* 70:Article 23 DOI 10.1186/s10152-016-0477-4.
- Saeedi H, Bernardino AF, Shimabukuro M, Falchetto G, Sumida PYG. 2019.** Macrofaunal community structure and biodiversity patterns based on a wood-fall experiment in the deep South-west Atlantic. *Deep Sea Research Part I: Oceanographic Research Papers* 145:73–82 DOI 10.1016/j.dsr.2019.01.008.
- Saeedi H, Costello MJ. 2019a.** The biology, ecology, and societal importance of razor clams. In: *Reference module in earth systems and environmental sciences*. Elsevier.
- Saeedi H, Costello MJ. 2019b.** A world dataset on the geographic distributions of Solenidae razor clams (Mollusca: Bivalvia). *Biodiversity Data Journal* 7:e31375 DOI 10.3897/BDJ.7.e31375.
- Saeedi H, Dennis TE, Costello MJ. 2017.** Bimodal latitudinal species richness and high endemism of razor clams (Mollusca). *Journal of Biogeography* 44:592–604 DOI 10.1111/jbi.12903.
- Saeedi H, Costello MJ, Warren D, Brandt A.** Latitudinal and bathymetrical species richness patterns in the NW Pacific and adjacent Arctic Ocean. *Scientific Reports*. In Press.
- Selig ER, Turner WR, Troeng S, Wallace BP, Halpern BS, Kaschner K, Lascelles BG, Carpenter KE, Mittermeier RA. 2014.** Global priorities for marine biodiversity conservation. *PLOS ONE* 9(1):e82898 DOI 10.1371/journal.pone.0082898.
- Serrano A, González-Irusta JM, Punzón A, García-Alegre A, Lourido A, Ríos P, Blanco M, Gómez-Ballesteros M, Druet M, Cristobo J, Cartes JE. 2017.** Deep-sea benthic habitats modeling and mapping in a NE Atlantic seamount (Galicia Bank). *Deep Sea Research Part I: Oceanographic Research Papers* 126:115–127 DOI 10.1016/j.dsr.2017.06.003.
- Shepherd E, Milner-Gulland EJ, Knight AT, Ling MA, Darrah S, Van Soesbergen A, Burgess ND. 2016.** Status and trends in global ecosystem services and natural capital: assessing progress toward aichi biodiversity target 14. *Conservation Letters* 9:429–437 DOI 10.1111/conl.12320.
- Stuart-Smith RD, Edgar GJ, Barrett NS, Bates AE, Baker SC, Bax NJ, Becerro MA, Berkhout J, Blanchard JL, Brock DJ. 2017.** Assessing national biodiversity trends for rocky and coral reefs through the integration of citizen science and scientific monitoring programs. *Bioscience* 67:134–146 DOI 10.1093/biosci/biw180.

- Tempestini A, Rysgaard S, Dufresne F. 2018.** Species identification and connectivity of marine amphipods in Canada's three oceans. *PLOS ONE* **13**:e0197174 DOI [10.1371/journal.pone.0197174](https://doi.org/10.1371/journal.pone.0197174).
- Thiel M, Penna-Díaz MA, Luna-Jorquera G, Salas S, Sellanes J, Stotz W. 2014.** Citizen scientists and marine research: volunteer participants, their contributions, and projection for the future. *Oceanography and Marine Biology: An Annual Review* **52**:257–314.
- Thomson SA, Pyle RL, Ah Yong ST, Alonso-Zarazaga M, Ammirati J, Araya JF, Ascher JS, Audisio TL, Azevedo-Santos VM, Bailly N. 2018.** Taxonomy based on science is necessary for global conservation. *PLOS Biology* **16**:e2005075 DOI [10.1371/journal.pbio.2005075](https://doi.org/10.1371/journal.pbio.2005075).
- Tittensor DP, Mora C, Jetz W, Lotze HK, Ricard D, Berghe EV, Worm B. 2010.** Global patterns and predictors of marine biodiversity across taxa. *Nature* **466**:1098–1101 DOI [10.1038/nature09329](https://doi.org/10.1038/nature09329).
- Tittensor DP, Walpole M, Hill SLL, Boyce DG, Britten GL, Burgess ND, Butchart SHM, Leadley PW, Regan EC, Alkemade R, Baumung R, Bellard C, Bouwman L, Bowles-Newark NJ, Chenery AM, Cheung WWL, Christensen V, Cooper HD, Crowther AR, Dixon MJR, Galli A, Gaveau V, Gregory RD, Gutierrez NL, Hirsch TL, Höft R, Januchowski-Hartley SR, Karmann M, Krug CB, Leverington FJ, Loh J, Lojenga RK, Malsch K, Marques A, Morgan DHW, Mumby PJ, Newbold T, Noonan-Mooney K, Pagad SN, Parks BC, Pereira HM, Robertson T, Rondinini C, Santini L, Scharlemann JPW, Schindler S, Sumaila UR, Teh LSL, Van Kolck J, Visconti P, Ye Y. 2014.** A mid-term analysis of progress toward international biodiversity targets. *Science* **346**:241–244 DOI [10.1126/science.1257484](https://doi.org/10.1126/science.1257484).
- Vianna G, Meekan M, Pannell D, Marsh S, Meeuwig J. 2012.** Socio-economic value and community benefits from shark-diving tourism in Palau: a sustainable use of reef shark populations. *Biological Conservation* **145**:267–277 DOI [10.1016/j.biocon.2011.11.022](https://doi.org/10.1016/j.biocon.2011.11.022).
- Woolley SNC, Tittensor DP, Dunstan PK, Guillera-Arroita G, Lahoz-Monfort JJ, Wintle BA, Worm B, O'Hara TD. 2016.** Deep-sea diversity patterns are shaped by energy availability. *Nature* **533**:393–396 DOI [10.1038/nature17937](https://doi.org/10.1038/nature17937).
- Yasuhara M, Hunt G, Dowsett HJ, Robinson MM, Stoll DK. 2012.** Latitudinal species diversity gradient of marine zooplankton for the last three million years. *Ecology Letters* **15**:1174–1179 DOI [10.1111/j.1461-0248.2012.01828.x](https://doi.org/10.1111/j.1461-0248.2012.01828.x).