



## Schistosomiasis vector snails and their microbiota display a phylosymbiosis pattern

Camille Huot, Camille Clerissi, Benjamin Gourbal, Richard Galinier, David Duval, Eve R Toulza

### ► To cite this version:

Camille Huot, Camille Clerissi, Benjamin Gourbal, Richard Galinier, David Duval, et al.. Schistosomiasis vector snails and their microbiota display a phylosymbiosis pattern. *Frontiers in Microbiology*, inPress, 10.3389/fmicb.2019.03092 . hal-02440210

**HAL Id: hal-02440210**

**<https://hal.science/hal-02440210>**

Submitted on 10 Mar 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# *Schistosomiasis vector snails and their microbiota display a phyllosymbiosis pattern*

Camille Huot<sup>1</sup>, Camille Clerissi<sup>1,2</sup>, Benjamin Gourbal<sup>1</sup>, Richard Galinier<sup>1</sup>, David Duval<sup>1</sup> and Eve Toulza<sup>1\*</sup>

<sup>1</sup>IHPE, Univ. Montpellier, CNRS, Ifremer, Univ. Perpignan Via Domitia, Perpignan France.

<sup>2</sup>Present address: EPHE, UPVD-CNRS, USR3278 CRIOBE, PSL Research University, Perpignan, France.

**\* Correspondence:**

Eve Toulza

[eve.toulza@univ-perp.fr](mailto:eve.toulza@univ-perp.fr)

**Keywords: microbiota, phyllosymbiosis, metabarcoding, Planorbid snails, tripartite interactions, schistosomiasis. (Min.5-Max. 8)**

## **Abstract**

Planorbidae snails are the intermediate host for the trematode parasite of the *Schistosoma* genus which is responsible for schistosomiasis, a disease that affects both humans and cattle. The microbiota for *Schistosoma* has already been described as having an affect on host/parasite interactions, specifically through immunological interactions. Here we sought to characterize the microbiota composition of seven Planorbidae species and strains. Individual snail microbiota was determined using 16S rDNA amplicon sequencing. The bacterial composition was highly specific to host strain with limited inter-individual variation. In addition, it displayed complete congruence with host phylogeny, revealing a phyllosymbiosis pattern. These results were confirmed in common garden, suggesting that the host highly constrains microbial composition. This study presents the first comparison of bacterial communities between several intermediate snail hosts of *Schistosoma* parasites, paving the way for further studies on the understanding of this tripartite interaction.

## **1 Introduction**

A microbiota consists of microbial communities in association with a host. Here, we defined the microbiota as all microorganisms involved in a long lasting interaction with a host, excluding parasites and pathogen microorganisms (Bordenstein & Theis, 2015). The microbiota is involved in numerous functions, including nutrition (McCutcheon *et al.*, 2009), development (Fraune & Bosch, 2010; McFall-Ngai, 2002), reproduction (Perlman *et al.*, 2008; Werren *et al.*, 2008) and immunity (Hahn & Dheilly, 2016; Lee & Mazmanian, 2010). For example, the bacterial microbiota of the mosquito gut is involved in the immune response of its host against DENV pathogen virus, through direct inhibition of the virus towards bacterial metabolites as well as through indirect effects by stimulating its basal immunity (Saraiva *et al.*, 2016). This illustrates the importance of considering microbiota in host-pathogen interactions.

Numerous studies have already explored the factors shaping microbiota composition in several models and highlighted the role of neutral processes (Burns *et al.*, 2015), environment (Roder *et al.*, 2015), host genetic background (Brucker & Bordenstein, 2011) or host physiology/immunity (Chu & Mazmanian, 2013; Hahn & Dheilly, 2016). Results from these studies demonstrate the effect of the host immune system in microbiota homeostasis. In *Hydra*, the nature and combination of antimicrobial peptides belonging to the arminin family are involved in the species-specificity of host microbial communities that follow host phylogeny (Franzenburg *et al.*, 2013).

Here, we characterized the microbiota of several genera of Planorbidae, a family of freshwater snails. These snails are the intermediate hosts for the parasite *Schistosoma spp.*, a genus of trematode parasites which develop asexually in the snails before infecting vertebrates where sexual reproduction takes place. Human *Schistosoma* species, mainly *S. mansoni*, *S. haematobium* and *S. japonicum*, infect about 250 million people (Hotez *et al.*, 2014) annually, and each year, more than 200 000 people die as a result of the infection worldwide (WHO, 2019). While *Biomphalaria glabrata* and *Biomphalaria pfeifferi* snails can be infected with *Schistosoma mansoni* (responsible for human intestinal schistosomiasis), *Planorbarius metidjensis* is responsible for the transmission of *S. bovis*, and *Bulinus truncatus* snails are natural hosts for *S. bovis* as well as *S. haematobium* (agent of the human urinary schistosomiasis). Interestingly, it has been shown that within the *B. glabrata* species, some strains can be completely refractory to infection depending on the parasite strain, a phenomenon called compatibility polymorphism (Galinier *et al.*, 2017; Theron *et al.*, 2014). These snail-parasite interactions, resulting from coevolution dynamics, reflect differences in host immune capacities or differences in immunobiological interactions between different host-parasite combinations.

The snail immune response in this interaction is complex with a specificity according to the parasite strain (Portela *et al.*, 2013). Indeed, snails immune effectors and receptors seem to be specific to the parasite and the type (cellular or humoral) and efficiency of immune response is linked to the infection type (primo-infection or challenge (homologous or heterologous) (Pinaud *et al.*, 2016). A shift in microbiota composition following an infection was observed after an immune challenge, where humoral immunity took place (Portet *et al.*, 2018). This highlights the importance of further in-depth studies of the relationship between the host's immune and vectorial capacities and its microbiota composition. To do this, it is essential to first characterize the factors that shape microbial communities and their host specificity.

To identify the effect of host identity in bacterial microbiota composition, we used 16S rDNA amplicon sequencing to analyse the bacterial communities at the individual level (10 to 15 samples per condition/strain) for six snail strains: four different strains of *Biomphalaria glabrata*, one strain of *Biomphalaria pfeifferi*, one strain of *Planorbarius metidjensis*, and one strain of *Bulinus truncatus*. Our results provided the first characterization of microbiota for several strains of molluscs, the intermediate hosts of the *Schistosoma sp.* parasite.

## 2 Material & Methods

### 2.1 Rearing conditions

### 2.1.1 Individual tank experiment

To determine the bacterial microbiota composition and specificity, we used four strains of *Biomphalaria glabrata*, one from Guadeloupe (B. gla GH2), two from Brazil (B. gla BAR2 and B. gla BRE2) and one experimentally selected for reduced compatibility to different *S. mansoni* parasite strains (B. gla BS902) (Theron et al., 2014). In addition, *Biomphalaria pfeifferi* (Oman) as well as another Planorbinae genus, *Planorbarius metidjensis* (Spain), and a Planorbidae non-Planorbinae species, *Bulinus truncatus* (Spain) were used (Table 1).

All strains were reared in the same conditions: 20 individuals of each strain were maintained in separate tanks (3L) and fasted one week before sampling to avoid changes in microbiota composition associated with diet. Snail shell size (diameter 7-8 mm), which is directly correlated to age, was similar for each experimental group.

### 2.1.2 Common garden experiment

Thirty molluscs of each strain were maintained for two months within the same 8L tank, where perforated baskets separating the strains were used in order to avoid mixing and potential antagonistic interactions, but which favoured the potential exchange of microbiota as they were reared in the same tank. Molluscs were fed with lettuce every 2 days (and fasted one week before sampling) and 50% of the water was renewed weekly.

## 2.2 Sampling

The mollusc shells were cleaned with cotton buds soaked in bleach (to avoid transfer of contaminants), and molluscs were then removed from the shell by dissection and flash-frozen individually in liquid nitrogen before being kept at -80°C until DNA extraction. DNA extraction and sequencing

DNA was extracted with the Nucleospin® tissue extraction kit from Macherey-Nagel and quantified with Qubit 2.0 Fluorometer following the procedure described in the Qubit™ dsDNA HS Assay Kit, to check its purity and yield.

For samples with highest DNA yield and quality (11 to 15 depending on snail strains for individual tank experiment, and 10 per strain for the common garden experiment, Table S1), 16S rRNA gene (V3-V4 regions) (Klindworth et al., 2013) libraries were generated using PCR primers 341F (5'-CCTACGGGNGGCWGCAG-3') and 805R (3'-GACTACHVGGGTATCTAATCC-5') following the standard Illumina two-step procedure. Libraries were paired-end sequenced with 250 bp read length on three different flow cells using the MiSeq system (Illumina) at the Génome Québec Innovation Centre, McGill University Montréal, Canada. A blank sample was sequenced in each of the three runs but as very few sequences were obtained, this dataset was not further analyzed.

## 2.3 Analysis of 16S sequences

The FROGS (Find Rapidly OTU with Galaxy Solution) pipeline implemented on a galaxy instance (<http://sigenae-workbench.toulouse.inra.fr/galaxy/>) was used for data processing (Escudié et al., 2017). In brief, paired reads were merged using FLASH (Magoč & Salzberg, 2011). After denoising and primers/adapters removal with CUTADAPT (Martin, 2011), de novo clustering was performed using SWARM with a local clustering threshold (Mahé et al., 2014), with aggregation distance  $d = 3$  after denoising. Chimeras were removed using VSEARCH (Rognes et al., 2016). We filtered the

dataset for singletons and performed affiliation using Blast+ against the Silva database (release 128, September 2016) for 16S rRNA gene amplicons. Finally, OTU tables were produced in a standard BIOM format for subsequent analyses.

We then used the packages phyloseq 1.24.2 (McMurdie & Holmes, 2013) and vegan 2.5-4 (Oksanen et al., 2018) with RStudio (R Core Team (2017)). Sample B. gla BRE\_JC\_7 had too low coverage (155 reads) and was thus discarded from subsequent analyses. Non-bacterial sequences as well as singletons remaining after all the secondary filtering steps were discarded from the dataset. We rarefied the data according to the sample with fewer sequence numbers (18299 reads for the Individual Experiment and 15969 reads for the Common Garden) in order to normalize for sequencing coverage. We characterized the beta-diversity dissimilarities using Principal Coordinates Analyses (PCoA) and Hierarchical Clustering on Bray-Curtis distance matrix (ranging from 0 for identical communities to 1 for completely different communities).

## 2.4 Core microbiota

To determine the core microbiota, which is the most stable part of the microbiota, we identified the families and genera that were either present in 100% of individuals or absent from a maximum of one individual for each strain, and represented at least 0.5% of sequences for each strain.

## 2.5 Snail Phylogeny

Phylogenetic analysis was performed using 28S rRNA gene sequences from the NCBI database (*B. glabrata*: AF435694.1, *B. pfeifferi*: MG461588.1, *P. metidjensis*: AF435671.1 and *B. truncatus*: AF435659.1). The 28S rRNA gene sequence of a Physidae species, *Physa* sp. (Egypt) (sister family of the Planorbidae) was used as an outgroup (AF435654.1). The sequences were aligned using MUSCLE (v3.8.31) and the tree was reconstructed using the maximum likelihood method implemented in the PhyML program (v3.1/3.0 aLRT) with 500 bootstraps on <http://www.phylogeny.fr> (Dereeper et al., 2008; Dereeper et al., 2010).

## 2.6 Statistical analyses

We analysed the variance due to host effect on dissimilarity matrices using Permutational Multivariate Analysis of Variance (PERMANOVA). PERMANOVA were done with 999 permutations. For all analyses, the threshold significance level was set at 0.05.

We used an indicator value index and 999 permutations (multipatt, {indicspecies}) (Caceres and Legendre 2009) to identify OTUs associated with the different host species. P-values were corrected for multiple comparisons using Benjamini and Hochberg's method (Benjamini and Hochberg 1995) (p.adjust, {stats}).

# 3 Results

## 3.1 Bacterial composition

At the phylum level, the composition of bacteria was similar in all *Biomphalaria* samples (Figure 1), with Proteobacteria being the predominant phylum for the different strains, in which the Flavobacteriaceae, Rhodobacteraceae, Comamonadaceae and Xanthomonadaceae families were the most represented. This is consistent with the results found by Portet et al. (2018). *P. metidjensis* composition also displayed a high proportion of Proteobacteria, in particular Alphaproteobacteria,

120 represented by Rhodobacteraceae (Table S1). In the case of *B. truncatus*, more pronounced  
 121 differences were visible at the phylum level, where Proteobacteria and Tenericutes were dominant  
 122 (Figure 1), with the latter represented mainly by Mycoplasmataceae, and more specifically by the  
 123 genus *Mycoplasma* (Table S1).

124 The core microbiota was determined as all bacterial families that were either present in 100% of  
 125 individuals or absent from a maximum of one individual. The core microbiota was composed of 44  
 126 families, for all strains included (Table S2). The core microbiota composition varied between strains;  
 127 whereas seven bacterial families were common to all strains, where *Cloacibacterium* (a  
 128 Flavobacteriaceae genus) were found as part of the core microbiota in all *Biomphalaria* strains and  
 129 species, except for *B. glabrata* BS90, which was absent in the core microbiota of *P. metidjensis* and  
 130 *B. truncatus*.

### 131 3.2 Beta-diversity, ordination and clustering

132 An ordination using PCoA was performed on Bray-Curtis (BC) distance matrix to visualize the  
 133 similarities between individuals according to their bacterial composition (Figure 2). The first two  
 134 axes explained 30% of the variability observed. Individuals tended to group according to host  
 135 species. *Biomphalaria* strains were grouped at the exception of the *B. glabrata* BS902 strain.  
 136 Individuals of the two other species, *Planorbarius metidjensis* and *Bulinus truncatus* were separated  
 137 from *Biomphalaria* individuals.

138 The hierarchical clustering analysis based on BC distance on the core microbiota confirmed a  
 139 grouping between individuals of the same strain or species. Moreover, the dendrogram of bacterial  
 140 communities reflected host phylogeny (Figure. 3). *B. truncatus* and *P. metidjensis* were separated  
 141 from *Biomphalaria* species, and *B. pfeifferi* was separated from *B. glabrata* strains. The microbiota  
 142 specificity according to host genetic background was confirmed by Manova on Bray-Curtis  
 143 dissimilarity matrix on core microbiota ( $p < 0.001$ ).

144 The core microbiota beta-diversity was analysed using the same approach with a PCoA ordination  
 145 based on the Bray-Curtis dissimilarity index (Figure. 3). Individuals belonging to the same strain tend  
 146 to cluster together and *Biomphalaria* strains were grouped, except for, again, the BS902 strain. In  
 147 addition, individuals from the two other species, *Planorbarius metidjensis* and *Bulinus truncatus*  
 148 were separated from *Biomphalaria* individuals. This analysis of the core microbiota composition  
 149 confirmed the pattern obtained for the whole microbiota with specific core microbiota associations  
 150 for individuals belonging to the same phylogenetic group (strain, species or genus), suggesting a  
 151 phyllosymbiosis pattern, driven by host species among snail intermediate hosts of schistosomiasis.

152 We used indicator value index and permutation tests to identify OTUs significantly associated with  
 153 each host species. In average, each species had 37 specific OTUs belonging to 88 genera (Table S3).  
 154 Although the 88 genera were mostly (77%) specific of each host species, this analysis highlighted  
 155 that specific OTUs belonging to 5 genera (*Pirellula*, *Planctomyces*, *Candidates Odysella*,  
 156 *Mesorhizobium*, and *Pseudomonas*) were found in more than 50% of host species. Strikingly, specific  
 157 OTUs from *Mesorhizobium* and *Pseudomonas* showed identical distribution within host species  
 158 (presence in BRE, BS90, Pfe, and Plan), suggesting that these bacteria might cooperate within host  
 159 microbiome.

### 160 3.3 Common Garden

Environmental conditions and/or host genetics can both affect microbiota composition. To investigate the main key drivers for core microbiota composition, we performed an additional experiment with all strains raised together in the same water for two months. However, we couldn't include *B. pfeifferi* in this analysis because most individuals did not survive until the end of the experiment as they escaped their basket and were predated.

The PCoA ordination revealed a similar microbiota specificity by strain to those observed in the first experiment with a grouping by strain then species (Figure S1), as confirmed by Manova analysis on host effect on Bray-Curtis dissimilarities between host strains ( $p < 0.001$ ).

#### 4 Discussion

To understand the host effect in shaping microbiota in Planorbidae schistosomiasis vector snails, we characterized the individual bacterial communities associated with several strains of *Biomphalaria glabrata*, *B. pfeifferi*, *P. metidjensis* and *B. truncatus* snails. Working on individuals reared in lab conditions favoured the control of most of the parameters that can influence microbiota composition.

In the present study, the whole microbiota was characterized using 16S amplicon sequencing. We identified 31207 OTUs among the six different snail strains. Most of OTUs were not assigned to the species level and 63% were assigned to the genus level. This corresponds to the limitation of the 16S V3V4 marker resolution. In addition, the blast+-based pipeline we use for taxonomic affiliation avoids false affiliation when a sequence matches with several sequences in the database. If several blast results have identical scores for a given OTU, and these taxonomies differ across hits, the OTU is set to "Multi-affiliation" (Escudié et al. 2017).

A few studies have characterized the cultivable flora of *B. glabrata* and have identified Aeromonadaceae, Enterobacteriaceae, Moraxellaceae and Pseudomonadaceae as being the most prevalent bacterial families in this species (Ducklow et al., 1979; Ducklow et al., 1981; Silva et al., 2013). The dominant families described in the previous studies were also represented in our dataset. However, the relative composition of microbiota at the phylum level revealed that Proteobacteria were dominant for most of the different *Biomphalaria* strains, represented by three main families: Rhodobacteraceae (Alphaproteobacteria), Commamonadaceae (Betaproteobacteria) and Xanthomonadaceae (Gammaproteobacteria). Not all bacterial families can easily be cultivated; the MiSeq technology allows identifying the whole bacterial diversity. Our results are consistent with those found for *B. glabrata* BRE bacterial microbiota in Portet et al. (2018), in which these three families were the most abundant of the core microbiota. The microbiota of a Guadeloupian strain of *B. glabrata* was also described using a similar approach (Allan et al., 2018) and similarly, the dominant phyla were Proteobacteria and Bacteroidetes.

Proteobacteria have been described as key factor in marine bivalve digestion, like the great scallop *Pecten maximus*, as they are involved in the degradation of major alimentary components contained in their diet (Lasa et al., 2016). This phylum is also dominant in other molluscs, as is the case for oysters *Crassostrea corteziensis*, *C. gigas* and *C. sikamea* (Trabal et al., 2012). As this is the first study to characterize the bacterial microbiota of *B. glabrata* BS902, *P. metidjensis*, *B. truncatus*, it is not possible to compare with previous results and to draw any definitive conclusions.

Interestingly, the bacterial families that comprise the core microbiota for each mollusc strain were also among the most abundant taxa in the whole microbiota, which is consistent with previous studies on this model (Portet et al., 2018). In corals, for example, the OTUs belonging to core microbiota are among the rare taxa and are difficult to detect within the whole microbiota (Ainsworth et al., 2015).

Due to the high inter-individual variation, in some studies, the core microbiota in the coral model was defined by OTUs present in a limited proportion of individuals, 30% in Ainsworth *et al.* (2015) and 50% in Brener-Raffalli *et al.* (2018). In our study, the most impressive case of high abundance in core taxa concerns *B. truncatus*, with the Mycoplasmataceae family, and more precisely the Genus *Mycoplasma*, that represents more than 47% of the whole microbiota. The Genus *Mycoplasma* was originally described as an obligate vertebrate parasite and the causative agent of human genital and respiratory diseases with a high tissue specificity (Razin *et al.*, 1998). This genus has been described in other models including algae and several invertebrates such as oysters (Clerissi *et al.*, 2018; King *et al.*, 2012), abalone (Huang *et al.*, 2010) and Sacoglossans (Davis *et al.*, 2013). It has also been described as being one of the most abundant microorganisms in the deep-sea bone-eating snail, *Rubyspira osteovora* (Aronson *et al.*, 2016). Its role in these organisms remains unclear but some authors hypothesized that they may help with digestion (Aronson *et al.*, 2016; Duperron *et al.*, 2012; Fraune & Zimmer, 2008), notably because of its presence in the digestive tract. In the present study, bacteria belonging to the genus *Cloacibacterium* have been found in all *Biomphalaria* strains and species, and *C. haliotis* has been described in another mollusc, the sea snail *Haliotis discus* (Hyun *et al.*, 2014).

The results of dissimilarity between strains revealed that the bacterial microbiota of *B. truncatus* individuals was distinct from other species with most of the Bray-Curtis distance values ranging between 0.8 and 0.99. The phylogenetic distance of this species from the others could explain this difference. Indeed, this is the only species, in this study, belonging to Bulinae sub-family, whereas all the others are classified in Planorbinae sub-family. Moreover, the bacterial microbiota of this species seems to be very specific, as suggested by the high abundance of *Mycoplasma*. Interestingly, individuals of the strain Bg BS90 also displayed strong dissimilarities with the other species and even the other strains of *B. glabrata*, with most of the dissimilarity values also ranging from 0.8 and 0.99.

As the different strains were maintained in separate tanks, we performed a common garden experiment to circumvent potential biases due to mollusc maintenance and tested whether the same microbial environment would lead to a homogeneous distribution of the bacterial communities between snail strains. This result confirmed a specificity of the microbiota by strain/species, suggesting that the importance of the host effect in microbiota composition is higher than the effect of rearing conditions. The microbiota can nevertheless vary during the host lifespan, with an initial recruitment of bacterial communities occurring during early development. It would be interesting to test the possibility of microbiota transfers from the environment in different developmental stages when the definitive flora is not yet fully established. A recent study showed a loss of microbial communities from one generation to the next in laboratory reared mosquitoes (Akorli *et al.*, 2019), which presents another avenue for our model to be further investigated.

In both individual and common garden experiments, almost every individual of each strain grouped together in the dendrogram, supporting the specificity according to the host. Additionally, the topology of microbiota dissimilarities was congruent with the mollusc phylogeny, despite a limited number of strains but that covers species, genera and sub-families of Planorbidae. This revealing suggests a pattern of phyllosymbiosis ~~driven by~~at the host species level among snail intermediate hosts of schistosomiasis. This has already been described in other models, for both vertebrates and invertebrates. In vertebrates, for example, a loose phyllosymbiosis pattern was identified between 44 species of coral reef fishes and their skin microbiota (Chiarello *et al.*, 2018), possibly related to a plasticity in the immune system. Host immune genes and other factors like nutrient production by the host and vertical transmission have also been hypothesized to explain phyllosymbiosis between

several populations of American pika, *Ochotona princeps* (Kohl *et al.*, 2017). For invertebrates, this pattern was shown in three *Nasonia* species, in a controlled environment, with such a codiversification and coevolution that there is a lethality of hybrids from a breed between two *Nasonia* species (Brucker & Bordenstein, 2013). This codiversification as a mechanism leading to phylosymbiosis has also been hypothesized in a study comparing microbiota composition of 15 *Cephalotes* species (Sanders *et al.*, 2014), whereas it would not be the main driver of this phenomenon in corals, in which phylosymbiosis would be led by other mechanisms like biogeography or host traits (Pollock *et al.*, 2018). Similar findings of phylosymbiosis driven by the host have been identified between two different species of *Hydra* (Fraune & Bosch, 2007) and many studies have shown that the host genetic background shape the microbiota in numerous models (Chaston *et al.*, 2016; Coon *et al.*, 2016; Paniagua Voirol *et al.*, 2018; Parker *et al.*, 2017; Sánchez-Cañizares *et al.*, 2017). In our model, this correlation between host and microbiota indicates that host phylogeny highly constrains the microbiota composition and structuration (Brooks *et al.*, 2016; Chiarello *et al.*, 2018). However, this pattern may not be ubiquitous and a few studies on *Drosophila* (Chandler *et al.*, 2011), mosquitos (Osei-Poku *et al.*, 2012) or flea beetles (Kelley & Dobler, 2011) identified no correlation between host phylogeny and microbiota composition. Nevertheless, we could not assess the phylosymbiosis pattern at a lower phylogenetic level (i.e. the strain) as we cannot determine the genetic distance between the different *Biomphalaria glabrata* strains because of inbreeding in the laboratory and high differentiation between strains.

In our case, the phylosymbiosis pattern could not be considered as a hallmark of coevolution because we focused on the whole bacterial community of a host, with very complex interactions, and not on a specific symbiont. Here, we defined coevolution according to O'Brien and collaborators (2019), as a "reciprocal evolution of [...] a broad range of interactions such as predator-prey, host-symbiont and host-parasite interactions, or interactions among the members of a community of organisms such as a host and its associated microbiome". O'Brien *et al.* noticed that hosts and their symbiont phylogenies are often mirrored, which can be interpreted as a parallel divergence called a codivergence. This codivergence has often led to obligatory symbiosis, as is the case between pea aphids and bacteria from the genus *Buchnera* (Baumann *et al.*, 2006) and is notably found in mutualistic symbiosis (O'Brien *et al.*, 2019). In this case, the protagonists have a very close interaction, with participation in each other's physiological mechanisms.

~~We considered the phylosymbiosis pattern to~~ The host-microbiota specificity illustrates the high interaction between the snails and their ~~microbiota~~ bacterial communities, suggesting a impact of the latter on its host fitness toward several functions like nutrition, development, reproduction and immunity. Given that, in this model, Planorbidae snails are intermediate hosts of *Schistosoma* parasites, it would be interesting to study the tripartite interaction between the trematodes, the molluscs and their microbiota.

Indeed, previous studies highlighted a variation in the compatibility phenotype between different combinations of *B. glabrata* strains and *S. mansoni* parasites (Galinier *et al.*, 2017; Theron *et al.*, 2014). Moreover, *P. metidjensis* and *B. truncatus* are not compatible with the same *Schistosoma* species. This compatibility polymorphism can be seen as a hallmark of differences in immune capacities. As the phylosymbiosis pattern suggests a strong link between host and microbiota, the hypothesis of a relationship between the snails' immune capacities and the composition of their microbiota can be made.

The protective role of whole microbiota (or gut microbiota), has indeed been shown in numerous models like the mosquitoes against dengue virus (Ramirez *et al.*, 2012) or the honey bees with the

augmentation in antimicrobial peptide production (Li *et al.*, 2017). Another example is the microbiota of *Dysdercus fasciatus* that acts as a physical barrier to prevent the entry or attachment of a parasite (Onchuru *et al.*, 2018). Chiu and collaborators (2017) also reviewed several examples of microbiota actions against pathogens, such as slowing or preventing the entry, installation, development and expansion of pathogens. In some models, the microbiota has a direct effect against their hosts' pathogens, producing effectors like ROS (Cirimotich *et al.*, 2011), or an indirect effect, promoting some immune pathway. Concerning the interaction between Planorbidae and *Schistosoma*, the immune mechanisms have been well studied, however, there are very few informations concerning the tripartite interactions. Some immune genes located in a Guadeloupe resistance to parasite complex (GRC) region have been shown previously to contribute in shaping microbiota (Allan *et al.*, 2018) highlighting a link between microbiota composition and host immunity.

The present study highlighted a ~~phylosymbiosis pattern~~strong host-microbiota specificity, which confirms the link between host genetics, immune capacity and microbiota composition. However, more information are needed to understand if there is a direct or indirect impact of microbiota on the host-parasite interaction.

The interaction between microorganisms and the host immune system can be complex. The microbiota stability can be affected upon parasite primo-infestation and challenge suggesting a tight control of immune system on bacterial composition (Portet *et al.*, 2018). The next step will be to compare the microbiota dynamics during an infection kinetic with several host/parasite combinations with different immunobiological interactions. Although a shift in microbiota composition during an infection associated with changes in snail immune gene expression was clearly established according to the host/parasite combination (sympatric/allopatric) (Portet *et al.*, 2018), further studies are needed to clarify the link between microbiota and snail host immunity. Phylosymbiosis pattern is a hallmark of tight interactions between host and microbiota, suggesting the role of microbial communities on different host physiological functions, including immunity. This study thus paves the way for future studies to decipher the role of microbiota in host fitness, including the development and transmission of parasites.

## Declarations

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author Contributions: CH, BG, RG, DD and ET were involved in the study concept and design. CH was involved in sampling and data acquisition. CH, CC and ET performed data analysis. CH and ET drafted the manuscript and all authors contributed to critical revisions and approved the final manuscript.

Funding: This work was supported by the DHOF program of the IHPE lab (<http://ihpe.univ-perp.fr/en/ihpe-transversal-holobiont/>). This study is set within the framework of the "Laboratoires d'Excellence (LABEX)" TULIP (ANR-10-LABX-41) and CeMEB (ANR-10-LABX-04-01).

Acknowledgments: We thank Nathalie Arancibia for and Jean-François Allienne for technical assistance. We are grateful to the genotoul bioinformatics platform Toulouse Midi-Pyrenees and Sigenae group for providing help and computing resources thanks to Galaxy instance <http://sigenae-workbench.toulouse.inra.fr>.

335 Data Availability Statement: The datasets generated for this study can be found in the Sequence read  
336 Archive repository under BioProject PRJNA554540 (sequence data to be released upon publication).

337 **References**

- 338 Ainsworth, T. D., Krause, L., Bridge, T., Torda, G., Raina, J.-B., Zakrzewski, M., Gates, R. D.,  
 339 Padilla-Gamiño, J. L., Spalding, H. L., Smith, C., Woolsey, E. S., Bourne, D. G., Bongaerts, P.,  
 340 Hoegh-Guldberg, O., & Leggat, W. (2015). The coral core microbiome identifies rare bacterial taxa  
 341 as ubiquitous endosymbionts. *The ISME Journal*, 9, 2261–2274.  
 342 <https://doi.org/10.1038/ismej.2015.39>
- 343 Akorli, J., Namaali, P. A., Ametsi, G. W., Kwesi Egyirifa, R., Adjoa, N., & Pels, P. (2019).  
 344 Generational conservation of composition and diversity of field-acquired midgut microbiota in  
 345 *Anopheles gambiae* (sensu lato) during colonization in the laboratory. *Parasites & Vectors*, 12(27),  
 346 1–9. <https://doi.org/10.1186/s13071-019-3287-0>
- 347 Allan, E. R. O., Tennessen, J. A., Sharpton, T. J., & Blouin, M. S. (2018). Allelic variation in a single  
 348 genomic region alters the microbiome of the snail *Biomphalaria glabrata*. *Journal of Heredity*, 1–6.  
 349 <https://doi.org/10.1093/jhered/esy014>
- 350 Aronson, H. S., Zellmer, A. J., & Goffredi, S. K. (2016). The specific and exclusive microbiome of  
 351 the deep-sea bone-eating snail, *Rubyspira osteovora*. *FEMS Microbiology Ecology*, 93(3), 1–13.  
 352 <https://doi.org/10.1093/femsec/fiw250>
- 353 Baumann, P., Moran, N. A., & Baumann, L. (2006). The evolution and genetics of aphid  
 354 endosymbionts. *BioScience*, 47(1), 12–20. <https://doi.org/10.2307/1313002>
- 355 Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to  
 356 multiple testing. *J. R. Stat. Soc. Ser. B*. 1995;289–300.
- 357 Bordenstein, S. R., & Theis, K. R. (2015). Host biology in light of the microbiome: ten principles of  
 358 holobionts and hologenomes. *PLoS Biol*, 13(8), 1–23. <https://doi.org/10.1371/journal.pbio.1002226>
- 359 Brener-Raffalli, K., Clerissi, C., Vidal-Dupiol, J., Adjerdoud, M., Bonhomme, F., Pratlong,  
 360 M., Aurelle, D., Mitta, G., & Toulza, E. (2018). Thermal regime and host clade, rather than  
 361 geography, drive Symbiodinium and bacterial assemblages in the scleractinian coral *Pocillopora*  
 362 *damicornis* sensu lato. *Microbiome*, 6(39), 1–13. <https://doi.org/10.1186/s40168-018-0423-6>
- 363 Brooks, A. W., Kohl, K. D., Brucker, R. M., van Opstal, E. J., Bordenstein, S. R., & Relman, D.  
 364 (2016). Phyllosymbiosis: relationships and functional effects of microbial communities across host  
 365 evolutionary history. *PLoS Biol*, 14(11), 1–29. <https://doi.org/10.1371/journal.pbio.2000225>
- 366 Brucker, R. M., & Bordenstein, S. R. (2013). The hologenomic basis of speciation: gut bacteria cause  
 367 hybrid lethality in the genus *Nasonia*. *Science*, 341(6146), 667–669.  
 368 <https://doi.org/10.1126/science.1240659>
- 369 Brucker, Robert M., & Bordenstein, S. R. (2011). The roles of host evolutionary relationships (genus:  
 370 *Nasonia*) and development in structuring microbial communities. *Evolution*, 66(2), 349–362.  
 371 <https://doi.org/10.1111/j.1558-5646.2011.01454.x>
- 372 Burns, A. R., Stephens, Z., Stagaman, K., Wong, S., Rawls, J. F., Guillemin, K., & Bohannan, B. J.  
 373 (2015). Contribution of neutral processes to the assembly of gut microbial communities in the

- 374 zebrafish over host development. The ISME Journal, 10, 655–664.  
375 <https://doi.org/10.1038/ismej.2015.142>
- 376 Cáceres MD, Legendre P. Associations between species and groups of sites: indices and statistical  
377 inference. Ecology. 2009;90:3566–3574.
- 378 Chandler, J. A., Lang, J. M., Bhatnagar, S., Eisen, J. A., & Kopp, A. (2011). Bacterial communities  
379 of diverse *Drosophila* species: ecological context of a host-microbe model system. PLoS Genetics,  
380 7(9), e1002272. <https://doi.org/10.1371/journal.pgen.1002272>
- 381 Chaston, J. M., Dobson, A. J., Newell, P. D., & Douglas, A. E. (2016). Host genetic control of the  
382 microbiota mediates the *Drosophila* nutritional phenotype. Applied and Environmental  
383 Microbiology, 82(2), 671–679. <https://doi.org/10.1128/AEM.03301-15>
- 384 Chiarello, M., Auguet, J. C., Bettarel, Y., Bouvier, C., Claverie, T., Graham, N. A. J.,  
385 Rieuvilleneuve, F., Sucré, E., Bouvier, T., & Villéger, S. (2018). Skin microbiome of coral reef fish  
386 is highly variable and driven by host phylogeny and diet. Microbiome, 6(1).  
387 <https://doi.org/10.1186/s40168-018-0530-4>
- 388 Chiu, L., Bazin, T., Truchetet, M. E., Schaefferbeke, T., Delhaes, L., & Pradeu, T. (2017). Protective  
389 microbiota: From localized to long-reaching co-immunity. Frontiers in Immunology, 8(1678).  
390 <https://doi.org/10.3389/fimmu.2017.01678>
- 391 Chu, H., & Mazmanian, S. K. (2013). Innate immune recognition of the microbiota promotes host-  
392 microbial symbiosis. Nat Immunol, 14(7), 668–675. <https://doi.org/10.1038/ni.2635>
- 393 Cirimotich, C. M., Dong, Y., Clayton, A. M., Sandiford, S. L., Souza-Neto, J. A., Mulenga, M., &  
394 Dimopoulos, G. (2011). Natural microbe-mediated refractoriness to *Plasmodium* infection in  
395 *Anopheles gambiae*. Science, 332(6031), 855–858. <https://doi.org/10.1126/science.1201618>
- 396 Clerissi, C., de Lorgeril, J., Petton, B., Lucasson, A., Gueguen, Y., Mitta, G., & Toulza, E. (2018).  
397 Diversity and stability of microbiota are key factors associated to healthy and diseased *Crassostrea*  
398 *gigas* oysters. BioRxiv. <https://doi.org/10.1101/378125>
- 399 Coon, K. L., Brown, M. R., & Strand, M. R. (2016). Mosquitoes host communities of bacteria that  
400 are essential for development but vary greatly between local habitats. Molecular Ecology, 25(22),  
401 5806–5826. <https://doi.org/10.1111/mec.13877>
- 402 Davis, J., Fricke, W. F., Hamann, M. T., Esquenazi, E., Dorrestein, P. C., & Hill, R. T. (2013).  
403 Characterization of the bacterial community of the chemically defended Hawaiian sacoglossan *Elysia*  
404 *rufescens*. Applied and Environmental Microbiology, 79(22), 7073–7081.  
405 <https://doi.org/10.1128/AEM.01568-13>
- 406 Dereeper, A., Guignon, V., Blanc, G., Audic, S., Buffet, S., Chevenet, F., Dufayard, J.-F., Guindon,  
407 S., Lefort, V., Lescot, M., Claverie, J.-M., & Gascuel, O. (2008). Phylogeny.fr: robust phylogenetic  
408 analysis for the non-specialist. Nucleic Acids Research, 36, 465–469.  
409 <https://doi.org/10.1093/nar/gkn180>

- 410 Dereeper, Alexis, Audic, S., Claverie, J.-M., & Blanc, G. (2010). BLAST-EXPLORER helps you  
411 building datasets for phylogenetic analysis. BMC Evolutionary Biology, 10(8), 1–6.  
412 <https://doi.org/10.1186/1471-2148-10-8>
- 413 Ducklow, H. W., Boyle, P. J., Mangel, P. W., Strong, C., & Mitchell, R. (1979). Bacterial flora of the  
414 schistosome vector snail *Biomphalaria glabrata*. Applied and Environmental Microbiology, 38(4),  
415 667–672.
- 416 Ducklow, H. W., Clausen, K., & Mitchell, R. (1981). Ecology of bacterial communities in the  
417 schistosomiasis vector snail *Biomphalaria glabrata*. Microbial Ecology, 7(3), 253–274.  
418 <https://doi.org/10.1007/BF02010308>
- 419 Duperron, S., Pottier, M. A., Léger, N., Gaudron, S. M., Puillandre, N., Le Prieur, S., Sigwart, J. D.,  
420 Ravaux, J., & Zbinden, M. (2012). A tale of two chitons: Is habitat specialisation linked to distinct  
421 associated bacterial communities? FEMS Microbiology Ecology, 83(3), 552–567.  
422 <https://doi.org/10.1111/1574-6941.12014>
- 423 Escudié, F., Auer, L., Bernard, M., Mariadassou, M., Cauquil, L., Vidal, K., Maman, S., Hernandez-  
424 Raquet, G., Combes, S., & Pascal, G. (2017). FROGS: Find, Rapidly, OTUs with Galaxy Solution.  
425 Bioinformatics, 34(8), 1287–1294. <https://doi.org/10.1093/bioinformatics/btx791>
- 426 Franzenburg, S., Walter, J., Künzel, S., Wang, J., Baines, J. F., Bosch, T. C. G., & Fraune, S. (2013).  
427 Distinct antimicrobial peptide expression determines host species-specific bacterial associations.  
428 Proceedings of the National Academy of Science, 110(39), E3730–E3738.  
429 <https://doi.org/10.1073/pnas.1304960110>
- 430 Fraune, S., & Bosch, T. C. G. (2007). Long-term maintenance of species-specific bacterial  
431 microbiota in the basal metazoan *Hydra*. PNAS, 104(32), 13146–13151. Retrieved from  
432 [www.pnas.org/cgi/content/full/](http://www.pnas.org/cgi/content/full/)
- 433 Fraune, S., & Bosch, T. C. G. (2010). Why bacteria matter in animal development and evolution.  
434 BioEssays, 32(7), 571–580. <https://doi.org/10.1002/bies.200900192>
- 435 Fraune, S., & Zimmer, M. (2008). Host-specificity of environmentally transmitted *Mycoplasma*-like  
436 isopod symbionts. Environmental Microbiology, 10(10), 2497–2504. <https://doi.org/10.1111/j.1462-2920.2008.01672.x>
- 438 Galinier, R., Roger, E., Moné, Y., Duval, D., Portet, A., Pinaud, S., Chaparro, C., Grunau, C.,  
439 Genthon, C., Dubois, E., Rognon, A., Arancibia, N., Dejean, B., Théron, A., Gourbal, B., & Mitta, G.  
440 (2017). A multistrain approach to studying the mechanisms underlying compatibility in the  
441 interaction between *Biomphalaria glabrata* and *Schistosoma mansoni*. PLoS Neglected Tropical  
442 Diseases, 11(3). <https://doi.org/10.1371/journal.pntd.0005398>
- 443 Hahn, M. A., & Dheilly, N. M. (2016). Experimental models to study the role of microbes in host-  
444 parasite interactions. Frontiers in Microbiology, 7(1300), 1–8.  
445 <https://doi.org/10.3389/fmicb.2016.01300>
- 446 Hotez, P. J., Alvarado, M., Basá, M.-G., Bolliger, I., Bourne, R., Boussinesq, M., Brooker, S. J.,  
447 Brown, S. A., Buckle, G., Budke, C. M., Halasa, Y. A., Jasrasaria, R., Johns, N. E., Keiser, J., King,

- 448 C. H., Lozano, R., Murdoch, M. E., O'hanlon, S., Bastien, S., Pion, D. S., Pullan, R. L., Ramaiah, K.  
449 D., Roberts, T., Shepard, D. S., Smith, J. L., Stolk, W. A., Undurraga, E. A., Utzinger, J., Wang, M.,  
450 Murray, C. J. L., & Naghavi, M. (2014). The global burden of disease study 2010: interpretation and  
451 implications for the neglected tropical diseases. *PLoS Neglected Tropical Diseases*, 8(7), 1–9.  
452 <https://doi.org/10.1371/journal.pntd.0002865>
- 453 Huang, Z. Bin, Guo, F., Zhao, J., Li, W. D., & Ke, C. H. (2010). Molecular analysis of the intestinal  
454 bacterial flora in cage-cultured adult small abalone, *Haliotis diversicolor*. *Aquaculture Research*,  
455 41(11), e760–e769. <https://doi.org/10.1111/j.1365-2109.2010.02577.x>
- 456 Hyun, D.-W., Shin, N.-R., Kim, M.-S., Kim, Y., Kim, S., Sei, J. O., Whon, T. W., & Bae, J.-W.  
457 (2014). *Cloacibacterium haliotis* sp. nov., isolated from the gut of an abalone, *Haliotis discus hannai*.  
458 *International Journal of Systematic and Evolutionary Microbiology*, 64, 72–77.  
459 <https://doi.org/10.1099/ij.s.0.054585-0>
- 460 Kelley, S. T., & Dobler, S. (2011). Comparative analysis of microbial diversity in *Longitarsus* flea  
461 beetles (Coleoptera: Chrysomelidae). *Genetica*, 139(5), 541–550. [https://doi.org/10.1007/s10709-](https://doi.org/10.1007/s10709-010-9498-0)  
462 [010-9498-0](https://doi.org/10.1007/s10709-010-9498-0)
- 463 King, G. M., Judd, C., Kuske, C. R., & Smith, C. (2012). Analysis of stomach and gut microbiomes  
464 of the eastern oyster (*Crassostrea virginica*) from coastal Louisiana, USA. *PloS One*, 7(12), e51475.  
465 <https://doi.org/10.1371/journal.pone.0051475>
- 466 Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., & Glöckner, F. O. (2013).  
467 Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation  
468 sequencing-based diversity studies. *Nucleic Acids Research*, 41(1), e1.  
469 <https://doi.org/10.1093/NAR/GKS808>
- 470 Kohl, K. D., Denise Dearing, M., Bordenstein, S. R., & Author, C. (2018). Microbial communities  
471 exhibit host-species distinguishability and phyllosymbiosis along the length of the gastrointestinal  
472 tract. *Mol Ecol*, 27(8), 1874–1883. <https://doi.org/10.1111/mec.14460>
- 473 Kohl, K. D., Varner, J., Wilkening, J. L., & Dearing, M. D. (2017). Gut microbial communities of  
474 American pikas (*Ochotona princeps*): Evidence for phyllosymbiosis and adaptations to novel diets.  
475 *Journal of Animal Ecology*, 87(2), 323–330. <https://doi.org/10.1111/1365-2656.12692>
- 476 Lasa, A., Mira, A., Camelo-Castillo, A., Belda-Ferre, P., & Romalde, J. L. (2016). Characterization  
477 of the microbiota associated to *Pecten maximus* gonads using 454-pyrosequencing. *International*  
478 *Microbiology*, 19, 93–99. <https://doi.org/10.2436/20.1501.01.267>
- 479 Lee, Y. K., & Mazmanian, S. K. (2010). Has the microbiota played a critical role in the evolution of  
480 the adaptive immune system? *Science (New York, N.Y.)*, 330(6012), 1768–1773.  
481 <https://doi.org/10.1126/science.1195568>
- 482 Li, J. H., Evans, J. D., Li, W. F., Zhao, Y. Z., DeGrandi-Hoffman, G., Huang, S. K., Li, Z. G.,  
483 Hamilton, M., & Chen, Y. P. (2017). New evidence showing that the destruction of gut bacteria by  
484 antibiotic treatment could increase the honey bee's vulnerability to *Nosema* infection. *PLoS ONE*,  
485 12(11), 1–18. <https://doi.org/10.1371/journal.pone.0187505>

- 486 Magoč, T., & Salzberg, S. L. (2011). FLASH: fast length adjustment of short reads to improve  
487 genome assemblies. *Bioinformatics* (Oxford, England), 27(21), 2957–2963.  
488 <https://doi.org/10.1093/bioinformatics/btr507>
- 489 Mahé, F., Rognes, T., Quince, C., de Vargas, C., & Dunthorn, M. (2014). Swarm: robust and fast  
490 clustering method for amplicon-based studies. *PeerJ*, 2, e593. <https://doi.org/10.7717/peerj.593>
- 491 Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.  
492 *EMBnet.Journal*, 17(1), 10. <https://doi.org/10.14806/ej.17.1.200>
- 493 McCutcheon, J. P., McDonald, B. R., & Moran, N. A. (2009). Convergent evolution of metabolic  
494 roles in bacterial co-symbionts of insects. *Proceedings of the National Academy of Sciences of the*  
495 *United States of America*, 106(36), 15394–15399. <https://doi.org/10.1073/pnas.0906424106>
- 496 McFall-Ngai, M. J. (2002). Unseen forces: the influence of bacteria on animal development.  
497 *Developmental Biology*, 242, 1–14. <https://doi.org/10.1006/DBIO.2001.0522>
- 498 Mcmurdie, P. J., & Holmes, S. (2013). phyloseq: an R package for reproducible interactive analysis  
499 and graphics of microbiome census data. *PloS One*, 8(4), 1–11.  
500 <https://doi.org/10.1371/journal.pone.0061217>
- 501 O'Brien, P. A., Webster, N. S., Miller, D. J., & Bourne, D. G. (2019). Host-microbe coevolution:  
502 applying evidence from model systems to complex marine invertebrate holobionts. *MBio*, 10(1), 1–  
503 14. <https://doi.org/10.1128/mbio.02241-18>
- 504 Oksanen, J., Blanchet, G. F., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R.,  
505 O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, H. M. H., Szoecs, E., and Wagner, H. (2019).  
506 *vegan: Community Ecology Package*. R package version 2.5-4. [https://CRAN.R-](https://CRAN.R-project.org/package=vegan)  
507 [project.org/package=vegan](https://CRAN.R-project.org/package=vegan)
- 508 Onchuru, T. O., Martinez, A. J., & Kaltenpoth, M. (2018). The cotton stainer's gut microbiota  
509 suppresses infection of a cotransmitted trypanosomatid parasite. *Molecular Ecology*, 27(16), 3408–  
510 3419. <https://doi.org/10.1111/mec.14788>
- 511 Osei-Poku, J., Mbogo, C. M., Palmer, W. J., & Jiggins, F. M. (2012). Deep sequencing reveals  
512 extensive variation in the gut microbiota of wild mosquitoes from Kenya. *Molecular Ecology*,  
513 21(20), 5138–5150. <https://doi.org/10.1111/j.1365-294X.2012.05759.x>
- 514 Paniagua Voirol, L. R., Frago, E., Kaltenpoth, M., Hilker, M., & Fatouros, N. E. (2018). Bacterial  
515 symbionts in Lepidoptera: their diversity, transmission, and impact on the host. *Frontiers in*  
516 *Microbiology*, 9(556), 1–14. <https://doi.org/10.3389/fmicb.2018.00556>
- 517 Parker, B. J., Mclean, A. H. C., Hrček, J., Gerardo, N. M., Charles, H., & Godfray, J. (2017).  
518 Establishment and maintenance of aphid endosymbionts after horizontal transfer is dependent on host  
519 genotype. *Biology Letters*, 13, 1–5. <https://doi.org/10.1098/rsbl.2017.0016>
- 520 Perlman, S. J., Kelly, S. E., & Hunter, M. S. (2008). Population biology of cytoplasmic  
521 incompatibility: maintenance and spread of *Cardinium* symbionts in a parasitic wasp. *Genetics*,  
522 178(2), 1003–1011. <https://doi.org/10.1534/genetics.107.083071>

- 523 Pinaud, S., Portela, J., Duval, D., Nowacki, F. C., Olive, M. A., Allienne, J. F., Galinier, R., Dheilly,  
524 N. M., Kieffier-Jaquinod, S., Mitta, G., Théron, A., & Gourbal, B. (2016). A shift from cellular to  
525 humoral responses contributes to innate immune memory in the vector snail *Biomphalaria glabrata*.  
526 PLoS Pathogens, 12(1), 1–18. <https://doi.org/10.1371/journal.ppat.1005361>
- 527 Pollock, F. J., Mcminds, R., Smith, S., Bourne, D. G., Willis, B. L., Medina, M., Thurber, R. V., &  
528 Zaneveld, J. R. (2018). Coral-associated bacteria demonstrate phyllosymbiosis and cophylogeny.  
529 Nature Communications, 9(4921). <https://doi.org/10.1038/s41467-018-07275-x>
- 530 Portela, J., Duval, D., Rognon, A., Galinier, R., Boissier, J., Coustau, C., Mitta, G., Théron, A., &  
531 Gourbal, B. (2013). Evidence for specific genotype-dependent immune priming in the  
532 lophotrochozoan *Biomphalaria glabrata* snail. Journal of Innate Immunity, 5(3), 261–276.  
533 <https://doi.org/10.1159/000345909>
- 534 Portet, A., Toulza, E., Lokmer, A., Huot, C., Duval, D., Galinier, R., & Gourbal, B. (2018).  
535 Dysbiosis of the *Biomphalaria glabrata* vector snail microbiota following infection by *Schistosoma*  
536 parasites. BioRxiv. <https://doi.org/10.1101/386623>
- 537 Ramirez, J. L., Souza-Neto, J., Cosme, R. T., Rovira, J., Ortiz, A., Pascale, J. M., & Dimopoulos, G.  
538 (2012). Reciprocal tripartite interactions between the *Aedes aegypti* midgut microbiota, innate  
539 immune system and dengue virus influences vector competence. PLoS Neglected Tropical Diseases,  
540 6(3), 1–11. <https://doi.org/10.1371/journal.pntd.0001561>
- 541 Razin, S., Yogev, D., & Naot, Y. (1998). Molecular biology and pathogenicity of Mycoplasmas.  
542 Microbiology and Molecular Biology Reviews (Vol. 62). Retrieved from  
543 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC98941/pdf/mr001094.pdf>
- 544 Roder, C., Bayer, T., Aranda, M., Kruse, M., & Voolstra, C. R. (2015). Microbiome structure of the  
545 fungid coral *Ctenactis echinata* aligns with environmental differences. Molecular Ecology, 24(13),  
546 3501–3511. <https://doi.org/10.1111/mec.13251>
- 547 Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. (2016). VSEARCH: a versatile open  
548 source tool for metagenomics. PeerJ, 4, e2584. <https://doi.org/10.7717/peerj.2584>
- 549 Sánchez-Cañizares, C., Jorrín, B., Poole, P. S., & Tkacz, A. (2017). Understanding the holobiont: the  
550 interdependence of plants and their microbiome. Current Opinion in Microbiology, 38, 188–196.  
551 <https://doi.org/10.1016/j.mib.2017.07.001>
- 552 Sanders, J. G., Powell, S., Kronauer, D. J. C., Vasconcelos, H. L., Frederickson, M. E., & Pierce, N.  
553 E. (2014). Stability and phylogenetic correlation in gut microbiota: lessons from ants and apes.  
554 Molecular Ecology, 23(6), 1268–1283. <https://doi.org/10.1111/mec.12611>
- 555 Saraiva, R. G., Kang, S., Simões, M. L., Angleró-Rodríguez, Y. I., & Dimopoulos, G. (2016).  
556 Mosquito gut antiparasitic and antiviral immunity. Developmental and Comparative Immunology,  
557 64, 53–64. <https://doi.org/10.1016/j.dci.2016.01.015>
- 558 Silva, T. M., Melo, E. S., Lopes, A. C. S., Veras, D. L., Duarte, C. R., Alves, L. C., & Brayner, F. A.  
559 (2013). Characterization of the bacterial microbiota of *Biomphalaria glabrata* (Say, 1818) (Mollusca:  
560 Gastropoda) from Brazil. Letters in Applied Microbiology, 57(1), 19–25.  
561 <https://doi.org/10.1111/lam.12068>

- 562 Theron, A., Rognon, A., Gourbal, B., & Mitta, G. (2014). Multi-parasite host susceptibility and  
563 multi-host parasite infectivity: A new approach of the *Biomphalaria glabrata*/*Schistosoma mansoni*  
564 compatibility polymorphism. *Infection, Genetics and Evolution*, 26, 80–88.  
565 <https://doi.org/10.1016/j.meegid.2014.04.025>
- 566 Trabal, N., Mazón-Suástegui, J. M., Vázquez-Juárez, R., Asencio-Valle, F., Morales-Bojórquez, E.,  
567 & Romero, J. (2012). Molecular analysis of bacterial microbiota associated with oysters (*Crassostrea*  
568 *gigas* and *Crassostrea corteziensis*) in different growth phases at two cultivation sites. *Microbial*  
569 *Ecology*, 64, 555–569. <https://doi.org/10.1007/s00248-012-0039-5>
- 570 Werren, J. H., Baldo, L., & Clark, M. E. (2008). Wolbachia: master manipulators of invertebrate  
571 biology. *Nature Reviews Microbiology*, 6(10), 741–751. <https://doi.org/10.1038/nrmicro1969>
- 572 WHO. (2019). Schistosomiasis. Retrieved from [https://www.who.int/en/news-room/fact-](https://www.who.int/en/news-room/fact-sheets/detail/schistosomiasis)  
573 [sheets/detail/schistosomiasis](https://www.who.int/en/news-room/fact-sheets/detail/schistosomiasis)
- 574

575 **Tables**

576 Table 1: Origin of snail strains used in this study

Species	Strain	Strain code	Strain origin
<i>Biomphalaria glabrata</i>	BAR2	B. gla BAR2	Belo Horizonte, Brazil (G. Oliveira, 2013)
<i>Biomphalaria glabrata</i>	BRE2	B. gla BRE2	Recife, Brazil (A. Théron, 1975)
<i>Biomphalaria glabrata</i>	GH2	B. gla GH2	Dans Fond, Guadeloupe (2005)
<i>Biomphalaria glabrata</i>	BS902	B. gla BS902	Salvador, Brazil (1960)
<i>Biomphalaria pfeifferi</i>	/	B. pfe	Anakhar, Oman (H. Moné, G. Moné, 2015)
<i>Planorbarius metidjensis</i>	/	P. met	Salamanca, Spain (S. Mas-Comà, 2014)
<i>Bulinus truncatus</i>	/	B. tru	Almeria, Spain (A. Olega, 2015)

577

578 **Figure legends**

579 Figure 1: Relative composition for the eight most abundant phyla in the microbiota for each strain of  
580 mollusc.

581 Figure 2: Principal Coordinate Analysis (PCoA) on Bray-Curtis dissimilarity matrix for bacterial  
582 microbiota composition. Each dot is an individual and each colour, a strain. The labels are displayed  
583 at the barycentre.

584 Figure 3: A) Hierarchical clustering based on Bray-Curtis dissimilarity matrix and Ward linkage for  
585 OTUs of the core microbiota. Each colour represents a strain. B) Phylogenetic tree of host species  
586 based on 28S rRNA gene sequence and using Maximum Likelihood with 500 bootstraps (%) for  
587 node support. *Physa sp.* was used as an outgroup. The red numbers are the bootstrap values for the  
588 nodes.

589 **Supplementary files**

590 Table S1: OTU table with taxonomic affiliation and read numbers for each sample.

591 Table S2: Core microbiota by strain/species at the family level (representing at least 0.6% of the total  
592 sequence read number by strain). The families highlighted in blue are common to all the strains.

593 Table S3: OTUs significantly associated with each single host species using indicpecies.

594 Supplementary Figure S1: Principal Coordinate Analysis (PCoA) of every individual from each  
595 strain in the common garden experiment. Each dot is an individual and each colour a strain. The  
596 labels are displayed at the barycentre.

597